

COMPARATIVE GENETIC CHARACTERIZATION BETWEEN ISOLATED POX VIRUS FROM DOVE AND SOME AVIAN POX STRAINS

MANAL AWAD*;NASSIF, S.A**; NAGDIA, Y.EL-NAGDI*** and AMAL, I.MOHAMED**.

* Serum and Vaccine Research Institute (SVRI)

** Central Laboratory for Evaluation of Vet. Biologics (CLEVB)

***Animal Health Research Institute.

Received: 19.2.2009

Accepted: 8.3. 2009

SUMMARY

Avipoxviruses from different geographic regions of the world have been characterized to study their genetic and biological properties, but so far, no such work has been performed on Egyptian isolates. Lesions suggestive of avian pox; found on Egyptian wild dove; were used for isolation of pox virus in a previous study. The resulting virus was propagated in chorio-allantoic membrane (CAM) of specific pathogen free (SPF) Embryonated chicken eggs (ECE).

PCR was carried out on the DNA of the dove poxvirus (DPV), pigeon poxvirus (PPV) and a vaccinal strain of fowl pox virus (FPV), then restriction fragment length polymorphism (RFLP) assay was carried out on the resulting amplicons of 578 bp length; using EcoRV and NlaIII re-

striction enzymes. The restriction profile revealed that the dove pox virus is identical to the PPV and both are different from FPV.

The results of immunoblotting analysis of the 3 pox viruses against chicken anti FPV revealed that in spite of the minor antigenic differences observed between them the DPV is closely related to the PPV. In conclusion the Egyptian wild doves are found to play a serious role in the epidemiology of PPV among pigeon flocks.

INTRODUCTION

Natural pox virus infections have been reported in more than 200 species of wild and domestic birds (Bolte et al., 1999). The pox viruses which infect birds belong to the genus Avipoxvirus of the Poxviridae family. Although a few members

(fowl pox, turkey pox, pigeon pox, canary pox, and quail pox) of the genus have been characterized.

Vaccination was found to be an effective method for control of this disease and thus, commercial fowl pox and pigeon pox virus vaccines were developed (Tripathy and Hanson, 1975).

The disease produces lesions on the skin, mouth, esophagus and trachea, although viruses may be found in other organs of the affected birds, resulting in substantial economic losses in commercial poultry (Tripathy et al,2003). However, there is no unified view point yet ,as to the exact number of species ,strains, or variants that exist within the genus Avipoxvirus (Bolte et al,1999) further more, information on genomic characterization as well as pathogenicity and effect of avipox virus isolates from wild birds on commercial chickens is limited (Weli et al 2004). So commercial poultry may be susceptible to various members of the genus Avipoxvirus (Tripathy and cunningham ,1984).

In Egypt the role of the local wild birds such as doves, sparrows and others in the epidemiology of the disease in Egypt is unknown.

In the present study we undergo to clarify the genomic characterization of an avipox virus isolated from an Egyptian dove compared with a fowl pox vaccinal strain and pigeon pox viruses , to in-

vestigate the role of doves in the epidemiology of the disease in Egypt.

MATERIALS AND METHODS

Virus propagation and purification:

In the present study the following viruses were obtained from the department of Pox virus, Serum and Vaccine Research Institute (SVRI): FPV vaccinal strain of the SVRI; PPV of the SVRI and DPV which was isolated in a previous work (Nagdia et al. 2005).

Each virus was propagated by inoculation on dropped CAM of 12 day old embryos (Schnitzlein et al 1988), the CAMs were collected 5 days post inoculation, homogenized in 1/10 (wt / vol) pbs containing 10µg / ml gentamicin , 10 mg / ml streptomycin and 1000 IU / ml penicillin G sodium.

The homogenate was centrifuged at 551 xg for 10 min. and finally the supernates were collected. The viruses were pelleted at 21,525 xg for 60 min at 4°C the suspended pellet was sonicated to 6 cycles, for 15 sec, and placed on ice for 40 sec. between each cycle (Tadese and Reed 2003). The sonicated resuspension was centrifuged through a 9 ml cushion of 36% (wt/vol) sucrose at 31,000 xg for 90 min at 4°C. The pellet was resuspended in TE (10 mM Tris-1 mM EDTA) of pH 9.0 and sonicated again to 2 cycles. The resulting resuspended pellet was layered on a 20%-50% (wt/vol)

continuous sucrose gradient and centrifuged at 31,000 xg for 60 min at 4°C. The band containing the purified virus particles was collected and pelleted by centrifuging at 86,100 xg for 60 min at 4°C. The pellet was then resuspended in TE and stored at -80°C. Virus suspensions were also purified by sodium diatrizoate gradient centrifugation (Esposito et al 1978).

Viral DNA Isolation:

Viral DNA was isolated from the infected and homogenized CAMs by a modification of the technique Schnitzlein et al (1988). In brief the homogenized CAMs were clarified by centrifugation at 551 xg for 10 min. the supernatants were centrifuged at 21,525 xg for 60 min, and the pellets were resuspended in TE. The resuspended pellets were further suspended in 0.5% 2-mercaptoethanol (2-ME) and 10% Triton X-100. The suspension was agitated on ice for 10 min and centrifuged at 1,240 xg for 5 min; the resulting supernatant was centrifuged at 31,000 xg for 60min at 4°C. The pellet was reconstituted in TE buffer containing; 0.25% 2-ME, Proteinase K (10 mg/ml) and 20% N-lauryl sarcosinate. After 30 min of agitation on ice, 54% of sucrose in water

and 20% sodium dodecyl sulfate (SDS) were added and incubated at 55°C for 2 hr. The resulting mixtures were extracted twice with 25:24:1 (vol/vol) phenol:chloroform:isoamyl alcohol and once with 24:1 (vol/vol) chloroform: isoamyl alcohol and concentrated by ethanol precipitation.

PCR Amplification:

The oligonucleotide primers were designed according to (Hofmann, 2006). The primers were designed on the basis of published DNA sequences of the 4 bp core protein gene sequence of FPV (HP444) (Binns et al 1989; Ghildyhal et al 1989). The primers sequence for the forward and reverse primers were illustrated in table (1). PCR was carried out in a final volume of 50µl containing 10 X PCR buffer, 2.5 µM Mgcl2, 200 µM of each of deoxynucleotide ,1 µM of each primer , 2.5 units Taq DNA polymerase and 0.2µg template DNA. The reaction mixture was subjected to 94°C for 7 min. followed by 35 cycles of 94°C for 1 min., 50°C for 1 min. and 72°C for 2 min. and final extension at 72°C for 7 min. The PCR products were separated on 1% agarose gel in 1X TAE .The PCR products are purified using PCR product purification kit of Qiagen Company.

Table(1): The sequences of the primer set

Primer	Sequence	Nucleotide position	Amplicons
Forward	CAG CAG GTG CTA AAC AAC AA	458 – 477	578
Reverse	CGG TAG CTT AAC GCC GAA TA	1016 - 1035	

RFLP assay:

RFLP assay was carried out on the 578- bp amplicons of FPV, DPV and PPV using EcoR V and Nla III restriction enzymes; according to the manufacturer instructions and Hoffmann, 2006. the resulted fragments for each virus were separated on 2% agarose gel in 1 x TAE.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis and immunoblotting:

This technique was carried out according to Tadese and Reed, 2003. Purified virus suspensions were solubilized in equal volumes of lysis buffer (50 mM Tris-HCl, pH 6.8; 5% 2-ME; 10% glycerol; 1% SDS; 0.001% bromophenol blue) and heated in a microwave oven (power output 725 W) for 60 sec. Supernatant extract from uninfected cells was also prepared and used as control. The protein concentrations of purified virus parti-

cles were determined photometrically using a protein assay kit of Bio-Rad Company, and samples of equal protein concentrations were loaded in a stacking gel of 4% polyacrylamide and a resolving gel of 12% polyacrylamide. After 1 hr of electrophoresis at 120 V and 4 hr at 200 V, proteins on the gel were transferred to nitro cellulose filters. The filters were blocked with 5% nonfat dry milk in phosphate-buffered saline (PBS), pH 7.4, washed 4 times with PBS, and made to react with 1:200 chicken anti-FPV serums for 24 hr at room temperature. After washing 4 times with 0.02% Tween-20 in PBS, peroxidase-labeled goat anti-chicken IgG (1:2,000) was added, and incubation was performed for 2 hr at room temperature. Following 4 washes with PBS, the filters were exposed to 4- chloro-1-naphtol, as substrate, containing 0.02% hydrogen peroxide. The relative molecular mass (rmm) of the reacted proteins was determined by comparison with 180 KD protein maker of Sigma Aldrich Company.

Table (2):The restriction fragment profile of DPV, PPV and FPV with either EcoR V or Nla III restriction enzymes.

EcoR V			Nla III		
DPV	PPV	FPV	DPV	PPV	FPV
252	252	252	446	446	185
177	177	177	132	132	132
149	149	149			112
					96
					53

RESULTS

PCR and RFLP analysis of the FP, PP and DP viruses:

The primer set designed from the 4b core protein gene of FPV amplified a 578 bp fragment for the 3 viruses. There was no difference in size of the amplified products. There was no amplification in the control sample from non infected CAM. (Fig.1).

PCR fragments of 578 bp of the FPV, PPV and DPV were digested with the EcoR V & NlaIII restriction enzymes (Table 2, Fig 2,3). The RFLP profile of the DPV was identical to that of the PPV and FPV after EcoRV digestion which yields 3 fragments of 252, 177 and 149 bp lengths for each virus. On the other hand the FPV showed a different RFLP profile after NlaIII digestion which yield 5 fragments of 185, 132, 112,

96 and 53 pb lengths, while the RFLP profile of both DPV and PPV were still identical and yield 2 fragments of 446 and 132 bp lengths.

Immunoblotting analysis of the FPV, PPV and DPV antigens:

In order to determine if FPV, PPV and DPV could also be differentiated antigenically i.e. on the level of immunogenic proteins, purified virus suspensions were first standardized and after electrophoreses on polyacrylamide and transfer to nitrocellulose filters, immunoblotted using chicken anti FPV serum. When antiserum against FPV was used (Fig. 4 and Table 3), antigens with approximate molecular weights of 65, 60, 36, 35.5 and 25 KD were detected in the 3 Pox viruses. Two other antigens of relative molecular mass (rmm) of 73 and 37 KD were unique to both PPV and DPV. On the other hand a 38 KD antigen were detected in the FPV only, while an antigen of rmm of 34 KD was detected in both FPV and DPV.

Table (3):Antigen profile of DPV compared with PPV and FPV using anti-serum against FPV.

Strain	Relative molecular mass (KD)								
	73	65	60	38	37	36	35.5	34	26
DPV	+	+	+	-	+	+	+	+	+
PPV	+	+	+	-	+	+	+	-	+
FPV	-	+	+	+	-	+	+	+	+

DPV = Dove pox virus; PPV = Pigeonpox virus; FPV = Fowlpox virus
KD = Kilo dalton; + = Presence of the specific antigen;
- = Absence of the specific antigen.




Fig. (3): Restriction profile of the 578-bp amplicons of the 3 avipox viruses using NlaIII. lane M, 100 bp DNA ladder; lane 1, DPV; lane 2, PPV and lane 3, FPV.




Fig. (4) Immunoblotting of purified lysates from non-infected CAM or infected with avipoxviruses with antiserum against FPV. Lane 1, DPV; lane 2, PPV; lane 3, FPV; lane 4, lysate of non infected CAM; lane M, pre-stained molecular weight marker.




Fig. (1) Amplification of DNA from FPV (lane 1) DPV (lane 2) and PPV (lane 3). lane M, 100 bp DNA ladder (of Fermentas company).




Fig. (2) Restriction profile of the 578-bp amplicons of the 3 avipox viruses using EcoR V. lane M, 100 bp DNA ladder; lane 1, FPV; lane 2, DPV and lane 3, PPV.

DISCUSSION

This study was designed to characterize an avipoxvirus isolated from local wild dove in Egypt, and to compare it with the fowl pox virus and PPV. Since such local wild birds like doves may play an important role in the epidemiology of avian pathogens. Via transmission of such pathogens either from wild migratory birds to the domestic birds (commercial chickens, Turkey, Pigeons or in between the domestic birds.

A trial to characterize the isolated DPV in comparison with the available avipoxviruses (FPV and PPV) to determine if such DPV is either related to any of them or even represent another species of avipox viruses. The result of the PCR/RFLP analysis for species differentiation in the genus avipox revealed that the DPV is closely related and identical to the PPV, and both of them are different from the FPV.

The 4 b core protein gene amplifying a 578-bp fragment provided the target for PCR/RFLP analysis for species differentiations in the genus avipoxvirus (Hoffman 2006).

When the FPV, PPV and DPV are compared by immunoblotting analysis, The PPV and DPV showed a similar immunogenic protein profile which was distinct from that of FPV.

In spite of the obtained results, there are many common antigens which were present in the FPV, PPV and DPV. However, through the use of SDS-PAGE, the immunogenic viral proteins were resolved and minor antigenic differences between the 3 viruses were observed. Since the uncommon viral proteins are detected by antiserum against FPV, at least some epitops or portion of these proteins have been conserved. In this regard the genetically similar DPV and PPV contained antigens common to FPV. The antigenic relatedness of FPV and PPV has previously been indicated by their ability to cross-immunize chickens (Winterfield and Hitchner, 1965).

The results of both PCR/RFLP assay and immunoblotting analysis revealed that DPV is closely related to PPV and could not be considered a new species.

From the epidemiological point of view, this DPV might have been acquired from infected pigeon, so doves may be considered as a biological vector for PPV and could play a serious role in the spread of the disease among non vaccinated or even vaccinated pigeon flocks.

So application of biosecurity measures in the pigeon farms and chicken farms will prevent entrance and contact of such wild birds with the commercial flocks and subsequently prevent spread of FPV, PPV or any avian pathogens among our poultry flocks.

REFERENCES

- Binns, M.M.; Boursnell, M.E.G.; Tomley, F.M. and Campbell, J.: 1989, Analysis of the fowlpox virus gene encoding the 4b core polypeptide and demonstration that it possesses efficient promoter sequences. *Virology* 170:288-291.
- Bolte, A.L.; Meurer, J. and Kaleta, E.F.: 1999, Avian host Spectrum of avipoxviruses. *Avian Pathol*; 28, 415-432.
- Esposito, J.J.; Nakano, J.H. and Obijeski, J.F.: 1978, Orthopoxvirus DNA: strain differentiation by electrophoresis of restriction endonuclease fragmented virion DNA. *Virology* 89:53-66.
- Ghildyhal, N.; Schnitzlein, W.M. and Tripathy DN: 1989, Genetic and antigen differences between fowlpox and quailpox viruses. *Arch Virol* 106:85-92.
- HoffmannTorge (2006). Molecular biological investigations for detection, epidemiological studies and differentiation of avian pox viruses PhD. Vet., Freie Uni. Berlin, Germany.
- Nagdia, Y. El-Nagdi; Amal El-Yosef and SH. Gheith : 2005, The role of free living dove in transmission of pox virus to some domesticated fowl birds. *Egypt J. Comp. Path. & Clinic. Path*: 18(1): 61-75.
- Schnitzlein, W.M.; Ghildyhal, N. and Tripathy, D.N.: 1988, Genomic and antigenic characterization of avipoxviruses. *Virus Res.* 10: 65-76.
- SimonChioma Weli; MalachyIfeanyi Okeke; Morten Tryland; (ivind Nilssen and Terje Traavik :2004, Characterization of avipoxviruses from wild birds in Norway. *The Canadian J. Vet. Res.* 68: 140-145.
- Theodros Tadese and Willie, M. Reed :2003: Use of restriction fragment length polymorphism, immunoblotting and polymerase chain reaction in the differentiation of avian poxviruses. *J. Vet. Diagn. Invest* 15: 141-150.
- Tripathy, D.N. and Cunningham, C.H.: 1984, Avian Pox. In: Hofstad M.S.; Barnes, H.J.; Calnek, B.W.; Reid, W.M.; Yoder, J. H.W (eds) *Diseases of poultry*, Iowa State University Press, Ames, IA, PP 524-534.
- Tripathy, D.N. and Hanson, L.E.: 1975, Immunity to fowl pox. *Am. J. Vet. Res.* 36: 541-544.
- Winterfield, R.W. and Hitchner, S.B.: 1965, The response of chickens to vaccination with different concentrations of pigeon pox and fowl pox viruses. *Avian Dis.* 9: 237-240.

مقارنة التوصيف الجيني لفيروس الجدري المعزول من اليمام وبعض عترات جدري الطيور

منال عوض ميخائيل * سمير عبد المعز ناصف ** نجية يوسف النجدي **
أمل اسماعيل محمد **

* معهد بحوث الأمصال واللقاحات البيطرية - العباسية - القاهرة

** المعمل المركزي للرقابة علي المستحضرات الحيوية البيطرية - العباسية - القاهرة

*** معهد بحوث صحة الحيوان - الجيزة

لقد تم دراسة الخصائص الجينية والبيولوجية لمجموعات فيروسات جدري الطيور وذلك في مختلف بقاع العالم، لكن والى حد بعيد لم تتم هذه الدراسات بالشكل الكافي على المعزولات المصرية من هذه الفيروسات. في هذه الدراسة تم لأول مرة محاولة توصيف وتعريف لاحدى معزولات فيروس الجدري والتي تم عزلها في دراسة سابقة من يمام مصرى مصاب بمرض الجدري،

وفي هذه الدراسة تم عمل اختبار تفاعل البلمرة المتسلسل لفيروس جدري اليمام وفيروس جدري الحمام والعتره المستخدمة في لقاح جدري الطيور بمعهد الأمصال واللقاحات البيطرية. ثم عمل اختبار RFLP على ناتج تفاعل البلمرة المتسلسل لكل فيروس باستخدام انزيم القطع EcoRV وانزيم القطع NlaIII حيث افادت النتائج ان فيروس جدري اليمام متشابه تماما مع فيروس جدري الحمام كما ان الأثنين مختلفين عن فيروس جدري الطيور. وبالرغم من أن نتائج الـ Immunoblotting أفادت أنه يوجد فروق ضئيلة جداً بين الفيروسات الثلاث إلا أن فيروس جدري اليمام متشابه إلى حد أكبر مع فيروس جدري الحمام. وخلصت الدراسة إلى أهمية الدور الوبائي الذي يلعبه اليمام البرى في إنتقال فيروس الجدري بين قطعان الحمام في مصر.