

**COMPLEMENTED EFFECT OF HUMIC ACID AND BIOFERTILIZERS ON
 WHEAT (*TRITICUM AESTIVUM* L.) PRODUCTIVITY
 BY**

Abou-Aly, H.E. and Mady, M. A.
 Botany Dept. Fac. of Agric., Moshtohor, Benha Univ., Egypt

ABSTRACT

The role of humic acid for enhancing biofertilization performance was studied on growth and yield of wheat in newly sand clay soil. Application of arbuscular mycorrhiza (AM) (*Glomus mosseae*) and plant growth promoting rhizobacteria (*Azotobacter chroococcum*) in combination with humic acid was evaluated. The results indicated that mycorrhizal root infection percentage significantly increased by application of humic acid with AM fungus. Inoculation with the biofertilizer agents increased phosphatase and dehydrogenase activity in wheat rhizosphere especially with AM inoculation. The highest values of enzymes activity were observed when the plants were treated by humic acid in the presence of biofertilizers especially with the dual inoculation. There were remarkable increases in available nutrients in rhizosphere of plants those inoculated with any of the two biofertilizers in combination with humic acid. Application of *A. chroococcum* and AM either alone or dual inoculation in the presence of humic acid gave considerable improvement in growth characteristics, photosynthetic pigments as well as nutrients uptake, total carbohydrates and crude protein of wheat plants when compared with either inoculated or uninoculated treatments without humic acid. Concerning endogenous phytohormones in wheat shoots, inoculation with *A. chroococcum* individually gave maximum value of auxins, while application of humic acid especially with dual inoculation of biofertilizers didn't have positive effect on auxins content. On the other hand, humic acid enhanced the effect of biofertilizers on increasing of cytokinins and gibberellins content in wheat shoots and reducing of abscisic acid. Moreover, application of humic acid gave the highest values of grain and straw yield when associated with dual inoculation or *A. chroococcum* individually. Also, maximum values of grain quality were obtained from plants those treated with dual inoculation and humic acid. Therefore, application of humic acid can be considered as a good approach in enhancement of biofertilizers performance in newly reclaimed soil.

Key words: Humic acid, biofertilizers, mycorrhizae, *A. chroococcum*, wheat, chlorophyll, endogenous hormones, grain yield

INTRODUCTION

Wheat enjoys a privileged position amongst food grain crops in the world in general and particularly in Egypt where it serves as a staple food for the majority of the population. Hence, under the prevailing circumstances, restoration and maintenance of soil fertility is a basic and critical problem, particularly in the newly reclaimed soil. This can be accomplished by adding organic material, biological active substances and plant growth-

promoting microorganisms, in addition to other field practices (Akhtar *et al.*, 2007). Soil organic contents are one of the most important parts that they directly affected the soil fertility and textures as well as increasing the microbial activities in the soil (Tejada *et al.*, 2006).

In recent years, humic substances can be added to the soil for improvement the crop yield. From the point of view of producers,

these chemical preparations have been perceived and accepted as a kind of hormone promoting the growth rather than improving the chemical and physical conditions of the soil (Cacco and Dell Agnolla, 1984). A benefit of humic acid is its ability to complex metal ions and can form aqueous complexes with micronutrients. It is the subject of studies in various areas of agriculture, such as soil chemistry, fertility, plant physiology as well as environmental sciences, because the multiple role by these materials can greatly improve plant growth and the plant nutrient uptake and was particularly important for the transport and availability of micronutrients (Bohrne and Lua, 1997 and Turkmen *et al.*, 2004). Also, humic acid may form an enzymatically active complex which can carry on reactions that are usually assigned to the metabolic activity of living microorganisms (Sellamuthu and Govindaswamy, 2003).

Microorganisms are important for agriculture in order to promote the circulation of plant nutrients and reduce the need for chemical fertilizers. Plant growth-promoting rhizobacteria (PGPR) can affect plant growth directly by the synthesis of phytohormones

and vitamins, inhibiting plant ethylene synthesis, enhancing stress resistance, improving nutrient uptake, fixing atmospheric nitrogen, solubilizing inorganic phosphate and mineralizing organic phosphate (Lucy *et al.*, 2004 and Cakmakci *et al.*, 2007). One of the most often reported PGPR is *A. chroococcum*. The beneficial effect of these bacteria is attributed to IAA production and to some extent to non-symbiotic N₂-fixation. So, these bacteria can potentially be used to improve wheat nutrition of micronutrients (Rajaei *et al.*, 2007). Arbuscular mycorrhizae (AM) are symbiotic associations formed between plants and soil fungi that benefit both partners. The role of AM in acquisition and sorption of nutrients from the soil has been recognized. Pronounced response had been obtained in the solubility of micronutrients in newly reclaimed soil when mycorrhiza was accompanied with organic substrates (Habashy *et al.*, 2008).

The present work is designed to evaluate integration between humic acid as soil enhancer and biofertilizers with *A. chroococcum* and mycorrhiza for improving the growth and yield of wheat in newly reclaimed soil.

MATERIALS AND METHODS

The study was conducted on newly soil cultivated with wheat (*Triticum aestivum* L. c.v. Sakha 93) at El-Bostan region, El-Behera Govern., Egypt during winter seasons of 2006/2007 and 2007/2008. Interaction effects between humic acid and endomycorrhizal fungi (*Glomus mosseae*) combined

with *A. chroococcum* on growth and yield of wheat were studied. Some physical and chemical properties of the experimental soil were estimated according to Jackson (1973) and Black *et al.* (1982), respectively (Table A).

Table (A): Physical and chemical analyses of the experimental soil.

Particle size distribution %				Soil chemical properties									
Sand	Silt	Clay	Texture class	pH	CaCO ₃ %	OM %	EC (ds/m ⁻¹)						
65.25	10.21	24.54	Sandy clay loam	8.16	14.27	0.97	1.43						
Soluble cations and anions in mol/L							Available nutrients (ppm)						
Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	CO ₃ ²⁻	HCO ₃ ⁻	Cl	N	P	K	Fe	Mn	Zn	Cu
7.92	4.25	9.19	0.63	0.00	2.96	11.81	25.8	3.1	13.7	2.1	0.62	0.39	0.36

Humic acid

Humic acid (85%) which contain 56% C, 4.5% H, 31% O and 4.5% N was

obtained from Sphinx for International trade Company, Cairo, Egypt.

Mycorrhizal inoculation

Arbuscular mycorrhizal fungus (*Glomus mosseae*) was obtained from Agric. Microbiol. Dept., Soils, Water and Environment Res. Inst., Agric. Res. Center, Giza, Egypt. Mycorrhizal inoculum consisted of root, hyphae, spores and growth media from a pot culture of onion plants which was previously infected with *Glomus mosseae* and grown for 4 months in pot culture. The standard inoculum (400 kg/fed.) contained about 270 spores/g. Spores of the fungus were measured by a wet-sieving and decanting technique (Gerdemann and Nicolson, 1963).

Azotobacter chroococcum

Growth regulators producing *Azotobacter chroococcum*—previously isolated and identified by El-Mehiy (2007) in Botany Dept., Fac. of Agric., Benha Univ., Egypt—was used as seed inoculants. The tested bacteria were grown on modified Ashby's medium (Abdel-Malek and Ishac, 1968) at 30°C for 7 days just before seed inoculation to reach the final density of 25×10^8 cfu/ml. Grains of wheat were mixed with the suspension for 30 min. Arabic Gum (16%) was applied to the grains as an adhesive agent. The grains were left to air-drying in shade, and then the grains became ready for sowing.

Experimental design

Grains of wheat (Sakha 93) were successfully washed with water and air-dried. Then, grains were soaked in solution of humic acid (2g/L) for 2 hrs and/or cell suspension of *A. chroococcum*. The grains were sown on the 15th and 17th of November in the two growing seasons, respectively. The experiments were arranged in randomized complete block design with three replicates. The plot area was 10.5 m² (3x 3.5m). All plots except N₂-fixer treatments received nitrogen fertilizers at the rate of 200 kg/fed urea (46 % N) in two equal doses (before the first and second irrigation). While, *A. chroococcum* treatments were supplemented with a half dose of inorganic N-fertilizer. Calcium super phosphate (15.5% P₂O₅) and potassium sulphate (48 % K₂O) were added before cultivation in both seasons at the rates of 150 and 100 kg/ fed., respectively. Humic acid was added at the rate of 5

kg/fed. after 30 and 60 days from sowing in two equal doses, while mycorrhiza was added just before sowing. The other required culture practices for growing wheat were followed as recommended.

This experiment included the following treatments:

- 1- Control.
- 2- Humic acid.
- 3- Arbuscular mycorrhiza (AM)
- 4- *A. chroococcum*.
- 5- Humic acid + AM
- 6- *A. chroococcum* + Humic acid
- 7- *A. chroococcum* + AM.
- 8- *A. chroococcum* + AM + Humic acid

Microbial activities

Microbial activities of the plants rhizosphere after 45 days from sowing were conducted. Mycorrhizal infection was microscopically estimated on a sample of fresh root as described by Giovannetti and Mosse (1980) after clearing and staining (Vierheilig *et al.*, 1998). The samples were analyzed for dehydrogenase activity according to the method described by Casida *et al.* (1964) while phosphatase activity was determined by the method given by Drobnikova (1961). Rhizosphere samples were analyzed for available nitrogen according to Page *et al.* (1982), available phosphorus was determined according to (A.P.H.A, 1992), available potassium according to Chapman and Pratt (1961) and available Fe and Zn were determined according to Page *et al.* (1982).

Sampling and collecting data

Nine plants of wheat from each treatment were randomly taken at 70 and 100 days after sowing to measure different morphological characteristics (plant height (cm), number of tillers/ plant, leaves dry weight (g/plant) and total leaf area (cm²/plant) using the disk methods according to Derieux *et al.* (1973).

Photosynthetic pigments

Chlorophyll a, b and carotenoids were colorimetrically determined in fresh leaves of wheat plants at 70 and 100 days after sowing during the two seasons according to the

methods described by Wettstein (1957) and calculated as mg/g fresh weight.

Chemical composition

Samples from wheat leaves at 70 and 100 days after sowing and grains at harvest were taken to determine total nitrogen (Horneck and Miller, 1998), phosphorus (Sandell, 1950), potassium (Horneck and Hanson, 1998). Also NPK uptake was calculated after determination of NPK according to (Chapman and Pratt, 1961). Total carbohydrate was determined according to (Dubois *et al.*, 1956). Crude protein was calculated according to the following equation: Crude protein = Total nitrogen x 5.75 (A.O.A.C., 1990).

Endogenous phytohormones

Endogenous phytohormones were quantitatively determined in wheat shoots at

80 days after sowing in the second season using High- Performance Liquid Chromatography (HPLC) according to Koshioka *et al.* (1983) for auxin (IAA), gibberellic acid (GA₃) and abscisic acid (ABA) while, cytokinins were determined according to Nicander *et al.* (1993).

Yield characteristics

At harvest, three plants were randomly taken /plot from each treatment for estimation of number of spikes/plant, grain yield (g)/plant, straw yield (g)/plant and weight of 1000 grains (g).

Statistical analysis

Data obtained in this study were statistically analyzed by using the least significant differences test (L.S.D) according to Senedecor and Cochran (1980).

RESULTS AND DISCUSSION

Mycorrhizal colonization and soil enzymes

Results of mycorrhizal colonization percent shown in Table (1) exhibited a gradual increase with inoculation by AM fungi, while it showed no significant increase with individual application of humic acid or *A. chroococcum* comparing to the control treatment. Mycorrhizal root infection was significantly increased by application of humic acid in combination with AM fungi. The results were in agreement with those obtained by Habashy *et al.* (2008) who reported that organic compounds significantly increased colonization of mycorrhiza. It was also noticed from Table (1) that individual application of humic acid or biofertilization with AM or *A. chroococcum* significantly increased phosphatase and dehydrogenase activity in wheat rhizosphere as compared to the control treatment. The combined inoculation with *A. chroococcum* and AM increased enzymes activity more than the individual inoculation. Also, the highest values of enzymes activity were recorded in rhizosphere of the plants that treated with humic acid in the presence of biofertilizer especially the dual inoculation. This may be due to the mechanisms of *Azotobacter* and AM on soil properties, also *Azotobacter*

require large amounts of available carbon for their survival in soil. Addition of humic acid may be of special importance in restoring optimal levels of organic matter for plant growth and for microbial activity which associated with enzymes activity (Karaca *et al.* (2006). These results showed a good agreement with Sellamuthu and Govindaswamy (2003) who reported an increase in enzymes activity with application of humic acid. They also reported that the microbial population and soil enzymes in the rhizosphere could be built up for the efficient utilization of nutrients.

Available nutrients in wheat rhizosphere

Data in Table (2) show that significant increases in available macronutrients (N, P and K) and some micronutrients (Fe and Zn) were observed when wheat plants received humic acid or individually inoculated with *A. chroococcum* or AM as compared to control plants. Application of humic acid with either AM or *A. chroococcum* exhibited values of available nutrients greater than the treatments of biofertilizers without humic acid. Also, application of humic acid with the dual inoculation gave the maximum values of available nutrients. This may be due to the

ability of humic acid to complex metal ions in agricultural systems, also humic acid can form aqueous complexes with soil nutrients, though not to the same extent as many synthetic chelating agents. Since humic acid binds to soil colloidal surfaces, it is not easily leached (Mackowiak *et al.*, 2001). On the other hand, the function of all mycorrhizal systems depends on the ability of the fungal symbiont to absorb inorganic and/or organic nutrients

available in soil (Turkmen *et al.*, 2005). Also, Habashy *et al.* (2008) found that a pronounced response had been obtained in the solubility of nutrients when mycorrhiza was accompanied with organic substances compared to AM inoculation or organic substances added alone. This may be due to that the addition of organic substances which improved the physical properties of the soil, and increased the supplying power of available nutrients to plants.

Table (1): Mycorrhizal colonization and activity of some soil enzymes in the rhizosphere of wheat as affected by humic acid and biofertilizers after 45 days from sowing during the two growing seasons (S₁ and S₂).

Characters Treatments	Mycorrhizal colonization %		Phosphatase (µg inorganic-P/g soil/day)		Dehydrogenase (mg of TPF/g soil/24 h)	
	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂
Control	9.3	15.8	26.7	32.1	23.4	31.1
Humic acid (HA)	11.6	23.2	39.7	46.8	38.4	36.2
Mycorrhiza (AM)	43.6	61.7	45.8	49.2	42.6	51.3
<i>A. chroococcum</i>	15.2	28.6	35.4	38.6	41.2	48.2
AM + HA	56.3	71.2	46.9	50.4	69.1	55.7
<i>A. chroococcum</i> + HA	12.8	25.3	37.9	49.7	51.4	57.8
<i>A. chroococcum</i> + AM	68.9	61.7	57.6	51.6	63.8	52.5
<i>A. chroococcum</i> + AM + HA	53.1	67.5	61.5	53.5	82.3	69.4
LSD at 5%	6.1	5.7	3.8	3.5	4.6	3.2

Table (2): Available nutrients of wheat rhizosphere as affected by application of humic acid and biofertilizers after 45 days from sowing during the two seasons (S₁ and S₂).

Characters Treatments	Available-N (ppm)		Available-P (ppm)		Available-K (ppm)		Available-Fe (ppm)		Available-Zn (ppm)	
	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂
Control	41.4	56.8	3.88	5.12	17.4	19.2	2.53	2.74	0.54	0.56
Humic acid (HA)	66.2	72.4	6.31	7.83	18.9	20.6	3.65	2.95	0.72	0.62
Mycorrhiza (AM)	71.4	65.9	6.21	9.15	21.5	23.7	3.19	3.62	0.53	0.61
<i>A. chroococcum</i>	69.3	71.6	5.70	6.83	20.6	19.4	2.46	3.37	0.58	0.51
AM + HA	89.2	96.5	8.64	9.98	21.7	23.4	4.26	4.83	0.78	0.84
<i>A. chroococcum</i> + HA	91.6	87.0	6.61	7.04	23.7	25.8	4.05	4.94	0.60	0.75
<i>A. chroococcum</i> + AM	87.3	80.7	8.23	9.70	25.0	23.0	4.52	4.02	0.72	0.67
<i>A. chroococcum</i> + AM + HA	98.4	106.3	9.34	9.61	25.8	23.7	4.71	4.28	0.89	0.77
LSD at 5%	10.8	11.3	1.12	0.8	1.06	0.83	0.66	0.37	0.07	0.05

Growth parameters

As shown in Table (3), the growth parameters of wheat plants as plant height, number of tillers/plant, dry weight of leaf/plant and total leaf area/plant were significantly

increased by individual application of humic acid and biofertilizers. Inoculation with *A. chroococcum* in the presence of humic acid or AM significantly increased number of tillers and total leaf area/plant at 70 and 100

days after sowing during the two seasons. In this regard, El-Mehiy (2007) reported that *A. chroococcum* possess a great variety of properties that are interest in the development of biofertilizers including production of growth promoting plant hormones (especially auxins, gibberellins and cytokinins) as well as N₂ fixation. Maximum stimulatory effect of the biofertilizers was obtained when they associated with humic acid application after 70 and 100 days from sowing in the two seasons. These results are in agreement with Turkmen *et al.* (2005) who reported that humic acid application positively affected the plant growth parameters. The mechanism of humic acid that is active in promoting plant growth is not completely known. However, increasing cell membrane permeability, oxygen uptake and root cell elongation are of plant growth factors which were reported. (Russo and Berlyn, 1990).

Photosynthetic pigments

Data in Table (4) indicated that different photosynthetic pigments i.e., chlorophyll a, b and carotenoids in wheat leaves were positively responded to application of humic acid, biofertilizers and their combinations at 70 and 100 days after sowing during the two seasons. Moreover, the interaction between humic acid and dual inoculation with *A. chroococcum* and AM gave the highest values of total pigments during the two seasons as compared with individual treatments and control plants. Generally, these results are to be considered as a good explanation to the obtained data regarding the favorable role of biofertilizers and humic acid on growth parameters (Table 3) that enhanced photosynthetic efficiency and increased dry matter accumulation. Ebrahim and Ali (2004) found that application of *Azotobacter* improved chlorophyll a, b and charotenoids content of wheat leaves.

Table (3): Growth characters of wheat as affected by humic acid and biofertilizers after 70 and 100 days from sowing during the two seasons (S₁ and S₂).

Characters Treatments	70 days after sowing							
	Plant height cm/ plant		No. of tillers /plant		Dry weight of leaf g/ plant		Total leaf area cm ² / plant	
	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂
Control	36.4	41.7	3.9	4.1	4.05	4.12	1250.3	1330.1
Humic acid (HA)	46.4	47.3	5.1	5.4	4.45	4.70	1380.1	1410.2
Mycorrhiza (AM)	43.5	45.7	5.6	5.5	4.90	5.10	1451.2	1490.4
<i>A. chroococcum</i>	50.4	46.9	5.0	5.4	5.22	5.38	1514.7	1540.3
AM + HA	49.7	52.4	5.9	6.0	5.88	5.72	1640.2	1690.4
<i>A. chroococcum</i> + HA	47.6	50.7	5.7	5.9	5.40	5.92	1701.3	1780.1
<i>A. chroococcum</i> + AM	44.7	49.2	5.2	5.7	5.42	5.60	1550.2	1617.1
<i>A. chroococcum</i> + AM + HA	40.6	43.4	6.1	6.3	6.59	6.44	1843.2	1820.2
LSD at 5%	3.12	3.23	0.5	0.62	0.42	0.60	81.2	97.43
100 days after sowing								
Control	86.2	89.1	5.2	5.7	5.95	6.25	1401.0	1480.7
Humic acid (HA)	98.5	97.2	7.7	7.4	9.20	9.62	1680.4	1885.9
Mycorrhiza (AM)	94.7	95.2	6.9	6.4	10.60	10.35	1835.2	1820.3
<i>A. chroococcum</i>	96.5	97.5	7.2	7.7	9.40	9.65	1620.1	1790.7
AM + HA	100.4	101.6	8.2	8.4	10.70	10.60	1875.2	1910.5
<i>A. chroococcum</i> + HA	98.7	99.4	7.9	7.3	10.25	10.75	1890.3	1905.1
<i>A. chroococcum</i> + AM	103.8	99.8	7.3	7.8	10.20	9.90	1870.2	1850.8
<i>A. chroococcum</i> + AM + HA	92.5	93.5	8.8	8.6	11.20	11.10	1980.0	1960.5
LSD at 5%	5.12	5.9	0.57	0.62	1.25	1.32	102.2	107.1

Table (4): Photosynthetic pigments, total carbohydrates and crude protein as affected by humic acid and biofertilizers after 70 and 100 days from sowing in the two seasons (S₁ and S₂).

Characters	70 days after sowing							
	Chlorophyll a mg/g F.W		Chlorophyll b mg/g F.W		Carotenoids mg/g F.W		Total pigments mg/g F.W	
	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂
Treatments								
Control	0.57	0.60	0.38	0.36	0.41	0.43	1.36	1.39
Humic acid (HA)	0.79	0.80	0.42	0.46	0.55	0.57	1.76	1.84
Mycorrhiza (AM)	0.67	0.71	0.44	0.48	0.51	0.52	1.62	1.71
<i>A. chroococcum</i>	0.78	0.81	0.51	0.50	0.54	0.56	1.83	1.87
AM + HA	0.87	0.84	0.50	0.52	0.73	0.77	2.10	2.13
<i>A. chroococcum</i> + HA	0.79	0.83	0.52	0.54	0.71	0.74	2.02	2.11
<i>A. chroococcum</i> + AM	0.75	0.77	0.49	0.51	0.62	0.69	1.86	1.97
<i>A. chroococcum</i> + AM + HA	0.99	0.94	0.57	0.60	0.75	0.72	2.31	2.26
LSD at 5%	0.21	0.23	0.19	0.17	0.15	0.12	0.36	0.39
	100 days after sowing							
Control	0.59	0.61	0.37	0.34	0.45	0.44	1.41	1.39
Humic acid (HA)	0.81	0.79	0.41	0.43	0.54	0.58	1.76	1.80
Mycorrhiza (AM)	0.71	0.74	0.51	0.54	0.57	0.59	1.79	1.87
<i>A. chroococcum</i>	0.71	0.73	0.58	0.59	0.61	0.62	1.91	1.94
AM + HA	0.87	0.89	0.63	0.62	0.72	0.74	2.21	2.25
<i>A. chroococcum</i> + HA	0.83	0.87	0.62	0.64	0.70	0.73	2.15	2.25
<i>A. chroococcum</i> + AM	0.72	0.78	0.61	0.59	0.64	0.63	1.97	2.00
<i>A. chroococcum</i> + AM + HA	0.92	0.90	0.63	0.62	0.77	0.76	2.32	2.28
LSD at 5%	0.22	0.19	0.12	0.15	0.21	0.18	0.32	0.34

Nutrients uptake and some bioconstituents in leaves

Table (5) clearly indicates that application of both humic acid and biofertilizers significantly increased NPK uptake, total carbohydrates and crude protein content in wheat leaves at 70 and 100 days after sowing during the two seasons as compared with control treatment. Moreover, combination between humic acid and dual inoculation with *A. chroococcum* and AM increased NPK uptake nearly more than two times at 70 days and nearly more than three times at 100 days compared with control treatment. Furthermore, the addition of humic acid associated with both biofertilizers increased nutrients uptake with an pronounced effect, and parallel trend for their increases in the soil (Table 2). This may be attributed to the enhancing effect of humic acid and mycorrhiza on soil physical properties to release nutrients in the rhizosphere which supply a power of available

nutrients to plants. The obtained data were in agreement with Turkmen *et al.* (2005) and Habashy *et al.* (2008). Also, Rajaei *et al.* (2007) reported that inoculation of wheat with *A. chroococcum* had a positive effect on nutrients uptake.

Regarding total carbohydrate and crude protein, the same positive trend was observed with application of humic acid and biofertilizers. All treatments showed a significant increase and the maximum one obtained by the interaction between *A. chroococcum* and AM in the presence of humic acid. In this respect, high content of total carbohydrates is a direct result for high rates of photosynthesis with great efficiency that was preceded with large photosynthetic area (Table 3) and high content of photosynthetic pigments (Table 4). The present results are in agreement with those of Ebrahim and Ali (2004).

Table (5): Nitrogen, phosphorus and potassium uptake in wheat leaves as affected by humic acid and biofertilizers after 70 and 100 days from sowing in the two seasons (S_1 and S_2).

Characters Treatments	70 days after sowing									
	N-uptake		P-uptake		K-uptake		Total carbohydrates mg/g D.W		Crude protein mg/g D.W	
	S_1	S_2	S_1	S_2	S_1	S_2	S_1	S_2	S_1	S_2
Control	86.6	84.4	11.5	11.9	98.4	85.2	487.3	492.7	123.1	117.9
Humic acid (HA)	163.5	163.5	18.6	20.4	152.4	156.9	560.7	571.4	211.3	200.1
Mycorrhiza (AM)	172.7	184.3	16.0	17.5	155.3	151.9	542.7	560.5	202.7	207.9
<i>A. chroococcum</i>	168.0	181.5	19.8	20.1	169.6	163.8	548.4	555.7	185.2	194.0
AM + HA	219.6	208.2	25.2	24.3	194.6	188.4	588.3	593.8	214.8	209.3
<i>A. chroococcum</i> + HA	199.2	208.6	22.4	25.3	168.2	193.2	590.6	611.7	212.2	202.7
<i>A. chroococcum</i> + AM	191.8	200.4	21.3	21.2	174.5	174.1	568.2	570.1	203.6	205.9
<i>A. chroococcum</i> + AM + HA	255.0	242.4	31.3	28.9	223.0	222.8	614.8	624.4	222.5	216.2
LSD at 5%	27.3	31.4	7.5	9.3	21.8	26.4	37.9	41.4	11.3	15.7
100 days after sowing										
Control	117.8	134.0	17.0	18.0	132.3	146.8	508.1	511.7	113.9	123.3
Humic acid (HA)	329.3	339.1	38.1	42.1	307.2	313.1	615.3	622.8	205.6	202.7
Mycorrhiza (AM)	366.7	365.8	36.0	37.7	343.9	323.7	591.4	598.4	199.0	203.3
<i>A. chroococcum</i>	308.7	325.2	32.3	33.7	311.6	311.2	602.1	610.7	188.9	193.8
AM + HA	410.8	394.3	44.0	43.4	333.8	340.2	630.8	637.4	220.8	213.9
<i>A. chroococcum</i> + HA	387.4	390.7	41.7	45.6	329.8	352.0	658.1	664.2	217.4	209.0
<i>A. chroococcum</i> + AM	366.1	360.8	37.3	39.1	342.7	334.6	622.4	625.3	206.4	209.6
<i>A. chroococcum</i> + AM + HA	443.2	426.2	52.0	49.4	377.4	375.7	673.2	680.4	222.7	220.8
LSD at 5%	55.3	43.1	8.5	9.4	12.3	17.6	38.6	34.1	13.5	12.4

Endogenous phytohormones

According to the data in Table (6), *A. chroococcum* gave maximum values of auxins in wheat shoots compared with all treatments, but inoculation of these bacteria with humic acid or humic acid with AM led to a decrease in auxins content compared to control. Gibberellins and cytokinins were improved by inoculation with *A. chroococcum* or AM and reached the highest values when the biofertilizers were supported by humic acid. Many investigators reported the role of plant growth promoting rhizobacteria such as *A. chroococcum* in the production of hormones such as gibberellins, auxins and cytokinins (El-Mehiy, 2007 and Rajaei *et al.*, 2007). On the other hand, abscisic acid, as growth inhibitor, was decreased with using AM or humic acid application while dual inoculation with AM

and *A. chroococcum* in the presence of humic acid recorded maximum reduction of abscisic acid content in wheat shoots.

Yield and its components

Data in Tables (7 & 8) showed that, number of spikes, grain yield, weight of thousand grains and straw yield of wheat as well as chemical composition of wheat grains significantly increased in response to any of the tested biofertilizer compared to control. Also, humic acid had positive effect on the same parameters. Moreover, humic acid application triggered and increased the positive effects of *A. chroococcum* and AM inoculation when wheat plants were inoculated with both biofertilizers in the presence of humic acid.

Table (6): Endogenous phytohormones in wheat shoots as affected by humic acid and biofertilizers applications at 80 days after sowing during second season.

Characters	Auxins		Gibberellins		Cytokinins		Total promoters		Cytokinins /Auxins		Abscisic acid (ABA)	
	µg/g F.W	±% Relative to control	µg/g F.W	±% Relative to control	µg/g F.W	±% Relative to control	µg/g F.W	±% Relative to control	µg/g F.W	±% Relative to control	µg/g F.W	±% Relative to control
Control	86.7	0.0	60.2	0.0	132.2	0.0	279.1	0.0	1.52	0.0	1.56	0.0
Humic acid (HA)	89.4	+3.1	87.9	+46.0	166.6	+26.0	343.9	+23.2	1.86	+22.4	1.25	-19.9
Mycorrhiza (AM)	110.9	+27.9	70.8	+17.6	170.5	+29.0	352.2	+26.2	1.54	+1.3	1.32	-15.4
<i>A. chroococcum</i>	132.7	+53.1	80.4	+33.6	168.4	+27.4	381.5	+36.7	1.27	-16.4	1.50	-3.8
AM + HA	90.3	+4.2	95.7	+59.0	172.4	+30.4	358.4	+28.4	1.91	+25.7	1.48	-5.1
<i>A. chroococcum</i> + HA	85.8	-1.1	90.3	+50.0	186.3	+40.9	362.4	+29.8	2.17	+42.8	1.20	-23.1
<i>A. chroococcum</i> + AM	114.7	+32.3	79.9	+32.7	168.5	+27.5	363.1	+30.1	1.45	-4.6	1.38	-11.5
<i>A. chroococcum</i> + AM+HA	82.4	-5.0	102.4	+70.1	193.7	+46.5	378.6	+35.6	2.35	+54.6	0.96	-38.5

Table (7): Yield components of wheat as affected by humic acid and biofertilizers applications during the two growing seasons (S₁ and S₂).

Characters	No. of spikes/ plant		Grain yield g/plant		Weight of 1000 grains		Straw yield g/plant	
	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂
Control	5.76	6.11	6.35	6.80	42.60	43.50	8.95	8.70
Humic acid (HA)	7.25	7.30	8.40	8.65	50.47	52.60	10.70	10.95
Mycorrhiza (AM)	6.70	6.40	7.75	7.90	48.80	49.20	10.50	10.25
<i>A. chroococcum</i>	6.59	6.96	8.25	8.45	51.20	50.80	10.40	10.35
AM + HA	7.15	7.22	9.45	9.15	52.80	52.85	11.60	11.23
<i>A. chroococcum</i> + HA	7.42	7.60	10.90	10.35	53.35	53.60	11.80	11.75
<i>A. chroococcum</i> + AM	6.90	6.93	9.30	9.70	50.20	51.30	11.20	10.90
<i>A. chroococcum</i> + AM + HA	8.59	8.70	11.72	11.20	55.70	54.42	12.45	12.80
LSD at 5%	0.46	0.31	0.45	0.52	2.05	1.76	0.59	0.64

Table (8): Chemical composition of wheat grains as affected by humic acid and biofertilizers applications during the two growing seasons (S₁ and S₂).

Characters	N mg/g D.W		P mg/g D.W		K mg/g D.W		Total carbohydrates mg/g D.W		Crude protein mg/g D.W	
	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂	S ₁	S ₂
Control	16.7	19.4	2.26	2.56	6.12	6.75	728.2	715.4	96.0	111.6
Humic acid (HA)	21.8	19.8	3.94	3.85	9.80	9.15	734.7	738.8	125.4	113.9
Mycorrhiza (AM)	20.7	20.9	2.35	2.41	8.75	8.20	730.5	744.2	119.0	120.2
<i>A. chroococcum</i>	21.4	19.9	2.82	2.74	7.25	7.40	752.1	794.5	123.1	114.4
AM + HA	21.8	20.6	2.48	2.33	7.90	7.88	760.4	755.2	125.4	118.5
<i>A. chroococcum</i> + HA	21.3	20.5	2.55	2.68	8.25	8.42	750.3	749.8	122.5	117.9
<i>A. chroococcum</i> + AM	20.3	21.2	2.71	3.66	9.95	10.15	769.6	763.5	122.5	121.9
<i>A. chroococcum</i> + AM + HA	22.4	21.7	4.12	3.95	10.70	10.80	787.1	770.4	128.8	124.8
LSD at 5%	1.05	0.95	0.12	1.07	1.25	2.11	12.88	18.50	7.41	5.20

The stimulatory effect of humic acid with dual inoculation on wheat yield would be expected since these applications promoted microbial activities (Tables 1 & 2), growth parameters (Table 3), increased photosynthetic pigments (Table 4), increased nutrients uptake and total carbohydrates (Table 5) as well as endogenous phytohormones (Table 6) as previously resulted and discussed in this work. These findings are supported by Turkmen *et al.* (2005) and Akhtar *et al.* (2007). They reported that the combined application of

Azotobacter or mycorrhiza with humic substances increased plant yield.

This study clearly indicated that humic acid could have positive effect on plant growth and yield by acting as soil enhancer and as well as by improving its physical properties. Also, the combined application of humic acid with the potent biofertilizers is a good tool for growth and yield promotion as well as improving soil health, particularly in newly soil.

REFERENCES

- A.O.A.C. (1990): Official methods of analysis, 15th Ed., Association of official analytical chemists, Inc., USA. manure. *Annals. Agric. Res.*, 24 (3): 466-473.
- A.P.H.A, American Public Health Association (1992): Standard methods for the examination of water and wastewater. Washington, D.C., USA.
- Abdel-Malek , Y. and Ishac, Y.Z. (1968): Evaluation of methods used in counting azotobacter. *J. Appl. Bact.*, 31: 267-275.
- Akhtar, M.J.; Asghar, H.N.; Asif, M. and Zahir, Z.A. (2007): Growth and yield of wheat as affected by compost enriched with chemical fertilizer, L-tryptophan and rhizobacteria. *Pak. J. Agri. Sci.*, 44(1):136-140.
- Black, C.A.; Evans, D.O.; Ensminger, L.E.; White, J.L; Clark, F.E. and Dinauer, R.C. (1982): *Methods of Soil Analysis*. Part 2. Chemical and microbiological properties. 2nd Ed. Soil Sci., Soc. of Am. Inch. Publ., Madison, Wisconsin, U.S.A.
- Bohme, M. and Lua, H.T. (1997): Influence of mineral and organic treatments in the rhizosphere on the growth of tomato plants. *Acta Hortic.*, 450:161-168.
- Cacco, G. and Dell Agnolla, G. (1984): Plant growth regulator activity of soluble humic substances. *Can. J. Soil Sci.*, 64:25-28.
- Cakmakci, R.; Donmez, M.F. and Erdogan, U. (2007): The effect of plant growth promoting rhizobacteria on barley seedling growth, nutrient uptake, some soil properties and bacterial counts. *Turk. J. Agric. Forest.*, 31:189-199.
- Casida, L.E.; Klein, D.A. and Santoro, T. (1964): Soil dehydrogenase activity. *Soil Sci.*, 98: 371-378.
- Chapman, H.D. and Pratt, F. P. (1961): *Methods of analysis of soil, plant and water*. Cal. Univ., 150-200.
- Derieux, M.; Krerrest, R. and Montalanty, Y. (1973): Etude dela surface foliaive et de lactivite photosynthetique chez kulkues hybrid de mais. *Ann. Amelior plants*, 23: 95-107.
- Drobnikova, V. (1961): Factors influencing the determination of phosphatase in soil. *Folia. Microbiol.*, 6, 260.
- Dubois, M.; Gilles, K.A.; Hamilton, J.K.; Rebens, P.A. and Smith, F. (1956): Colorimetric methods for determination sugars and related substances. *Annals. Chem. Soc.*, 46: 1662- 1669.
- Ebrahim, M.K. and Ali, M.M. (2004): Physiological response of wheat to foliar application of zinc and inoculation with some bacterial fertilizers. *J. Plant Nutrition*, 27(10): 1859-1874.
- El-Mehiy, R.M.M. (2007): Efficiency of some microorganisms in production of some plant growth stimulating substances. M.Sc. Thesis, Fac. of Agric. Benha Univ. Egypt.
- Gerdemann, J.W. and Nicolson, T.H. (1963): Spores of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. *Trans. Brit Mycol. Soc.*, 46:235-244.

- Giovannetti, M. and Mosse, B. (1980): An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection in roots. *New Phyt.*, **84** (3):489-500.
- Habashy, N.R.; Amal W. Abou El-Khair and Raafat N. Zaki (2008): Effect of organic and biofertilizers on phosphorus and some micronutrients availability in a calcareous soil. *Res. J. Agric. & Biol. Sci.*, **4**(5):454-552.
- Horneck, D.A. and Hanson, D. (1998): Determination of potassium and sodium by flame Emission spectrophotometry. In hand book of reference methods for plant analysis, e.d Kolra, Y. P.(e.d). 153-155.
- Horneck, D.A. and Miller, R.O. (1998): Determination of total nitrogen in plant tissue. In hand book of reference methods for plant analysis, e.d Kolra, Y.P.(e.d). 73.
- Jackson, M.L. (1973): *Soil Chemical Analysis*. Prentice-Hall of India, Private New Delhi.
- Karaca, A.; Tugay, O.C. and Tamer, N. (2006): Effects of a humic deposit (gyttja) on soil chemical and microbiological properties and heavy metal availability. *Biol Fertil Soils*, **42**: 585-592
- Koshioka, M.; Harada, J.; Noma, M.; Sassa, T.; Ogiama, K.; Taylor, J. S.; Rood, S. B.; Legge, R.L. and Pharis, R.P. (1983): Reversed phase C18 high performance liquid Chromatography of acidic and conjugated gibberellins. *J. Chromatogr.*, **256**: 101-115.
- Lucy, M.; Reed, E. and Glick, B.R. (2004): Application of free living plant growth-promoting rhizobacteria. *Antoni Van Leeuwenhoek*, **86**:1-25.
- Mackowiak, C.L.; Grossl, P.R. and Bugbee, B.G. (2001): Beneficial effects of humic acid on micronutrients availability to wheat. *Soil Sci. Soc. Am. J.*, **65**:1744-1750.
- Nicander, B.; Stahi, U.; Bjorkman, P.O. and Tillberg, E. (1993): Immunoaffinity copurification of cytokinins and analysis by high-performance liquid chromatography with ultraviolet spectrum-detection. *Planta*, **189**: 312- 320.
- Page, A.L.; Miller, R.H. and Keeny, D.R. (1982): *Methods of soil analysis*. Amer. Soc. Agron., Madison., Wisc.
- Rajace, S.; Alikhani, H.A. and Raiesi, F. (2007): Effect of plant growth-promoting potentials of *Azotobacter chroococcum* native strains on growth, yield and uptake of nutrients in wheat. *J. Sci.& Technol. Agric. & Natur.*, **11**(41): 297-306.
- Russo, R.O. and Berlyn, G.P. (1990): The use of organic bio-stimulants to help low input sustainable agriculture. *J. Sust.Agr.*, **1**:19-42.
- Sandell, R. (1950): *Colorimetric determination of traces of metal* 2nd Ed. Inter science. Pub. Inc. New. York
- Sellamuthu, K.M. and Govindaswamy, M. (2003): Effect of fertilizer and humic acid on rhizosphere microorganisms and soil enzymes at an early stage of sugar cane growth. *Sugar Tech.*, **5**(4):273-277.
- Snedecor, G.W. and Cochran, W.G. (1980): *Statistical methods*. 7 th Ed. Iowa state Univ. Press. Ames. Iowa, USA.
- Tejada, M.; Hernandez, M.T. and Garcia, C. (2006): Application of two organic amendments on soil restoration: Effects on the soil biological properties. *J. Environ. Qual.*, **35**:1010-1017.
- Turkmen, O.; Bozkurt, M.A.; Yildiz, M. and Cimrin, K.M. (2004): Effect of nitrogen and humic acid applications on the head weight, nutrient and nitrate contents in lettuce. *Adv. Food Sci.*, **26**:1-6.
- Turkmen, O.; Demir, S.; Sensoy, S. and Dursun, A. (2005): Effect of arbuscular mycorrhizal fungus and humic acid on the seedling development and nutrient content of pepper growth under saline conditions. *J. Biol. Sci.*, **5**(5):568-574.
- Vierheilig, H.; Coughlan, A.P.; Wyss, U. and Piche, Y. (1998): Ink and Vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Appl.& Environ. Microbiol.*, **64**(12): 5004-5007.
- Wettstein, D. (1957): Chlorophyll, letal und der submikroskopische formmech sell- der plastiden, *Exptl. Cell. Res.*, **12**:427.

التأثير المتكامل لحمض الهيوميك والأسمدة الحيوية على تحسين إنتاجية نبات القمح

حامد السيد أبوعلی ، محمد أحمد ماضی

قسم النبات الزراعى - كلية الزراعة بمشتر - جامعة بنها - مصر

فى هذه التجربة تم تقييم دور حمض الهيوميك فى تحسين كفاءة التسميد الحيوى على نمو وإنتاجية نبات القمح فى الأراضى الجديدة. لذلك استخدم نوعين من الأسمدة الحيوية وهما الميكوريزا الداخلية (*Glomus mosseae*) وبكتريا مشجعة للنمو (*Azotobacter chroococcum*) فى وجود حمض الهيوميك. وقد أظهرت النتائج أن استخدام حمض الهيوميك فى وجود الميكوريزا قد أدى إلى زيادة نسبة إصابة الميكوريزا للجذور. وكذلك فإن التلقيح بأى من الأزوتوباكتر أو الميكوريزا أدى إلى زيادة فى نشاط كل من انزيم الفوسفاتيز والديهيدروجينيز فى ريزوسفير نبات القمح وخاصة مع استخدام الميكوريزا. أما أعلى نشاط انزيمى فقد لوحظ فى ريزوسفير النباتات المضاف لها حمض الهيوميك مع التسميد الحيوى المزدوج. وقد كانت تلك المعاملة الأفضل أيضا فى زيادة نسبة العناصر الميسرة فى التربة. وقد أدى استخدام الأزوتوباكتر أو الميكوريزا فرديا أو بالاشتراك فى وجود حمض الهيوميك إلى تحسن ملحوظ فى صفات النمو لنبات القمح وكذلك فى صبغات البناء الضوئى والعناصر والكربوهيدرات الكلية والبروتين الموجود فى نباتات القمح. أيضا أدى التلقيح بالأزوتوباكتر بمفردها إلى زيادة محتوى النبات من الأكسجين، بينما أعلى محتوى من الجبريلين والسيتوكينين وأقل نسبة من حمض الأبسيسيك كانت عند إضافة حمض الهيوميك للنباتات الملقحة تلقىح مزدوج بالأسمدة الحيوية. علاوة على ذلك فإن استخدام حمض الهيوميك مع التلقيح المزدوج بالتسميد الحيوى بكل من الأزوتوباكتر والميكوريزا أعطى أعلى القيم فى محصول الحبوب ومحصول القش وكذلك فى محتوى الحبوب من العناصر والكربوهيدرات والبروتين. لذلك فإن هذه الدراسة تؤكد على استخدام حمض الهيوميك كطريقة فعالة فى تحسين كفاءة التسميد الحيوى وخاصة فى الأراضى المستصلحة الحديثة.