

EVALUATION OF SOME WHEAT GENOTYPES UNDER DIFFERENT SEEDING RATES BY

Mehasen, S.A.S.*; Ahmed, M.A. ** and Morsy, M.A.M.***

* Dept. of Agron., Fac. of Agric., Moshtohor, Benha Univ., Egypt.

** Dept. of Agron., Fac. of Nasser Agric Science, Aden Univ., Yemen.

*** Central Administration for Seed Certification, Ministry of Agric. and Land Reclamation, Egypt.

ABSTRACT

Two field experiments were conducted at the Experiment and Research Center, Fac. of Agric., Moshtohor, Benha Univ., Qalyoubia Governorate, Egypt, during 2007/08 and 2008/09 seasons, to study the effect of three seeding rates (40, 60 and 80 kg fed⁻¹) on growth, yield and yield components of seven wheat genotypes (Giza 168, Sakha 93, Sakha 94, Gemmiza 7, Gemmiza 9, Sids 1 and Sids 12). The most important results which were obtained from this study were as follows:

Gemmiza 7 gave the highest values of flag leaf area, spike length, No. of spikelets spike⁻¹, seed index and grain yield fed⁻¹, Gemmiza 9 gave the highest values of plant height and straw yield fed⁻¹ while, Sakha 94 gave the highest values of No. of tillers plant⁻¹ and No. of spikes/m², moreover, Sids 1, Gemmiza 7 and Sakha 93 genotypes were earlier heading, respectively compared with other genotypes.

The highest values of plant height, No. of spikes/m², grain and straw yields fed⁻¹ were produced by the highest seeding rate (80 kg fed⁻¹). On the other hand, the lowest seeding rate (40 kg fed⁻¹) producing the highest values of No. of tillers plant⁻¹, flag leaf area, spike length, No. of spikelets spike⁻¹ and seed index, while No. of days to 50% heading was not significantly affected by different seeding rates.

Effect of the interaction between wheat genotypes and seeding rates was significant on No. of tillers plant⁻¹, No. of days to 50% heading, flag leaf area, plant height, spike length, No. of spikelets spike⁻¹, No. of spikes/m², seed index, grain and straw yields fed⁻¹. Gemmiza 9 with 80 kg fed⁻¹ seed rate gave the highest values of grain and straw yields fed⁻¹.

Results indicate that No. of tillers plant⁻¹, plant height, spike length, number of spikes/m² and straw yield fed⁻¹ had the greatest influence on grain yield fed⁻¹. Flag leaf area was significantly correlated with grain yield fed⁻¹.

Key words: Wheat, Genotypes, Seeding rates, Growth, Yield and its components.

INTRODUCTION

Wheat is considered the main source of food in the world and Egypt. Raising wheat production through increasing productivity and increasing the cultivated area is an important national target to minimize the gap between the Egyptian production and consumption. In Egypt, the wheat cultivated area is about 2.9 million feddans producing 7.83 million tons (Agric. Statistic Bulletin, 2008).

Increasing wheat yield per unit area can be achieved by breeding high yielding varieties and improving the cultural treatments of the crop.

Several investigators showed that wheat cultivars differed in growth, yield and its components (Metwally *et al.*, 1998; Mehasen, 1999; El-Hawary, 2000; Abd El-

hameed, 2002; Ali *et al.*, 2004; Mehasen and Mohamed, 2005; Abu-Grab *et al.*, 2006; Omar, 2007, Hassan, 2008 and El-Nady, 2009).

Most of the previous studies have shown different effects for seeding rate on growth, yield and yield attributes of wheat. Toaima *et al.* (2000); Saleh (2002); Ramadan (2003); Ali *et al.* (2004); Abu-Grab *et al.*, (2006); Omar (2007); El-Afandy *et al.* (2007); El-Ganayni and Mahmoud (2008) and Ramadan and Awaad (2008) reported increases in plant height, No. of spikes/m², grain and straw yields with increasing seeding rates, while No. of tillers plant⁻¹, No. of days

to 50% heading, flag leaf area, length spike, No. of spikelets spike⁻¹ and 1000-kernels weight were decreased.

Grain yield fed⁻¹ was positively and significantly correlated with plant height, spike length, No. of spikes/m² and straw yield (El-Wakil and Abd-Alla, 2004; Hassan, 2008; Swelam, 2008 and Ramadan and Awaad, 2008).

Therefore, the present investigation was designed to study the performance and productivity of wheat genotypes under different seeding rates, in the Middle Delta of Egypt (Qalyobia Governorate).

MATERIALS AND METHODS

- Description of the studied materials and experimental treatments.

Two field experiments were conducted at the Experimental Research Center, Faculty of Agriculture at Moshtohor, Benha Univ., Qalyobia Governorate, Egypt, during the two winter successive growing seasons 2007/08 and 2008/09, to study the effect of three seeding rates (40, 60 and 80 kg fed⁻¹) on growth, yield and yield components for seven wheat genotypes (Giza 168, Sakha 93, Sakha 94, Gemmiza 7, Gemmiza 9, Sids 1 and Sids 12). The soil was clay in texture, pH values, organic matter content and available N were 7.81, 7.79, 1.67, 1.78% and 50, 58 ppm during the two growing seasons, respectively.

The treatments were assigned in a split-plot design with three replications. Genotypes of wheat were arranged at random in the main plots while, seeding rates occupied the sub-plots. The sub-plot area was 10.5 m² consisted of 15 rows of 3.5 m length and spaced 20 cm apart.

Wheat genotypes were cultivated on November 28th and 30th in the first and second seasons, respectively. N fertilizer with the rate of 80 kg N fed⁻¹ was splitted into two equal doses as urea form (46% N) applied before the first and the second irrigations in both season. P fertilizer with the rate of 15 kg P₂O₅ fed⁻¹ was one equal dose as calcium super

phosphate form (15% P₂O₅) applied before sowing during seedbed preparation. The preceding crop was maize in both seasons. The normal cultural practices were carried out as recommends.

- Collected data.

The studied traits were No. of tillers plant⁻¹, No. of days to 50% heading and flag leaf area (cm²). Random samples of 10 plants were taken from every sub-plots at harvesting time to determine the following characters: plant height (cm), spike length (cm), and No. of spikelets spike⁻¹. For determine No. of spikes/m² and seed index (g) a sample of one square meter from each sub-plot was taken. Grain and straw yields (kg fed⁻¹) were estimated on whole sub-plot basis.

- Statistical analysis.

Analysis of variance was done for the data of each season separately and combined analysis was performed for the data over the two seasons according to Snedecor and Cochran (1980) treatment means were compared using least significant difference test at 0.05 level of significance. Simple and multiple correlation and coefficient of determination were computed between the above mentioned characters as outlined by Steel and Torrie (1987). Using the MSTAT-C Statistical Software package (Michigan State University, 1983).

RESULTS AND DISCUSSION

Analysis of variances for all traits in each season as well as the combined analysis are presented in Table (1). Test of homogeneity revealed that the error variance for the two seasons were homogenous, therefore combined analysis was processed. Year's mean squares were highly significant for all the studied traits except for No. of spikelets spike⁻¹ and seed index were not significant. Wheat genotypes mean squares were highly significant for all traits in both seasons as well as the combined data. Seeding rates mean squares were significant for all traits in both seasons as well as the combined data except No. of days to 50% heading. The interaction between years and wheat genotypes mean squares was not significant for all of the studied characters except No. of tillers plant⁻¹ and straw yield fed⁻¹. The interaction between years and seeding rates mean squares was not significant for all of the studied characters except No. of tillers plant⁻¹, flag leaf area, No. of spikelets spike⁻¹ and seed index. The interaction between years, wheat genotypes and seeding rates mean squares were not significant for all of the studied characters except No. of tillers plant⁻¹ and straw yield fed⁻¹.

-Genotypes differences.

The results reported in Table (2) indicate clearly that, there were significant differences between the different wheat genotypes in all studied traits. Moreover, it is clear from Table (2) that Gemmiza 7 gave the highest values of flag leaf area (42.66 cm²), spike length (12.80 cm), No. of spikelets spike⁻¹ (21.62), seed index (44.11 g) and grain yield fed⁻¹ (3211 kg), Gemmiza 9 gave the highest values of plant height (108.61 cm) and straw yield fed⁻¹ (4343 kg) while, Sakha 94 gave the highest values of No. of tillers plant⁻¹ (4.07) and No. of spikes/m² (460.1) moreover, Sids 1, Gemmiza 7 and Sakha 93 genotypes were earlier heading compared with other genotypes.

It could be concluded that varietal differences among wheat genotypes may be due to genetical make up. The superiority of Gemmiza 7 and Gemmiza 9 in grain yield (kg fed⁻¹) over other genotypes might be due to the increase in yield components, namely, spike

length, No. of spikelets spike⁻¹ and seed index. The results were obtained by Metwally *et al.* (1998); Mehasen (1999); El-Hawary (2000); Abd El-hameed (2002); Ali *et al.* (2004); Mehasen and Mohamed (2005); Abu-Grab *et al.* (2006); Omar (2007); El-Ganayni and Mahmoud (2008); Hassan (2008) and El-Nady (2009) indicated marked differences among wheat varieties in growth, yield and yield components.

-Seeding rate effect:

Results in Table (2) indicated that, growth, yield and its attributes of wheat i.e. No. of tillers plant⁻¹, flag leaf area, plant height, spike length, No. of spikelets spike⁻¹, No. of spikes/m², seed index, grain and straw yields were significantly affected by different seeding rates, whereas, No. of days to 50% heading was insignificantly responded to different seeding rates, under investigation. It is clear that the significant highest values of plant height (105.57cm), No. of spikes/m² (478.9), grain yield fed⁻¹(3209 kg) and straw yield fed⁻¹(4418 kg) were produced by the highest seeding rate (80 kg fed⁻¹). On the other hand, the lowest seeding rate (40 kg fed⁻¹) producing the highest values of No. of tillers plant⁻¹(4.80), flag leaf area (45.12 cm²), spike length (12.12 cm), No. of spikelets spike⁻¹ (36.21) and seed index (43.33 g). The negative effect of highest seeding rate on some yield components could be attributed to the increase in population, as a result of higher seeding rate and consequently low penetration of light within wheat canopy, hence high competition between plants for water, minerals and other environmental factors.

Grain yield fed⁻¹ as a final result of all yield attributers had benefited from the progressive effects of highest seed rate on plant height and No. of spikes/m². Such positive effects were turn in grain yield fed⁻¹. The three seeding rates 40, 60, 80 kg fed⁻¹, yielded 3006, 3145 and 3209 kg, respectively. The present results are in full harmony with those of Toaima *et al.* (2000); Saleh (2002); Ramadan (2003); Ali *et al.* (2004); Abu-Grab *et al.* (2006); Omar (2007); El-Afandy *et al.* (2007); El-Ganayni and Mahmoud (2008); Hassan (2008) and Ramadan and Awaad (2008).

Table (1): Mean squares values and significance for growth, yield and yield components of wheat genotypes in 2007/08, 2008/09 seasons and their combined analysis

S.O.V	d.f	No. of tillers plant ⁻¹	No. of days to 50% heading	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	No. of spikelets spike ⁻¹	No. of spikes/m ²	Seed index (g)	Grain yield (kg fed ⁻¹)	Straw yield (kg fed ⁻¹)
2007/2008 season											
Rep.	2	0.192*	2.016	0.800	0.103	0.007	0.044	0.444*	0.291	7938.111	3594.048*
Geno.	6	0.395**	23.106**	68.660**	489.92**	11.62**	3.452**	454.92**	27.783**	118688.4**	36340.06**
Err.(a)	12	0.033	2.812	2.527	0.711	0.035	0.053	11.981	0.152	2838.481	590.103
SR.	2	18.94**	0.111	644.46**	368.22**	2.325**	4.488**	20100**	32.465**	223291.9**	584754.3**
G.xSR.	12	0.168**	0.630	5.166**	8.087**	0.052	0.056*	105.75**	0.181*	2303.680**	3381.611**
Err.(b)	28	0.027	0.698	1.270	1.894	0.035	0.023	26.071	0.078	734.024	565.357
C.V.%		4.38	0.91	2.90	1.37	1.60	0.73	1.15	0.67	0.87	0.56
2008/2009 season											
Rep.	2	2.90	0.762	0.970	1.968	0.004	0.089	18.143	0.136	3603.762	2401.921*
Geno.	6	0.092**	22.222**	61.449**	420.27**	11.65**	3.017**	681.98**	25.193**	88599.51**	37171.40**
Err.(a)	12	0.017	1.429	1.554	1.339	0.028	0.040	21.513	0.151	1076.188	482.550
SR.	2	14.80**	0.619	636.50**	468.63**	2.517**	7.888**	22844**	43.871**	226737.5**	646942.7**
G.xSR.	12	0.067**	1.452**	5.893**	5.394**	0.037	0.073	50.148	0.382**	1583.942**	1903.693*
Err.(b)	28	0.014	0.405	1.493	1.643	0.025	0.066	13.889	0.086	457.770	846.508
C.V.%		2.91	0.69	3.08	1.25	1.33	1.23	0.83	0.70	0.68	0.68
Combined analysis											
Year	1	1.904**	22.294**	25.976**	160.25**	0.731**	0.168	613.36**	0.446	30520.00**	268366.8**
RepxY	4	0.099**	1.389	0.885	1.036	0.006	0.066	9.294	0.213	5770.937*	2997.984**
Geno.	6	0.184**	43.442**	127.51**	907.98**	23.26**	6.431**	1096.7**	52.812**	205506.1**	71779.77**
YxG.	6	0.303**	1.886	2.590	2.216	0.015	0.038	40.180	0.164	1781.767	1731.698*
Err.(a)	24	0.025	2.120	2.040	1.025	0.032	0.046	16.747	0.152	1957.335	536.327
SR.	2	33.57**	0.103	1272.3**	833.65**	4.839**	12.13**	42897**	75.891**	449686.0**	1230423**
YxSR.	2	0.184**	0.627	8.591**	3.205	0.003	0.239**	47.484	0.446**	343.437	1273.198
G.xSR.	12	0.127**	1.529**	9.787**	12.102**	0.080**	0.105*	132.94**	0.456**	3479.481**	3825.967**
YxGxS	12	0.108**	0.553	1.272	1.379	0.009	0.023	22.966	0.107	408.140	1459.337*
Err.(b)	56	0.020	0.552	1.381	1.769	0.030	0.045	19.980	0.082	595.897	705.933
C.V.%		3.68	0.81	2.99	1.31	1.47	1.02	1.00	0.68	0.78	0.62

*and ** significant at 5% and 1% level of probability, respectively.

Table (2): Growth, yield and yield components of wheat as affected by genotypes and seeding rates (Combined analysis of 2007 and 2008 seasons)

Treatments	No. of tillers plant ⁻¹	No. of days to 50% heading	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	No. of spikelets/spike	No. of spikes/m ²	Seed index (g)	Grain yield (kg fed ⁻¹)	Straw yield (kg fed ⁻¹)
Genotypes										
Giza 168	3.94	94.11	34.22	87.40	9.34	20.02	446.6	39.81	2913	4183
Sakha 93	3.80	90.50	38.99	98.50	12.02	20.74	457.3	43.77	3147	4256
Sakha 94	4.07	92.16	40.22	99.93	12.20	21.11	460.1	42.92	3165	4281
Gemmiza 7	3.79	90.83	42.66	106.84	12.80	21.62	440.5	44.11	3211	4332
Gemmiza 9	3.88	93.83	39.26	108.61	12.46	21.47	444.0	42.42	3200	4343
Sids 1	3.81	90.27	38.21	103.40	11.73	20.25	440.5	40.45	3039	4195
Sids 12	3.84	91.66	41.13	104.63	12.03	20.6	446.8	40.67	3163	4224
L.S.D at 5%	0.11	0.99	0.98	0.69	0.12	0.15	2.8	0.27	30	16
Seeding rates										
40 kg/fed	4.80	91.85	45.12	96.69	12.12	21.36	415.1	43.33	3006	4078
60 kg/fed	3.81	91.92	38.38	101.73	11.83	20.86	449.8	42.09	3145	4281
80 kg/fed	3.02	91.95	34.22	105.57	11.44	20.28	478.9	40.65	3209	4418
L.S.D at 5%	0.06	N.S	0.51	0.58	0.08	0.09	2.0	0.13	11	12

-Interaction effect:

Significant effect of interaction between wheat genotypes and seeding rates was obtained for all growth, yield and yield components namely, No. of tillers plant⁻¹, of days to 50% heading, flag leaf area, plant height, spike length, No. of spikelets spike⁻¹, No. of spikes/m², seed index, grain and straw yields fed⁻¹ (Table 3). Gemmiza 9 with 80 kg fed⁻¹ seed rate gave the highest values of plant height (113.33 cm), No. of spikes/m² (479.8), grain yield fed⁻¹(3301 kg) and straw yield fed⁻¹(4522 kg) while, Gemmiza 7 with 40 kg fed⁻¹ seed rate gave the highest values of flag leaf area (49.05 cm²), spike length (13.20 cm), No. of spikelets spike⁻¹(22.11) and seed index (45.80 g). Moreover, Sakha 94 with 40 kg fed⁻¹ seed rate gave the highest value of No. of tillers plant⁻¹ (5.14) and sids 1 with 40 kg fed⁻¹ seed rate surpassed the other 6 genotypes in early heading (89.50 days). On the other hand, Giza 168 with 40 kg fed⁻¹ seed rate gave the lowest values of plant height (82.33 cm), grain yield fed⁻¹(2771 kg) and straw yield fed⁻¹(3958 kg). Giza 168 with 80 kg fed⁻¹ seed rate gave the lowest values of flag leaf area (31.65 cm²), spike length (9.10 cm), No. of spikelets spike⁻¹(19.36) and seed index (38.73 g) while, Gemmiza 7 with 80 kg fed⁻¹ seed rate gave the lowest value of No. of tillers plant⁻¹(2.84) and Gemmiza 9 with 80 kg fed⁻¹ seed rate gave the

lowest value of No. of spikes/m² (405.3). Similar results were also reported by Hassanein *et al.* (2001); El-Ganayni and Mahmoud (2008) and Ramadan and Awaad (2008).

-Simple correlation analysis:

Data of simple correlation coefficient matrix shown in Table (4). Results indicate that No. of tillers plant⁻¹, plant height, spike length, number of spikes/m² and straw yield fed⁻¹ had the greatest influence on grain yield fed⁻¹ with r values being -0.559, 0.826, 0.517, 0.620 and 0.804 respectively. flag leaf area, was significantly correlated with grain yield fed⁻¹ with r value of -0.198. Another correlation worthy of some attention is that between No. of tillers plant⁻¹ and flag leaf area, plant height, No. spikelets/spike⁻¹, number of spikes /m², seed index and straw yield fed⁻¹ with r values being of 0.798, -0.450, 0.554, -0.850, 0.514 and -0.754, respectively while, spike length was significantly associated with No. of tillers plant⁻¹ with r value being of 0.215. Also positive and highly significant relationships were found between flag leaf area and each of spike length, No. spikelets/spike⁻¹, number of spike/m², seed index and straw yield fed⁻¹ r values being of 0.619, 0.761, -0.781, 0.715 and -0.576, respectively, as well as, between plant height and each of spike length, number

of spike/m² and straw yield fed⁻¹ with r values being of 0.624, 0.381 and 0.651, respectively. Results clearly indicated that spike length showed highly significant and positive correlation with No. spikelets/spike⁻¹ and seed index (r values = 0.732 and 0.692, respectively), while it was significant and positive association number of spikes/m². On the other hand, No. spikelets/spike⁻¹ was highly significant and positive correlation with number of spike/m² and seed index (r values = -0.548 and

0.811, respectively) and significant and positive correlation with straw yield fed⁻¹. Number of spikes/m² revealed highly significantly and positively correlated with seed index and straw yield fed⁻¹ (r values = -0.461 and 0.841, respectively). These findings in most cases were in accordance with those obtained by El-Wakil and Abd-Alla (2004); Swelam (2008); El-Ganayni and Mahmoud (2008); Ramadan and Awaad (2008) and Hassan (2008).

Table (3): Effect of the interaction between genotypes and seeding rates on growth, yield and yield components of wheat (over the combined analysis)

Treatments		No. of tillers plant	No. of days to 50% heading	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	No. of spikelets/ spike	No. of spikelets/m ²	Seed index (g)	Grain yield (kg fed ⁻¹)	Straw yield (kg fed ⁻¹)
Gi. 168	40 kg/fed	4.63	94.50	37.75	82.33	9.71	20.63	415.0	40.85	2771	3958
	60 kg/fed	3.94	93.66	33.28	88.36	9.21	20.08	445.3	39.85	2926	4215
	80 kg/fed	3.24	94.16	31.65	91.50	9.10	19.36	479.5	38.73	3043	4378
Sa. 93	40 kg/fed	4.64	90.83	45.03	92.33	12.40	21.28	430.0	44.85	3054	4101
	60 kg/fed	3.81	90.50	38.10	99.66	12.16	20.83	459.3	43.90	3174	4290
	80 kg/fed	2.95	90.16	33.85	103.50	11.51	20.11	482.6	42.58	3213	4378
Sa. 94	40 kg/fed	5.14	92.50	46.38	95.83	12.55	21.78	433.6	43.98	3079	4118
	60 kg/fed	3.93	91.83	39.78	98.13	12.26	21.20	458.6	42.93	3191	4303
	80 kg/fed	3.16	92.16	34.51	105.83	11.80	20.36	488.0	41.86	3226	4423
Ge. 7	40 kg/fed	4.74	90.50	49.05	102.33	13.20	22.11	406.5	45.80	3105	4155
	60 kg/fed	3.78	91.16	42.86	107.33	12.91	21.68	440.3	44.15	3231	4355
	80 kg/fed	2.84	90.83	36.06	110.86	12.30	21.06	474.6	42.40	3297	4487
Ge. 9	40 kg/fed	5.03	94.16	46.38	102.33	12.73	21.96	405.3	44.00	3060	4141
	60 kg/fed	3.70	93.83	37.98	110.16	12.46	21.50	446.8	42.55	3240	4366
	80 kg/fed	2.91	93.50	33.43	113.33	12.18	20.95	479.8	40.73	3301	4522
Si. 1	40 kg/fed	4.60	89.50	43.95	99.83	12.00	20.60	405.4	41.80	2911	4015
	60 kg/fed	3.83	90.33	36.81	103.71	11.80	20.23	446.3	40.65	3070	4206
	80 kg/fed	3.00	91.00	33.86	106.65	11.41	19.93	469.8	38.91	3138	4365
Si. 12	40 kg/fed	83.4	91.00	47.35	101.83	12.26	21.13	410.0	42.06	3064	4060
	60 kg/fed	3.69	92.16	39.88	104.75	12.03	20.55	452.3	40.66	3182	4236
	80 kg/fed	3.02	91.83	36.16	107.33	11.81	20.20	478.1	39.31	3243	4376
L.S.D at 5%		0.16	0.86	1.36	1.54	0.20	0.24	5.2	0.33	28	31

Table (4): Correlation coefficient between grain yield and its attributes of wheat genotypes (combined data of the two seasons)

Characters	1	2	3	4	5	6	7	8	9
Y-Grain yield fed ⁻¹	-0.559**	-0.152	-0.198	0.826**	0.517**	0.141	0.620**	0.112	0.804**
1-No. of tillers plant ⁻¹	1.000	0.085	0.798**	-0.450**	0.215	0.554**	-0.850**	0.514**	-0.754**
2-No. days to heading		1.000	-0.178	-0.194	-0.373**	0.004	0.047	-0.251**	0.101
3-Flag leaf area (cm ²)			1.000	-0.067	0.619**	0.761**	-0.781**	0.715**	-0.576**
4-Plant height (cm)				1.000	0.624**	0.177	0.381**	0.038	0.651**
5-Spike length (cm)					1.000	0.732**	-0.219	0.692**	0.065
6-No. spikelets/spike ⁻¹						1.000	-0.548**	0.811**	-0.226
7-No. of spikes/m ²							1.000	-0.461**	0.841**
8-Seed index (g)								1.000	-0.238
9-Straw yield fed ⁻¹									1.000

REFERENCES

Abd El-Hameed, I.A. (2002): Effect of some agronomic practices on wheat. Ph.D. Thesis, Fac. Agric., Zagazig Univ., Egypt.

Abu-Grab, O.S.; Mousa, A.M. and El-Shaarawy, G.A. (2006): Photosynthetic and N-use efficiencies for some wheat cultivars in relation to planting density and nitrogen fertilization levels. *Egypt. J. Appl. Sci.*, 21(2B): 475-492.

Ali, G.A.; Zeiton, O.E.; Bassiouny, A.H. and El-Banna, A.R.Y.A. (2004): Productivity of wheat cultivars grown at El-Khattara and El-Arish under different levels of planting densities and N-fertilization. *Zagazig J. Agric. Res.*, 31(4A): 1225-1256.

El-Afandy, K.T.; Abdel-Ati, A.A. and Mohamed, M.A. (2007): Effect of nitrogen, phosphorus and seeding rates on wheat production and weed control in Siwa Oasis. *J. Agric. Sci., Mansoura Univ.*, 32(8): 6099-6111.

El-Ganayni, A.A. and Gamalat A. Mahmoud (2008): Productivity of two bread wheat cultivars under three seeding rates. *J. Agric. Sci., Mansoura Univ.*, 33(6): 3993-4000.

El-Hawary, M.A. (2000): Evaluation of some wheat varieties under water deficit conditions. *Zagazig J. Agric. Res.*, 27(4): 819-830.

El-Nady, Hanim, E.A. (2009): Breeding bread wheat of drought and earliness. Ph. D. Thesis, Fac. Agric., Moshtohor, Benha Univ., Egypt.

El-Wakil, Nadia A.R. and Maha M. Abd-Alla (2004): Influence of seeding rate and nitrogen fertilization level on yield and its attributes of some wheat cultivars. *Egypt. J. Appl. Sci.*, 19(2): 129-150.

Hassanein, M.S.; El-Hariri, D.M. and Ahmed, M.A. (2001): Effect of variety and seed rate on yield and yield components of wheat (*Triticum aestivum* L.). *Annals of Agric. Sci., Moshtohor*, 39(1): 1-13.

Hassan, Manal A. (2008): Effect of seeding rate and row spacing on productivity and resistance to powdery mildew of two bread wheat cultivars. *Egypt. J. Appl. Sci.*, 23 (10A): 169-182.

Mehasen, S.A.S and Mohamed, N.A. (2005): multivariate and response curve analyses for yield and its attributes in some wheat cultivars under nitrogen fertilization levels. *Egypt. J. Appl. Sci.*, 20(2): 93-109.

Mehasen, S.A.S. (1999): Response of some wheat varieties to agrispon foliar application and nitrogen rates. *Annals of Agric. Sci., Moshtohor*, 37(2): 853 – 864.

Metwally, I.O.E.; Abd El-All, A.M. and Leilah, A.A. (1998): Effect of preceding summer crops and N fertilizer levels on growth, yield and yield components of wheat. *Proc. 8th Conf Agron., Suez Canal Univ., Ismailia, Egypt*, 73-79.

Michigan State University, (1983). *MSTAT-C: Micro- computer Statistical Program*, Version 2.0. Michigan State University, East Lansing.

Omar, A.E.A. (2007): Productivity of some wheat cultivars as affected by sowing date and seeding rate. *Egypt. J. Appl. Sci.*, 22(3): 103-116.

- Ramadan, A.R. (2003): Breeding for improvement wheat plant characteristics. Ph. D. Thesis, Fac. Agric., Zagazig Univ., Egypt.
- Ramadan, A.R. and Awaad, S.S. (2008): Response of yield attributes of some bread wheat varieties to irrigation levels and seeding rates under old land conditions. J. Agric. Sci., Mansoura Univ., 33(7): 4717-4737.
- Saleh, M. E. (2002): Response of two wheat cultivars to seeding rate and nitrogen level. Zagazig J. Agric. Res., 29(5): 1343-1366.
- Snedecor, G.W. and Cochran, W.G. (1980): Statistical Methods, 7th Ed., Iowa State Univ. Press, Ames, Iowa, USA.
- Steel, R.G.D. and Torrie, J.H. (1987): Principles and procedures of statistics. A Biometrical Approach 2nd, 6th printing. Mc. Graw. Hill Book. Company: 272 - 277.
- Swelam, A.A. (2008): Mean performance, correlation and path coefficient analysis for grain yield and its components of four bread wheat cultivars grown under four nitrogen levels. Egypt. J. Agric. Res., 86: 561-573.
- Toaima, S.E.; El-Hofi, A.A. and Ashoub, H. (2000): Yield and technological characteristics of some wheat varieties as affected by nitrogen fertilizer and seed rates. J. Agric. Sci., Mansoura Univ., 25(5): 2449-2467.

تقييم بعض التراكيب الوراثية للقمح تحت معدلات تقاوي مختلفة

- صديق عبد العزيز صديق محيسن*، محسن على أحمد**، محمد عبد المنعم محمد مرسى***
 * قسم المحاصيل - كلية الزراعة بمشتهر - جامعة بنها - مصر.
 ** قسم المحاصيل - كلية ناصر للعلوم الزراعية - جامعة عدن - اليمن.
 *** الإدارة المركزية لفحص واعتماد التقاوي - وزارة الزراعة واستصلاح الاراضي - مصر.

- أقيمت تجربتان حقليتان بمركز البحوث والتجارب بكلية الزراعة بمشتهر - جامعة بنها - محافظة القليوبية - مصر خلال موسمي ٢٠٠٧/٢٠٠٨، ٢٠٠٨/٢٠٠٩ م لدراسة تأثير ثلاثة معدلات تقاوي (٤٠، ٦٠، ٨٠ كجم للفدان) علي النمو والمحصول ومكوناته لسبعة تراكيب وراثية من القمح (جيزة ١٦٨، سخا ٩٣، سخا ٩٤، جيزة ٧، جيزة ٩، سدس ١ وسدس ١٢)، ويمكن تلخيص أهم نتائج التحليل التجميعي للموسمين فيما يلي.
- سجل صنف جيزة ٧ أعلى متوسطات لكل من مساحة ورقة العلم، طول السنبل، عدد السنبيلات بالسنبل، وزن الحبة ومحصول الحبوب للفدان. وسجل صنف جيزة ٩ أعلى القيم لكل من طول النبات ومحصول القش للفدان. بينما سجل الصنف سخا ٩٤ أعلى القيم لكل من عدد الأشطاء للنبات وعدد السنابل في المتر المربع. علاوة علي ذلك كانت أفضل الأصناف في صفة التبيكير لطرده السنابل هي سدس ١ وجيزة ٧ وسخا ٩٣ علي الترتيب مقارنة بالتراكيب الوراثية الأخرى.
- أعطى أعلى معدل تقاوي (٨٠ كجم للفدان) أعلى قيم لكل من طول النبات، عدد السنابل في المتر المربع ومحصول الحبوب والقش للفدان. وعلي العكس من ذلك أعطي أقل معدل تقاوي (٤٠ كجم للفدان) أعلى قيم لكل من عدد الأشطاء للنبات، مساحة ورقة العلم، طول السنبل، عدد السنبيلات بالسنبل، ووزن الحبة. بينما لم تتأثر معنويا صفة عدد الأيام لطرده ٥٠% من السنابل بمعدلات التقاوي المختلفة.
- تأثر معنويا كل من عدد الأشطاء للنبات، عدد الأيام لطرده ٥٠% من السنابل، مساحة ورقة العلم، طول النبات، طول السنبل، عدد السنبيلات بالسنبل، وعدد السنابل في المتر المربع، وزن الحبة ومحصول الحبوب والقش للفدان بالتفاعل بين التراكيب الوراثية ومعدلات التقاوي. سجل الصنف جيزة ٩ بمعدل تقاوي ٨٠ كجم للفدان أعلى محصول حبوب وقش للفدان.
- أظهرت النتائج وجود ارتباط معنوي وعالي المعنوية بين محصول الحبوب للفدان و كل من صفات النمو ومكونات المحصول المدروسة ما عدا عدد الأيام لطرده ٥٠% من السنابل، عدد السنبيلات بالسنبل و وزن الحبة ١٠٠٠ حبة.