

Evaluation of Some Environmentally Safe Chemicals Against *Spodoptera littoralis*

Ismail, Seham M.,¹ M. Morshedy²

ABSTRACT

Toxicity effect of Dipel-2X, and three insect growth regulators IGRs: Diflubenzuron; Spiromesifen and Pyriproxyfen, were determined against 2nd larval instar of *Spodoptera littoralis*. The results showed that Diflubenzuron was the most potent toxicity followed by Spiromesifen; Pyriproxyfen, and Dipel-2X. The effect of LC₅₀ of the tested IGRs on the *in vivo* inhibition of chitinase from *Spodoptera littoralis* was assayed. The interaction effect of Dipel-2X with IGRs was investigated. Results proved that pretreated of Dipel-2X with IGRs caused more toxicity effect than single treatment. The sensitivity of chitinase activity to the three tested IGRs was measured by the I₅₀ values. The I₅₀ values of Diflubenzuron; Spiromesifen and Pyriproxyfen on Lab strain larval chitinase are 0.54, 0.60 and 0.73 μM respectively. While the I₅₀ values are 0.60, 0.72 and 0.81 μM respectively against Field strain. The results proved that chitinase was sensitive to the IGRs. Generally, Dipel-2X pretreated with IGRs will produce a new trend so as increase toxicity of the bioinsecticide, enhance the role of beneficial insects. The results of the present study may add some forward steps to use bioinsecticide as alternative to conventional insecticides especially against this insect. So, the tested compounds can be involved in important steps necessary for successful IPM programmes applied against *S. littoralis*.

INTRODUCTION

The Egyptian cotton leafworm, *Spodoptera littoralis* is one of the major pests in the middle east. It has quickly developed resistance to chemical pesticides (Chung and Cote, 1992 & Amin, *et al.*, 2001). Therefore, the cotton leafworm in Egypt exhibits multiple resistance to nearly all insecticides used (Keddis, *et al.*, 1988 & Ishaaya and Kleen, 1990). Owing to the endless and various problems that have been arisen by using insecticides (eg., the development of pest resistance, rapid resurgence of target species and outbreaks of secondary pests), the need to develop novel alternatives or functional combinations of pest control techniques is emphatically a product of this decade. Attention was therefore paid to control insects using different non traditional insecticides, e.g., insect growth regulators (IGRs) and *Bacillus thuringiensis* (BT). (Klein, *et al.*, 1996; Abdel-Halim, 1997 & Rizk, *et al.*, 1999).

IGRs show good effect against scale insects on cotton. Their effects have been observed in development, on behavior and several forms of diapauses. Also IGRs compounds which are considered nowadays one of the mainly component of IPM program term IGRs describe a new class of bio-rational compounds. (Mesbah, *et al.*, 1982; Abdel-Naby, *et al.*, 1990; Palma and Meola, 1993; Pawar, *et al.*, 1995 & Shiotsuki, *et al.*, 1999).

In recent years, much attention has been paid to the use of the *Bacillus thuringiensis* (B.t.) against *Spodoptera littoralis* (Lecadet and Martouret, 1987; Entwistle, *et al.*, 1993; Klein, *et al.*, 1996 & Abdel-Halim, 1997). *Bacillus thuringiensis* subsp. *Kurstaki* active vs. lepidopterous insects (Lereclus, *et al.*, 1989). The mixtures of Bt. with some insecticides have been evaluated against various insects. (Pree and Daly, 1996 & Abou-Taleb, 2000).

The present investigation aimed to study the efficiency of bioinsecticide (Diple-2X) either alone or in their combination with some IGRs (Diflubenzuron; Spiromesifen and Pyriproxyfen) on *Spodoptera* larvae. Also the study was directed to throw the light on the effect of these tested IGRs on the activity of chitinase.

MATERIALS AND METHODS

Insect:

Laboratory strain of cotton leafworm, *Spodoptera littoralis* was chosen for bioassays and biochemical assessment. This strain start as field strain reared under laboratory condition for several years in central lab. of pesticides, Agricultural Research Center (ARC) Cairo, Egypt. Field strain of *Spodoptera littoralis* egg masses were collected from cotton fields at Abeis area, the 2nd larval instar used for assessments.

Chemical:

Three IGRs insecticides: Diflubenzuron, 25% (WP) Novartis Co. (Syngenta). Spiromesifen, 24% (SC); was obtained from MyTrade Co., and Pyriproxyfen, 10% (EC) was obtained from Sumitomo Chemical Co.

Bioinsecticide *Bacillus thuringiensis* subsp. *Kurstaki*; Diple-2X 6.4% (WP) (32,000 International Units/mg). The product was produced by Abbott Laboratories. Chemical and Agricultural Products

¹ Central Laboratory of Pesticides, Sabahia, Alexandria, A.R.C.

² Pesticide Chemistry Dept., Fac. of Agric., Alex., Univ.

Received March 15, 2009, Accepted March 30, 2009

Division, North Chicago, USA, and Provided by Bayer Company.

Bioassay tests:

1-Toxicity of Bt.:

Second instar larvae were starved for 6hrs before exposed test. The selected larvae were bioassayed against commercial strain (Diple-2X). Using three replicates for each concentration with ten larvae in each replicate.

Disc dipping technique was used since it has been proved to be the most common procedure for assessing toxicity to commercial formulation of Bt. (Tabashnik and Cushing, 1987). Each castor leaves disc (2Cm²) was dipped into the suspension of tested formulation for 10s. Tested concentration were prepared in glass distilled water (GDW) (Toni and Fred, 1996) discs were held vertically to allow excess solution to drip off and placed on a rack to dry for at least 2hr. Treated discs were offered to starved larvae (on disc per cup) and left under constant conditions (27±2 °C). There after survivors were transferred with fresh castor oil plant leaves to clean cups and kept under the same conditions. Control larvae were allowed to fed on castor oil leave discs treated with distilled water. Mortality was percentage calculated for each concentration daily for 24; 48, and 72hrs and corrected according to Abbott (1925) and subjected to probit analysis using the computer program (Finney, 1971).

2-Toxicity of the Tested IGRs Against *S. littoralis*:

Diflubenzuron; Spiromesifen and Pyriproxyfen, were bioassayed against the 2nd larval instar *S. littoralis*. The castor leaves were dipped in different concentrations of the tested IGRs. All insecticides concentrations were prepared in distilled water. The treated leaves were placed in clean glass container at the laboratory conditions of (27±2°C) and 65-70%RH. Ten larvae (Lab and Field strains) were used for each test with three replicate. Mortality was recorded after 24; 48 and 72hr and subjected to probit analysis.

3-Toxicity of Tested Diple-2X in Presence of IGRs:

S. littoralis 2nd instar (Lab and Field strains) were treated with solution of Diflubenzuron; Spiromesifen and Pyriproxyfen at LC₅₀ values concentrations before 24; 48 and 72hr of feeding on discs of castor oil leaves discs treated with LC₅₀ of Diple-2X, joint action experiments have tow controls. Larvae of the first control were allowed to fed castor oil leaf discs treated with concentration equivalent LC₅₀ of Diple-2X alone, while larvae of the second control were fed with untreated discs. Mortality counted and recorded daily for 3days. Percentage of mortality were calculated

according to Abbott (1925) and subjected to probit analysis (Finney, 1971).

Enzyme Preparation and Activity Assay:

Chitinase was prepared from *S. littoralis*. 2nd instar larvae (Lab and Field strains) according to the method of Deul, *et al.*, (1978). Homogenate was prepared in 10⁻³M Cleland's reagent (dithiotheritol, DTT) (v/w=2). The homogenate was centrifuged for 15min at 12,000g. An equal volume of saturated ammonium sulfate solution was slowly added to the supernatant. After stirring for 1hr, the suspension was centrifuged for 10min at 10,000g. The precipitate was washed with half-saturated ammonium sulfate solution and was recentrifuged, after which it was suspended in a small volume of water, followed by dialysis 20hr. Any occasional precipitate was removed by centrifugation and was discarded as it proved to be enzymatically inactive. After dialysis, water was added to the original ratio (v/w=2). All manipulations were carried out at 0-2°C.

Chitinase activity was determined according to the method of Reissig, *et al.*, (1955) which modified by Andrew, *et al.*, (1982), using sodium acetate buffer instead of tris-HCl buffer and wave-length 416nm was used instead of 544nm. 25µl of chitin (20mg/ml), 100µl of enzyme prep and 225µl of sodium acetate buffer, (pH 4.5) in total volume 350µl. The enzyme substrate mixture was incubated at 35°C for 60min, then the reaction was stopped by adding 100µl of 0.8M borate buffer (pH10.0) followed by determination of N-acetylglucoseamine by the method of Reissig, *et al.*, (1955). By adding 1.5ml of p-dimethyl amino benzaldehyde (DMAB, reagent). The samples were incubated in a shaker water bath at 35°C for 20min, the samples were measured spectrophotometrically at λ416nm.

The protein content of *S. littoralis* 2nd instar larvae homogenates was assayed spectrophotometrically by the method of Lowery, *et al.*, (1951) at λ750nm using bovine serum albumin as a standard protein.

Inhibition of Chitinase Activity:

The inhibition of chitinase was determined in 2nd instar larvae *S. littoralis* using the LC₅₀ values of each of the three tested IGRs. The method of Dixon and Webb, (1964) was adopted to draw the Dixon-plots by plotting 1/V versus concentrations of the inhibitor at two concentrations of the substrate. Chitine (the substrate of chitinase) concentrations were 3.0 and 5.0mM. Estimation of I₅₀ value (the concentration of the inhibitor which inhibits 50% of the enzyme activity) was carried out by pre incubating the enzyme with the inhibitor for 30 min.

RESULTS AND DISCUSSIONS

Toxicity of Dipel-2X and Three IGRs:

The results of the toxicity of the Dipel-2X and IGRs in terms of LC_{50} are given in Table (1) for 2nd instar larvae of *S. littoralis*. LC_{50} values after 24hr were 7.31, 3.62, 4.25 and 5.38ppm for Dipel-2X, Diflubenzuron, Spiromesifen and Pyriproxyfen, respectively against *Spodoptera* Lab strain. For Field strain, LC_{50} values were 9.44, 4.39, 5.66 and 6.17ppm respectively. While LC_{50} values after 48hr were 5.40, 1.22, 3.41 and 4.00ppm respectively against Lab strain. For Field strain, the LC_{50} were 7.36, 2.16, 3.02 and 4.40ppm respectively. LC_{50} values after 72hr were 1.23, 0.23, 0.44 and 0.74ppm respectively against Lab strain. For Field strain, the LC_{50} were 2.13, 0.45, 0.68 and 0.91ppm respectively. According to the LC_{50} values, it is quite clear that the susceptibility of *Spodoptera* larvae to Dipel-2X, and the Lab strain of *Spodoptera* larvae is more susceptible to Dipel-2X in comparison to the Field strain. Also IGRs may act as growth distrupctor, it interferes with moulting by softening the larval endocuticle through reduction in its chitin content and by hardening of the exocuticle as result of enhanced phenoloxidase activity. These results are in agreement with many investigators, Dulmage, 1971; Ibrahim, 1974; Ascher and Nemny, 1979; Dimetry, *et al.*, 1979; Grosscurt and Anderson, 1980; Radwan, *et al.*, 1980; El-Sayed, 1981; El-Nockrashy, *et al.*, 1986; Lecadet and Martouret, 1987; Marguerre-M and Daniel, 1987; Chilcott and Ellar, 1988; Chung and Cote, 1992; Fisk and Wright, 1992; Tabashnik, 1992; Chandler, 1993; Palma and Meola, 1993; Forrester, 1994; El-Kordy, *et al.*, 1995; Pawar, *et al.*, 1995; Smagghe, *et al.*, 1997; Barker, 1998; Said, 1998; Abd-Allah, 2000; Abou-Taleb, 2000; Ali, 2001 & El-Aw, 2006.

Toxicity of Dipel-2X Alone or Pretreated with the LC_{50} Values of IGRs Against *S. littoralis* Larvae:

Data in Table (2) show the LC_{50} values of Dipel-2X are 7.31, 5.40 and 1.23ppm after 24; 48 and 72hr against Lab *Spodoptera* strain respectively, while the LC_{50} values are 9.44, 7.36 and 2.13ppm against Field *Spodoptera* strain respectively. The interaction of IGRs with Dipel-2X against Lab and Field strains of

Spodoptera larvae were studied. Larvae were allowed to feed on castor oil leave discs treated with LC_{50} of the different IGRs.

The LC_{50} values, of Dipel-2X pretreated with the LC_{50} values of Diflubenzuron; Spiromesifen and Pyriproxyfen on Lab and Field strains of *Spodoptera* larvae are presented in Table (2). The LC_{50} values of Dipel-2X when pretreated with IGRs was lower than LC_{50} of Dipel-2X alone in Lab or Field *Spodoptera* strains. The enhancement of toxicity is calculated as a Potentiation factor (P.f.) Table (2). Potentiation factor (P.f.) values for Diflubenzuron; Spiromesifen and Pyriproxyfen are 13.54, 11.97 and 9.14 respectively, after 24hr for Lab strain, while the P.f. values of three IGRs are 13.88, 12.26 and 10.04 respectively, after 24hr treatment, for Field strain. The P.f. values of three IGRs are 16.36, 13.17 and 8.57 respectively, after 48hr for Lab strain, while the P.f. values for Field strain are 13.63, 12.07 and 10.51 for three IGRs respectively. While the P.f. values of three IGRs are 9.68, 6.87 and 5.33 respectively, for Lab strain after 72hr treatment, while the P.f. values for Field strain are 13.33, 9.52 and 7.41 for three IGRs respectively. It is clear that the LC_{50} values concentrations of IGRs enhancement the toxicity of the Dipel-2X on *S. littoralis* larvae. The mixtures of Diflubenzuron+Dipel-2X were the most toxic treatments than Spiromesifen+Dipel-2X and Pyriproxyfen+Dipel-2X respectively.

In general, the susceptibility of *Spodoptera* larvae to Dipel-2X increases when treatment after IGRs. The IGRs+Dipel-2X caused more toxic effect than single treatment with Dipel-2X, it could be concluded that IGRs enhanced the toxicity effect of Dipel-2X. Based on P.f. values, the Lab strain of *Spodoptera* larvae is more susceptible to Dipel-2X in comparison to the Field strain. Generally, efficacy of IGRs have a very good additive toxicity for Dipel-2X either in Lab or Field *Spodoptera* strains. These results are agreement with finding (Salama, *et al.*, 1992; David and Joanne, 1996; Klein, *et al.*, 1996; Pree and Daly, 1996; Liburd, *et al.*, 2000 & Mona, *et al.*, 2004) whom found that when certain pairs of drugs or insecticides are administered

Table 1. LC_{50} values of Dipel-2X and three IGRs to 2nd instar *S. littoralis* larvae

Compounds	LC_{50} (ppm)					
	24hr		48hr		72hr	
	Lab strain	Field strain	Lab strain	Field strain	Lab strain	Field strain
Dipel-2X	7.31	9.44	5.40	7.36	1.23	2.13
Diflubenzuron	3.62	4.39	1.22	2.16	0.23	0.45
Spiromesifen	4.25	5.66	3.41	3.02	0.44	0.68
Pyriproxyfen	5.38	6.17	4.00	4.40	0.74	0.91

Table 2. Comparative toxicities of Dipel-2X alone or pretreated with three IGRs on *Spodoptera* larvae

Compounds	LC ₅₀ (ppm)											
	24hr				48hr				72hr			
	Lab strain	P.f.	Field strain	P.f.	Lab strain	P.f.	Field strain	P.f.	Lab strain	P.f.	Field strain	P.f.
Dipel-2X	7.31		9.44		5.40		7.36		1.23		2.13	
Diflubenzuron +Dipel-2X	0.54	13.54	0.68	13.88	0.33	16.36	0.54	13.63	0.22	9.68	0.30	13.33
Spiromesifen +Dipel-2X	0.61	11.97	0.77	12.26	0.41	13.17	0.61	12.07	0.31	6.87	0.42	9.52
Pyriproxyfen +Dipel-2X	0.80	9.14	0.94	10.04	0.63	8.57	0.70	10.51	0.40	5.33	0.54	7.41

*Potentiation factor (P.f.) = LC₅₀ Dipel-2X alone / LC₅₀ IGRs + Dipel-2X

together, the effects may be greater or less than might be expected from the sum of the activities of the components when administered separately. The phenomena involved, included under the term "synergism" "potentiation" and "antagonism", are becoming increasingly important in, for example, practical insect control and mammalian toxicology.

The observation that Dipel-2X had the lowest effect when applied alone but it was the best when mixed with IGRs. These findings may be resulted insect cuticle easily penetration which caused by IGRs in the mixture, and these results show that IGRs are act in similar manner in reducing chitin incorporation in the cuticle of *S. littoralis*. So these mixture are a good control of Lepidopterous larvae.

Generally, it could be concluded that the use of insect growth regulators (IGRs) and their mixtures with biological insecticides (Dipel-2X) instead of conventional hazardous insecticides; and these may reduce the environmental pollution and hazard effects on human health. Dipel-2X may play an important role in future insect pest management programs especially when mixed with IGRs.

***In Vivo* Inhibition of *S. littoralis* Chitinase Activity:**

The *in vivo* inhibition effect of the LC₅₀ values three IGRs against to the *Spodoptera* 2nd instar Lab and Field strains larval chitinase is shown in the data given in Table (3). The data declared that Diflubenzuron; Spiromesifen and Pyriproxyfen exhibited the high percentages of reduction of chitinase activity as values were 87.2, 80.5 and 75.4% respectively, for Lab strain, while values were 80.6, 76.9 and 70.5% respectively, for Field strain. Results indicated that Diflubenzuron in more effect on the chitinase activity than the Spiromesifen and Pyriproxyfen on 2nd instar. It is quite clear that the IGRs at LC₅₀ concentration acts as

potential inhibitors for *Spodoptera* larvae chitinase activity.

Table 3. *In vivo* inhibition of *Spodoptera* larvae 2nd instar Chitinase activity by LC₅₀ of three IGRs

IGRs	%Inhibition	
	Lab strain	Field strain
Diflubenzuron	87.2	80.6
Spiromesifen	80.5	76.9
Pyriproxyfen	75.4	70.5

***In Vitro* Inhibition of *S. littoralis* Chitinase Activity:**

Table (4) show the *in vitro* inhibition of IGRs on chitinase activity of *S. littoralis* 2nd instar. The I₅₀ values of Diflubenzuron; Spiromesifen and Pyriproxyfen for Lab strain larval chitinase are 0.54, 0.60 and 0.73 μM respectively. While the I₅₀ values are 0.60, 0.72 and 0.81 μM respectively against Field strain. To characterize in more details about the *in vitro* inhibition of chitinase by the inhibitor, the I₅₀ and Ki values of each inhibitor were estimated from the graphical method of Dixon and Webb, (1964) Table (4). The obtained data proved that IGRs competitive inhibition of chitinase activity and Ki values were 40.0 and 15.0 μM for Lab and Field strains respectively, in the case of Diflubenzuron. While these values were 52.0 and 18.0 μM for Lab and Field strains respectively, in the case of Spiromesifen. On the other hand, Ki values were 61.0 and 56.0 μM for Lab and Field strains respectively, in case of Pyriproxyfen.

In conclusion, Chitinase plays an essential role during ecdysis chitin. This enzyme is vital to moulting in insects, and may also affect gut physiology through their involvement in peritrophic membrane turnover. The exoskeleton of insects might constitute a useful target site for insecticidal chemicals.

Table 4. *In vitro* inhibition of *Spodoptera* larvae Chitinase activity by some IGRs

IGRs	I_{50} (μ M)		Ki(μ M)	
	Lab strain	Field strain	Lab strain	Field strain
Diflubenzuron	0.54	0.60	24	30
Spiromesifen	0.60	0.72	33	44
Pyriproxyfen	0.73	0.81	42	56

REFERENCES

- Abbott, W. S. (1925). A method for computing the effectiveness of an insecticide. *J. Econ. Entomol.*, *18*: 265-267.
- Abd-Allah, S. M. A. (2000). *Bacillus thuringiensis* (B.t.), Bioinsecticidal toxin agent: protein purification and study mode of action. Ph. D. Thesis. Faculty of Science, Alex. University.
- Abdel-Halim, S. M. (1997). Efficacy and residual effect of some microbial and chemical insecticides against the cotton leafworm, *Spodoptera littoralis*. (Boisd.). *Egyptian J. of Biological Pest Control*. *7*: 71-76.
- Abdel-Naby, L. M.; M. R. Farrag, and M. F. El-Sheikh (1990). Biological effects of two insect growth regulators on the Egyptian cotton leafworm *Spodoptera littoralis* (Boisd.). *J. of Agri-Sci. Mansoura Univ. (Egypt)*. *13*: 328-333.
- Abou-Taleb, H. K. (2000). Corn integrated insect management, Managing corn borers by *Bacillus thuringiensis* in the field. M. Sci. Thesis. Faculty of Agriculture, Alex. University.
- Ali, G. F. (2001). Toxicological studies on pesticide alternatives. M. Sc. Thesis submitted to Alex. University.
- Amin, A. A. A.; M. Gergis, and M. El-Naggar (2001). Alternative in field refuge strategies for controlling certain cotton key pests in middle Egypt. The ESA 2001 Annual Meeting-December 2001: An Entomological Odyssey of ESA, San Diego, CA.
- Andrew, C. C.; Mayer, R. T., and De-loach, J. R. (1982). Purification and characterization of chitinase from the stable fly *Stomoxys calcitrans*. *Arch. Biochem. Biophys.*, *216*: 314-321.
- Ascher, K. R. S., and N. E. Nemny (1979). Toxicity of chitin synthesis inhibitors diflubenzuron and its dichlorobenzoyl analogue to *Spodoptera littoralis* larvae. *Pestic., Sci.*, *7*: 1-9.
- Barker, J. F. (1998). Effect of *Bacillus thuringiensis* supsp. *Kurstaki* toxin on the mortality and development of the larval stages of the banded sunflower moth (Lepidoptera: Cochylidae). *J. Econ. Entomol.*, *9*: 1084-1088.
- Chandler, L. D. (1993). Use of feeding stimulants to enhance insect growth regulator induced mortality of the fall armyworm (Lepidoptera: Noctuidae) larvae. *Florida Entomol.*, *76*: 316-326.
- Chilcott, C. N., and D. J. Ellar (1988). Comparative toxicity of *Bacillus thuringiensis* var *israelensis* crystal proteins *in vivo* and *in vitro*. *J. Gen. Microbiol.* *134*: 2552-2558.
- Chung, Y. S., and C. J. Cote (1992). Crystal of *Bacillus thuringiensis* two mutants of show increased toxicity against larvae of *Spodoptera littoralis*. *Entomophga*, *37*:193-196.
- Deul, D. H.; De-Jong, B. J., and kortenbach. J. A. M. (1978). Inhibition of chitin synthesis by two 1-(2,6-disubstituted benzoyl)-3-phenylurea insecticides. *Pesticide Biochemistry and Physiology*. *8*: 98-105.
- Dimetry, Nadia Z., and M. A. Ramadan (1979). Effect of insect growth regulator on the cotton leafworm, *Spodoptera littoralis* (Boisd.). 3rd Pest. Conf. Tanta Univ., *1*: 85-91.
- Dixon, M., and E. C. Webb (1964). *Enzymes*. Academic press, Inc., New York. 2nd edition., 328-330.
- Dulmage, H. T. (1971). A proposed standardized bioassay for formulations of *Bacillus thuringiensis* based on the international unit. *J. Invertebr. Pathol.*, *18*: 240-345.
- El-Aw, A. M. (2006). Efficacy of two biorational insecticides, emamectin benzoate and spinosad, against the immature stages of the cotton leafworm, *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *Alex. J. Agric. Res.*, *51*: 55-68.
- El-Kordy, M. W.; A. I. Gadallah; M. G. Abas, and S. A. Mostafa (1995). Effect of Pyriproxyfen; flufenoxuron and teflubenzuron on some biochemical aspects of *S. littoralis*. A1-Azhar. *J. Agric. Res.*, *21*: 223-238.
- El-Nockrashy, A. S.; H. S. Salama, and F. Taha (1986). Influence of bait formulations on the effectiveness of *Bacillus thuringiensis* against *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *J. Appl. Entomol.*, *101*: 389-299.
- El-Sayed, E. I. (1981). Effect of diflubenzuron and diflubenzuron hostathion combinations on larvae and adults of the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.). *Bull. Ent. Soc. Egypt, Econ. Ser.*, *12*: 195-201.
- Entwistle, P. M.; J. Baily; J. Cory, and S. Higgs (1993). *Bacillus thuringiensis* an environmental biopesticide wiley, New York. 172-185.
- Finney, D. J. (1971). *Probit analysis*, 3rd ed Cambridge Univ. Press, Cambridge, England.

- Fisk, T., and D. J. Wright (1992). Response of *Spodoptera exempta* (Walk.) larvae to stimulated field spray applications of acylurea insect growth regulators with observations on cuticular uptake of acylureas. *Pesticide Since.*, 35: 331-337.
- Forrester, N. W. (1994). Use of *Bacillus thuringiensis* in integrated control, especially on cotton pestes Agriculture, Ecosystems and Environment, 49: 77-83.
- Grosscurt, A. C., and S. O. Anderson (1980). Effects of diflubenzuron on some chemical and mechanical properties of the eiyra *Leptinatarsa decemlineata*. *Proc. Kon. Ned. Acad. Wetenschappen.* 83: 143-150.
- Ibrahim, A. A. (1974). Studies on the biological control of *Spodoptera littoralis* (Boisd.) in A.R.E., Ph.D. Thesis Faculty of Agriculture. Cairo University. 282pp.
- Ishaaya, I., and M. Kleen (1990). Response of susceptible laboratory and resistant field strains of *Spodoptera littoralis* (Lepi. Noct.) to teflubenzuron. *J. Econ. Entomol.*, 83: 59-62.
- Keddis, M. E.; A. F. Ayad; S. M. Abdel-Fattah, and A. M. El-Guindy (1988). Studies of resistance to urea derivatives and their mixtures with insecticides in field strains of cotton leafworm, *Spodoptera littoralis* (Boisd.) during the cotton seasons 1983, 1984 and 1985. *Bull. Ent. Soc. Egypt, Econ. Ser.*, 15: 229-234.
- Klein, C. D.; R. D. Johnson; B. H. Myers, and M. L. Page (1996). Evaluation of *Bacillus thuringiensis* and ovicide combinations for bollworm and tobacco budworm. *Proc. Beltwide Cotton Conf., Nashville, TN, USA*, 2: 789-799.
- Lecadet, M. M., and D. Martouret (1987). Host specificity of the *Bacillus thuringiensis* α -endotoxin toward lepidopteran species: *Spodoptera littoralis* Bdv. and *Pieris brassicae* L. *J. Invert. Pathol.*, 49: 37-48.
- Lereclus, D.; C. Bourguoin; M. M. Lecader; A. Klier, and G. Rapoport (1989). Role structure, and molecular organization of the genes coding for the parasporal δ -endotoxins of *Bacillus thuringiensis*. In Regulation of prokaryotic development. (Smith, R. A. Slepecky, and Setlow, P. ed.). American Society for Microbiology, Washington. D. C. p.225-276.
- Lourd, E. O.; J. E. Funderburk, and S. M. Olson (2000). Effect of biological and chemical insecticides on *Spodoptera* species (Lep., Noctuidae) and marketable yield of tomatoes. *J. of Applied Entomol.*, 124: 19-25.
- Lowery, H. O.; Rosbrough, N. J.; Farr, A. L., and Ranball, R. J. (1951). Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Marguerre-M, L., and M. Daniel (1987). Host specificity of the *Bacillus thuringiensis* δ -endotoxin toward lepidopteran species: *Spodoptera littoralis* Bdv. and *Pieris brassicae* L. *J. Invert. Pathol.*, 49: 37-48.
- Mesbah, H. A.; A. N. Hassan; A. S. M. Radwan; A. R. M. Abdel-Mohymen, and A. S. M. Abdel-Fattah (1982). Synergism between certain adjuvants and two insect growth regulators against the larvae of the cotton leafworm *Bull. Ent. Soc. Egypt. Econ. Ser.*, 3: 25-30.
- Mona, B. R. El-Mandarawy; Moustafa, A. Z. El-Naggar, and Saadia, A. Abdel-Samae (2004). Evaluation of Feeding stimulant mixed with chemical and bio insecticides on certain Lepidopterous pests and their natural enemies in cotton and soybean fields. *J. Appl. Sci. Environ. Mgt.*, 8: 77-83.
- Palma, K., and R. Meola (1993). Mode of action of pyriproxyfen and smethoprene on cat flea (*Ctenocephalides felis*) eggs. *J. Med. Entomol.*, 421-426.
- Pawar, P. V.; P. S. Pisale, and N. R. Sharma (1995). Effect of some new insect growth regulators on metamorphosis & reproduction of *Aedes aegypti*. *Indian J. Med. Res.*, 101: 8-13.
- Pree, J. D., and C. J. Daly (1996). Toxicity of mixtures of *Bacillus thuringiensis* with endosulfan and other insecticides to the cotton bollworm *Helicoverpa armigera*. *J. Pestic. Sci.*, 48: 199-204.
- Radwan, H. S. A.; G. A. M. Rizk, and D. M. Assal (1980). The dose toxicity of certain insect growth regulators on cotton leafworm in relation to development inhibitory effect. (*Proc. Arab Pest Conf., Tanta Univ.*, 1: 415-422).
- Reissig, J. L.; Strominger, J. L., and Leloir, L. F. (1955). A modified colorimetric method for the estimation of N-acetylamine sugars. *J. Biol. Chem.*, 217: 959-966.
- Rizk, M. A.; A. G. El-Sisi; N. A. Badr, and S. M. Abdel-Halim (1999). Controlling of cotton sucking pests using safe materials. 2nd Int. Conf. of Pest Control, Mansoura, Egypt, Sept., Special Proc., 211-221.
- Said, A. M. (1998). Bioefficacy of certain insect growth regulators on the black cutworm, *Agrotis ipsilon* (Hufu.) *Egypt. J. Appl. Sci.*, 13: 113-120.
- Salama, H. S.; M. S. Abd El-Baset, and M. Ragaei (1992). Mode of action of chemical additives in enhancing the potency of *Bacillus thuringiensis* against Lepidopterous insects. *J. Appl. Entomol.*, 114: 167-173.
- Shiotsuki, T.; F. Yukuhiro; M. Kiuchi, and E. Kuwano (1999). Effect of 1-(4-phenoxyphenoxy) imidazole (KS-175) on larval growth in the silkworm *Bombyx mori*. *J. Insect Physiol.*, 45: 1040-1047.
- Smagghe, G.; M. Auda; K. Laecke, and D. Van Degheele (1997). Significance of penetration and metabolism on topical toxicity of diflubenzuron in *Spodoptera littoralis* and *Spodoptera exigua*. *Entomol. Exp. Et Applicata.*, 82: 255-260.
- Tabashnik, B. E. (1992). Evaluation of synergism among *Bacillus thuringiensis* toxins. *Appl. Environ. Microbiol.*, 58: 3343-3346.
- Tabashnik, B. E., and L. N. Cushing (1987). Leaf residue vs. topical bioassays for assessing insecticide resistance in the diamond back moth *Plutella ylostella*. *F. A. O. Plant Prot. Bull.*, 35: 11-14.
- Toni, M., and G. Fred (1996). Effect of surfactants, *Bacillus thuringiensis* formulations, and plant damage on oviposition by diamond back moth (Lepidoptera: plutellidae). *J. Econ. Entomol.*, 89: 891-897.

الملخص العربي

تقييم بعض المركبات الآمنة بيئياً ضد دودة ورق القطن

سهام منصور إسماعيل، عمود مرشدى

وهو أنزيم الكيتينيز ولقد أوضحت النتائج أن في حالة الدايفلوبيتريون كانت النسبة المئوية للتشيط هي ٨٧,٢ و ٨٠,٦% للسلاطة العملية والحقلية على الترتيب، بينما في حالة السبرومييسيفين كانت النسبة المئوية للتشيط هي ٨٠,٥ و ٧٦,٩% لكلا من السلاطة العملية والحقلية على التوالي، ولقد كانت النسبة المئوية للتشيط بواسطة البيريروكسيفين هي ٧٥,٤ و ٧٠,٥% للسلاطة العملية والحقلية على الترتيب. وكذلك تم دراسة تأثير منظمات النمو الحشرية المخترة على قيم I_{50} أوضحت النتائج حدوث زيادة في الفعل الأبادى للدابليل-٢ أكس على يرقات العمر الثانى للدودة ورق القطن المعاملة من قبل منظمات النمو الحشرية. ومن هذه النتائج نجد أن الخلاط أعطت تأثير أكبر من الدابليل-٢ أكس ومنظمات النمو الحشرية عند تطبيقهم بصورة فردية مما يوضح أن منظمات النمو الحشرية تنشيط عمل الدابليل-٢ أكس ولذلك تعتبر هذه الدراسة خطوة في اتجاه إستخدام هذه المخالط كأحد عناصر المكافحة المتكاملة لدودة ورق القطن حيث إنها أكثر أماناً للإنسان والبيئة.

الهدف من البحث هو تقييم التأثير الأبادى للمبيد الحيوى دايبيل-٢ أكس مع ثلاث تجهيزات من منظمات النمو الحشرية هى الدايفلوبيتريون؛ سبرومييسيفين وبيريروكسيفين ومخالطهما وذلك على يرقات العمر الثانى لدودة ورق القطن للسلاطة العملية والحقلية بهدف تلاشى تأثير المبيدات التقليدية الضار على البيئة. وقد تم تسجيل قيم التركيزات النصف مميتة (LC_{50}) لكلا من الدابليل-٢ أكس ومنظمات النمو الحشرية تحت الدراسة بصورة فردية. ثم تم معاملة اليرقات العمر الثانى لدودة ورق القطن بتركيزات مختلفة (LC_{50}) من منظمات النمو الحشرية تحت الدراسة ثم معاملة هذه اليرقات بتركيز (LC_{50}) من الدابليل-٢ أكس بعد ٢٤ و ٤٨ و ٧٢ ساعة من المعاملة. منظمات النمو الحشرية تحت الدراسة، فأوضحت النتائج أن قيم LC_{50} بعد المعاملة أنخفضت بدرجة ملحوظة ويتضح ذلك من قيم معامل التنشيط (P.f.) الذى تم حسابها. وكذلك تم دراسة المقدرة التشيطية لمنظمات النمو الحشرية المخترة على النشاط الأنزيمى لأنزيم هام وحيوى بالنسبة للحشرة