



## PRODUCTIVITY AND NUTRITIVE VALUE OF NATURAL VEGETATION AT WADI-WATEER REGION IN SINAI

[7]

Baumi<sup>1</sup>, N.H.M. and Nassar, Zienab<sup>1</sup>, M.

1- Range management Unit, Desert Research Center, Matariya, Cairo

**Keywords:** Wadi wateer, Aridity, Plant associations, Natural vegetation, Pasture measurements

### ABSTRACT

The present study was carried out in 2004/2005 – 2005/2006 seasons to make survey of plant vegetation at Wadi Wateer (East southern of Sinai) (a bout 100 km length) at 9 successive sites. The study was carried through two years (two wet and dry seasons) to evaluate the vegetation structure throughout 20 chart quadrates (2 m x 2 m) at each site. The results showed that most of the pasture measurements such as plant density (plants / 4 m<sup>2</sup>), coverage %, frequency %, abundance % and fresh, and dry yields (ton/fad.) reached its maximum values during the wet seasons. Sites 5, 6, 7, 8 and 9 were more suitable for growth of the plant associations than other sites under studies. Many plant species such as *Astragalus spinosus*, *Medicago laciniata*, *Artemisia Judica*, *Artemisia monosperma*, *Legum spartum*, *Haloxylon salicornicum*, *Calotropis procera*, *Lycium shawii* and *Paronychia sinaica* were capable to grow under the aridity conditions at Wadi Wateer area.

### INTRODUCTION

Natural vegetation are mainly dependent on the prevailing environmental conditions in the desert, such conditions play a great role in controlling the seasonal aspects of the floral vegetation.

East southern of Sinai, Egypt is considered the minimum district in its annual winter rainfall, were the average 16.17 mm at Nowibaa and 17.6 mm at Nikhil with uneven distribution, low and medium relative humidity through 5 months, were various plant communities can be grow during such limited rainy seasons. Natural vegetation in most countries of the world consider as main resource for supporting the wild and domestic animals with its basic forage requirements. The productivity and

quality in relation to prevailing environmental conditions need to be assessed and clarified in order to get more Justification for the optimum and adequate use for better range management.

Ayyad and Ghabbour (1977) revealed that the existing environmental conditions have an important role in range flourishing productivity and nutritive value.

Moreover, moisture availability for vigorous growth of native plants is limited, under arid land condition, however profit feasible biomass production is associated with adequate precipitation Abou Dyea and Salem (1990 a) and Habbs et al (1984).

Abou Dyea and Salem (1990 b) in El-Negila area they found that the genera of legumes and grasses had the greatest number of species. Density (plant/ m<sup>2</sup>), coverage %, abundance % and frequency % varied widely from one season to another, where different vegetation characters attained its maximum in spring and winter seasons. Reiad et al (1996b) studied the effect of locations and environmental conditions in Sidi- Barrani and El- Negila area on vegetative characters. They reported that the vegetative measurements varied according to the studied locations and growing seasons and the highest averages values were recorded at Sidi- Barrani due to its favorable environmental conditions. Hendawy (2002) studied the effect of eight different sites at Gabal El- Maghara region in middle Sinai on the natural vegetation. He found that the highest values of the most parameters were recorded during the spring seasons. This is could be due to the increase in available relative humidity which result in more growth activity. El-Toukhy et al (2002) studied the productivity of some associations and seasonal locations as well as climatic factors on growth parameters at wadi El- Natron , Al- Alameen road in the North Western Coast (NWC) of Egypt under 5 different sites condition. The results showed that sites 5 and

(Received November 15, 2008)

(Accepted January 10, 2009)

1 produced higher yields during winter seasons, where these sites more correlated by the availability of rains as well as the other favourable climatic factors.

Ahmed and Nassar (1999) studied the chemical composition of 108 samples of various range plants belonging to 19 families from different sties along the NWC of Egypt. The results indicated that the chemical composition of the concerned range plants varied within the different species under different habitat conditions.

#### MATERIALS AND METHODS

Natural range vegetation depends mainly on the amount of rainfall which widely varies in quantity and frequency from year to year. Moreover, various types of range plants are closely associated with the edaphic and prevailing climatic conditions of the studied areas. This investigation was conducted to asses the productivity and the nutritive values for the present range plants in Wadi Wateer area in east south Sinai. This study was conducted in two seasons of nine successive sites during the two years (from autumn 2004 up to spring 2006).

Nine sites were determined inside Wadi Wateer beginning from the 260 km at the international road at middle Sinai to Nowibaa city, each site was determined after each consecutive 10 km distance in the direction of Nowibaa. The different sites are located at 8, 19, 27, 36, 48, 55, 66, 78, 95 km from the beginning of the wadi, respectively.

Means of some meterological data of Nowibaa and Nikhl city during the period extended from autumn 2004 up to spring 2006 are presented in Table (2).

Twenty chart quadrates of 2 m x 2m tool were used to determine the plant density (plants/ 4 m<sup>2</sup>), coverage %, frequency %, abundance % of the natural vegetation. Number of individual species and the area occupied by them were determined from twenty stand sampling over the nine sties during both wet (spring) and dry (autumn) seasons.

Hanson and Churchill (1965) equations were used to calculate the previous measurements as the following

$$\text{Density (plant/m}^2\text{)} = \frac{\text{Number of the individual species}}{\text{Total area (in units)}}$$

$$\text{Coverage \%} = \frac{\text{The area occupied by the species}}{\text{The whole investigated area (In units)}} \times 100$$

$$\text{Frequency \%} = \frac{\text{Number of occurrence of the individual species}}{\text{Number of occurrence of the whole species}} \times 100$$

$$\text{Abundance \%} = \frac{\text{Number of the individual species}}{\text{Total number of the whole species}} \times 100$$

#### Fresh and dry production

Fresh production was determined by weighting each species in a square meter and for whole plant species in unit area, then calculated in (ton/fad).

#### Dry production

Samples were randomly taken from the fresh matter of each species and then dried in an air forced drying oven at 70°C for a constant weight (dry weight) during the two seasons.

#### Plant identification

Plant identification was done according to Täckholm (1974) and Boulos (2005). The list of the collected plants included family name, scientific name, palatability and life duration Aslan, (1959).

#### Chemical composition

Chemical composition of dry matter of each of the palatable plant species were determined as follows

Crude protein (CP) was estimated by using modified micro kjeldahl method as described by Peach and Tracey (1956) to determine total nitrogen and then multiplied by 6.25.

Crude fiber (CF) and total ash were determined according to A.O.A.C. (1970).

Ether Extract (EE) was extracted by Hexane in soxhelt according to A.O.A.C (1970).

Sodium and potassium contents of plant samples were estimated using flame- photometer in an acid digestion solution according to A.O.A.C. (1980).

Total digestible nutrient (TDN) was determined according to Khafagi (1977).

#### Soil physical and chemical properties

Soil samples were taken from the different sites from two depths (0- 20) and (20-40) ; physical analysis was conducted by using the international pipette method described by Piper (1950). Chemical determinations were carried out according to Jackson (1958) as presented in Table (1).

Table 1. Soil physical and chemical properties of nine studied sties in Wadi Water during the period extended from autumn 2004 up to spring 2006

Site	Depth (cm)	pH	E.C m.mhos/cm	Cations (me/L)				Anions (me/L)				CaCO <sup>3</sup> %	Texture
				Ca <sup>++</sup>	Mg <sup>++</sup>	Na <sup>+</sup>	K <sup>+</sup>	Cl <sup>-</sup>	So <sub>4</sub> <sup>-</sup>	HCO <sub>3</sub> <sup>-</sup>	CO <sub>3</sub>		
1	0-20	7.80	1.15	2.45	6.14	2.81	1.13	8.40	0.71	2.08	-	27.81	sandy
	20-40	7.74	1.19	2.56	6.18	1.19	1.82	8.34	0.88	2.81	-	21.46	sandy
2	0-20	7.70	1.20	2.63	7.00	1.30	1.91	9.14	0.91	1.89	-	31.40	Loamy sandy
	20-40	7.65	1.13	2.34	6.99	1.45	1.95	9.80	0.89	2.00	-	22.14	Loamy sandy
3	0-20	7.62	1.40	2.14	7.18	1.22	1.53	8.20	0.75	2.03	-	27.33	Loamy sandy
	20-40	7.79	1.55	3.33	7.81	2.40	2.67	10.95	0.66	2.91	-	30.01	Loamy sandy
4	0-20	7.56	1.30	2.18	6.94	3.5	1.13	10.88	0.95	2.13	-	30.41	Loamy sandy
	20-40	7.61	1.42	3.88	6.14	3.01	1.01	10.77	0.69	2.18	-	24.50	Loamy sandy
5	0-20	7.80	1.81	4.00	7.19	4.22	2.18	14.21	0.95	3.20	-	26.18	Loamy sandy
	20-40	7.55	1.94	4.81	6.90	5.21	2.11	14.99	0.78	4.03	-	28.29	Loamy sandy
6	0-20	7.40	1.90	4.85	7.34	4.88	2.10	13.80	0.88	4.80	-	33.15	Loamy sandy
	20-40	7.88	1.93	4.70	6.93	5.00	2.81	12.30	0.99	5.00	-	35.19	Loamy sandy
7	0-20	7.69	1.40	2.88	5.70	2.71	2.73	8.85	0.69	3.00	-	31.45	Loamy sandy
	20-40	7.67	1.48	2.74	5.37	3.13	2.18	7.40	0.94	5.71	-	30.80	Loamy sandy
8	0-20	7.43	2.90	10.19	8.14	6.55	4.30	20.94	0.83	6.84	-	30.18	Loamy sandy
	20-40	7.52	2.40	8.80	6.92	5.17	2.22	20.33	0.76	2.45	-	31.44	Loamy sandy
9	0-20	7.49	2.30	9.71	6.87	5.18	2.19	19.82	0.63	2.84	-	35.15	Loamy sandy
	20-40	7.50	2.17	8.81	6.00	5.14	2.01	2.01	0.74	3.07	-	35.13	Loamy sandy

Table 2. Means of some meteorological data of Nowibaa and Nikhl city during the period extended from autumn 2004 up to spring 2006

month	Nowibaa											
	Jan.	Feb.	Mar.	April	May.	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
Temperature (°C)												
Maximum	21.0	21.8	24.5	28.3	32.5	35.7	37.5	37.0	34.4	30.5	26.2	22.6
Minimum	13.6	14.0	16.3	19.2	22.7	24.8	26.8	27.3	25.6	23.1	19.2	15.5
Relative humidity % (R.H.)	49	47	49	50	49	48	50	52	55	55	51	50
Rainfall (mm)	5.8	1.4	0.7	1.2	1.03	0	0.04	0	0	2.7	2.9	0.4
Wind speed (km/h)	15.7	15.3	15.5	14.8	13.5	13.8	10.9	11.1	13.1	11.2	13.1	14.8
	Nikhl											
Temperature (°C)												
Maximum	18.0	19.5	23.0	28.2	32.4	34.2	37.0	35.8	32.8	28.9	24.3	19.4
Minimum	1.4	2.4	5.1	8.7	12.0	14.4	16.9	17.5	15.1	12.8	7.2	2.4
Relative humidity % (R.H.)	63	58	55	48	45	50	49	55	59	64	62	63
Rainfall (mm)	3.7	2.4	0.4	0.2	1.1	0	0	0	0	5.9	0.3	3.6
Wind speed (km/h)	10.5	11.4	12.0	11.6	10.5	8.8	7.4	9.62	9.2	7.9	7.4	8.5

Source: Meteorological authority, Cairo.

## RESULTS AND DISCUSSION

### Vegetative growth traits

### Botanical composition

Data presented in Table (3) showed that the natural vegetation components of Wadi Wateer consisted of 33 plant species belonged to 15 families. Most of the existing plant species in this area 27 species were perennials. Whereas, the annuals species were 6 plants species, nineteen species were palatable and the other 14 were unpalatable. The highest number of the following plant species i.g., *Artemisia Judica*, *Artemisia monosperma*, *Peganum harmala*, *Zygophyllum album*, *Haloxylon salicornicum*, *Zilla spinosa* and *Calotropis procera* were noticed in both wet and dry seasons at most sites. The dominant of these plant species compared with other plant species plant could be due to its wide adaptability and capability to grow under the prevailing.

### Density (Plant / 4 m<sup>2</sup>)

Data in Table (4) indicated that the density of the native plant species were greatly higher in wet season compared with that of dry season. The total density of the nine investigated sites amounted 35.6 (plant / 4m<sup>2</sup>) in wet season compared with 17.65 (plant/ 4m<sup>2</sup>) in dry season. The density of wet season were recorded with sites, 9, 4, 7 and 5 and with sites 9 and 8 in dry season. The density of these sites were 10.95, 6.15, 5.50, 3.20 (plant / 4 m<sup>2</sup>) in wet season and in dry season 6.10, 5.25 (plant / 4 m<sup>2</sup>), respectively. The above results proved plant density at that Wadi Wateer in wet season at sites 4, 6, 7, 9 showed that superiority over of sites 1, 2, 3, 5, 8. This finding means that the vegetative growth activities are more dependable on the suitability and convenience of the prevailing environmental conditions, since it grow well under the adequacy of the available water, moderate temp. and dew.... etc. as presented in Table (2), this could be attributed mainly to many edaphic factors as soil texture, water holding capacity, organic matter, salt content Table (1).

Similar results were previously obtained by Ibrahim (1995) and Reiad et al (1996 a) in North Western Coast of Egypt. They found that Sidi - Barrani associations had the highest plant density over all seasons than El-Negila area. The plant density was the highest value during the winter period (wet season) at Sidi - Barrani and El-

Negila than dry season. El-Toukhy et al (2002) in Al-Alameen road (NWC), El-Morsy (2002) in NWC of Egypt and Ibrahim et al (2006) in North Eastern Coast of Egypt.

### Coverage %

Data in Table (5) showed that the highest plant coverage percentages in Wadi Wateer were recorded at sites 4, 5, 6, 7, 8, 9 which were (11.88, 11.37, 12.65, 11.48, 12.75 and 14.27%) in wet season and sites (4, 5, 6, 7, 8,9) which were (9.74, 7.91, 12.54, 10.04, 9.25, 9.94) in dry season. The superiority of coverage % in wet season than that of dry season may be attributed to the precipitation and environmental effects and the relatively appropriate soil (Tables 1, 2). Similar results were reported by Ibrahim (1995) in NWC of Egypt ; Reiad et al (1996 a & b) in NWC of Egypt; Hendawy (2002) in Gabal El Maghara region, Middle Sinai, El- Morsy (2002) in Wadi Magid and Wadi Mehgun at NWC of Egypt and Ibrahim et al (2006) In North Eastern Coast of Egypt. Plant species of the highest coverage percentage at Wadi Wateer were *Artemisia Judica*, *Artemisia monosperma*, *Haloxylon salicornicum*, *Zilla spinosa*, *Calotropis procera* and *Lyceum shawii*.

### Frequency %

Results in Table (6) represent the effect of sites and growing seasons on the frequency of the naturally growing plant species. It is obviously clear that the site 6 in wet seasons was of the highest frequency compared with other sites in wet and dry seasons. The highest frequencies of plant species were for *Artemisia Judica*, *Artemisia monosperma* and *Haloxylon salicornicum* in dry and wet seasons. This finding revealed that these plant species are dominant in this area and this could be due to the effect of edaphic factors which act as promoter for the richest and most adaptive to the dominant environment conditions. Abd-Alla (1999) in North Western Coast of Egypt; Hendawy (2002) in Gabal El Maghara region, Middle Sinai and Ibrahim et al (2006) In North Eastern Coast of Egypt had similar results.

### Abundance %

Data in Table (7) showed the effect of sites and growing seasons on the abundance of the naturally grown plant species. The highest abundance % of natural plant species were noticed in sites 5, 6, 7, 8, 9 in both dry and wet seasons. This may be due

Table 3. Effect of sites and growth seasons on palatability and duration of all plant species recorded in wadi Water district during autumn 2004 up to spring 2006

Family name	No.	Scientific name	Palatability	Duration	S1		S2		S3		S4		S5		S6		S7		S8		S9	
					Au	Sp	Au	Sp	Au	Sp	Au	Sp	Au	Sp	Au	Sp	Au	Sp	Au	Sp	Au	Sp
Leguminosae	1	<i>Acacia tortilis sub sp. raddiana</i>	P	per																	+	
	2	<i>Acacia nilotica</i>	P	per								+									+	+
	3	<i>Alhagi graecorum</i>	P	per						+							+				+	+
	4	<i>Astragalus spinosus</i>	P	per						+	+	+	+		+	+		+			+	+
	5	<i>Retama raetam</i>	P	per												+	+				+	+
	6	<i>Medicago laciniata</i>	P	Ann										+		+		+	+	+	+	+
	7	<i>Melilotus indica</i>	P	Ann																		+
	8	<i>Melilotus sulcatus</i>	P	Ann																		+
	9	<i>Trifolium tomentosum</i>	P	Ann																		+
Compositae	10	<i>Achilla fragrantissima</i>	up	per										+			+			+	+	
	11	<i>Artemisia judaica</i>	P	per				+	+			+	+			+	+	+	+	+	+	+
	12	<i>Artemisia monosperma</i>	P	per				+	+		+	+	+		+	+		+	+	+	+	+
Gramineae	13	<i>Glebionis coronaria</i>	up	per										+		+		+	+	+	+	+
	14	<i>Legum spartum</i>	P	per								+			+	+		+	+	+	+	+
Zygophyllaceae	15	<i>Fagonia bruguieri</i>	up	per						+				+								
	16	<i>Fagonia glutinosa</i>	up	per											+						+	
	17	<i>Peganum harmala</i>	up	per	+			+	+	+	+	+	+	+		+				+	+	+
	18	<i>Zygophyllum album</i>	up	per				+	+	+		+	+	+	+	+	+	+	+	+	+	+
	19	<i>Zygophyllum decumbens</i>	up	per														+	+			
	20	<i>Zygophyllum simplex</i>	up	Ann											+				+			
Chenopodiaceae	21	<i>Anabasis setifera</i>	up	per						+	+		+	+	+	+				+		
	22	<i>Haloxylon salicornicum</i>	up	per	+	+		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Tamaraceae	23	<i>Tamarix aphylla</i>	p	per											+					+		
	24	<i>Tamarix decumbens</i>	p	per								+	+	+								+
Nitrariaceae	25	<i>Nitraria retusa</i>	p	per											+		+	+				+
Cruciferae	26	<i>Zilla spinosa</i>	up	per				+	+		+	+		+	+	+		+	+	+	+	+
Asclepiadaceae	27	<i>Calotropis procera</i>	up	per				+	+	+	+	+	+	+	+	+		+	+		+	+
Cleomaceae	28	<i>Cleome Africana</i>	up	per												+	+	+				
Solanaceae	29	<i>Lyceum shawii</i>	p	per								+	+	+		+		+			+	+
Capparaceae	30	<i>Capparis spinosa var. canescens</i>	up	per													+	+		+	+	
Polygonaceae	31	<i>Calligonum comosum</i>	p	per						+						+		+				
Caryophyllaceae	32	<i>Paronychia sinacia</i>	p	per								+						+				
Labiatae	33	<i>Phlomis aurea</i>	up	per										+			+					+

Table 4. Plant density (plant /4m<sup>2</sup>) of plant species in Wadi Wateer as affected by sites and seasons during the period extended from autumn 2004 up to spring 2006

No.	Dry seasons										Wet seasons										
	1	2	3	4	5	6	7	8	9	Total	1	2	3	4	5	6	7	8	9	Total	
1								0.05		0.05											
2								0.05	0.05	0.10			0.05					0.05	0.10		
3			0.10					0.05		0.15					0.05		0.05		0.10		
4			0.05	0.1		0.05	0.05	0.05		0.30			0.05	0.05	0.05				0.05	0.20	
5						0.05		0.05		0.10					0.05		0.05		0.10		
6								2.50	5.00	7.50				0.50	1.00	2.50	2.00	1.50	7.50		
7									0.25	0.25											
8																		1.50	1.50		
9																		2.50	2.50		
10								0.05	0.05	0.10				0.05			0.05		0.10		
11		0.05			0.15		0.05			0.25			0.15	0.20	0.05	0.05	0.15	0.20	0.80		
12		0.05		0.05				0.05	0.05	0.20			0.05	0.10	0.10	0.05	0.10	0.15	0.05	0.65	
13								0.70		0.70				0.50	100	1.00	1.50	4.50	8.50		
14				0.05		0.05		0.05		0.15					0.05	0.05	0.05	0.05	0.20		
15													0.05	0.10					0.15		
16						0.05				0.05							0.05		0.05		
17	0.2	0.05	0.20	0.25	0.30			0.05	0.05	1.10			0.10	0.25	0.25	0.15	0.05	0.05	0.05	0.90	
18		0.30	0.15	0.15	0.15	0.20	0.15		0.15	1.25			0.15		0.05	0.05	0.20	0.05	0.10	0.65	
19							0.10			0.10							0.05		0.05		
20														0.05		0.05			0.10		
21			0.05			0.05		0.05		0.15			0.15	2.50	0.25	0.20				3.10	
22	0.8	0.4	0.40	0.25	0.40	0.35	0.35	0.45	0.30	3.7		1.00	0.40	0.3	0.20	0.15	0.30	0.10	0.15	2.75	
23						0.05		0.05		0.10											
24				0.05	0.05					0.10				0.05	0.05				0.05	0.15	
25						0.05	0.05			0.10				0.05		0.05		0.05	0.05	0.15	
26		0.15		0.05		0.05		0.05	0.05	0.35			0.15	0.05	0.05	0.05	0.05	0.05	0.05	0.50	
27			0.10	0.05	0.05	0.05			0.05	0.30			0.05	0.15	0.05	0.05	0.05		0.05	0.45	
28							0.05			0.05						0.10	0.05			0.15	
29				0.05					0.05	0.10				0.05	0.05	0.10	0.05		0.05	0.30	
30							0.20		0.05	0.25							0.10	0.05		0.15	
31													0.1		0.05	0.05				0.2	
32														2.50		1.00				3.50	
33							0.05		0.05	0.10				0.05						0.05	
T	1.00	1.00	1.05	1.00	1.15	1.00	1.10	4.25	6.10	17.65		1.00	1.05	1.20	6.15	2.15	3.20	5.50	4.40	10.95	35.6

T. Total

Table 5. Coverage percentage of plant species in Wadi Water as affected by sites and seasons during the period extended from autumn 2004 up to spring 2006

No.	Dry seasons										Wet seasons									
	1	2	3	4	5	6	7	8	9	M.	1	2	3	4	5	6	7	8	9	M.
1								+												
2								1.25	1.10	0.26				1.25					1.20	0.27
3			0.35						0.25	0.06						0.70		0.20		0.10
4			0.43	0.26		1.12	0.43	0.25		0.27			0.30	0.40	0.25				0.70	0.18
5						2.00		1.25		0.36						2.10		1.20		0.37
6								0.25	0.56	0.09					0.10	0.25	0.25	0.45	0.25	0.14
7									0.10	0.01										
8																			0.70	0.08
9																			0.90	0.10
10								0.15	0.07	0.02					0.43			0.20		0.07
11		0.43			0.48		0.25			0.13	0.87		0.90		0.18	0.40	2.20	0.65	0.58	
12		0.15		0.80				0.18	1.25	0.26	0.18	0.50	0.50	0.26	0.40	0.70	2.30	0.45	0.59	
13								0.07		0.01					0.25	1.10	0.25	1.10	1.00	0.41
14				1.25		0.15		0.90		0.25						0.30	0.60	1.00	0.80	0.30
15												0.25		0.80						0.12
16						0.80				0.08								0.15		0.02
17	0.29	0.18	0.97	1.10	1.40			0.52	0.40	0.54	0.06	0.80	0.90	1.50	0.37		0.50	0.56	0.59	
18		1.33	0.88	0.48	0.53	0.70	0.23		0.48	0.51	1.41		1.00	0.37	0.56	0.50	0.05	1.10	0.55	
19							0.15			0.02							0.30			0.03
20															0.18		0.25			0.05
21			0.35			0.18		0.15		0.07			0.25	0.70	0.40	0.15				0.17
22	2.92	1.80	1.61	3.17	2.25	3.02	3.02	2.06	3.70	2.61	7.35	5.35	2.80	1.70	2.21	3.90	1.05	1.50	2.40	3.14
23						1.80		1.87		0.40										
24				1.25	1.25					0.27			1.25	1.25					1.20	0.41
25						1.25	1.25			0.27			0.43			1.12		1.20	0.31	
26		0.51		0.18		0.40		0.18	0.25	0.16	1.10	1.45	0.90	0.87	0.18	0.60	0.70	0.26	0.67	
27			0.40	1.25	0.80	1.12			0.30	0.43	0.70	1.15	0.50	1.00	0.70	0.43		0.45	0.55	
28							0.43			0.04						0.40	0.60			0.11
29					1.20				0.37	0.17			1.10	1.10	1.10	1.20		0.45	0.55	
30							4.06		1.25	0.59							2.30	1.20		0.39
31													0.56		0.26	0.50				0.15
32														0.35		0.43				0.09
33							0.07		0.18	0.02					0.40					0.04
T	3.21	4.40	4.99	9.74	7.91	12.54	10.04	9.25	9.94	8.00	7.35	9.67	8.06	11.88	11.37	12.65	11.48	12.75	14.27	11.05

T. Total

Table 6. Frequency percentage of plant species in Wadi Water as affected by sites and seasons during the period extended from autumn 2004 up to spring 2006

No.	Dry seasons										Wet seasons									
	1	2	3	4	5	6	7	8	9	M.	1	2	3	4	5	6	7	8	9	M.
1								5		0.56										
2								5	5	1.11			5					5		1.11
3								5		1.11					5		5			1.11
4			5	5		5	5	5		2.78			5	5				5		2.22
5						5		5		1.11					5		5			1.11
6								5	5	1.11				5	5	5	5	5		2.78
7									5	0.56										
8																		5		0.56
9																		5		0.56
10								5	5	1.11				5			5			1.11
11		5			15		5			2.78		15	5		5	5	15	10		6.11
12		5		5				5	5	2.22		5	5	5	5	5	15	5		5.56
13								5		0.56				5	5	5	5	5		2.78
14				5		5		5		1.67					5	5	5	5		2.22
15													5	5						1.11
16						5				0.56							5			0.56
17	20	5	20	25	20			5	5	11.11		5	25	20	15	5		5	5	8.89
18		30	15	15	15	20	15		15	13.89		15		5	5	10	10	5	5	6.11
19							5			0.56								5		0.56
20														5		5				1.11
21			5			5		5		1.67			5	5	5	5				2.22
22	80	40	40	25	35	35	30	30	30	38.33	100	40	30	20	15	30	10	15	15	30.56
23							5		5	1.11										
24				5	5					1.11				5	5				5	1.67
25						5	5			1.11				5			5		5	1.67
26		15		5		5		5	5	3.89		15	5	5	5	5	5	5	5	5.56
27			10	5	5	5			5	3.33		5	15	5	5	5	5		5	5.00
28							5			0.56						5	5			1.11
29					5				5	1.11				5	5	10	5		5	3.33
30							20		5	2.78							10	5		1.67
31													5		5	5				1.67
32														5			5			1.11
33							5		5	1.11				5						0.56
T	100	100	100	95	100	100	100	100	95	98.91	100	100	100	100	100	115	100	100	100	101.67

T. Total



Table 7. Abundance percentage of plant species in Wadi Water as affected by sites and seasons during the period extended from autumn 2004 up to spring 2006

No.	Dry seasons											Wet seasons										
	1	2	3	4	5	6	7	8	9	M.	1	2	3	4	5	6	7	8	9	M.		
1								1.2		0.1												
2								1.8	0.8	2.9			1.3					0.5	0.2			
3			9.5					1.2		1.2					1.5		1.1		0.3			
4			4.8	5.0		5.0	4.8	1.2		2.3			4.2	1.3	2.3			0.5	0.9			
5						5.0		1.2		0.7					1.5		1.1		0.3			
6								58.8	81.3	15.6					23.3	30.8	45.5	44.4	13.6	17.5		
7									4.6	0.5												
8																			13.6	1.5		
9																			22.9	2.5		
10							4.8	1.2		0.7				2.3			1.1		0.4			
11		4.8			13.0		4.8			2.5			14.3		5.1	1.5	0.9	3.3	0.5	2.8		
12		4.8		5.0				1.2	0.8	1.3			4.8	8.3	2.6	2.3	1.5	1.8	5.6	0.5	3.0	
13								16.5		1.8					23.4	30.8	18.2	33.3	41.0	16.3		
14				5.0		5.0		1.2		1.2						1.5	0.9	1.1	0.5	0.4		
15													4.2		4.7					1.0		
16						5.0				0.6								1.1		0.1		
17	20.0	4.8	19.0	25.0	26.1			1.2	0.8	10.8			9.5	20.8	6.4	7.0	1.5		1.1	0.5	5.2	
18		28.6	14.3	15.0	13.0	2.0	14.3		3.34	12.1			14.3		1.3	2.3	3.5	3.6	1.1	0.9	3.0	
19							9.5			1.1								0.9		0.1		
20														2.3			1.9			0.5		
21			4.8			5.0		1.2		1.2				12.5	6.4	11.6	6.2				4.1	
22	80.0	42.7	38.1	30.0	34.8	35.0	28.6	9.7	4.9	33.8			100.0	38.1	25.0	5.1	7.0	9.2	0.9	3.3	1.4	21.0
23						5.0		1.2		0.7												
24				5.0	4.3					1.0					1.3	2.3				1.6	0.6	
25						5.0	4.8			1.1					1.3			0.9		0.5	0.3	
26		14.3		5.0		5.0		1.2	0.8	2.9			14.3	4.2	1.3	2.3	1.5	0.9	1.3	0.5	2.9	
27			9.5	5.0	4.3	5.0			0.8	2.7			4.7	12.5	1.3	2.3	1.5	0.9		0.5	2.6	
28							4.6			0.5						3.00	0.9				0.4	
29					4.5				0.8	0.6					1.3	2.3	3.0	0.9		0.5	0.9	
30							19.0		0.8	2.2								1.8	1.1		0.3	
31															8.3		1.5	0.9				1.2
32																64.0		18.2				9.1
33							4.8		0.8	0.6						2.3						0.3
T	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100

T. Total

Table 8. Fresh and dry foliage yields (ton/fad) of plant species in Wadi Water as affected by sites and seasons during the period extended from autumn 2004 up to spring 2006

Seasons productivity	Dry seasons								
	1	2	3	4	5	6	7	8	9
Fresh yield (ton/fed)	0.161	0.300	0.402	0.548	0.532	0.604	0.604	0.571	0.619
Dry yield (ton/fed)	0.068	0.134	0.188	0.270	0.250	0.270	0.263	0.251	0.311
Seasons productivity	Wet seasons								
	1	2	3	4	5	6	7	8	9
Fresh yield (ton/fed)	0.180	0.380	0.377	0.668	0.681	0.846	0.797	0.770	0.795
Dry yield (ton/fed)	0.101	0.201	0.192	0.297	0.285	0.391	0.403	0.332	0.396

to both edaphic and climatic factors particularly precipitation and wind stress effect. These results confirmed what obtained by Reiad *et al* (1996 a); Hendawy (2002) and Ibrahim *et al* (2006). In general, the greatest abundance of grown plant species were, *Medicago laciniata* in both wet and dry season, *Glebionis coronaria* wet season, *Haloxylon salicornicum* dry and wet seasons, *Paronychia sinaica* wet season.

#### Fresh and dry productivity

Fresh and dry yields of native plants are presented in Table (8). Results showed that sites 6, 7 and 9 produced the highest fresh yields 0.846, 0.797, 0.795 (ton/fad) respectively, during wet season. This indicated that the yields of this three sites was correlated much by availability of soil moisture as well as climatic and edaphic factors which considered as suitable for most plant species in this area.

In respect of the dry foliage yield results in Table (8) showed that the highest dry yields (0.403 and 0.396 ton/fad) were obtained in sites 7, 9 respectively, in wet season and in site 9 in dry season (0.311 ton/fad).

Moreover, the following plant species, i.e., *Artemisia Judica*, *Artemisia monosperma* and *Haloxylon salicornicum* plants produced the highest fresh and dry weights. The obtain results are in agreement with those of Ibrahim (1995) and Reiad *et al* (1996 b & c) in North Western Coast of Egypt; Hendawy (2002) in Gabal El Maghara region Middle Sinai El-Morsy (2002) in NWC of Egypt and Ibrahim *et al* (2006) in North Eastern Coast of Egypt.

#### Chemical composition

Data of chemical composition of some native plant species are presented in Table (9). The results revealed that the all chemical composition traits, i.e., CP, CF, Ash, EE, TDN, Na and K were higher in wet season than that in dry season. The highest chemical composition content of plants in wet seasons compared with that in dry season indicated that chemical composition content of plants were much positively correlated with the available moisture in the soil as well as the surrounding favourable atmosphere for mineral absorption and accumulation in the grown plant species. These results are in harmony with what was reported earlier by Ibrahim (1995) and Reiad *et al* (1996 b & c) noticed that the superiority of surpassed native plants with respect to the chemical composition (CP, CF, EE and total carbohydrates) in Sidi-Barrani in compare with El- Negila site these results were truth in autumn season only in Sidi-Barrani and El- Negila . Ahmed and Nassar (1999) studied the chemical composition of 108 samples from different sites along the North Western Coast of Egypt. They found that the highest average of CP was concerned to *Fabaceae* compared to *Poaceae* and *Chenopodiaceae* in the North Western Coast of Egypt. Average of ash and mineral composition reached to us maximum in *Chenopodiaceae*. El-Toukhy *et al* (2002) studied effect of the variations in climatic on growth parameters of most of the grown plants species during survey of the plant vegetation at Wadi El-Natron - El-Alameen road at five sites. They noticed no clear trend for the chemical constituents for the different studied sites except for Sodium which was higher in some plants for some of the studied sites. Fiber content was higher in *Zilla spinoza*, *Artemisia inculata*, *Pituranthos tortuosus*, *Carduncellus eriocephalus* and *Lycium eurpeum*. and

Table 9. Mean values of chemical content of different plant species at different sites in Wadi Wateer in wet and dry seasons

Site	No.	Dry season							Wet season						
		CP	CF	Ash	Fat	TDN	Na	K	CP	CF	Ash	Fat	TDN	Na	K
2	11	5.21	20.10	6.81	3.40	25.18	0.13	0.20	6.81	20.13	7.12	5.30	30.14	0.73	0.27
	12	6.14	21.80	5.14	2.20	28.30	0.91	0.74	6.13	22.18	8.13	5.20	31.20	0.15	0.34
3	3	12.70	23.11	5.40	6.80	18.19	0.12	0.10	-	-	-	-	-	-	-
	4	13.40	25.18	6.20	5.04	33.14	0.15	0.73	13.18	26.10	5.02	6.71	32.0	0.17	0.56
	12	-	-	-	-	-	-	-	8.14	30.18	2.18	4.10	30.11	0.33	0.34
4	31	-	-	-	-	-	-	-	4.05	30.80	3.17	4.05	14.18	0.20	0.43
	2	-	-	-	-	-	-	-	18.14	29.19	8.19	3.02	27.11	0.81	0.93
	4	12.41	28.02	2.15	3.30	20.99	0.10	0.25	14.13	22.18	3.14	4.35	21.13	0.19	0.33
	11	-	-	-	-	-	-	-	5.80	23.19	6.22	6.01	28.29	0.30	0.32
	12	4.13	30.18	4.19	3.20	23.12	0.31	0.82	6.13	29.90	5.00	5.14	29.00	0.21	0.34
	14	4.15	28.31	6.90	2.80	30.19	0.75	0.84	-	-	-	-	-	-	-
	24	6.18	30.19	5.33	4.31	18.19	0.14	0.39	7.00	29.40	6.20	5.14	28.00	0.33	0.57
	25	-	-	-	-	-	-	-	6.19	25.18	8.9	4.15	20.18	0.74	0.98
	29	-	-	-	-	-	-	-	4.80	29.90	7.14	3.19	28.29	0.21	0.17
	32	-	-	-	-	-	-	-	6.80	19.8	4.14	2.33	29.17	0.33	0.35
5	4	-	-	-	-	-	-	-	9.38	25.80	6.08	6.14	33.70	0.12	0.31
	6	-	-	-	-	-	-	-	10.10	20.91	5.50	5.37	29.10	0.37	0.55
	11	6.01	22.18	5.80	2.14	30.31	0.19	0.40	-	-	-	-	-	-	-
	12	-	-	-	-	-	-	-	7.80	30.14	6.71	3.19	37.0	0.17	0.80
	24	5.44	31.14	3.81	3.11	22.07	0.25	0.45	8.19	30.80	5.19	4.90	22.0	0.47	0.79
29	5.05	28.19	7.55	3.51	29.11	0.74	0.71	6.18	30.19	6.02	5.80	30.30	0.15	0.93	
6	3	-	-	-	-	-	-	-	11.71	25.18	8.14	3.33	30.70	0.18	0.80
	4	10.30	26.10	7.14	3.90	29.80	0.25	0.22	-	-	-	-	-	-	-
	5	8.80	26.81	6.60	4.19	22.14	0.28	0.19	9.11	30.74	7.15	5.37	20.00	0.70	0.33
	6	-	-	-	-	-	-	-	14.4	23.1	5.3	5.7	25.19	0.30	0.35
	11	-	-	-	-	-	-	-	5.3	31.6	7.0	3.1	19.6	0.86	0.23
	12	-	-	-	-	-	-	-	6.9	33.8	8.4	2.9	31.14	0.90	0.59
	14	5.21	28.0	8.9	3.3	25.5	0.74	0.82	6.6	35.4	8.3	3.7	30.4	0.66	0.71
	23	6.6	29.1	7.4	2.2	23.4	0.70	0.11	-	-	-	-	-	-	-
	25	6.5	30.7	6.8	5.4	28.8	0.54	0.64	-	-	-	-	-	-	-
	29	-	-	-	-	-	-	-	6.2	29.1	5.5	2.4	21.7	0.17	0.91
	31	-	-	-	-	-	-	-	8.4	23.4	3.2	3.7	18.8	0.27	0.61
7	4	11.8	20.7	8.4	3.9	22.14	0.11	0.74	-	-	-	-	-	-	-
	6	-	-	-	-	-	-	-	10.5	24.10	7.7	4.5	35.9	0.81	0.52
	11	6.8	29.90	7.7	2.3	26.4	0.71	0.64	6.53	30.20	7.8	4.3	30.18	0.89	0.71
	12	-	-	-	-	-	-	-	8.9	31.1	8.0	4.7	30.7	0.68	0.50
	14	-	-	-	-	-	-	-	7.7	25.9	5.1	2.8	28.8	0.19	0.37
	25	7.4	29.13	5.0	6.6	22.17	0.33	0.78	8.9	30.14	7.9	6.8	30.81	0.84	0.91
	29	-	-	-	-	-	-	-	6.3	29.9	6.8	3.9	22.9	0.54	0.69
	30	8.01	30.13	6.7	3.9	25.9	0.76	0.27	8.4	31.10	6.8	4.00	26.3	0.68	0.81
	31	-	-	-	-	-	-	-	7.4	25.8	5.10	2.2	21.7	0.21	0.37
	32	-	-	-	-	-	-	-	5.9	21.7	3.0	3.05	19.7	0.20	0.81
	8	1	16.8	27.20	8.9	5.7	22.9	0.71	0.77	-	-	-	-	-	-
2		15.1	30.7	7.8	5.9	23.1	0.65	0.89	-	-	-	-	-	-	-
3		12.3	29.1	5.2	3.4	21.9	0.54	0.73	12.1	30.2	6.1	2.9	23.8	0.51	0.11
5		10.3	26.4	6.7	2.1	24	0.71	0.65	9.14	35.1	5.9	4.10	29.9	0.84	0.97
6		13.7	28.1	9.7	5.2	35.7	0.19	0.81	14.0	29.8	7.20	6.0	34.7	0.71	0.52
11		-	-	-	-	-	-	-	7.7	31.7	5.7	4.8	20.3	0.81	0.21
12		8.2	31.3	7.8	3.3	25.0	0.91	0.72	8.3	30.9	6.00	4.5	26.7	0.57	0.55
14		5.5	29.1	6.6	2.9	20.4	0.71	0.52	5.4	27.1	5.7	5.3	23.5	0.81	0.25
23		9.3	30.8	8.7	5.1	29.1	0.29	0.54	-	-	-	-	-	-	-
30		-	-	-	-	-	-	-	7.7	35.8	8.3	5.5	23.3	0.87	0.74

Table 9. Cont.

Site	No.	Dry season							Wet season						
		CP	CF	Ash	Fat	TDN	Na	K	CP	CF	Ash	Fat	TDN	Na	K
9	2	15.35	29.7	9.1	6.4	29.2	0.91	0.11	18.0	37.8	9.8	7.0	24.8	0.84	0.88
	4	-	-	-	-	-	-	-	9.8	29.9	5.4	4.8	29.5	0.94	0.44
	6	10.5	29.7	6.6	5.4	29.5	0.65	0.51	11.0	28.8	7.8	6.2	30.7	0.71	0.72
	7	9.9	24.0	4.8	2.51	25.4	0.21	0.35	-	-	-	-	-	-	-
	8	-	-	-	-	-	-	-	9.8	22.4	6.8	5.1	33.8	0.81	0.77
	9	-	-	-	-	-	-	-	13.4	20.9	7.8	6.2	38.9	0.74	0.58
	11	-	-	-	-	-	-	-	8.5	31.0	8.0	3.2	20.7	0.85	0.52
	12	7.3	29.9	5.20	4.1	19.4	0.22	0.29	8.0	33.4	6.7	4.0	18.5	0.84	0.41
	14	-	-	-	-	-	-	-	6.7	30.8	5.9	2.1	25.4	0.71	0.88
	24	-	-	-	-	-	-	-	9.9	31.9	7.8	5.8	22.8	0.71	0.94
	29	6.3	30.4	6.1	3.2	22.9	0.68	0.85	8.5	33.8	7.0	4.1	20.7	0.51	0.34
	30	10.4	28.8	9.4	6.2	24.8	0.53	0.81	-	-	-	-	-	-	-

ash content was higher for *Anabasis articulata* followed *Arthocnemon glaucum* and *Carduncellus eriocophalus*. However, they reported no clear trend for the rest of chemical contents in wet and dry seasons.

#### REFERENCES

- Abd-Alla, Saida, O.M. (1999). Effect of Afforestation *Acacia saligna* on the Development and Improvement of Some Plant Association at the North Western Coast of Egypt. p. 72, Ph. D.Thesis. Fac. Agric., Ain Shams Univ., Cairo, Egypt.
- Abou-Deya, I.B. and M.O. Salem (1990 a). Seasonal changes in the natural vegetation at El-Mathany area. Proc 4<sup>th</sup> Conf. Agron., Cairo, 11: 627-642.
- Abou- Deya, I.B. and M.O. Salem (1990 b). Seasonal variation in the vegetation structure of protected area at El- Nigeala. Proc 4<sup>th</sup> Conf. Agron., Cairo, 15-16 Spt. (11): 676-691.
- Ahmed, A.M. and Zienab M. Nassar (1999). Chemical composition of some natural range plants from the North Western Coast of Egypt. VI<sup>th</sup> National Conf. for Environmental Studies & Research. Institute of Environmental Studies & Research. Ain Shams Univ. Cairo Egypt. pp. 551 - 568.
- A.O.A.C. (1970). Association of Official Agriculture Chemists. Official Methods of Analysis. 11<sup>th</sup> Ed., p. 382, Washington, D.C.
- A.O.A.C. (1980). Association of Official Analytical Chemists. Official Methods of Analysis. 13<sup>th</sup> Ed., p. 414, Washington, U.S.A.
- Aslan, M.H. (1959). Adaptability and Palatability of Some Forage Plants in the Western Desert of Egypt. pp. 35- 60, M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt.
- Ayyad, M.A. and S.I. Ghabbour (1977). Systems analysis of Mediterranean desert ecosystems of Northern Egypt. samdene. Environmental Conservation L. 2: 91- 102.
- Boulos, L. (2005). Flora of Egypt. (4), pp. 325, Al-Hadara publishing, Cairo, Egypt.
- El-Morsy, M.H. (2002). Studies on Forage Plants in Wadi Magid and Wadi Mehgun in the North Western Coast of Egypt. pp. 42-56. Ph.D. Thesis, Fac. Agric., Cairo Univ. Egypt.
- El-Toukhy, Salwa, A.M.; K.W. Ahmed and S.H. Hendawy (2002). Productivity and nutritive value of some association at Wadi El- Natom - El-Alameen road in the North Western Coast. J. Agric. Sci. Mansoura Univ., 27(1): 233- 244.
- Habbs, T.J.; A.D. Sparrow and J.L. Landsbery (1984). A model of soil moisture balance and herbage growth in arid rangelands of central Australia, J. Aride, Environ. 28: 281- 298.
- Hanson, H.C. and E.D. Churhcill (1965). The Plant Community. pp. 255-280, Reinhold publishing crop. New York.

- Hendawy, S.H. (2002). Diversity of natural vegetation of Gabal- El-Maghara region, Middle Sinai. *Minufiya. J. Agric. Res.* 27(3): 475-486.
- Ibrahim, K.A. (1995). Productivity and Nutritive Value of Some Range Plants of the North Western Coast. pp. 45-84, M.Sc. Thesis, Fac. Agric., Ain Shams Univ., Cairo, Egypt.
- Ibrahim, K.M.; Abd-Alla, Saida. O.M. and Salwa, A.M. El-Toukhy (2006). Potentiality of native range plants under different locations at North Eastern Coast of Egypt. *Annals Agric. Sci., Moshtohor*, 44(1): 1-14.
- Jackson, M.L. (1958). Soil Chemical Analysis Advanced Course. pp. 302-320, Published by the Auther. Dept. Soils, Wisconsin Univ. Madison U.S.A.
- Khafagi, E.A. (1977). Comparative Studies on Evaluating Roughages Including the Markers and Prediction Methods with Reference to Legnin. pp. 36-62, Ph.D. Thesis Fac. Agric. Cairo Univ. Giza , Egypt.
- Peach, K. and M.R. Tracey (1956). *Modern Methods of Plant Analysis*. Springer Verlag, Berlin.
- Piper, C.S. (1950). *Soil and Plant Analysis*, pp. 163-177, Inter. Pub., New York.
- Reiad, M.SH.; M.A. Ashoub; I.B. Abou-Deya; M.S. El-Hakeem and K.M. Ahmed (1996 a). Effect of edaphic and climatic factors on soil and vegetation characters of different associations representative to the North Western Coast of Egypt. *J. Appl. Sic.*; 11 (1): 216- 231
- Reiad, M.SH.; M.A. Ashoub; I.B. Abou-Deya; M.S. El-Hakeem and K.M. Ahmed (1996 b). Environmental effect on yield and quality of five plant associations of the North Western Coast of Egypt. *J. Appl. Sci. Moshtohor*, 34(1): 37- 52.
- Reiad, M.SH.; M.A. Ashoub; I.B. Abou-Deya; M.S. El-Hakeem and K.M. Ahmed (1996 c). Productivity and nutritive value of some range communities at the North Western Coast of Egypt. 1. *Thymelaea hishirsuta* (L.) Endle . *Egypt. J. Appl. Sci.*, 11 (3): 146-163.
- Täckholm. V. (1974). *Student Flora of Egypt*. 2<sup>nd</sup> Ed. Published by Cairo Univ. Cairo, Egypt.



## الانتاجية والقيمة الغذائية للكسء الخضرى الطبيعى بمنطقة وادى وتير بسيناء

[٧]

ناجى حسين محمود بيومى<sup>١</sup> - زينب محمود نصار<sup>١</sup>

١- وحدة المراعى - مركز بحوث الصحراء - المطرية- القاهرة

### الملخص العربى

قسمت النباتات بعد تعريفها إلى مجاميع حسب العائلة، النباتات المستساغة وغير المستساغة ، الحولية والمعمرة. كما تم تحليل عينات التربة فى المواقع المختلفة وكذلك بيانات الأرصاد الجوية متمثلة فى الرطوبة النسبية ، نسبة الأمطار ، درجة الحرارة، سرعة الرياح فى منطقتين نوبيع ممثلة لنهاية الوادى ومنطقة نخل ممثلة لبداية الوادى، وذلك لمعرفة تأثير الظروف الجوية والأرضية على التركيب النباتى بالموقع.

أوضحت النتائج أن معظم القياسات الخضرية للتغطية والوفرة والتكرار والكثافة أو الحاصل الغض والجاف الناتج وكذلك التحليل الكيماوى للنباتات كانت جميعها أعلى فى موسم الربيع عن موسم الخريف . احتوت المنطقة على ثلاثة وثلاثون نوعا نباتيا تتبع خمسة عشر عائلة نباتية والأغلبية كانت للنباتات المعمرة (العادر- الرمث - الشيح- الكبارس) حيث ان لها السيادة نتيجة لتأقلمهم للجفاف. كما أوضحت النتائج ان الموقع ٦، ٧، ٨، ٩ كانت أعلى فى أغلب القياسات الخضرية والمحصولية والتركيب الكيماوى نظراً لكونها فى نهاية مسرى الوادى حيث تستقر مياه الأمطار نتيجة للجريان السطحى فى نهاية الوادى كما كانت المواقع ٨، ٩ أعلى عن غيرها فى أغلب القياسات.

اجرى هذا البحث خلال الموسمين ٢٠٠٤/٢٠٠٥ و٢٠٠٥/٢٠٠٦ خلال موسمى الخريف والربيع لدراسة تأثير المواقع، والمواسم على التركيب النباتى الطبيعى بمنطقة وادى وتير يبدأ وادى وتير عند الكيلو ٢٦٠ من الطريق الدولى وحتى مدينة نوبيع حيث يبلغ طول الوادى حوالى ١٠٠ كم. تم تقسيم الوادى بداية من الطريق الدولى وحتى مدينة نوبيع إلى ٩ مواقع حددت . وكانت كالتالى عند الكيلو ٨ ، ١٩ ، ٢٧ ، ٣٦ ، ٤٨ ، ٥٥ ، ٦٦ ، ٧٨ ، ٩٥ بواقع تقريبا بين كل موقعين ١٠ كم وبدأت الدراسة من خريف ٢٠٠٤ بغرض دراسة التركيب النباتى وخواص الغطاء الخضرى فى المواسم المختلفة وذلك باستخدام طريقة القطع المربعة (٢م × ٢م). حيث تم أخذ ٢٠ قراءة فى كل موقع وقد تم حساب متوسط الكثافة (نبات /م<sup>٢</sup>) ، التغطية % ، التكرار % ، الوفرة % لكل موقع ، الحاصل الفض والجاف (طن / فدان) لكل موقع، التركيب الكيماوى للنباتات المستساغة حيث تم إجراء التحليلات الأتية وحساب نسبة (البروتين- الالياف- الرماد - الزيوت- TDN - نسبة الصوديوم والبيوتاسيوم) .