



IMPACT OF MANAGEMENT SYSTEMS ON SOME SOILS QUALITIES IN EGYPT

[33]

Hoda, A. Elia¹; M.Talha²; M.Y. Afifi¹ and Al-Hassana Abu Gabal²

1- Desert Research center, Mattariya, Cairo, Egypt

2- Faculty of Agriculture, Ain Shams University, Shoubra El-Kheima, Cairo, Egypt

Keywords: Soil quality, Minimum data set (MDS), Principal components analysis, Mean weight diameter (MWD), Soil organic matter, Soil quality index

ABSTRACT

The current work aims to evaluate the effect of different periods of organic farming on some soils qualities compared with the traditional system. Several farms varied in their soil nature and periods of practicing organic farming system were chosen. The farms are located in Belbes, El-Manayef, El-Fayoum and El-Behera areas. Moreover, the study also involved adjacent traditional managed farms. The evaluation was based on the weighted values of the physical, chemical and biological properties of the 0-50 cm and 0-100 cm soil depths. The considered soil properties were soil organic matter (SOM); bulk density; porosity; available water; penetration resistance, mean soil particles weight diameter (MWD), cation exchange capacity and total microbial count. The obtained results indicated that soil biological parameters were less important than the physical or the chemical factors. Principle component₁ (PC₁) scores indicated that 79.97% and 78.05 % of total variance are attributed to the 0-50 cm soil depth of the organic and conventional managed soils, respectively. In the meantime, 75.72 and 71.94 % are related to the 0-100 cm soil depth for the two farming systems, (Organic and conventional) in the same sequence. The only significant factor contributing to PC₂ was total count; PC₂ indicated that (10.29 and 10.87% of the total variance are accounted for 0-50 and 0-100 cm soil depth of organic farming as well as 15.3 and 16.68% of the total variance for 0-50 and 0-100 cm soil depth of conventional farming system, respectively.

INTRODUCTION

It is well known that the inherent soil characteristics are those related to the soil forming factors; i.e., climate, parent material, ... etc, whereas the dynamic soil properties are influenced by the management practices imposed by mankind.

Both are very important to sustainability. Evaluating inherent soil properties and their effect on land use and the suitability of each soil for various uses are the basic concern of soil surveyors. On the other hand, dynamic soil quality focuses on the status of specific soil due to the relatively recent management practices (Karlen *et al* 1997; Doran and Parkin 1994).

Doran and Parkin (1994) defined soil quality as "the capacity of a soil to function". Meanwhile a more complete definition of soil quality was offered by a committee appointed by the Soil Science Society of America (SSSA) as "soil quality is the capacity of a specific kind of soil to function, within natural or managed ecosystem boundaries, to sustain plant and animal productivity, maintain or enhance water and air quality and support human health and habitation".

Doran and Parkin (1994) also stated that soil quality assessment is conducted by evaluating indicators. These indicators can be physical, chemical and biological properties, processes or characteristics of soils.

Jawson (2001) mentioned that physical properties include soil structure, aggregate stability, wind and water erosion. Chemical properties include pH, total plant nutrients, and salinity. Biological properties include root microbial count and other organism-driven processes such as respiration, mineralization, immobilization and denitrification.

(Received June 8, 2009)

(Accepted June 24, 2009)

Since it is impractical to measure every ecosystem or soil property, many researchers proposed a minimum data set which is the smallest set of soil properties or indicators needed to measure or characterize soil quality.

Larson and Pierce (1994), Doran and Parkin (1996) proposed a minimum data set (MDS) consisting of an array of soil chemical, physical, and biological characteristics as basic indicators of soil quality. The soil physical indicators include soil texture, rooting depth, infiltration, water holding capacity and bulk density. Soil chemical indicators embraces electrical conductivity, pH and extractable N, P, K. Soil biological indicators evolve microbial biomass, C and N, potentially mineralizable; N and soil respiration. However, **Seybold et al (1996)** reported that the proposed indicators should be related to soil function and quality.

On the other hand, **Vance (2000) and Doran (2002)** stated that because of its positive influence on several soil processes, crop productivity and environmental quality; therefore soil organic matter (SOM) is often considered the single most important indicator of soil quality and sustainable land management. Moreover, **Van Noordwijk et al (1997)** stated that SOM is a soil property that is generally most sensitive to crop management.

The current work aims at evaluating the impact of organic farming system on soil properties, especially the physical ones that determine soil quality.

MATERIALS AND METHODS

Investigation sites

Due to the fact that organic farming in Egypt still in transition from conditions developed under the conventional system, this study has focused on the beneficial impact of organic farming, if any, on soil properties which enhance soil quality, compared to those obtained under the adjacent conventional farming conditions. The impact of soil type and time – scale on assessing soil quality under organic farming conditions. The chosen farms were selected to represent two soil textural categories, i.e. sandy soils present in Sharkia and Ismailia governorates and the relatively fine soils present in El- Fayoum and El-Behera governorates and to represent different periods from starting organic cultivation. The soil samples depths were 0-50 cm and 0-100 cm for both organic and conventional farming systems were statistically analyzed as well as soil quality index. SQI, was obtained.

Three main steps were followed to assess soil quality index (SQI)

- (i) Select a minimum data set (MDS) of indicators that best represent soil function.
- (ii) Score the MDS indicators based on performance of soil functions.
- (iii) Integrate the indicator scores into a comparative index of soil quality.

A) To select a representative minimum data set (MDS) (**Doran and Parkin 1994**) for the organic and conventional systems, the first performed standardized principal component analysis (PCA) of all untransformed data that showed statistically significant differences between management systems using ANOVA. Principal components (PCs) for a data set are based on their linear combinations of the variables that account for maximum variance within the set by describing vectors of closest fit to the n observations in p -dimensional space, subject to being orthogonal to one another. PCs are assumed receiving high values best represent system attributes. Therefore, the PCs were examined only with eigenvalues ≥ 1 (**Brejda et al 2000**).

For a particular PC, each variable received weight or factor loading that represents its contribution to the PC. Only the highly weighted variables from each PC were retained for the MDS. Highly weighted is defined as that within 10% of the highest factor loading using absolute values. When more than one variable was retained within a PC, linear correlations were calculated to determine whether the variables could be considered redundant and, therefore, eliminated from the MDS (**Andrews 1998**). If the highly weighted variables were not correlated (assumed to have a correlation coefficient < 0.60), then each was considered important and was retained in the MDS. Among well-correlated variables within a PC, the variable with the highest sum of correlation coefficients (absolute values) was chosen for the MDS (**Andrews and Carroll 2001**).

B) Scored each of the MDS variables based on their performance of soil functions using SPSS ver. 10.0 for windows (**SPSS 2002**). The following classical statistic parameters were calculated: minimum, maximum, mean, standard deviation and coefficient of variation (**Webster 1977 and Wilding & Drees 1983**).

Once transformed, the MDS variables for each observation were weighted using the PCA results. Each PC explained a certain amount (%) of the variation in the total data set. This percentage,

divided by the total percentage of variation explained by all PCs with eigenvalues >1 , provided the weighting factor for variables chosen under a given PC.

C) The weighted MDS variable scores were summed for each observation in the following formula: $SQI = \sum Wi \times S$

Where Wi is the PC weighting factor and S is the indicator score. It was compared the calculated SQI organic and conventional means for 0-50 cm and 0-100 cm soil depths. Higher index scores was assumed to mean better soil quality or greater performance of soil functions.

RESULTS AND DISCUSSION

Soil layers physical, chemical, and biological characteristics were determined for the chosen organic and conventional farming sites the obtained data are given in **Tables from (1 to 4)**.

The results in **Tables (1 and 2)** indicate that Galvina and El-Adlia (Belbes area) as well as El-Manayef (Ismailia area) represent varying periods of organic farming. The soils of the chosen farms are generally, non saline except for few cases, and sandy to sandy loam in texture. Organic matter content, except for the uppermost layers, is less than 1%. Thereby, CEC, total aggregates and their MWD is considerably low. Meanwhile, the basic infiltration rate is appreciably high. The data also reveal that the OM content, MWD, Total aggregates, CEC and total microbial count in the organically managed farms, are relatively higher than their corresponding values in the traditionally managed ones.

On the contrary, data obtained for El- Fayoum and El- Behera farms, presented in **Tables (3 and 4)** indicates that their texture are finer than the previous ones. Their textural classes range from sandy loam to clay. Also these soils are non-saline, exceptional for Abu El Matameer site as it is considered slightly saline. Due to the fine texture of such soils, CEC, Total aggregate, and their MWD as well as the retained soil moisture at 0.1 bar and the amount of available water for plant growth are appreciably high for organic farming system especially in the surface soil layer. Therefore, the soil bulk density is low. The data delineate in all chosen sites that the values organic matter content, BD, total aggregates, MWD and total microbial content are considerably higher in the organic managed soils compared to that of the conventional managed ones.

Principal Components Analysis (PCA) for soil properties

To explore the multivariate relationships among soil properties for organic and conventional farming systems, a Principal Components Analysis (PCA) for the physical, chemical, and biological characters was performed using for the whole profiles of the organic system soil variables separately for each soil depth. The same analysis was also performed using all of the weighted attributes for 0-50 cm and 0-100 cm soil depth, under both organic and conventional farming systems.

From the PCA of the standardized values of the determined eleven soil variables for both soil depths under organic and conventional farming; it is found that the characteristics of the first two principal components are significant. In all cases, it reached from 86.6 to 93.4 % of the total variation.

Following univariate analysis, it retained parameters were assessed using PCA. The PCA output for the physical, chemical, and biological characters is given in **Tables (5 & 6)**, Only PCs with eigenvalues >1 explained that at least 5% of the total variance, were retained for interpretation. Ninety five and ninety four percent of the variance, indicated by the first three PCs, are related to 0-50 cm and 0-100 cm soil depth for organic farming system. Also ninety seven and ninety five percent of the variance are attributed to 0-50 cm and 0-100 cm soil depth for conventional farming system respectively.

The PC1 explained that 79.97 and 78.05% are related to the total variance of 0-50 cm organic and conventional farming systems, and 75.72 and 71.94 % are attributed to 0-100 cm soil depth for organic and conventional farming systems, respectively. There were nine variables that had significant response on PC1, eight scores that were > 0.98 , positive weighted (OM %, porosity, moist at 0.1 bar, available water, penetration resistance, MWD, silt+ clay and CEC) and two negatively weighted (Bulk density, basic infiltration rate). These finding align with those for **Harris et al (1996)** who stated that soil quality decreased with increasing bulk density and increased with total stable aggregates (mean weight diameter). It is diminished by increasing the time required for water to infiltrate.

The PC1 had significant positive response on soil penetration resistance variable. The strong dependence of penetration resistance suggests that soils with higher PC1 scores were more consolidated. **Hussain et al (1999)** found penetration

Table1. Soil physical, chemical, and biological characteristic for organic farming and conventional cultivation site, Belbes area

| Pro. No. | Farming type & Age | Dep. cm | T.S ¹ | Silt % | Clay % | Tex ² | O.M.% | B.D ³ | Porosity ⁴ | 0.1 bar ⁵ | Avai. W ⁶ | T.Agg ⁷ | MWD ⁸ | P.R Mpa ⁹ | CEC ¹⁰ | Ec dS/m | Total bact. count | I _{bas} ¹¹ |
|---|--------------------------|---------|------------------|--------|--------|------------------|-------|------------------|-----------------------|----------------------|----------------------|--------------------|------------------|----------------------|-------------------|---------|---------------------|--------------------------------|
| Organic farming (Galvina area) | | | | | | | | | | | | | | | | | | |
| 1 | Annual crops 25 year | 0: 25 | 76.06 | 3.37 | 20.57 | S.C.L. | 2.67 | 1.44 | 44.40 | 33.17 | 17.50 | 28.04 | 0.49 | 1.50 | 16.30 | 2.40 | 24*10 ⁵ | 11.03 |
| | | 25:45 | 80.38 | 2.74 | 16.88 | S.L. | 0.73 | 1.53 | 42.06 | 24.86 | 11.05 | 24.12 | 0.35 | 2.11 | 7.88 | 2.48 | 12*10 ⁵ | |
| | | 45:68 | 86.01 | 0.20 | 13.79 | L.S. | 0.40 | 1.59 | 39.77 | 20.56 | 9.51 | | | 2.61 | 7.34 | 2.03 | | |
| | | 68:120 | 85.8 | 1.46 | 12.74 | L.S. | 0.14 | 1.63 | 38.26 | 25.69 | 11.16 | | | | 7.17 | 1.78 | | |
| 2 | orchards 25 year | 120:150 | 56.83 | 6.42 | 36.75 | S.C. | 0.16 | 1.32 | 50.19 | 42.18 | 18.09 | | | | 33.70 | 2.03 | | 7.48 |
| | | 0: 15 | 71.65 | 9.87 | 18.48 | S.L. | 2.51 | 1.41 | 45.98 | 27.34 | 14.79 | 27.65 | 0.49 | 1.999 | 16.74 | 8.85 | 17*10 ⁵ | |
| | | 15:50 | 78.32 | 8.95 | 12.73 | S.L. | 0.61 | 1.56 | 40.68 | 25.24 | 12.30 | 27.36 | 0.43 | 2.24 | 8.70 | 2.81 | 15*10 ⁶ | |
| | | 50:110 | 85.32 | 4.22 | 10.46 | L.S. | 0.95 | 1.59 | 39.77 | 17.44 | 10.17 | | | 3.099 | 8.42 | 3.79 | | |
| | | 110:135 | 88.11 | 3.25 | 8.64 | S:L.S | 0.94 | 1.59 | 39.77 | 16.82 | 9.32 | | | | 16.30 | 2.41 | | |
| | | 135:160 | 45.57 | 19.47 | 34.96 | S.C.L | 0.98 | 1.38 | 48.31 | 36.58 | 15.88 | | | | 35.65 | 2.33 | | |
| Conventional cultivation (Galvina area) | | | | | | | | | | | | | | | | | | |
| 3 | Annual crops 25 year | 0: 20 | 78.37 | 8.46 | 13.18 | S.L | 2.04 | 1.49 | 43.35 | 24.66 | 14.19 | 23.08 | 0.33 | 2.44 | 8.26 | 3.82 | 98*10 ⁵ | 3.91 |
| | | 20:70 | 81.2 | 4.36 | 14.44 | L.S. | 0.60 | 1.58 | 39.69 | 18.07 | 10.08 | 14.16 | 0.26 | 2.996 | 9.89 | 3.13 | 15*10 ⁵ | |
| | | 70:135 | 78.2 | 6.59 | 15.21 | S.L. | 0.29 | 1.62 | 38.40 | 22.70 | 11.29 | | | 2.55 | 7.17 | 5.79 | | |
| | | 135:160 | 63.38 | 15.61 | 21.00 | S.C.L. | 0.52 | 1.46 | 44.70 | 28.78 | 16.45 | | | | 10.87 | 6.23 | | |
| Organic farming (El-Adliya area) | | | | | | | | | | | | | | | | | | |
| 4 | Annual crops 7 year | 0: 15 | 71.75 | 11.15 | 17.10 | S.L. | 2.04 | 1.49 | 43.35 | 31.88 | 14.59 | 25.47 | 0.34 | 2.25 | 11.30 | 2.50 | 133*10 ⁵ | 6.76 |
| | | 15: 42 | 73.6 | 11.69 | 14.71 | S.L. | 0.41 | 1.52 | 41.31 | 22.45 | 10.07 | 19.11 | 0.30 | 3.11 | 2.61 | 4.00 | 149*10 ⁴ | |
| | | 42: 80 | 74.14 | 11.89 | 13.98 | S.L. | 0.13 | 1.52 | 42.64 | 26.10 | 12.19 | | | 2.71 | 3.48 | 3.22 | | |
| 5 | orchards 7 years | 80:130 | 87.48 | 4.76 | 7.76 | S:L.S | 0.06 | 1.65 | 36.54 | 16.89 | 7.81 | | | | 2.61 | 3.08 | | 6.898 |
| | | 0: 15 | 73.4 | 10.59 | 16.01 | S.L | 1.22 | 1.45 | 44.23 | 23.52 | 12.99 | 22.59 | 0.32 | 2.42 | 9.78 | 4.65 | 113*10 ⁵ | |
| | | 15: 60 | 73.65 | 10.47 | 15.89 | S.L. | 0.23 | 1.5 | 42.08 | 22.94 | 12.97 | 20.73 | 0.26 | 2.56 | 5.22 | 1.21 | 84*10 ⁴ | |
| 6 | Annual crops one year | 60:115 | 69.4 | 13.42 | 17.17 | S.L. | 0.17 | 1.52 | 41.98 | 17.61 | 9.98 | | | 2.996 | 5.22 | 3.08 | | |
| | | 115:140 | 75.53 | 10.40 | 14.07 | S.L | 0.06 | 1.65 | 37.02 | 21.16 | 13.34 | | | | 4.35 | 2.91 | | |
| | | 0: 10 | 76.45 | 10.88 | 12.68 | S.L | 1.04 | 1.58 | 40.15 | 22.78 | 12.50 | 22.30 | 0.22 | 2.62 | 8.26 | 9.42 | 100*10 ⁵ | |
| | | 10: 47 | 79.71 | 8.30 | 11.99 | S.L | 0.39 | 1.61 | 39.25 | 16.55 | 8.92 | 13.31 | 0.14 | 3.10 | 5.87 | 2.73 | 64*10 ⁴ | |
| 6 | Annual crops one year | 47:65 | 74.4 | 7.49 | 18.11 | S.L | 0.09 | 1.61 | 39.02 | 22.35 | 11.38 | | | 2.59 | 5.87 | 2.56 | | |
| | | 65:90 | 74.9 | 7.07 | 18.03 | S.L | 0.09 | 1.65 | 37.50 | 20.03 | 10.43 | | | 2.84 | 5.22 | 3.37 | | |
| | | 90:135 | 74.00 | 6.67 | 19.33 | S.L | 0.19 | 1.54 | 41.89 | 21.98 | 10.24 | | | | 9.13 | 3.09 | | |

1= total sand

2 = texture

3= bulk density (Mg/m³)

4 = porosity %

5= moisture content at 0.1 bar

6= available water

7 = total aggregate

8 = mean weight diameter

9= penetration resistance

10 = CEC meq/100 gm soil

11 = Basic infiltration rate (cm/hr.)

Table 2. Soil physical, chemical, and biological characteristic for organic farming and conventional cultivation site, El-Manayef area

| Pro. No. | Farming type & Age | Depth cm | T.S % ¹ | Silt% | Clay% | Tex. ² | O.M.% | B.D. ³ Mg/m ³ | Porosity % ⁴ | 0.1 bar ⁵ | Avai. W. ⁶ | T.Agg ⁷ | MWD ⁸ | P.R Mpa ⁹ | CEC ¹⁰ | Ec dS/m | Total bact count | I _{bas} ¹¹ |
|---------------------------------|-------------------------|----------|--------------------|-------|-------|-------------------|-------|-------------------------------------|-------------------------|----------------------|-----------------------|--------------------|------------------|----------------------|-------------------|---------|---------------------|--------------------------------|
| Organic farming | | | | | | | | | | | | | | | | | | |
| 7 | Annual crops 12 year | 0:15 | 82.88 | 3.76 | 13.36 | S.L. | 2.82 | 1.49 | 40.87 | 26.23 | 14.18 | 36.72 | 0.29 | 1.37 | 9.78 | 7.66 | 153*10 ⁵ | 8.44 |
| | | 15:30 | 79.45 | 4.66 | 15.89 | S.L. | 3.03 | 1.45 | 42.69 | 25.68 | 11.68 | 21.41 | 0.22 | 1.09 | 11.30 | 2.99 | 104*10 ⁴ | |
| | | 30:67 | 91.19 | 4.34 | 4.47 | S | 0.57 | 1.76 | 33.08 | 15.93 | 9.70 | 21.41 | 0.22 | 2.54 | 6.30 | 2.15 | | |
| | | 67:130 | 88.4 | 2.91 | 8.69 | L.S. | 1.52 | 1.65 | 36.54 | 18.28 | 10.20 | | | 2.11 | 8.26 | 4.88 | | |
| 8 | orchards 12 year | 0:15 | 79.55 | 4.29 | 16.16 | S.L. | 3.09 | 1.45 | 42.46 | 25.23 | 12.77 | 33.63 | 0.28 | 1.29 | 9.78 | 0.99 | 220*10 ⁵ | 7.14 |
| | | 15:35 | 88.79 | 1.56 | 9.65 | L.S | 1.47 | 1.68 | 35.88 | 22.05 | 11.13 | 23.08 | 0.25 | 1.84 | 5.87 | 0.69 | 90*10 ⁴ | |
| | | 35:60 | 84.98 | 2.67 | 12.35 | L.S. | 0.55 | 1.64 | 37.40 | 22.72 | 10.28 | 23.08 | 0.25 | 1.62 | 8.69 | 0.71 | | |
| | | 60:120 | 87.62 | 2.43 | 9.95 | L.S. | 0.53 | 1.74 | 33.33 | 17.27 | 8.49 | | | 2.24 | 7.83 | 0.60 | | |
| Conventional cultivation | | | | | | | | | | | | | | | | | | |
| 9 | Annual crops | 0:15 | 88.52 | 2.47 | 9.01 | L.S | 1.79 | 1.59 | 38.37 | 22.14 | 11.30 | 30.97 | 0.25 | 1.75 | 7.98 | 3.84 | 99*10 ⁵ | 15.89 |
| | | 15:38 | 95.96 | 0.52 | 3.52 | S | 0.52 | 1.74 | 33.59 | 19.49 | 8.18 | 21.21 | 0.23 | 1.93 | 2.17 | 0.85 | 80*10 ⁴ | |
| | | 38:70 | 82.44 | 3.44 | 14.12 | S.L | 0.92 | 1.7 | 34.62 | 20.15 | 10.48 | 21.21 | 0.23 | 2.04 | 6.30 | 2.65 | | |
| | | 70:85 | 92.32 | 1.28 | 6.40 | S | 1.33 | 1.68 | 35.63 | 15.29 | 8.07 | | | 2.04 | 5.22 | 2.68 | | |
| | | 85:135 | 95.47 | 0.40 | 4.13 | S | 0.61 | 1.79 | 32.45 | 15.12 | 8.38 | | | | 2.61 | 2.87 | | |
| 10 | Orchards | 0:15 | 84.58 | 4.71 | 10.71 | L.S | 1.06 | 1.59 | 40.00 | 21.14 | 11.02 | 30.62 | 0.24 | 1.86 | 3.91 | 0.79 | 142*10 ⁵ | 10.51 |
| | | 15:35 | 94.96 | 0.35 | 4.69 | S | 0.59 | 1.66 | 37.12 | 17.38 | 9.21 | 17.74 | 0.20 | 2.24 | 3.04 | 0.62 | 31*10 ⁴ | |
| | | 35:50 | 88.92 | 2.14 | 8.94 | L.S | 1.08 | 1.66 | 36.64 | 16.10 | 9.21 | 17.74 | 0.20 | 2.4 | 6.30 | 1.24 | | |
| | | 50:64 | 88.72 | 2.56 | 8.72 | L.S | 1.56 | 1.67 | 36.02 | 17.40 | 9.47 | | | 2.28 | 6.74 | 0.74 | | |
| | | 64:84 | 90.41 | 0.99 | 8.60 | S | 1.08 | 1.71 | 34.23 | 17.97 | 8.25 | | | 2.10 | 5.87 | 1.24 | | |
| | | 84:135 | 85.78 | 3.98 | 10.24 | L.S | 0.76 | 1.75 | 32.95 | 19.68 | 10.64 | | | | 5.22 | 0.82 | | |

1 = total sand

2 = texture

3 = bulk density (Mg/m³)

4 = porosity %

5 = moisture content at 0.1 bar

6 = available water

7 = total aggregate

8 = mean weight diameter

9 = penetration resistance

10 = CEC meq/100 gm soil

11 = Basic infiltration rate (cm/hr.)

Table 3. Soil physical, chemical, and biological characteristic for organic farming and conventional cultivation site, El-Fayoum area

| Pro. No. | Farming Type& age | Depth cm | T. S. ¹ % | Silt% | Clay% | Text ² | O.M.% | B.D ³ Mg/m ³ | Poro % ⁴ | 0.1 bar ⁵ | Aval W. ⁶ | T.Ag ⁷ | MWD ⁸ | P.R Mpa ⁹ | CEC ¹⁰ | Ec dS/m | Total bact. count | I _{bas} ¹¹ |
|---------------------------------|-----------------------|----------|----------------------|-------|-------|-------------------|-------|------------------------------------|---------------------|----------------------|----------------------|-------------------|------------------|----------------------|-------------------|---------|---------------------|--------------------------------|
| Organic farming | | | | | | | | | | | | | | | | | | |
| 11 | Annual crops (15year) | 0 : 30 | 46.23 | 19.09 | 34.68 | S.C.L. | 3.12 | 1.17 | 53.75 | 41.11 | 20.08 | 51.03 | 0.59 | 2.43 | 49.56 | 0.89 | 176*10 ⁵ | 7.28 |
| | | 30 : 60 | 61.13 | 14.51 | 24.36 | S.C.L. | 1.30 | 1.36 | 47.29 | 38.61 | 18.01 | 31.31 | 0.34 | 2.94 | 35.87 | 1.73 | 156*10 ⁵ | |
| | | 60 : 140 | 80.19 | 7.51 | 12.30 | L.S. | 0.30 | 1.55 | 39.69 | 39.55 | 21.07 | | | | 20.43 | 1.63 | | |
| 12 | orchards (15 year) | 0 : 21 | 47.01 | 20.32 | 32.67 | S.C.L. | 4.58 | 1.19 | 52.59 | 41.44 | 20.51 | 54.33 | 0.63 | 2.45 | 44.35 | 1.15 | 197*10 ⁵ | 3.03 |
| | | 21 : 50 | 50.68 | 18.92 | 30.40 | S.C.L. | 2.44 | 1.23 | 50.85 | 38.24 | 16.08 | 32.75 | 0.48 | 2.79 | 34.78 | 3.24 | 175*10 ⁵ | |
| | | 50 : 70 | 62.92 | 17.37 | 19.71 | S.L. | 0.65 | 1.48 | 42.41 | 39.68 | 20.82 | 30.13 | 0.19 | 3.11 | 29.56 | 6.30 | | |
| | | 70 : 140 | 66.41 | 15.31 | 18.28 | S.L. | 0.50 | 1.52 | 39.68 | 35.87 | 16.01 | | | | 23.50 | 6.53 | | |
| Conventional cultivation | | | | | | | | | | | | | | | | | | |
| 13 | Annual crops | 0 : 25 | 55.85 | 14.74 | 29.41 | S.C.L. | 2.56 | 1.28 | 49.61 | 40.92 | 20.29 | 42.6 | 0.51 | 2.71 | 31.52 | 1.69 | 158*10 ⁵ | 2.16 |
| | | 25 : 50 | 76.22 | 10.13 | 13.65 | S.L. | 0.59 | 1.47 | 41.90 | 41.06 | 18.40 | 14.11 | 0.27 | 2.98 | 21.74 | 1.81 | 91*10 ⁴ | |
| | | 50 : 57 | 59.36 | 15.86 | 24.78 | S.C.L. | 0.65 | 1.26 | 48.78 | 40.57 | 17.70 | 19.52 | 0.24 | 2.68 | 32.60 | 1.77 | | |
| | | 57 : 80 | 79.28 | 8.24 | 12.48 | S.L. | 0.53 | 1.28 | 50.58 | 30.17 | 15.12 | | | 3.78 | 19.57 | 1.68 | | |
| | | 80 : 95 | 48.81 | 21.45 | 29.74 | S.C.L. | 1.02 | 1.25 | 50.00 | 38.12 | 16.06 | | | | 34.78 | 1.19 | | |
| | 95 : 150 | 77.8 | 8.19 | 14.01 | S.L. | 0.59 | 1.54 | 40.54 | 39.97 | 23.34 | | | | 20.65 | 2.38 | | | |
| 14 | orchards | 0 : 35 | 55.2 | 18.86 | 25.94 | S.C.L. | 2.87 | 1.24 | 50.40 | 42.09 | 19.29 | 40.72 | 0.43 | 2.81 | 38.26 | 2.08 | 162*10 ⁵ | 1.495 |
| | | 35 : 67 | 56.15 | 14.77 | 29.08 | S.C.L. | 1.02 | 1.20 | 51.61 | 33.99 | 16.25 | 19.35 | 0.21 | 2.96 | 28.26 | 1.45 | 110*10 ⁵ | |
| | | 67 : 90 | 72.03 | 8.47 | 19.50 | S.L. | 0.45 | 1.24 | 52.49 | 32.21 | 15.73 | | | 3.62 | 23.69 | 4.51 | | |
| | | 90 : 140 | 80.12 | 8.00 | 11.88 | S.L. | 0.33 | 1.60 | 39.62 | 37.18 | 19.02 | | | | 20.00 | 6.29 | | |

1 = total sand
7 = total aggregate2 = texture
8 = mean weight diameter3 = bulk density (Mg/m³)
9 = penetration resistance4 = porosity %
10 = CEC meq/100 gm soil5 = moisture content at 0.1 bar
11 = Basic infiltration rate (cm/hr.)

6 = available water

Table 4. Soil physical, chemical, and biological characteristic of an organic farming and conventional cultivation profiles of El-Behera area

| Pro. No. | Farming Type & age | Depth cm | T. S.% ¹ | Silt % | Clay % | Tex. ² | O.M% ³ | B.D. ³ Mg/m ³ | Por.% ⁴ | 0.1 bar ⁵ | Ava. W ⁶ | T.Aggr ⁷ | MWD ⁸ | P.R Mpa ⁹ | CEC ¹⁰ | Ec dS/m | Total bact. count | I _{bas} ¹¹ |
|--|-------------------------|----------|---------------------|--------|--------|-------------------|-------------------|-------------------------------------|--------------------|----------------------|---------------------|---------------------|------------------|----------------------|-------------------|---------|----------------------|--------------------------------|
| Organic farm (Abo El-Matameer area) | | | | | | | | | | | | | | | | | | |
| 15 | Annual crops (10 years) | 0:30 | 26.67 | 27.22 | 46.11 | C. | 3.89 | 1.11 | 55.78 | 51.38 | 23.38 | 55.86 | 0.76 | 2.49 | 45.28 | 3.15 | 170*10 ⁶ | 2.757 |
| | | 30:50 | 21.1 | 29.18 | 49.72 | C. | 1.48 | 1.14 | 52.72 | 50.12 | 24.18 | 57.62 | 0.64 | 2.70 | 55.22 | 4.29 | 212*10 ⁴ | |
| | | 50:68 | 42.72 | 35.00 | 22.28 | L. | 0.90 | 1.08 | 55.93 | 54.85 | 24.24 | 54.02 | 0.54 | 2.59 | 32.39 | 4.50 | 104*10 ⁴ | |
| | | 68:105 | 30.57 | 36.12 | 33.31 | C.L. | 0.90 | 1.12 | 53.63 | 53.10 | 23.68 | | | 2.53 | 35.65 | 4.46 | | |
| | | 105:135 | 42.41 | 27.61 | 29.98 | C.L. | 0.84 | 1.24 | 50.88 | 48.16 | 21.62 | | | | 14.78 | 4.83 | | |
| Conventional cultivation(Abo El-Matameer area) | | | | | | | | | | | | | | | | | | |
| 16 | Annual crops | 0:10 | 29.16 | 17.86 | 52.98 | C. | 2.64 | 1.14 | 54.31 | 52.02 | 22.53 | 50.73 | 0.55 | 2.62 | 44.16 | 6.97 | 128*10 ⁵ | 1.667 |
| | | 10:25 | 33.44 | 25.24 | 41.32 | C. | 1.48 | 1.16 | 52.24 | 49.13 | 24.84 | 47.78 | 0.57 | 2.94 | 31.52 | 8.22 | 180*10 ⁴ | |
| | | 25:65 | 26.44 | 27.97 | 45.59 | C. | 0.90 | 1.11 | 54.25 | 51.40 | 25.14 | 60.06 | 0.50 | 2.59 | 44.57 | 9.47 | | |
| | | 65:95 | 24.71 | 27.05 | 48.24 | C. | 0.58 | 1.09 | 56.09 | 54.46 | 22.52 | | | 2.29 | 46.09 | 8.81 | | |
| | | 95:135 | 52.33 | 29.96 | 17.71 | L:S:L | 0.81 | 1.24 | 50.93 | 47.02 | 21.12 | | | | 14.13 | 7.84 | | |
| Organic farm (Etai El- Barood area) | | | | | | | | | | | | | | | | | | |
| 17 | Annual crops (10 years) | 0:30 | 35.86 | 24.44 | 39.70 | C.L | 3.92 | 1.13 | 54.78 | 50.04 | 21.81 | 50.77 | 0.71 | 2.47 | 49.26 | 0.90 | 182.*10 ⁵ | 5.888 |
| | | 30:60 | 43.3 | 26.40 | 30.30 | C.L. | 0.99 | 1.16 | 52.40 | 49.78 | 19.25 | 60.08 | 0.55 | 2.57 | 67.39 | 1.05 | 196*10 ⁴ | |
| | | 60:80 | 22.36 | 27.16 | 50.48 | C. | 0.96 | 1.12 | 52.54 | 44.45 | 20.48 | | | 2.88 | 67.39 | 0.71 | | |
| | | 80::135 | 46.99 | 28.01 | 25.00 | L:SCL | 0.49 | 1.24 | 50.25 | 44.65 | 20.50 | | | | 75.00 | 0.76 | | |
| | | | | | | | | | | | | | | | | | | |
| Conventional cultivation (Etai El- Barood area) | | | | | | | | | | | | | | | | | | |
| 18 | Annual crops | 0:35 | 37.53 | 20.03 | 42.44 | C | 2.32 | 1.20 | 50.95 | 48.91 | 19.19 | 46.21 | 0.62 | 2.79 | 44.57 | 0.58 | 139*10 ⁵ | 1.994 |
| | | 35:55 | 61.05 | 13.30 | 25.65 | S.C.L | 0.96 | 1.27 | 50.49 | 47.81 | 17.21 | 33.77 | 0.43 | 2.96 | 48.91 | 0.66 | 141*10 ⁴ | |
| | | 55:85 | 67.83 | 12.97 | 19.20 | S.L. | 0.32 | 1.48 | 42.80 | 40.90 | 15.07 | | | 3.67 | 19.57 | 0.93 | | |
| | | 85:135 | 83.58 | 5.41 | 11.01 | L.S | 0.17 | 1.61 | 38.74 | 30.94 | 17.30 | | | | 10.87 | 0.80 | | |
| | | | | | | | | | | | | | | | | | | |

1= total sand 2 = texture 3 = bulk density (Mg/m³) 4 = porosity % 5 = moisture content at 0.1 bar 6= available water
 7 = total aggregate 8 = mean weight diameter 9 = penetration resistance 10 = CEC meq/100 gm soil 11 = Basic infiltration rate (cm/hr.)

Table 5. The Principal Components (PC) from the principal component analysis (PCA) of the standardized values of eleven soil variables and their weights based on eleven simultaneous correlated variables from all minimum data set of organic farming system (Only principal components with eigenvalues >1)

| Measurements | Organic farming system, 0-50 cm | | | Organic farming system, 0-100 cm | | |
|---|------------------------------------|--------|--------|-------------------------------------|--------|--------|
| | Principal component | | | | | |
| | PC 1 | PC2 | PC3 | PC 1 | PC2 | PC3 |
| Eigenvalue | 8.797 | 1.132 | 0.562 | 8.33 | 1.195 | 0.800 |
| Proportion of variance | 79.97 | 10.289 | 5.112 | 75.724 | 10.865 | 7.273 |
| Cumulative proportion | 79.97 | 90.259 | 95.371 | 75.724 | 86.590 | 93.863 |
| Rotated scores of two retained eigenvectors | | | | | | |
| OM% | .897 | | -.924 | .669 | -.131 | -.948 |
| Bulk Density (Mg/m ³) | -.990 | | .789 | -.991 | | .711 |
| Porosity % | .994 | | -.754 | .997 | | -.678 |
| Moisture content at 0.1bar | .989 | -.138 | -.734 | .982 | | -.728 |
| Available water | .986 | -.112 | -.724 | .965 | | -.746 |
| Penetration resistance (Mpa) | .899 | .214 | -.702 | .674 | .588 | -.608 |
| MWD | .950 | .154 | -.641 | .948 | | -.484 |
| Infiltration rate (cm/hr.) | -.683 | .218 | .956 | -.640 | | .946 |
| CEC (meq/100 gm soil) | .974 | | -.751 | .931 | | -.683 |
| Total bacterial count | -.114 | .992 | .209 | -0.355 | .889 | .392 |
| Silt +clay% | .983 | | -.804 | .976 | -.146 | -.767 |

PC1= the first principal component PC2 = the second principal component PC3 = the third principal component

Table 6. The Principal Components (PC) from the PCA of the standardized values of eleven soil variables and their weights based on eleven simultaneous correlated variables from all minimum data set of conventional farming system (Only principal components with eigenvalues >1)

| Measurements | Conventional farming system 0-50 cm | | | Conventional farming system 0-100 cm | | |
|---|--|--------|--------|---|--------|--------|
| | Principal component | | | | | |
| | PC 1 | PC2 | PC3 | PC 1 | PC2 | PC3 |
| Eigenvalue | 8.585 | 1.683 | 0.419 | 7.941 | 1.835 | 0.72 |
| Proportion of variance | 78.049 | 15.301 | 3.806 | 71.941 | 16.678 | 6.546 |
| Cumulative proportion | 78.049 | 93.35 | 97.156 | 71.941 | 88.619 | 95.165 |
| Rotated scores of two retained eigenvectors | | | | | | |
| OM% | .726 | .820 | -.140 | .426 | .322 | .991 |
| Bulk Density (Mg/m ³) | -.986 | -.354 | .281 | -.961 | -.308 | -.527 |
| Porosity % | .983 | .367 | -.318 | .959 | .336 | .535 |
| Moisture content at 0.1 bar | .983 | .254 | -.142 | .991 | .154 | .423 |
| Available water | .959 | .203 | -.274 | .983 | .165 | .431 |
| Penetration resistance (Mpa) | .706 | .662 | -.758 | .568 | .847 | .475 |
| MWD | .947 | | -.174 | .892 | | .136 |
| Infiltration rate (cm/hr.) | -.817 | -.540 | .700 | -.805 | -.668 | -.371 |
| CEC (meq/100 gm soil) | .981 | .368 | -.149 | .987 | .226 | .499 |
| Total bacterial count | | .944 | -.455 | | .937 | .267 |
| Silt +clay% | .986 | .166 | -.214 | .978 | | .306 |

PC1= the first principal component PC2 = the second principal component PC3 = the third principal component

Table 7. Formula of calculated soil quality index (SQI) using weighting factors for each scored of minimum data set (MDS) variable

| Soil depth | Farming system | Equation |
|------------|------------------|---|
| 0-50 cm | Organic cultiva. | $SQI = \sum [0.89(S_{OM\%} - S_{BD} + S_{poros.\%} + S_{Pene.Resis.} + S_{MWD} + S_{CEC} + S_{silt+clay} - S_{IR}) + (0.11 \times S_{Total\ count})]$ |
| 0-100 cm | Organic cultiva. | $SQI = \sum [0.81(S_{OM\%} - S_{BD} + S_{poros.\%} + S_{Pene.Resis.} + S_{MWD} + S_{CEC} + S_{silt+clay} - S_{IR}) + (0.116 \times S_{Total\ count})]$ |
| 0-50 cm | Conven. Cultiva | $SQI = \sum [0.836(S_{OM\%} - S_{BD} + S_{poros.\%} + S_{Pene.Resis.} + S_{MWD} + S_{CEC} + S_{silt+clay} - S_{IR}) + (0.16 \times S_{Total\ count})]$ |
| 0-100 cm | Conven. Cultiva | $SQI = \sum [0.76(S_{OM\%} - S_{BD} + S_{poros.\%} + S_{Pene.Resis.} + S_{MWD} + S_{CEC} + S_{silt+clay} - S_{IR}) + (0.175 \times S_{Total\ count})]$ |
| 0-50 cm | Total areas | $SQI = \sum [0.833(S_{OM\%} - S_{BD} + S_{poros.\%} + S_{Pene.Resis.} + S_{MWD} + S_{CEC} + S_{silt+clay} - S_{IR}) + (0.166 \times S_{Total\ count})]$ |
| 0-100 cm | Total areas | $SQI = \sum [0.75(S_{OM\%} - S_{BD} + S_{poros.\%} + S_{CEC} + S_{MWD} + S_{silt+clay} - S_{IR}) + (0.15 \times S_{Total\ count}) + (0.09 \times S_{Pene.Resis.})]$ |

S is the score for the subscript variable and the coefficients is the weighting factors derived from the PCA

Table 8. Soil quality and their weights and classes for evaluating studied areas soil qualities

| Indicators | Weight (organic) | | Weight (conven.) | | Scores (S) | | | | | The weights *the marks of the indicator (W _i x S) | | | |
|-----------------------------|-------------------|---------|------------------|---------|--------------|-------------|-------------|---------|-------|--|----------|---------|----------|
| | 0-50cm | 0-100cm | 0-50 cm | 0-100cm | I | II | III | IV | V | 0-50 cm | 0-100 cm | 0-50 cm | 0-100 cm |
| OM% | 2.24 | 1.47 | 1.47 | 1.08 | >1.5 | 1.5-1 | 1-0.5 | .5-0.25 | <0.25 | 11.20 | 5.90 | 5.88 | 4.33 |
| B.D. (Mg/m ³) | 1.37 | 1.43 | 1.40 | 1.43 | <1.3 | 1.3-1.8 | 1.8-2.1 | >2.1 | | 4.10 | 4.28 | 4.21 | 4.29 |
| Porosity % | 46.63 | 44.32 | 45.04 | 44.21 | 50 | 50-60;50-40 | 60-80;40-20 | | | 93.26 | 88.64 | 90.07 | 88.41 |
| A.W | 16.48 | 15.87 | 16.00 | 15.35 | >18 | 18-14 | 14-10 | 10-6 | <6 | 65.93 | 63.50 | 64.00 | 61.40 |
| T. Aggre | 36.71 | 35.66 | 32.49 | 29.43 | >60 | 60-45 | 45-30 | <30 | | 73.41 | 71.32 | 64.99 | 29.43 |
| CEC meq/100g soil | 28.57 | 25.88 | 23.82 | 21.34 | >30 | 30-15 | 15-10 | 10-5 | <5 | 114.27 | 103.53 | 95.29 | 85.34 |
| Silt+Clay % | 38.78 | 34.37 | 34.89 | 32.27 | <35 | 35-50 | 50-60 | 60-80 | >80 | 155.12 | 137.47 | 139.57 | 129.07 |
| Soil quality index | | | | | | | | | | 517.27 | 474.64 | 464.01 | 402.27 |
| Relative soil quality index | | | | | | | | | | 73.90 | 67.81 | 66.29 | 57.47 |

resistance is negatively correlated with the rooting relation function of soil quality. At the same time, soils with high PC1 scores had more surface residue and so may be less susceptible to erosion. Soils in the 0-50 cm soil depth for organic farming system had significant greater PC1 scores than the others.

It is concluded that soil physical, chemical and biological conditions increased with PC1 scores. Among organical and conventional cultivation systems, PC1 is represented by the following order: (0-50 cm) organic > (0-50 cm) conventional > (0-100 cm) organic > (0-100 cm) conventional. This order is due to the increase in the organic matter content of 0-50 cm soil depth for the conventional farming system from 0-100 cm of the organic.

The PC2 represented by 10.29, 10.87% the total variance for farming system for 0-50 and 0-100cm soil depths of the organic farming system and 15.3 and 16.68% for the 0-50 and 0-100cm soil depths for conventional farming system respectively. The PC2 had significant positive results for the total microbial count. The strong dependence of PC2 on the total microbial count suggests that soils with higher PC2 scores are more consolidated. According to Harris *et al* (1996), total biomass affects the environmental and production functions of soil quality.

The PCA of indicators that exhibited significant differences between management systems for the four areas, two PCs had eigenvalues >1. Highly weighted variables under PC1 included OM %, porosity, soil moisture at 0.1 bar, available water, penetration resistance, MWD, silt+ clay, IR and CEC. Correlation coefficients among these variables revealed that IR is not correlated with the other highly weighted variables. Hence, IR was retained for the MDS of the remaining eight well-correlated variables. SOM was the most highly correlated and was chosen for the MDS as the most representative for that group.

This result indicates that organic matter, biological and physical aspects of soil quality were the most sensitive indicators of soil quality. Farming systems aspect of soil quality increased with PC1 scores.

The data in Table (7) reveal that, the SQI was calculated using weighting factors for each scored MDS variable according to the following formula:

$$SQI = \sum W_i \times S$$

Where S is the score for the subscript variable and the coefficients are the weighting factors derived from the PCA. Weights were determined by the percent of variation in the data set explained by

the PC that contributed the indicated variable divided by the total percentage of variation explained by all PCs with eigenvalues > 1.

The data in Table (8) reveal for organic and conventional farming systems that soil quality index for 0-50 cm soil depth in organic farming system > 0-100 cm soil depth in organic farming system > or = 0-50 cm soil depth in conventional farming system > 0-100 cm soil depth in conventional farming system.

REFERENCE

- Andrews, S.S. (1998). *Sustainable Agriculture Alternatives: Ecological and Managerial Implications of Poultry Litter Management Alternatives Applied to Agronomic Soils*. pp. 68-109. Ph.D. Diss., Univ. of Georgia, Athens.
- Andrews, S.S. and C.R. Carroll (2001). Designing a decision tool for sustainable agro ecosystem management: Soil quality assessment of a poultry litter management case study. *Ecol. Applic.* 11(6): 1573-1585.
- Brejda, J.J.; T.B. Moorman; D.L. Karlen and T.H. Dao (2000). Identification of regional soil quality factors and indicators: I. Central and Southern High Plains. *Soil Sci. Soc. Am. J.* 64: 2115-2124.
- Doran, J.W. (2002). Soil health and global sustainability: translating science into practice. *Agriculture, Ecosystem & Environment*. 88: 119-127.
- Doran, J.W. and T.B. Parkin (1994). Defining and Assessing Soil Quality. For a sustainable environment. In: Doran J.W.; D.C. Coleman; D.F. Bezdicek and B.A. Stewart (eds). *Defining Soil Quality for a Sustainable Environment*. *Soil Sci. Soc. Am. Special Publication No. 35*. 44: 450-457. Madison, Wisconsin.
- Doran, J.W. and T.B. Parkin (1996). Quantitative indicators of soil quality: A minimum data set. In: Method for Assessing Soil Quality, Doran J.W. and A.J. Jones (eds.). *Soil Sci. Soc. Am. Special Publication No. 49*, pp. 25-37. Madison, Wisconsin, USA.
- Harris R.F.; D.L. Karlen and D.J. Mulla (1996). A conceptual framework for assessment and management of soil quality and soil health. In: Doran J.W., Jones A.J., eds. *Methods for Assessing Soil Quality*. *Soil Sci. Soc. Am. Special Publ.* 49: 61-82. Madison, WI.
- Hussain, I.; K.R. Olson; M.M. Wander and D.L. Karlen (1999). Adaptation of soil quality indices and application to three tillage systems in Illinois. *Soil Till. Res.* 50: 237-249.

- Jawson, Michael, D. (2001).** *Soil Management and Conservation Chapter 4.2 p. 10.* www.ers.usda.gov/publications/arei/ah722/arei4_2/AREI4_2soilmgmt.pdf
- Karlen, D.L.; M.J. Mausbach; J.W. Doran; R.G. Cline; R.F. Harris and G.E. Schuman (eds) (1997).** Soil quality: a concept definition and framework for evaluation. *Soil Sci. Soc. Am. J.* **61:** 4-10.
- Larson, W.E. and D.J. Pierce (1994).** The dynamics of soil quality as a measure of sustainable management in. Doran, J.W. Coleman, D.C. Bezdicsek D.F. and Stewart B.A. (Eds). *Defining soil quality for a sustainable environment. Soil Sci. Soc. Am Special Publication No. 35, 44:* 450-457. Madison. Wisconsin.
- Seybold, C.A.; M.J. Mausbach; D. L. Karlen and H.H. Rogers (1996).** Quantification of soil quality. *Adv. Soil Sci.* **50:** 229-236.
- SPSS. (2002).** SPSS statistical package for windows. Users manual. Suhaedi E., Metternicht G., and Lodwick G. 2002 Geographic information systems and multiple goal analysis for spatial land use modeling in Indonesia. [www. Gisdevelopment. Net/ aars/ acrs/2002/ luc/ luc002. Shtml.](http://www.Gisdevelopment.Net/aars/acrs/2002/luc/luc002.Shtml)
- Van Noordwijk, M.; C. Cerri; P.L. Woomer and K. Nugroho (1997).** Soil carbon dynamics in the humid tropical forest zone. *Geoderma*, **79:** 187-225.
- Vance, E.D. (2000).** Agricultural site productivity: principles- derived from long- term experiments and their implications for intensively managed forest. *Forest Ecology and Management.* **138:** 369-396.
- Webster, R. (1977).** *Quantitative and Numerical Methods in Soil Classification and Survey.* **52:** 1162-1167. Clarendon Press, Oxford.
- Wilding, L.P. and L.R. Drees. (1983).** Spatial variability and pedology. In: *Pedogenesis and Soil Taxonomy. I: Concepts and Interpretation.* pp. 83-116, Elsevier, Amsterdam.



أثر نظم الإدارة الزراعية على جودة التربة فى مصر

[٣٣]

هدى عبده ايليا^١ - محمود طلحة^٢ - محمود يوسف عفيفي^١ - الحسنا أبو جبل^٢

١- مركز بحوث الصحراء - المطرية - القاهرة - مصر

٢- قسم الأراضى - كلية الزراعة - جامعة عين شمس - شبرا الخيمة - القاهرة - مصر

ومعدل الرشح الأساسى والسعة التبادلية الكاتيونية
والعد الميكروبي الكلى.

ودلت النتائج أن خواص التربة البيولوجية أقل
أهمية من الخواص الطبيعية والخواص الكيمبأوية.
وأن التأثير النسبى للخواص من التباين الكلى للمكون
الأساسى الأول ٧٩,٩٧ و ٧٨,٠٥ ٪ وذلك للعمق
(صفر - ٥٠سم) لكلا من الزراعة العضوية والزراعة
التقليدية على التوالى. وفى نفس الوقت كان التأثير
النسبى ٧٥,٧٢ و ٧١,٩٤ ٪ للعمق (صفر - ١٠٠
سم) لكلا من النظامين (الزراعة العضوية والزراعة
التقليدية) على نفس المنوال.

وكان العامل الوحيد ذو التأثير المعنوى للمكون
الأساسى الثانى هو العد الميكروبي الكلى. وأن التأثير
النسبى له يمثل ١٠,٢٩ و ١٠,٨٧ ٪ من التباين
الكلى للخواص لكلا من العمقين (صفر - ٥٠سم)
(صفر - ١٠٠ سم) للزراعات العضوية. بينما كان
التأثير النسبى يمثل ١٥,٣ و ١٦,٦٨ ٪ من التباين
الكلى للخواص لكلا من العمقين (صفر - ٥٠ سم)
(صفر - ١٠٠ سم) للزراعات التقليدية على
التوالى.

الموجز

الهدف من هذه الدراسة هو تقييم أثر الأعمار
المختلفة للزراعة العضوية على جودة التربة مقارنة
بنظام الزراعة التقليدية.

ولإتمام هذه الدراسة تم إختيار مواقع عديدة
مختلفة فى القوام وعمر الزراعة العضوية. وكانت
المواقع المختارة هى بلبس بمحافظة الشرقية
والمنايف بمحافظة الإسماعيلية وسكران بمحافظة
الفيوم و(إيتاى البارود وابو المطامير) بمحافظة
البحيرة

وأىضا تم إختيار مواقع أخرى للزراعات التقليدية
مجاورة للزراعات العضوية للمقارنة.

تم التقييم على أساس مدى التأثير فى الخواص
الطبيعية والخواص الكيمبأوية والخواص البيولوجية
للأعماق (صفر - ٥٠ سم) و (صفر - ١٠٠ سم)
وكانت الخواص المأخوذة فى الأعتبار ذات التأثير
المعنوى هى: محتوى التربة من المادة العضوية
والكتافة الظاهرية والمسامية والماء الميسر ومقاومة
التربة لإختراق الجذور ومتوسط القطر الوزنى

تحكيم: أ.د التونى محمد على

أ.د سمير على محمد