

## **Suppressive Effect of Sulphadimidine on Major Histocompatibility Complex as an Immune Parameter in Broiler Chicken** **A. El-Mahmoudy<sup>1\*</sup>, M. E. Mahmoud<sup>2</sup> and K. Farah<sup>3</sup>**

Departments of Pharmacology<sup>1</sup> and Clinical Pathology<sup>3</sup>, 13736 Moshtohor Faculty of Veterinary Medicine, Benha University, Egypt;

<sup>2</sup>Department of Animal Hygeine and Behaviour, Faculty of Veterinary Medicine, Assiut University, Egypt.

\*Coressponding author E-mail: [a.elmahmoudy@yahoo.com](mailto:a.elmahmoudy@yahoo.com)

### **Abstract**

The present study aimed at evaluation of the immuomodulatory effect of sulphadimidine by investigating its ability to affect constitutive expression of MHC-II molecules on B-cells, which are one determinant of antigen presentation and the vigor of immunity. Three groups of broiler chicken were used; the first two groups received 1 g/kg and 2 g/kg sulfadimidine, respectively, in drinking water; while the third group was left as control. Bursal samples were taken after 7 and 14 days of treatment and checked for MHC-II gene expression by RT-PCR technique. The obtained results demonstrate that oral administration of sulphadimidine inhibited the expression of MHC-II mRNA in bursal of Fabricius of broiler chicken in a dose and age dependent manner. This finding is of clinical importance as many farmers routinely add sulpha drugs to drinking water as a prophylactic measure against some infections; yet the data presented in this study doesn't recommend this managing behavior.

Key words: Major histocompatibility complex class II, (MHC-II), Sulphadimidine, Bursa of Fabricius, B-cells, Broiler chickens.

### **Introduction**

Major histocompatibility complex class II (MHC-II) molecules are considered all-or-non gate rule for exogenous antigen presentation and consequent initiation of adaptive immunity by the virtue of their ability to present non-self peptides to the receptors of helper lymphocytes. Presentation of antigen is necessary for lymphocyte activation and differentiation (23 and 18). According to the type of antigen, the MHC molecules presented as class I or class II (6). Hence, MHC molecules play important roles in the regulation of the immune response by

*Third Inter. Sci. Conf., 29 Jan.- 1 Feb./ 2009, Benha & Ras Sudr, Egypt  
Fac. Vet. Med. (Moshtohor), Benha Univ*

communicating among different cellular components of the immune system: T cells, B cells, and antigen-presenting cells (14).

Chicken MHC-II (called blood linked complex; B<sub>L</sub>-complex) genes are  $\alpha$  and  $\beta$  heterodimeric transmembraneous glycoproteins. The main component responsible for antigen presentation is  $\beta$ -molecule, while  $\alpha$ -molecule is of relatively lower function. In a general sense, the bursa of Fabricius of chicken is a unique organ that provides only B-lymphocytes (B-cells), initiates the adaptive (specific) immune response and an equivalent to the bone marrow in mammals, at least with respect to B-cell development. Also the bursa of Fabricius is critical only within the first 2-3 weeks after hatching. As about 5% of bursal cells leave the bursa per day to the periphery after that the bursa will be functionless (15 and 17). MHC-II genes are one marker of B-cell receptors. Although B-cells constitutively express these surface molecules to counteract pathogens, their expression can be induced by infections or pro-inflammatory cytokines IFN- $\gamma$  and TNF- $\alpha$ .

Evidence from several laboratories in animals suggested that the level of MHC-II mRNA reflects its surface expression (22; 21; 16; 18; 7; and 27)

In spite of these facts, there is no available information regarding the effect of sulphonamides on adaptive (MHC-II) components in chicken. Therefore, the ultimate goal of this study is to elucidate the effect of sulpham drug, exemplified by sulphadimidine, treatment on MHC-II-gene expression in chickens.

## **Material and Methods**

### **1. Experimental birds**

A day old, newly hatched male chicks (Ross strain, body weight of 40 $\pm$ 5 g), each was permitted free access for standard nipple drinker and feed trough. Chicks were maintained on 12:12 light-dark cycle and kept in thermostatically controlled cages (Hokkaido, Japan) to match chick requirements. Starting ambient temperature was 35°C and decreased 1°C each other day.

### **2. Experimental design**

Birds were divided to three groups each group consisted of 5 birds. Group 1 kept without any treatment (0 g sulphadimidine/L); group 2 was

administered 1 g sulphadimidine/L for 14 days; while group 3 was administered 2 g sulphadimidine/L for 14 days. Samples were taken on day 7 (2 samples from each group) and day 14 (2 samples from each group) of treatment. The experiment was repeated 3 times.

### **3. Measurement of MHC-II mRNA expression by RT-PCR**

Bursa of Fabricius was dissected from chicks at 7 and 14 days of treatment with sulphadimidine (0 g/L, 1 g/L, and 2 g/L). For detection of target genes (MHC-II), 14 samples from bursa of Fabricius were run together in each reverse transcription and polymerase chain reaction (RT-PCR) trial; 4 from 1 g/L-treated chicks, 2 from 7 & 2 from 14 day-treatment periods; 4 from 2 g/L-treated chicks, 2 from 7 & 2 from 14 day-treatment periods; and 4 from untreated ones (0 g/L). Two additional samples (1 from 1 g/L group and the other from 2 g/L group) that were run with no reverse transcriptase (RT) added (called 'RT' controls).

The bursa of Fabricius was cooled in ice-cold distilled water and homogenized. Total cellular RNA was extracted from tissue homogenates by using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) (2). The RNA concentration was determined on the basis of absorbance at 260 nm, and its purity was evaluated by the ratio of absorbance at 260/280 nm (>1.8). After digestion of genomic DNA by treatment with deoxyribonuclease I (Invitrogen), first-strand cDNA was synthesized from 2 µg of total RNA by using Superscript II RNase, reverse transcriptase and oligo (dT) primer (Invitrogen). To validate successful deoxyribonuclease I treatment, the reverse transcriptase was omitted in control reactions. The absence of PCR-amplified DNA fragments in these samples indicated the isolation of RNA free from genomic DNA contamination.

For performing PCR, 1 µl of template cDNA was added to a PCR cocktail consisting of 10 mM Tris-HCl, 50 mM KCl, 1.25 mM MgCl<sub>2</sub>, 10 mM deoxy-NTPs, 200 nM primers, and 0.05 U Taq DNA polymerase (Takara Bio, Otsu, Japan) for a final volume of 10 µl. Primers set for house keeping β-actin gene (GenBank accession number; NM\_031144) were as follows: sense, 5'-gtg ttc cac cag gag atg ttg-3' and anti-sense, 5'-ctc ctg ccc atc gag ttc gtc-3' (predicted size = 434 bp); BL-alpha (GenBank accession number; AY357253), sense 5'-ggg ccg tcc tga agccgc ac-3' and antisense, 5'-ttc cgg ctc cca cat cct ct- 3' (predicted size = 410 bp). BL-beta (GenBank

*Third Inter. Sci. Conf., 29 Jan. - 1 Feb./ 2009, Benha & Ras Sudr, Egypt*

*Fac. Vet. Med. (Moshthor), Benha Univ*

accession number; AY510090), sense; 5'-gcg ggg gcc gtg ctg gtg gc-3' and antisense; 5'-acg tag cac gcc aga cgg tc-3' (predicted size = 560 bp). PCR was performed with a thermal cycler (GeneAmp. PCR system 2700, Applied Biosystems, Foster City, CA, USA). A first incubation at 94 °C for 3 min was performed for initial denaturation. This was followed by 25 cycles of the following sequential steps: 30 s at 94 °C (denaturation), 1 min at 55 °C or 51 °C for  $\beta$ -actin or Bl-alpha, and at 57°C for BL-beta respectively (annealing), and 1 min at 72 °C (extension); the final incubation was at 72 °C for 7 min (final extension). The reaction products were electrophoresed on 1% agarose gels and stained with ethidium bromide (0.2  $\mu$ g/ml). The gels were exposed to UV with a UV transilluminator (UVP Laboratory Products, Upland, CA, USA), and the images were picked up using a camera and analyzed only qualitatively.

## Results

To match doses of the drug routinely used by farmers, 2 doses (1 and 2 g/L) were selected in drinking water along their concept that they protect their chicks against possible infections. Because the genes encoding MHC-II molecules are regulated in tight fashion at mRNA level to meet with the local requirements for the adaptive immune response, measurement of MHC-II by RT-PCR was adopted. As shown in Fig (1), administration of sulphadimidine for first 7 days of age resulted in suppression of MHC-II molecules indicated by absence of the bands representing  $\square$ -component of the molecule with non-significant effect on  $\square$ -component. The effect was observed in both doses of sulfadimidine used (1 and 2 g/L).

Unexpectedly, as shown in Fig. (2) the suppressive effect of the sulpha drug used on MHC-II molecules after 14 days of treatment in drinking water, was relatively less than that recorded after 7 days of treatment, however the suppressive effect was still evident. Also,  $\square$ -component was not affected while the whole effect was only on the main  $\square$ -component. The effect was unrelated the dose of sulphadimidine used as the effect was similar in 1 and 2 g/L groups.

The effect of sulphadimidine is specific to the drug as the control  $\square$ -actin bands were intact in both treated and control groups (Fig. 3).

*Third Inter. Sci. Conf., 29 Jan.- 1 Feb./ 2009, Benha & Ras Sudr, Egypt  
Fac. Vet. Med. (Moshtohor), Benha Univ*




Figure 1: Effect of administration of sulphadimidine in drinking water (1g/L and 2g/L) to chicks for 1 week and 2 weeks on expression of MHC-II ( $\beta$ -component) by bursal cells.




Figure 2: Effect of administration of sulphadimidine in drinking water (1g/L and 2g/L) to chicks for 1 week and 2 weeks on expression of MHC-II ( $\alpha$ -component) by bursal cells.




Figure 3: The non-effect of administration of sulphadimidine in drinking water (1g/L and 2g/L) to chicks for 1 week and 2 weeks on expression  $\beta$ -actin by bursal cells; indicating specific action of the drug used.

### Discussion

Long term ago, some farming problems as coccidiosis is being controlled by routine supplying sulpha drugs, such as sulphadimidine, into drinking water to chicks. Although this prophylactic measure was successful sometimes, and unsuccessful other times, findings in this study strongly recommend avoiding sulpha drug administration to chicks especially in newly hatching chicks. This recommendation comes from data of this manuscript which clearly showed the immunosuppressive effect of sulphadimidine by inhibiting MHC class II mRNA expression. As a result, this drug could suppress the antigen presentation function of bursal immune cells. It is well established that any antigen is not processed as it is in immune pathways in the body; any antigen must be firstly presented by antigen presenting cells such as macrophages,

lymphocytes and fibroblasts to molecules recognizable by lymphocytes that are MHC-II molecules (23; 6; 20; 18 and 27). This crucial function of the MHC in the immune response makes these molecules a promising candidate for genetic selection to improve avian immunity.

Failure to present antigens to immune cells, results in weakness or even absence of immune defense mechanisms adopted upon antigen invading with serious outbreaks and economic losses. In addition, the immunological response to a given vaccine will be also drastically affected. In fact, there is an association between the chicken MHC and antibody production against a variety of antigens (5; 13; 14; 12 and 24), since long-term selection for antibody response to SRBC was accompanied by a correlated change in MHC allelic frequencies in the chicken (4; 19 and 25). Furthermore, genetic selection of optimal MHC genotypes was used for improvement of humoral immunity (26). Interestingly, chicken MHC is roughly 20 fold smaller than the human MHC, therefore, this crucial property (smaller size and simplicity) makes it possible to study the striking MHC-determined pathogen specific disease resistance at the molecular level (1 and 9). Among markers of B-cells is MHC-II which is expressed on their membranes to counteract pathogens, and also can be induced by infection, pro-inflammatory cytokines IFN- $\gamma$  and TNF- $\alpha$ . MHC molecules could be measured at genetic level in terms of their mRNA expression (22; 21 and 7). Unlike viral infection which down-regulates the surface expression of MHC class I or destroys B-cells in busa of Fabricius cells in chicken early life (3 and 8), sulphadimidine inhibits the constitutive expression of MHC-II in chicken.

The classical chains alpha and beta of chicken MHC-II are not necessarily to be linked genetically and the beta chains are highly variable between individuals (11). In mammalian species,  $\beta$ -component is the main component while  $\alpha$ -component comes in the second rank. The reverse is true in chicken where the main component responsible for antigen presentation is  $\alpha$ -molecule, while  $\beta$ -molecule is of relatively lower function. Therefore, the more conserved L-alpha chain was the target of (11) to investigate the regulation procedures of these genes in chicken. The fact that BL-alpha chains are more conserved has been reproduced in

present study (Fig. 2). On the other hand, BL-beta was affected by our treatment constantly in 7 and 14 day-old chicken as shown in Fig. (1).

The bursa of Fabricius of chicken provides only B-lymphocytes that initiate the adaptive (specific) immune response upon invading antigen or vaccination. It strongly determines the resistance and susceptibility to infectious diseases (9). Because role of the bursa of is critical only within the first 2-3 weeks after hatching; after that that it becomes functionless (15 and 17), therefore, we focused our research on the first 2 weeks after hatching. In the present study, the constitutive expression of MHC-II (BL- $\beta$ ) mRNA by bursal cells was severely inhibited upon administration of sulphadimidine for 7 and 14 days at 1 and 2 g/L drinking water. However, the suppressive effect of sulphadimidine on MHC-II mRNA expression was not only dose dependent (more evident at high dose) but also age dependent (1 week-old chicks were affected more than 2-weeks old chicks).

In fact, the effectiveness of sulphonamides against some microbes is out of debate, and the use of these drugs could be inevitable. However, our finding strongly recommends avoiding sulphadimidine administration to chicks especially in the first week after hatching as this may lead, on the other hand, to drastic immune effect which makes chicks more liable to microbes to which sulpha is not a protector.



## References

1. Briles, W., Briles, R., Taffs, R., Stone, H. (1983): Resistance to malignant lymphoma in chicken is mapped to a subregion of major histocompatibility (B-complex). *Science*, 219, 977-979.
2. Chomczynski, P. and Sacchi N. (1987): Single step method of RNA isolation by acid guanidinium thiocyanate-phenol chloroform extraction. *Anal. Biochem.*, 162, 156-159.
3. Cohen, J. I. (1998): Infection of cells with varicella-zoster virus downregulates surface expression of class I major histocompatibility complex antigens. *J. Infect. Dis.*, 177(5), 1390-1393.
4. Dunnington, E.A., Briles, R.W., Briles, W.E., Gross, W.B., Siegel, P.B. (1984): Allelic frequencies in eight alloantigen systems of chickens selected for high and low antibody response to sheep red blood cells. *Poultry Sci.*, 63: 1470-1472.
5. Dunnington, E.A., Larsen, C.T., Gross, W.B., Siegel, P.B. (1992): Antibody responses to combinations of antigens in white Leghorn chickens of different background genomes and major histocompatibility complex genotypes. *Poultry Sci.*, 71: 1801-1806.
6. Glimcher, L.H. and Kara, C.J. (1992): Sequences and factors: A guide to MHC class II transcription. *Annu. Rev. Immunol.*, 10-13.
7. Hobart, M. V; Ramassar, N; Goes, J; Urmson, P. and Halloran, F. (1997): IFN regulatory factor-1 plays a central role in the regulation of the expression of class I and II MHC genes *in vivo*. *J. Immunol.*, 158: 4260.
8. Hunt, H.D., Lupiani, B., Miller, M.M., Gimeno, I., Lee, L.F., Parcels, M. S. (2001): Marek's Disease Virus Down-Regulates Surface Expression of MHC (B complex) Class I (BF) Glycoproteins during Active but not Latent Infection of Chicken Cells. *Virology*, 282, 198-205.
9. Kaufman, J., Lamont, S. (1996): The chicken major histocompatibility complex. In: Schook LB, Lamont SJ (Ed) 1996, The major histocompatibility region in domestic animal species. CRC, Boca Raton, Fla, pp 35-64.
10. Kaufman, J., Milne, S., Gobel, T.W., Walker, B.A., Jacob, J.P., Auffray, C., Zoorob, R., Beck, S. (1999): The chicken B locus is a minimal essential major histocompatibility complex. *Nature* 401: 923-925.
11. Kumanovics, A; Takada, T. and Lindahl, K.F. (2003): Genomic organization of the mammalian MHC. *Annu. Rev. Immunol.*; 21: 629-6257.
12. Karaca, M., Johnson, E., Lamont, S.J. (1999): Genetic line and major histocompatibility complex effects on primary and secondary antibody responses to T-dependent and T-independent antigens. *Poultry Sci.*, 78: 1518-1525.
13. Lakshmanan, N., Gavora, J.S., Lamont, S.J. (1997): Major histocompatibility complex class II DNA polymorphisms in chicken strains selected for Marek's disease resistance and egg production or for egg production alone. *Poultry Sci.*, 76: 1517-1523.
14. Lamont, S.J. (1998): Impact of genetics on disease resistance. *Poultry Sci.*, 77: 1111-1118.

*Third Inter. Sci. Conf., 29 Jan.- 1 Feb./ 2009, Benha & Ras Sudr, Egypt  
Fac. Vet. Med. (Moshtohor), Benha Univ*

15. Lassila, O. (1989): Emigration of B cells from the chicken bursa of Fabricius. *Eur J. Immunol.*, 19, 955-958.
16. Lee, Y.J; Han, Y; Lu, H.T; Nguyen, V; Qin, H; Howe, P H; Hocevar, B. A; Boss, J. M; Ransohoff, R. M. and Benveniste, F. N. (1997): TGF- $\beta$  suppresses IFN- $\gamma$  induction of class II MHC gene expression by inhibiting class II transactivator messenger RNA expression. *J. Immunol.*, 158: 2065.
17. Paramithiotis, E. and Ratcliffe, M.J. (1994): B cell emigration directly from the cortex of lymphoid follicles in the bursa of Fabricius. *Eur. J. Immunol.*, 24, 458-463.
18. Pieters, T. (1997): MHC class II restricted antigen presentation. *Curr. Opin. Immunol.*, 9: 89-96.
19. Pinard, M.H., Janss, L.L., Maatman, R, Noordhuizen, J.P., van der Zijpp, A.J. (1993): Effect of divergent selection for immune responsiveness and of major histocompatibility complex on resistance to Marek's disease in chickens. *Poultry Sci.*, 72:391-402.
20. Riley, J.L.; Westerheide, S. D.; Price, J. A.; Brown J. A. and Boss, J. M. (1995): Activation of class II MHC genes requires both the X box region and the class II transactivator (CIITA). *Immunity*, 2: 533.
21. Rohn, W.; Lee, Y., Benveniste, F. (1996): Regulation of class II expression. *Crit. Rev. Immunol.*, 16:311-330.
22. Sicher, S.C; Chung, G.C; Vazquez, M.A. and Lu, C.Y. (1995): Augmentation or inhibition of IFN- $\gamma$  induced MHCII expression by Lipopolysaccharides. The roles of TNF- $\alpha$  and nitric oxide, and the importance of the sequence of signaling. *J. Immunol.*, 155: 5826-5834.
23. Unanue, E.R. (1992): Cellular studies on antigen presentation by class II MHC molecules. *Curr. Opin. Immunol.*, 4: 63.
24. Weigend, S., Matthes, S., Solkner, J., Lamont, S.J. (2001): Resistance to Marek's disease virus in White Leghorn chickens: effects of avian leucosis virus infection genotype, reciprocal mating, and major histocompatibility complex. *Poultry Sci.*, 80: 1064-1072.
25. Yonash, N., Kaiser, M.G., Heller, E.D., Chaner, A., Lamont, S.J. (1999): Major histocompatibility complex (MHC) related cDNA probes associated with antibody response in meat-type chickens. *Anim. Genet.*, 30: 92-101.
26. Zhou, H., and Lamont, S.J. (2003): Chicken MHC class I and II gene effects on antibody response kinetics in adult chickens. *Immunogenetics*, 55:133-140.
27. Zhu, S.X; Pai, R.K; Askew, D; Boom, W.H. and Hardinng, C.V. (2002): Regulation of class II MHC expression in APCs: Roles of types I, III and IV class II transactivator. *J. Immunol.*, 169:1326-1333.

التأثير المثبط لعقار السلفاديميدين على إمتش سي كعنصر مناعي في دجاج التسمين

ابوبكر المحمودي<sup>١</sup>, معتمد محمود<sup>٢</sup>, خالد فراره<sup>٣</sup>  
<sup>١</sup> قسم الفارماكولوجيا والباثولوجيا الإكلينيكية بكلية الطب البيطري - جامعة بنها  
<sup>٢</sup> قسم صحة وطبائع الحيوان بكلية الطب البيطري - جامعة أسيوط

### الملخص العربي

تقوم جزيئات مركب التوافق النسيجي الثاني (إم إتش سي الصنف الثاني) بأدوار حاسمة في التفاعل المناعي التكيفي إستناداً إلى قدرتهم لتقديم الأنتيجينات الغير ذاتية إلى الخلايا المناعية المساعدة (CD4T-Helper cells) و من ثم تؤدي إلى تنشيطها وتفاضلها. وان الجينات التي تُشفر جزيئات إم إتش سي الصنف الثاني تكون بالتالي مسنولة بشكل كبير عن نمو التفاعل المناعي التكيفي. ولذا وجود هذه الجينات بصورة كافية ضرورة لضمان سلامة الجهاز المناعي التكيفي. وحيث أن حويصلة فابريسي تقوم بدور هام في تطور المناعة المكتسبة في الطيور (مشابه لدور لنخاع العظام في الثدييات), لكن يتضاءل دورها بعد الثلاث الأسابيع الأولى من العمر. ولدراسة تأثير مركبات السلفا على الجهاز المناعي التكيفي تم قياس جينات إم إتش سي الصنف الثاني عند ٧ و ١٤ يوماً من عمر الكتاكيت بعد معالجتها بعقار السلفاديميدين. توضح النتائج التي تم الحصول عليها بواسطة استخدام تقنية RT-PCR أن إعطاء عقار السلفاديميدين في مياه الشرب (١ جم و ٢ جم لكل لتر) يمنع تعبير mRNA الخاص بـ إم إتش سي الثاني في دجاج التسمين وذلك على حسب الجرعة المعطاة. هذه النتائج تبين التأثير المثبط للمناعة لعقار السلفاديميدين ولذلك فهي ذات أهمية حقلية كبرى حيث أن الكثير من المزارعين يستخدمون مركبات السلفا في الكتاكيت بشكل دوري كإجراء وقائي ضد بعض الإصابات, ولكن البيانات التي قدمت في هذه الدراسة توصي بعدم استخدام عقار السلفا لهذا الغرض.