

## SEASONAL FLUCTUATION OF *HELICOVERPA ARMIGERA* (HUBNER) (LEPIDOPTERA: NOCTUIDAE) ON COTTON AND OKRA AND HEAT UNITS RELATED

EL-SAYED, A.A.A, MERVAT A. KANDIL AND A.E. AMER

Plant Protection Research Institute, ARC, Doki, Giza

(Manuscript received 14 November 2009)

---

### Abstract

The seasonal fluctuation of the American bollworm (ABW), *Helicoverpa armigera* (Hub.), (moths and larvae) was studied on cotton, *Gossypium barbadense* and okra, *Hibiscus esculentus*, in Dakahlia Governorate during 2007 and 2008 cotton seasons. Results revealed that the population of ABW male moths occurring three peaks each season. The first peak had highest average number of male moths than others during the two seasons. Populations of ABW larvae infest cotton and okra plants appeared two peaks during the same period. The average numbers of larvae on okra was higher than cotton during the two seasons and average. The probability regression values revealed highly significant between accumulated heat units and each of ABW male moths and larval infestation in cotton and okra. The calculated  $R^2$  values ranged between 0.8148 and 0.9619. The same trend was found between accumulated ABW male moths and larval infestation in cotton and okra. The calculated  $R^2$  values ranged between 0.9185 and 0.9793. Also, between larval infestation in okra and cotton plants were highly significant. The calculated  $R^2$  values ranged between 0.6030 and 0.9899. Consequently, accumulated heat units can be used to predict pheromone traps catch and larval infestation on okra and cotton. Pheromone traps catch can be used to predict larval infestation on okra and cotton also can be used larval infestation on okra to predict larval infestation on cotton.

### INTRODUCTION

The American bollworm *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) is a polyphagous pest causing extensive damage on many crops such as cotton, okra and tomato (Sharma, 2001). Intercropping cotton with okra reduced the boll damage and okra acts as a good trap and highly preferred for feeding more than cotton for *H. armigera* (DuraimuruJan and Regupathy 2005 a and b). Eggs numbers of *H. armigera* were significantly higher on sunflower, okra and tomato than cotton, but larval numbers were not significantly differed from cotton at comparable time (Ravi *et al.* 2005). The host plants sequence and host suitability play an important role in population dynamic of *H. armigera* male moths (Fitt, 1989). Mean survival, development and reproductive performance of *H. armigera* varies on different food plants (Pretorius, 1976 and Jallow *et al.* 2001). The pheromone traps are convenient and powerful tools which gives reliable information for monitoring of population

density and intensity and for forecasting of *H. armigera* infestation (Gawand *et al.*, 2006). Pheromone traps had been used to determine the seasonal dynamics (Slosser *et al.*, 1987). Monitoring the population of *H. armigera* is greatly important in the programs of integrated pest management in crops.

Common in Egypt, okra is planted in small area adjacent cotton fields. Thus, the aim of this work was to study seasonal fluctuation of *H. armigera*, relationships between accumulated heat units, male moths and larval infestation on cotton and okra plants.

## MATERIALS AND METHODS

The present study was carried out at Aga district in Dkahlia Governorate during two cotton seasons 2007 and 2008. The cotton variety Giza 86 and okra were planted on the second week of April during the two seasons. The total area planted with cotton at Aga district was about 150 and 75 feddan in 2007 and 2008 cotton season, respectively. The experimental area was 2200 m<sup>2</sup> during the two seasons. The cotton crop occupied in area of 2000 m<sup>2</sup> and okra about 175 m<sup>2</sup> on the border of the cotton area. The okra was normally planted on border of cotton plants in Aga districts. The agronomic practices were carried out as usual. The experimental area was sprayed according to the recommendation ministry of agriculture program.

Three funnel pheromone traps were fixed in cotton fields throughout a period extended from fourth week of May to the 3<sup>rd</sup> week of September during the two cotton seasons 2007 and 2008. The pheromone capsule was changed every two weeks. The numbers of *H. armigera* male moths caught was recorded weekly. Daily maximum and minimum temperatures were obtained from the Meteorological Laboratory, Agricultural Research Center. The degree days were calculated according to Seaver *et al.* 1990 and the lower and upper temperature thresholds were 11.73 and 30 °C. The accumulated heat units were calculated from 1<sup>st</sup> January in both seasons.

### Samples

Weekly twenty plants from each cotton and okra plants were chosen week per randomly and investigated carefully to estimate the larvae of *H. armigera* from the fourth week of May to the 3<sup>rd</sup> week of September.

### Statistical analysis

Data were statistically analyzed using 'Costat' program a product of Cohort software Inc., Berkeley, California.

## RESULTS AND DISCUSSION

### 1. Seasonal fluctuation of male moths

The population of ABW, *H. armigera*, male moths captured per trap fluctuated during the two seasons showing up three peaks each (Figures 1). The average number of ABW captured increased gradually from the fourth week of May to reach the first peak on 21<sup>st</sup> June and 14<sup>th</sup> June with 135 and 175 moth/trap/week during 2007 and 2008 cotton seasons, respectively. The population declined gradually and increased to reach the second peak at 26<sup>th</sup> and 19<sup>th</sup> July with 25 and 55moths/trap/week during the first and second season, respectively. The population decreased and after that, the mean numbers of male moths captured increased till the third peak on 30<sup>th</sup> August during the two seasons. The first peak had the highest number of male moths captured than the others during the two seasons. The total numbers of male moths captured was higher during the second season than the first one.

### 2. Seasonal fluctuation of larval infestation in cotton and okra

The population of ABW larval infestation in cotton and okra crops started earlier one week on okra than cotton during the two seasons (Figures 1). On okra, larval infestation began at the fourth week of May with 1 and 2 larva/20 plants, while on cotton plants began at the first week of June with 6 and 5 larva/20 plants during 2007 and 2008 cotton seasons, respectively. Larval populations fluctuated on cotton and okra crops occurring two peaks during each season. On okra plants, the infestation with the ABW larvae increased gradually to reach the first peak with 17 and 20 larvae/20 plants at the fourth week of June and the first week of July during the two seasons, respectively. And after that, the population decreased and fluctuated during July and August till the second peak with 11 and 13 larvae/20 plants on second and first week of September during the 2007 and 2008 seasons, respectively. On cotton plants, the infestation with the ABW larvae increased gradually to reach the first peak with 13 and 17 larvae/20 plants at fourth week of June during the two seasons, respectively. And after that, the population decreased and fluctuated during July and August till the second peak on second week of September (6 and 7 larvae/20 plants). The total numbers ABW larval infestation was higher on okra than cotton plants.

### **3- Heat units and male moths captured & larval infestation related**

The probability regression values between accumulated heat units (ACHu) and weekly average number of ABW male moths captured in sex pheromone funnel taps revealed that the slope values were insignificant during 2007 cotton season while, it was significant during 2008 season and average for the two seasons. However,  $R^2$  values were 0.1984, 0.4709 and 0.3948, for the two seasons and average, respectively (Tables 1-3 and Figures 2). Also, relationships between accumulated heat unit and larval infestation in cotton and okra were insignificant for the two seasons and average, while highly significant at accumulated male moths, accumulated larval infestation in cotton and okra for the two seasons and average. The calculated  $R^2$  values ranged between 0.8148 and 0.9619 (Tables 1-3).

#### **Relation between male moths captured and larval infestation**

The probability regression values between male moths captured and larval infestation in cotton and okra were highly significant during the two seasons and average, except in case of okra during the second season which was insignificant. The obtained  $R^2$  values were (0.4959 & 0.4269), (0.4084 & 0.1769) and (0.5614 & 0.3953) for the two seasons and average, respectively. The probability regression values between accumulated male moths and accumulated larvae infested cotton and okra were highly significant during the two seasons and average. The calculated  $R^2$  values ranged between 0.9185 and 0.9793 (Tables 1-3 and Figures 3).

#### **Relation between okra and cotton larval infestation**

The probability regression values between larval infestation on okra and cotton plants were highly significant during the two seasons and average. Also, the regression value between accumulated larvae infested okra and accumulated larvae infested cotton were highly significant during the two seasons and average. The obverse  $R^2$  values ranged between 0.6030 and 0.9899 (Tables 1-3 and Figures 4).

Gawande *et al.* (2006) reported that the male moth trap catches of *H. armigera* had significant and positive correlation with its egg population in the field and infestation in green fruiting, whereas it had non significant correlation with larval population. Egg numbers were significantly higher on okra than on cotton (Ravi *et al.*, 2005). Survival of larval stage is very much dependent on the appropriate responses by females when selecting host plants (Fitt and Bayan, 1991 & Fitt, 1991).




Figure 1. Weekly average number of ABW caught moths, larvae infested cotton & Okra plants and Accumulated heat units during 2007, 2008 seasons and the average.

914 SEASONAL FLUCTUATION OF *HELICOVERPA ARMIGERA* (HUBNER) (LEPIDOPTERA: NOCTUIDAE) ON COTTON AND OKRA AND HEAT UNITS RELATED




Figure 2. Weekly accumulated average number of ABW caught moths (accmoth), larvae infested cotton (Acccotton) & Okra (Accokra) plants and accumulated heat units related during 2007, 2008 season and average.




Figure 3. Weekly accumulated number of ABW larvae infested cotton ( Accotton) and okra (Accokra) plants and accumulated moths caught related during 2007, 2008 seasons and average.

916 SEASONAL FLUCTUATION OF *HELICOVERPA ARMIGERA* (HUBNER) (LEPIDOPTERA: NOCTUIDAE) ON COTTON AND OKRA AND HEAT UNITS RELATED




Figure 4. Weekly accumulated number of ABW larvae infested cotton and okra plants related during 2007, 2008 seasons and average.



Table 1. Regression equations between accumulated heat units, weekly average numbers of ABW moths, larvae infest cotton and Okra plants during 2007cotton season

| Independent                      | dependant                          | Equation parameters |        |                           | R <sup>2</sup> |
|----------------------------------|------------------------------------|---------------------|--------|---------------------------|----------------|
|                                  |                                    | a                   | bx     | probability of regression |                |
| Accumulated heat units           | Mean number of moth                | 80.05               | -0.027 | .0640 ns                  | 0.1984         |
| Accumulated heat units           | Larvae infested cotton             | 9.63                | -0.003 | .0962 ns                  | 0.1634         |
| Accumulated heat units           | Larvae infested okra               | 10.26               | -0.002 | .4109 ns                  | 0.0427         |
| Accumulated heat units           | Accumulated moth                   | -168.33             | 0.310  | .0000 ***                 | 0.8864         |
| Accumulated heat units           | Accumulated larvae infested cotton | -25.54              | 0.043  | .0000 ***                 | 0.8796         |
| Accumulated heat units           | Accumulated larvae infested okra   | -46.11              | 0.066  | .0000 ***                 | 0.9289         |
| Mean number of moth              | larvae infested cotton             | 2.01                | 0.081  | .0011 **                  | 0.4959         |
| Mean number of moth              | larvae infested okra               | 4.04                | 0.094  | .0033 **                  | 0.4269         |
| Accumulated moth                 | Accumulated larvae infested cotton | -1.57               | 0.139  | .0000 ***                 | 0.9793         |
| Accumulated moth                 | Accumulated larvae infested okra   | -6.90               | 0.206  | .0000 ***                 | 0.9702         |
| Larvae infested okra             | Larvae infested cotton             | -0.61               | 0.742  | .0000 ***                 | 0.8492         |
| Accumulated larvae infested okra | Accumulated larvae infested cotton | 3.59                | 0.669  | .0000 ***                 | 0.9887         |

Table 2. Regression equations between accumulated heat units, weekly average numbers of ABW moths, larvae infest cotton and Okra plants during 2008 cotton season.

| Independent                      | dependant                          | Equation parameters |       |                           | R <sup>2</sup> |
|----------------------------------|------------------------------------|---------------------|-------|---------------------------|----------------|
|                                  |                                    | a                   | bx    | probability of regression |                |
| Accumulated heat units           | Mean number of moth                | 160.41              | -0.06 | .0017 **                  | 0.4709         |
| Accumulated heat units           | Larvae infested cotton             | 11.89               | 0.00  | .0851 ns                  | 0.1739         |
| Accumulated heat units           | Larvae infested okra               | 11.48               | 0.00  | .3057 ns                  | 0.0654         |
| Accumulated heat units           | Accumulated moth                   | -137.04             | 0.45  | .0000 ***                 | 0.8148         |
| Accumulated heat units           | Accumulated larvae infested cotton | -43.33              | 0.06  | .0000 ***                 | 0.9210         |
| Accumulated heat units           | Accumulated larvae infested okra   | -60.50              | 0.08  | .0000 ***                 | 0.9619         |
| Mean number of moth              | Larvae infested cotton             | 3.25                | 0.06  | .0043 **                  | 0.4084         |
| Mean number of moth              | Larvae infested okra               | 6.10                | 0.04  | .0822 ns                  | 0.1769         |
| Accumulated moth                 | Accumulated larvae infested cotton | -17.21              | 0.13  | .0000 ***                 | 0.9574         |
| Accumulated moth                 | Accumulated larvae infested okra   | -22.55              | 0.15  | .0000 ***                 | 0.9185         |
| Larvae infested okra             | Larvae infested cotton             | -0.11               | 0.78  | .0002 ***                 | 0.6030         |
| Accumulated larvae infested okra | Accumulated larvae infested cotton | 2.91                | 0.79  | .0000 ***                 | 0.9885         |

918 SEASONAL FLUCTUATION OF *HELICOVERPA ARMIGERA* (HUBNER) (LEPIDOPTERA: NOCTUIDAE) ON COTTON AND OKRA AND HEAT UNITS RELATED

Table 3. Regression equations between accumulated heat units, weekly average numbers of ABW moths, larvae infest cotton and okra plants (mean of two seasons), 2007- 2008 cotton seasons.

| Independent                      | dependant                          | Equation parameters |       |                           | R <sup>2</sup> |
|----------------------------------|------------------------------------|---------------------|-------|---------------------------|----------------|
|                                  |                                    | a                   | bx    | probability of regression |                |
| Accumulated heat units           | Mean number of moth                | 120.73              | -0.04 | .0052 **                  | 0.3948         |
| Accumulated heat units           | Larvae infested cotton             | 10.75               | 0.00  | .0810 ns                  | 0.1782         |
| Accumulated heat units           | Larvae infested okra               | 10.84               | 0.00  | .3166 ns                  | 0.0626         |
| Accumulated heat units           | Accumulated moth                   | -154.79             | 0.38  | .0000 ***                 | 0.8478         |
| Accumulated heat units           | Accumulated larvae infested cotton | -34.69              | 0.05  | .0000 ***                 | 0.9076         |
| Accumulated heat units           | Accumulated larvae infested okra   | -53.48              | 0.07  | .0000 ***                 | 0.9513         |
| Mean number of moth              | Larvae infested cotton             | 2.18                | 0.08  | .0003 ***                 | 0.5614         |
| Mean number of moth              | Larvae infested okra               | 4.73                | 0.07  | .0052 **                  | 0.3953         |
| Accumulated moth                 | Accumulated larvae infested cotton | -10.04              | 0.13  | .0000 ***                 | 0.9779         |
| Accumulated moth                 | Accumulated larvae infested okra   | -15.95              | 0.18  | .0000 ***                 | 0.9518         |
| Larvae infested okra             | Larvae infested cotton             | -1.28               | 0.89  | .0000 ***                 | 0.8470         |
| Accumulated larvae infested okra | Accumulated larvae infested cotton | 3.02                | 0.74  | .0000 ***                 | 0.9899         |

Generally, the larval infestation of *H. armigera* on okra was more than cotton plants during the whole two seasons and started earlier on okra than cotton with one week during the two seasons. The average number of larvae was decreased during July and August on cotton and okra during the two seasons, it may be due to climate changes and pesticides used to control cotton bollworm. The pesticide application could causes changes in the distribution pattern of the *H. armigera* (Balla and Singh, 1997).

Therefore, accumulated heat units can be used to forecast pheromone traps catch and larval infestation on cotton & okra. Pheromone traps catch can be used to forecast of *H. armigera* larval infestation on cotton & okra, also larval infestation on okra can be used to forecast larval infestation on cotton.

Khan *et al.* (2003) found positive correlation between infestation and weather factors (temperature and relative humidity). Nada *et al.* (2004) found that the population of *H. armigera* at Dakahlia had three and four peaks and the days from peak to peak ranged between 30-33 days, also, heat units between 440-486 units.

## REFERENCES

1. Balla, C. R. and S. P. Singh, 1997. Effect of pesticides applications on pheromone trap catches and distribution pattern of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) in pigeon pea ecosystem. Proceedings the First National Symposium on Pest Management in Horticulture Crops: Bangalore, India, 15-17 October, pp. 124-128.
2. DuraimuruJan, P. and A. Regupathy. 2005a. Mitigation of insecticide resistance in of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) by conjunctive use of trap crops, neem and *Trichogramma chilonis* Ishii in cotton. International J. of Zoological Research, 1(1): 53-58.
3. DuraimuruJan, P. and A. Regupathy. 2005b. Stimulo-deterrent diversionary strategy with conjunctive use of trap of insecticide resistant *Helicoverpa armigera* (Hubner) in cotton. J. Biological Sciences, 5(6): 681-686.
4. Fitt, G. P. 1991. Host selection in Heliiothina. In Reproductive Behaviour of Insects: Individuals and populations (eds W.J. Bailey and Ridsdill-Smith), pp. 172-201. Chapman and Hall, London, UK.
5. Fitt, G. P. 1989. The ecology of Heliiothis species in relation to agroecosystems. Ann. Rev. Entomol., 34:17-52.
6. Fitt, G. P. and G. S. Bayan. 1991. Methods for studying behaviour. In: *Helicoverpa Research Methods and Prospects* (ed. MP Zalucki), PP. 122-150. Springer Verlag, New Yourk, USA.

920 SEASONAL FLUCTUATION OF *HELICOVERPA ARMIGERA* (HUBNER) (LEPIDOPTERA: NOCTUIDAE) ON COTTON AND OKRA AND HEAT UNITS RELATED

7. Jallow, M. F. A., M. Matsumura and Y. Suzuki , 2001. Oviposition preference and reproductive performance of Japanese *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). *Applied Entomology and Zoology*, 36(4):419-429.
8. Gawand, G. V., B. S. Ghodki, H.U. Khambadkar and V. Farkade. 2006. Correlation studies on pheromone trap catches of *Helicoverpa armigera* with its field infestation. *Journal of Soils and Crops*, 16(1):250-253.
9. Khan, B. S., M. Afzal and M. A. Murtaza. 2003. Effect of abiotic factors against the infestation of American bollworm *Helicoverpa armigera* (Hubner) on different varieties of cotton under unsprayed conditions. *Journal of biological Sciences*, 3(1):82-85.
10. Nada, M. A., M. G. Ragab and E. S. Mansour. 2004. Seasonal abundance and prediction possibility of American bollworm, *Helicoverpa armigera* (Hub.) in relation to pheromone trap catches and heat unit accumulated. *Egypt. J. Appl. Sci*, 19 (11): 294-409.
11. Pretorius, L. M. 1976. Laboratory studies on the development and reproductive performance of *Heliothis armigera* (Hubn.) on various food plants. *J. Entomol. Soc. Sth. Afr.*, 39:337-343.
12. Ravi, K. C., K. S. Mohan, T. M. Manjumth, G. Head, B. V. Patil, D. P. A. Greba, K. Premalatha, J. Peter and N. G. V. Rao. 2005. Relative abundance of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) on different host crops in India and the role of these crops as natural refuge for *Bacillus thuringiensis* cotton. *Envi. Entomol.*, 34(1):59-69.
13. Seaver, D., J. Strand and A. J. Strawn. 1990. DDU degree-day utility users guide version 2.0, Univ. of California Division of Agriculture and Natural Resources UCIPM Publication 9, Integrated Pest Management Program, Univ. of California, Davis.
14. Sharma, H. C. 2001. Cotton bollworm/ legume pod borer, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) biology and management. International Institute for Semi-Arid Tropics, India.
15. Slosser, J.E., J.A. Witz, G. I. Putcrka, J. I. Price and A. N. Hartostock. 1987. Seasonal changes in bollworm (Lepidoptera: Noctuidae) moths catches in traps in a large area. *J. Environ. Entomol.*, 16(6): 1296-1301.

## التقلبات الموسمية لدودة اللوز الأمريكية في القطن والبامية وعلاقتها بالوحدات الحرارية

علي احمد السيد ، مرفت عبد البرقنديل ، عادل السيد عامر

معهد بحوث وقاية النباتات مركز البحوث الزراعية - الدقى - الجيزة

تم دراسة التقلبات الموسمية لفرشات ويرقات دودة اللوز الأمريكية في القطن و البامية في محافظة الدقهلية خلال موسمي ٢٠٠٧ و ٢٠٠٨. أوضحت النتائج أن لفرشات دودة اللوز الأمريكية المصادة بواسطة المصائد الفرمونية ثلاثة قمم خلال كل موسم وكانت القمة الأولى أعلاها في التعداد. وكان لإعداد يرقات دودة اللوز الأمريكية علي القطن و البامية قمتين خلال الموسم وكانت القمة الأولى هي الأعلى في التعداد وكان تعداد اليرقات علي البامية اعلي من أعدادها الموجودة علي القطن خلال الموسمين. ووجد ان هناك علاقة ارتباط معنوية بين الوحدات الحرارية المتجمعة وتعداد اليرقات في المصائد وعدد اليرقات في القطن و البامية وتراوحت قيمة  $R^2$  بين ٠,٨١٤٨ و ٠,٩٦١٩. ووجد أن هناك علاقة مابين التعداد المتجمع لفرشات دودة اللوز الأمريكية وتعداد اليرقات في القطن و البامية وكانت قيمة  $R^2$  ٠,٩١٨٥ و ٠,٩٧٩٣ وكذلك وجدت علاقة عالية المعنوية مابين تعداد اليرقات في البامية و القطن وكانت قيمة  $R^2$  ٠,٦٠٣٠ و ٠,٩٨٩٩. وبصفة عامه يمكن استخدام الوحدات الحرارية المتجمعة للتنبؤ بتعداد اليرقات في المصائد الفرمونية و الإصابة في القطن و البامية. كما يمكن استخدام تعداد اليرقات في القطن و البامية وكذلك استخدام تعداد اليرقات في البامية للتنبؤ بتعداد يرقات دودة اللوز الأمريكية في القطن.