

EFFECT OF SITOFEX (CPPU) ON FRUIT SET, FRUIT QUALITY OF ANNA APPLE TREES

Fatma I. I. Abou Grah, Abd El-Megeed Nagwa and El-Shereif H.

Hort. Res. Institute, Agric. Res. Center Giza, Egypt .

ABSTRACT

Efficiency of the synthetic cytokinin (CPPU) (N- (2-chloro-4-pyridyle-N-phenylurea) was studied on Anna apple trees at full bloom, 14 days after full bloom and two time spraying (at full bloom and 14 days after full bloom). CPPU tended significantly to increase the percentage of fruit set, yield and fruiting, while it decreased fruit drop of all treatments. The best results were obtained by 20 ppm CPPU at full bloom and 10 ppm CPPU two time spraying in the first season. While, the 15 ppm CPPU at full bloom was the best during the second season.

Highest fruit weight, fruit size and fruit dimensions were obtained by 15 ppm concentrations at 14 days after full bloom in the two seasons. Concerning fruit shape index (L/D), an increase in fruit diameter than its length due to all conducted CPPU treatments in both seasons. As well as, acidity and TSS/acid ratio were significantly improved as a result of all studied treatments when compared to the control.

It could be concluded that, most of studied treatments resulted in a positive and significant effect on most studied characteristics, since (CPPU at 15 ppm at full bloom and 20 ppm at 14 days after full bloom were the most effective treatments for increasing fruit set and yield as well as improving the most fruit properties.

Key words: Cytokinin (CPPU), Anna apple trees, Fruit set and Fruit drop

INTRODUCTION

Apple is considered one of the major and the most important deciduous fruit trees in Egypt. Many investigators reported that, yield and quality of Anna apple fruits depended upon several factors, one of the most vital factor which affects and plays an important role in this concern is spraying with some growth regulators which enhance fruit set, reduce fruit drop, consequently increase productivity. Moreover, both concentration and date of application are very important factors which in true reflect in increasing and improving fruit yield and fruit characteristics.

Several investigations mentioned that, spraying deciduous fruit trees with Sitofex (CPPU) different concentrations enhanced cell division, increased cell size, increased fruit weight, size and fruit yield. Furthermore, application of the abovementioned growth regulators improved the most fruit properties. Nickell (1986), Rizk (1998), Feng *et al.*, (1999), Al-Ashkar (2000) Ranpise *et al.*, (2000) and Marwad (2001) on grapes; El-Barkooky (1985), Greene (1989) and Khurshid *et al.*, (1997), on apple; Biasl *et al.*, (1991) and Lowes and Woolley (1992) on kiwi; Jindal and Sharma (1986) on plum; Kabeel (1999) on persimmon; Kabeel and Fawaaz (2005) on pear.

Due to the little information currently available about the effect of CPPU on apple fruit, this study was carried out to explore the effect of concentration and application time of CPPU on apple, fruit set, drop and quality.

MATERIAL AND METHODS

The current investigation was under taken in the experimental farm at El-Kanater, Horticultural Research Station, Kalyubia, Governorate, Egypt. This study has been extended for the two consecutive seasons of 2007 and 2008 on 10- year-old-apple trees. Anna apple trees were budded on Malling Maritton 106 rootstock, planted at 2.5 meters apart and grown in clay loamy soils. Selected trees were healthy, nearly uniform as possible in their vigour and subjected to the similar fertilization, irrigation, pruning and pest control programs usually done at this region.

Different foliar sprays with Sitofex (CPPU) treatments used in this study were as follows:

- 1- CPPU at 5 ppm.
- 2- CPPU at 10 ppm
- 3- CPPU at 15 ppm
- 4- CPPU at 20 ppm
- 5- Control

These treatments were sprayed at: a) Full bloom stage, b) At full bloom and two weeks after full bloom and c) Two weeks after full bloom.

Forty five trees were devoted and complete randomized design was used, since each treatment was replicated by a two trees. Four main branches well distributed around the periphery of tree (one branch on each direction) were tagged and the following measurements were determined:

1- Fruiting measurements:**1-a. Percentages of fruit set and fruit drop:**

Both number of flowers and set fruitlets on the tagged branches were counted and recorded for all treatments, then percentage of fruit set was calculated by the following equation according to **Westwood (1978)**

$$(\%) \text{ Fruit set} = \frac{\text{Number of set fruitlets}}{\text{No. of opened flowers}} \times 100$$

Furthermore, number of dropped fruits were recorded till harvest time, then estimated as percentage on the basis of initial number of fruitlets according to this equation:

$$(\%) \text{ Fruit drop} = \frac{\text{Number of dropped fruits}}{\text{Number of set fruitlets}} \times 100$$

1-b. Yield and percentage of yield increment than control:

The average of tree yield in kgs for each treatment was determined at harvest time (at maturity stage). Furthermore, the yield increment percentage for each treatment as compared to the control was estimated according to the following equation:

$$(\%) \text{ Yield incr.} = \frac{\text{Yield / treatment} - \text{yield / control}}{\text{Yield / control}} \times 100$$

At picking date, number of fruits/tree were used to calculate yield monetary value = Fruit yield (kg)/tree x farm – gate price (L.E.1.5).

2- Fruit quality:

At the time of harvest (at maturity stage), ten fruits from each replicate were randomly sampled and the following fruit characteristics were determined including average fruit weight (gm.), fruit firmness (lb/inch²) using a **Magness and Tayler** pressure tester with 7/18 inch plunger. Furthermore, fruit chemical

properties were also determined including the average fruit juice TSS percentage using handy refractometer, fruit juice acidity percentage as malic acid (mgs/100 gms fruit juice) according to **A.O.A.C. (1985)** and **Vogel (1968)**, TSS/acid ratio was calculated.

All the obtained data were statistically analyzed of variance method according to **Snedecor and Cochran (1990)** using L.S.D. values at 0.5 % level. However, means were compared according to Duncan's multiple range test (**Duncan, 1955**).

RESULTS AND DISCUSSION

1. Fruiting measurements:

1-a. Percentages of fruit set and fruit drop:

Data in Table (1) displayed clearly that, all treatment sprays resulted in a significant increase in fruit set % as compared to the control. Moreover, trees sprayed with 20 ppm concentration of CPPU were statistically the superior as exhibited significantly the highest value fruit set (18.59 %). Meanwhile, the opposite trend was observed with the control which was statistically the inferior as exhibited the least value of fruit set (11.58 %). On the other hand, the best time to treat spraying was at full bloom and after two weeks (15.72 % and 20.71) this result was detected during both 2007 and 2008 seasons. In additions, the best interaction in this respect was obtained by 20 ppm concentration of CPPU with the two time applications during 2007 and 2008 seasons, respectively (22.5 and 26.87 %).

With regard to the percentage of fruit drop, data in the same Table showed obvious trend where all treatment concentrations under study decreased significantly percentage of fruit drop as compared to the control in the two experimental seasons. Data pointed out that, the highest percentage of fruit drop was always concomitant to the control (85.5 and 80.83 %) whereas either CPPU at 5 ppm in the first season (65.22 %) and CPPU at 10 ppm in the second season (68.53 %) were the most effective treatments regarding reducing fruit drop. Since they resulted in statistically the lowest values in this concern.

On the other hand, the application time after two weeks of full bloom had the least value of fruit drop (70.19 and 68.15 %) during the two seasons. Whereas either CPPU at 20 ppm or at 10 ppm application after two weeks of full bloom or the two application (at full bloom and after two weeks) were the most effective treatments regarding reducing of fruit drop. The obtained results are in conformity with those previously reported by **Nickell (1986)** and **Feng et al., (1999)** on grapes, **El-Barkouky (1985)** and **Khurshid et al., (1997)** on apple; **Kabeel (1999)** on persimmon, **Guirguis et al., (2003)** and **Kabeel and Fawaaz (2005)** on pear trees.

Table (1): Effect of Sitofex spray at different concentrations (A) and at different dates (B) on percentage of fruit set and fruit drop.

2007 season								
Date	Fruit set (%)				Fruit drop (%)			
Treat	A*	B	C	Ave.(A)	A	B	C	Ave. (A)
5 ppm	13.33d-f	13.37ef	11.83ef	12.84D	54.50f	67.83de	73.33cd	65.22CD
10 ppm	16.67c	15.07c-e	13.00d-f	14.91C	74.23cd	76.27bc	66.70e	72.48B
15 ppm	12.87d-f	16.93bc	20.00b	16.60B	78.23bc	79.23bc	68.90de	75.46B
20 ppm	17.20bc	22.50a	16.07cd	18.59A	81.63ab	73.67cd	55.03f	70.11C
Control	12.67ef	10.73f	11.33f	11.58E	82.23ab	87.33a	86.97a	85.50A
Ave. (B)	14.55B	15.72A	14.45B		74.17B	76.87A	70.19C	
2008 season								
5 ppm	20.70b	22.43b	15.87c	19.67A	77.20bc	71.83cd	72.33cd	73.79B
10 ppm	14.57ed	21.00b	10.63e	15.40C	76.20c	63.80e	65.60e	68.53C
15 ppm	15.13ed	21.70b	20.33b	19.05B	77.33bc	72.67cd	67.83de	72.61B
20 ppm	20.50b	26.87a	15.17cd	20.84A	71.60cd	82.53ab	62.33e	72.15B
Control	12.60de	11.53e	10.53e	11.55D	84.83a	85.00a	72.67cd	80.83A
Ave. (B)	16.70B	20.71A	14.51C		77.43A	75.17A	68.15B	

L.S.D. at 5 %:

A =	1.315	1.125	2.600	2.520
B =	1.697	1.453	3.357	3.258
A x B =	2.94	2.517	5.814	5.649

* A: Spray at full bloom.

B: Spray at full bloom and two weeks after full bloom.

C: Spray at two weeks after full bloom.

1-b. Yield and yield increment % in relation to the control:

Data tabulated in Tables (2 & 3) clear that, both number of fruits/tree, yield/tree, yield/feddan and yield increment percentage were responded significantly to all used concentrations as compared to the control during the two studied seasons. Furthermore, the greatest statistically values of yield parameters were resulted from Anna apple trees being sprayed with CPPU at 20 ppm (450 fruits/tree, 59.27 kg fruits/tree, 15.53 ton fruits/feddan and 2.7% increment) followed by 15 ppm, 10 ppm and 5 ppm treatments as compared to the control, which reflected significantly the lowest value of yield parameters (170.6 fruits/tree, 16.01 kg/fruits/tree, 4.29 ton fruits/feddan and 0.0% increment). On the other hand, the best results in this respect were obtained by two application time (at full bloom and after two weeks) during the two seasons. While, the lowest value were obtained by sprayed with CPPU after two weeks from full bloom. Moreover, the interaction 20 ppm CPPU sprayed at full bloom and after two weeks was better than the other treatments.

These results are completely agreed with those being mentioned by many investigators Greene (1989) on apple, Kabeel (1999) Kabeel and Fawaaz (2005) on pear, Feng *et al.*, (1999), Al-Ashkar (2000), Marwad (2001) on grapes.

Fayoum J. Agric. Res. & Dev., Vol.23, No.1,(B). January, 2009

Table (2): Effect of Sitofex spray at different concentrations (A) and at different dates (B) on number of fruit/tree and fruit yield/tree.

2007 season												
Date Treat	Numbr of fruits/tree				Fruit yield/tree				(L.E.)			
	A*	B	C	Ave. (A)	A	B	C	Ave. (A)	A	B	C	Ave. (A)
5 ppm	176.7g	200.0f	200.0f	192.2D	20.13d-g	22.19d	21.65d-f	21.33C	30.2gh	33.3gh	32.5gh	32.0D
10 ppm	210.0ef	223.3e	200.0f	211.1C	24.33d	25.10d	21.73d-f	23.72C	36.5fg	37.7f	32.6gh	35.6C
15 ppm	300.0d	320.0c	290.0d	303.3B	36.44c	39.37c	38.37c	38.06B	54.7e	59.1d	57.6de	57.1B
20 ppm	466.7a	483.3a	400.0b	450.0A	58.69b	63.57a	55.64b	59.27A	87.9b	95.4a	83.5c	88.9A
Control	173.3g	173.3g	165.0g	170.6E	15.45g	16.08g	16.50e-g	16.01D	23.2i	24.1i	24.8i	24.0E
Ave. (B)	265.30B	280.0A	251.0C		30.99AB	33.26A	30.78B		46.5B	49.9A	46.2B	
2008 season												
5 ppm	213.3e	266.7d	186.7f	222.2D	20.35fg	26.10e	20.06fg	22.17D	30.5fg	39.2e	30.1fg	33.3D
10 ppm	216.7e	303.3c	206.0e	242.0C	25.09e	33.47d	22.03ef	26.87C	37.6e	50.2d	33.0f	40.3C
15 ppm	305.0c	423.3a	303.3c	343.9B	37.31cd	52.28a	39.76c	43.12B	56.0c	78.4a	59.6c	64.7B
20 ppm	376.7b	433.3a	320.0c	376.7A	46.60b	54.91a	44.19b	48.57A	69.9b	82.4a	66.3b	72.9A
Control	176.7f	180.0f	170.0f	175.6E	16.55g	18.21fg	16.83g	17.19E	24.8h	27.3gh	25.2h	25.3E
Ave. (B)	257.7B	321.3A	237.2C		29.18B	36.99A	28.57B		43.8B	55.5A	42.8B	

L.S.D. at 5 %:

A =	8.131	7.191	2.324	1.735	1.970	1.798
B =	10.500	9.283	3.000	2.240	2.543	2.321
A x B =	18.180	16.080	5.196	3.879	4.075	4.021

* A: Spray at full bloom.

B: Spray at full bloom and two weeks after full bloom.

C: Spray at two weeks after full bloom.

It is also noticeable that, yield monetary value (Table 2) was parallel to the increase of CPPU concentration from 0.0 ppm (LE 24.0), to 5 ppm (LE 32.0) to 10 ppm (LE 35.6) to 15 ppm (LE 37.1) and to 20 ppm (LE 88.9). this trend was clear throughout the two studied seasons and statistically confirmed. We can also say that CPPU sprays have better return when sprayed at full bloom and next after two weeks (LE 49.9 and 55.5) than the other treatments. However, the best interaction was 20 ppm at full bloom and after two weeks (LE 95.4 and 82.4).

Table (3): Effect of Sitofex spray at different concentrations (A) and at different dates (B) on percentage of yield increment than control and fruit yield/feddan.

2007 season								
Date	Yield increment than control				Yield/feddan (ton)			
Treat	A*	B	C	Ave. (A)	A	B	C	Ave. (A)
5 ppm	0.317de	0.400de	0.307de	0.341C	5.083h	5.817f-h	5.670gh	5.52D
10 ppm	0.583d	0.613d	0.313de	0.503C	6.373fg	6.573f	5.690gh	6.21C
15 ppm	1.363c	1.487c	1.333c	1.394B	9.543e	10.310d	10.050de	9.97B
20 ppm	2.777ab	3.033a	2.370b	2.727A	15.350b	16.650a	14.570c	15.53A
Control	0.000e	0.000e	0.000e	0.00D	4.223i	4.327i	4.323i	4.29E
Ave. (B)	1.008AB	1.107A	0.865B		8.11B	8.74A	8.06B	
2008 season								
5 ppm	0.237de	0.423d	0.197de	0.286C	5.333fg	6.830e	5.250fg	5.81D
10 ppm	0.483d	0.533d	0.310de	0.442C	6.573e	8.760d	5.770ef	7.03C
15 ppm	1.317c	1.907ab	1.363c	1.529B	9.770cd	13.700a	10.410c	11.29B
20 ppm	1.917ab	2.017a	1.636bc	1.854A	12.210b	14.340a	11.580b	12.71A
Control	0.000e	0.000e	0.000e	0.00D	4.330g	4.760fg	4.400g	4.50E
Ave. (B)	0.791B	0.976A	0.700B		7.64B	9.68A	7.48B	

L.S.D. at 5 %:

A =	0.209	0.148	0.324	0.497
B =	0.270	0.191	0.419	0.642
A x B =	0.467	0.330	0.725	1.112

* A: Spray at full bloom.

B: Spray at full bloom and two weeks after full bloom.

C: Spray at two weeks after full bloom.

2. Fruit characteristics:

2-1. Fruit physical characteristics:

2-1-a. Fruit weight, size and firmness:

As shown in Table (4), fruit weight was increased by CPPU applications after two weeks of full bloom (117.4 and 116.4 g) as compared with the other treatments (113.2, 113.3, 109.6 and 114.0 g.) through 2007 and 2008 seasons, respectively.

It is also noticeable that, fruit weight gradually and significantly increased with increasing CPPU concentration from 5 ppm (104.0g.) to 10 ppm (110.2g.) to 15 ppm (125.3g.) to 20 ppm (129.4g.) comparing to control (97.8g.).

It is clear from the same Table that, foliar applications of the deferent treatments of CPPU, resulted in significant increases in fruit size. The highest value was obtained by CPPU at 20 ppm (138.1 cm³) followed by 15 (125.0 cm³), 10 (123.3 cm³) 5 ppm (105.2 cm³) compared to the control (86.0 cm³). The present data are in accordance with those mentioned by **El-Barkouky (1985); Jindal and Sharma (1986); Biasl et al., (1991); Lowes and Woolley (1992); Rizk (1998); Kabeel (1999) and Guirguis et al., (2003)** on some fruit deciduous trees.

Data of fruit flesh firmness in the two seasons as shown in the same Table obviously indicate that, it was significantly increased by increasing CPPU concentrations. The results also indicated that, all the after full bloom application gave higher values than the other treatments and showed a positive relation as the values increased by increasing CPPU concentrations. These results are in agreement with the fact that, at maturity, firmer fruits easily tolerate post harvest treatments. Moreover, previous reports of **Khurshid et al., (1997); Kabeel (1999) and Guirguis et al., (2003)** on apple, persimmon and pear trees have supported this trend.

2-1-b. Fruit dimensions:

It is clear in Table (5) that, fruit length and diameter gradually increased as CPPU concentration increased from 0.0 ppm (4.76 and 4.90 cm.) to 5 ppm (4.86 and 5.06 cm.) to 10 ppm (5.5 and 5.22 cm.) to 15 ppm (5.87 and 5.60 cm.) to 20 ppm (5.97 and 5.81 cm.).

Meanwhile, CPPU spray was more effective when sprayed at full bloom and after two weeks (5.57 and 5.46 cm.) than the other treatments. Moreover, 20 ppm CPPU spray at full bloom and after two weeks show better interaction in this respect.

The results of the two seasons indicated that, all the CPPU treated fruits resulted in an increase in length than in diameter as all obtained shape index values were less than the control and the increase in length values has positively linked with a parallel increase in CPPU concentration. Obtained results concerning the response of fruit dimensions and fruit shape index were generally supported by finding of **Nickell (1986), Biasl et al., (1991), Lowes and Woolley (1992) Kabeel (1999) and Guirguis et al., (2003)** on kiwi, grapes and pear fruits.

2-2. Fruit chemical properties:

2-2-a. Fruit juice TSS %:

Regarding the response of fruit juice TSS % of tested treatments, data in Table (6), indicates that TSS responded significantly to the most of treatments. Moreover, the richest fruits in their content of TSS % was achieved by trees sprayed with 5 ppm CPPU treatments (13.39 and 11.78%). Meanwhile, juice TSS was better when CPPU sprayed after two weeks from full bloom. Meanwhile, the lowest significant values of fruit juice TSS was the interaction 10, 15 ppm CPPU applied twice (at full bloom and after two weeks).

On the other hand, total soluble solids (T.S.S.) percentage results of the first season showed that, the most CPPU treatments increased values than the control either applied after two weeks from full bloom are applied twice (at full bloom and after two weeks).

Table (4): Effect of Sitofex spray at different concentrations (A) and at different dates (B) on fruit weight, size and firmness.

2007 season												
Date	Fruit weight (g.)				Fruit size (cm ³)				Fruit firmness (lb/inch ²)			
Treat	A*	B	C	Ave. (A)	A	B	C	Ave. (A)	A	B	C	Ave. (A)
5 ppm	110.0d	110.9d	107.9d	109.6C	102.3ef	110.0cd	103.3d	105.2C	10.27e	13.67cd	10.67e	11.53B
10 ppm	114.8d	111.1d	108.0d	111.3C	141.7a	113.3c	115.0c	123.3B	10.73e	9.73e	14.33b-d	11.60B
15 ppm	122.9c	123.0c	132.2b	126.0B	110.0cd	140.0a	125.0b	125.0B	10.00e	10.17e	16.27ab	12.14B
20 ppm	125.0bc	127.2bc	139.1a	130.5A	140.7a	131.7b	145.0a	138.1A	12.03de	13.33cd	12.00de	12.46B
Control	93.2e	94.4e	100.0e	95.86D	80.7g	81.7g	98.3f	86.9D	12.00de	14.67bc	18.00a	14.89A
Ave. (B)	113.2A	113.3B	117.4A		114.9A	115.33A	115.7A		11.01C	12.31B	14.25A	
2008 season												
5 ppm	95.3fg	109.8d	106.9de	104.0D	109.0cd	110.0c	102.3de	107.1C	9.97d	13.37c	10.37d	11.23B
10 ppm	113.7d	110.0d	106.9de	110.2C	140.7a	112.3c	114.0c	122.3B	10.43d	9.43d	14.03bc	11.30B
15 ppm	121.8c	122.9c	131.1b	125.3B	101.3e	139.0a	124.0b	121.4B	9.70d	9.97d	15.97b	11.88B
20 ppm	121.9c	126.16bc	138.0a	129.4A	139.7a	130.7b	144.0a	138.1A	11.73cd	13.03c	11.70cd	12.16B
Control	93.3g	101.3ef	99.0fg	97.86E	80.0f	93.3e	81.0f	86.11D	13.00c	15.67b	19.00a	15.89A
Ave. (B)	109.6B	114.0A	116.4A		114.14B	117.87A	113.06B		10.97C	12.29B	14.21A	

L.S.D. at 5 %:

A =	3.074	2.943	3.285	3.043	0.959	0.965
B =	3.969	3.800	4.240	3.928	1.238	1.245
A x B =	6.875	6.581	7.345	6.804	2.144	2.157

Table (5): Effect of Sitofex spray at different concentrations (A) and at different dates (B) on fruit length, diameter and shape index.

2007 season												
Date	Fruit length (cm.)				Fruit diameter (cm.)				Fruit shape index			
	A*	B	C	Ave. (A)	A	B	C	Ave. (A)	A	B	C	Ave. (A)
Treat												
5 ppm	4.93fg	5.067e-g	4.57g	4.86C	4.93e-g	5.43b-d	4.80fg	5.06CD	1.02a-c	0.94bc	0.95bc	0.97A
10 ppm	5.23d-f	5.80b-d	5.47b-f	5.50B	5.07d-g	5.37b-d	5.23c-e	5.22C	1.04a-c	1.08ab	1.04a-c	1.06A
15 ppm	5.93a-c	5.97ab	5.70b-e	5.87A	5.50a-c	5.53a-c	5.73ab	5.60B	1.08ab	1.08ab	0.99a-c	1.05A
20 ppm	5.30c-f	6.07ab	6.53a	5.97A	5.90a	5.80ab	5.73ab	5.81A	0.90c	1.06a-c	1.14a	1.03A
Control	4.33fg	4.97fg	4.47g	4.76C	4.83e-g	5.17c-f	4.70g	4.90D	1.02a-c	0.96bc	0.95bc	0.98A
Ave. (B)	5.25B	5.57A	5.35B		5.25B	5.46A	5.24B		1.01A	1.02A	1.02A	
2008 season												
5 ppm	5.13fg	5.37d-g	5.27e-g	5.26C	5.30b-d	5.27b-d	5.23b-d	5.27CD	0.97e	1.03a-e	1.01b-e	1.00B
10 ppm	5.70b-e	5.57c-f	5.77b-e	5.68B	5.50b-d	5.57a-d	5.36b-d	5.46BC	1.04a-e	1.00b-e	1.11a	1.05A
15 ppm	5.97a-c	5.80a-e	5.83a-d	5.87AB	5.73ab	5.83a	5.53a-d	5.70AB	1.04a-e	0.99c-e	1.06a-d	1.03AB
20 ppm	5.97a-c	6.33a	6.13ab	6.14A	5.63a-d	5.90a	5.67a-c	5.73A	1.06a-d	1.07a-c	1.08ab	1.07A
Control	5.03fg	5.27e-g	5.00g	5.10C	5.20b-d	5.17cd	5.10d	5.16D	0.97e	1.03a-e	0.98de	0.99B
Ave. (B)	5.56A	5.67A	5.60A		5.47A	5.55A	5.37A		1.02A	1.02A	1.05A	

L.S.D. at 5 %:

A =	0.260	0.219	0.172	0.204	0.067	0.033
B =	0.336	0.283	0.222	0.263	0.086	0.042
A x B =	0.582	0.491	0.385	0.455	0.150	0.075

Table (6): Effect of Sitofex spray at different concentrations (A) and at different dates (B) on fruit juice TSS, acidity and TSS/acidity ratio.

2007 season												
Date Treat	TSS (%)				Acidity (%)				TSS/acidity ratio			
	A*	B	C	Ave. (A)	A	B	C	Ave. (A)	A	B	C	Ave. (A)
5 ppm	12.00bc	12.17b	13.00a	12.39A	0.359bc	0.390a-c	0.333c	0.361B	31.19cd	31.19cd	41.99a	35.89A
10 ppm	11.50c-e	10.83fg	12.33b	11.56BC	0.307c	0.472a	0.472a	0.416A	23.58b	23.90f	28.64de	30.04B
15 ppm	11.50c-e	10.83fg	11.17e-g	11.17D	0.465a	0.478a	0.478a	0.462A	24.84ef	24.65f	23.65f	24.38C
20 ppm	11.83b-d	11.50c-e	11.83b-d	11.72B	0.333c	0.468a	0.468a	0.420A	36.18b	25.00ef	25.44ef	28.87B
Control	12.00bc	10.67g	11.33d-f	11.33CD	0.466a	0.479a	0.479a	0.471A	25.84ef	23.46f	23.98f	24.43C
Ave. (B)	11.77A	11.20B	11.93A		0.386B	0.446A	0.446A		31.79A	25.64C	28.74B	
2008 season												
5 ppm	11.50bc	11.83ab	12.00a	11.78A	0.349bc	0.380a-c	0.323c	0.351C	34.12cd	31.14de	40.15a	35.14A
10 ppm	11.33cd	11.00d	12.17a	11.50B	0.297c	0.458a	0.462a	0.406A	38.30ab	24.76fg	28.91ef	30.66B
15 ppm	11.33cd	11.00d	11.33cd	11.22C	0.455a	0.432ab	0.468a	0.452A	25.06fg	25.58fg	24.17g	24.94C
20 ppm	11.33cd	11.83ab	12.00a	11.72AB	0.323c	0.450ab	0.458a	0.410A	35.76bc	26.31fg	26.93fg	29.67B
Control	11.33cd	11.50bc	12.00a	11.61AB	0.455a	0.459a	0.469a	0.461A	25.06fg	25.61fg	25.89fg	25.52C
Ave. (B)	11.37B	11.43B	11.90A		0.376B	0.436A	0.436A		31.66A	26.68C	29.21B	

L.S.D. at 5 %:

A =	0.232	0.175	0.071	0.042	1.535	1.679
B =	0.299	0.227	0.053	0.054	1.982	2.168
A x B =	0.518	0.392	0.092	0.093	3.433	3.754

* A: Spray at full bloom.

B: Spray at full bloom and two weeks after full bloom.

C: Spray at two weeks after full bloom.

2-2-b. Fruit juice total acidity %:

Obtained results in Table (6) show clearly that, CPPU treatments decreased values than the control specially 5 ppm (0.361 and 0.351 % during the two studied seasons, respectively. Moreover, CPPU spray at full bloom resulted in lower acidity (0.386 and 0.376 % in 2007 and 2008 seasons, respectively) than the other treatments. However, 10 ppm CPPU spray at full bloom consider better interaction in this respect (0.307 and 0.297 %, respectively). Meanwhile, TSS/acidity ratio did not show clear trend.

CONCLUSION

The present results of 5, 10, 15 and 20 ppm CPPU sprays at: A) full bloom, B) full bloom and after two weeks, and C) after two weeks of full bloom showed a positive effect than control. However, 20 ppm concentration was superior where it increased percentage of fruit set, number of fruits/tree, fruit yield per tree and per feddan, yield increment than control and subsequently yield monetary value. Also 20 ppm has a benefit effect (than the other treatments) on fruit characteristics (fruit weight, size, dimensions, fruit shape index and juice TSS). Moreover, CPPU spray at full bloom and after two weeks has better effect on the former fruit and yield attributes. So, we can recommended apple (Anna cv.) growers to spray 20 ppm CPPU at full bloom and next after two weeks to increase the yield, fruit quality and yield monetary value.

REFERENCES

- Al-Ashkar, R.A. (2000).** Effect on some GA₃ and CPPU treatments on yield and fruit quality of Ruby seedless and Flame seedless grapes. *Egy. J. Appl. Sci.*, 15 (7): 491-510.
- Association of Official Agriculture Chemists A.O.A.C. (1985).** "Official Methods of Analysis" Benjamin Franklin Station, Washington D.C. U.S.A., 495-510.
- Biasl, R.; G. Costa; R. Glulioni; F. Sacci and S. Sansavini (1991).** Effects of CPPU on kiwi fruit performance. *Acta. Hort.*, 297, 367-373.
- Duncan, D.B. (1955):** Multiple range and multiple F. tests. *Biometrics*, 11: 1-42.
- El-Barkouky, F.M.Z. (1985).** Effect of some growth regulators on flowering fruit setting and fruit quality of apple. Ph. D. Thesis Fac. of Agric. Ain Shams Univ.
- Feng, X.H.; H.F. Song; Y.M. Qian and G.J. Liu (1999).** Effect of treatment with CPPU and GA₃ after flowering on the production and berry quality of kyoga grapes. *South china fruits*, 28: 2, 41. (C.F. comp. Research).
- Greene, D.W. (1989).** CPPU influences "Mcintosh" apple crop load and fruit characteristics. *Hort. Scienc*, 24, 94-96.
- Guirguis, N, S.; E. S. Attala and M.M. Ali (2003).** Effect of sitofex (CPPU) on fruit set, fruit quality of "Le-Conte" pear cultivar. *Annals of Agric. Sci. Moshtohor*, 41 (1): 271-282.
- Jindal, K.K. and N.S. Sharma (1986).** Effect of some growth retardants in combination with nutrients on fruit maturity and quality of Japanese plum (*Prunus salicina lindl*) cv. Santa Rosa. *Advances in research on Temperate fruits proceedings of the National Symposium on Temperate Fruits 15-18 March (1984).* Hemichall Pradesh Agricultural University, Sloan, India, 281-285, 14 ref.
- Kabeel, H. (1999).** Effect of some growth regulators on fruit set, yield and fruit quality of "Costata" persimmon trees, *Minufiya Jour. Agric. Res.*, 24 (5): 1727-1739.

- Kabeel, H. and S.A.A. Fawaaz (2005).** Effect of spraying some growth regulators on "Le-Conte" pear trees on I- productivity, fruit quality and leaf mineral content. Hort. Res. Inst, Agric. Res. Cen., Giza, Egypt., 3 (3): 173-193.
- Khurished, T.; D.J.Mc. Neiland and M.C. Trought (1997).** Effect of foliar applied gibberellins and soil applied paclobutrazol on reproductive and vegetative growth of "Braebun" apple trees growing under a high density planting system. New Zealand Jour. Of Crop and Hort. Sci., Lincoln Univ. (C.F. Hort. Abst. 67-6637).
- Lowes, G. S. and Woolley, D. J. (1992).** A new way to grow bigger kiwi fruit. Department of plant science, Marsey University, April. The Orchardist pp. 35-37.
- Marwad, I.A. (2001).** Effect of some sitofex (CPPU) and gibberellin (GA₃) treatments on yield fruit quality of Thompson seedless grapes. Egy. J. Appl. Sci.; 16 (10): 210-232.
- NickeH, L.G. (1986).** Effects of N (2-chloro-4-pyridyl- (N-phenylurea) on grapes and other crops. Proc. of Plant Growth Regulator Society of America. 13: 236-241.
- Ranpise, S.A.; B.T. Pat. L and T.K. Ghure (2000).** Effect of 4-chloro fernuron (CPPU) and gibberellic acid on physiochemical properties of Thompson seedless grapes. Journal of Maharashtra, India, No. 526: 293-299.
- Rizk, M.H. (1998).** Effect of sitofex (CPPU), GA₃ and hand thinning yield and fruit quality of Thompson seedless grapes. J. Agric. Sci. Mansoura Univ., Egypt. 23 (1): 397-404.
- Snedecor, G.W. and W.G. Cochran (1990):** "Statistical Methods". 7th ed., Iowa State Univ. Press Amer. Iowa, USA pp.85-86.
- Vogel, A. (1968):** A Text Book of Quantitative Inorganic Analysis. Longmans, New York, pp. 1216.
- Westwood, M.N. (1978):** Temperate Zone Pomology W. H. Freeman and Company. San Francisco.

تأثير السيتوفكس CPPU على عقد ومحصول وجودة ثمار التفاح صنف آنا

فاطمة إبراهيم إبراهيم أبو جرة - نجوى عبد المجيد - هناء الشريف
معهد بحوث البساتين - مركز البحوث الزراعية

تم دراسة تأثير رش مادة السيتوفكس (CPPU) على أشجار التفاح صنف آنا في ثلاث مواعيد: (١) عند تمام التزهير. (٢) عند تمام التزهير وبعد ١٤ يوم من تمام التزهير، (٣) بعد ١٤ يوم من تمام التزهير. وذلك خلال الموسمين ٢٠٠٧، ٢٠٠٨. وأوضحت الدراسة أن رش اشجار التفاح بمركب السيتوفكس أدى إلى زيادة نسبة العقد والإثمار وتقليل النسبة المئوية لتساقط الثمار وزيادة المحصول والعائد النقدي مقارنة بالكنترول وكانت أفضل النتائج عند استخدام التركيز ٢٠ جزء في المليون عند الرش عند تمام التزهير و ١٥ جزء في المليون عند الرش بعد ١٤ يوم من تمام التزهير وذلك خلال الموسم الأول بينما كانت أفضل النتائج في الموسم الثاني عند الرش بتركيز ١٥ جزء في المليون عند تمام التزهير.

كما أوضحت الدراسة إلى أن كل من الصفات الطبيعية للثمار مثل (وزن - حجم - أبعاد - معامل الشكل - الصلابة) وكذلك الصفات الكيماوية للثمار مثل (النسبة المئوية للمواد الصلبة الذائبة الكلية - نسبة الحموضة - النسبة بين المواد الصلبة الذائبة / الحموضة) قد تحسنت معنويًا نتيجة رش معظم تلك المعاملات تحت الدراسة بمادة السيتوفكس مقارنة بالأشجار الغير معاملة (الكنترول) في كل من موسمي الدراسة.

ونتيجة لذلك فإنه يمكن الإشارة إلى أن رش أشجار التفاح بالسيتوفكس قد أدت إلى تأثير إيجابي ومعنوي لمعظم الصفات المدروسة وكانت المعاملة (السيتوفكس ٢٠ جزء في المليون عند الرش عند تمام التزهير)، السيتوفكس ١٥ جزء في المليون عند الرش بعد ١٤ يوم من تمام التزهير هي أفضل المعاملات فاعلية في زيادة عقد الثمار والمحصول وكذلك تحسين معظم صفات الثمار الطبيعية والكيماوية.