

ALLEVIATE THE HARMFUL EFFECTS OF SALINE RECLAIMED SOIL CONDITIONS ON GROWTH AND PRODUCTIVITY OF ONION PLANTS USING ASCORBIC ACID AND ORGANIC FERTILIZER

Ashraf Sh. Osman and Mofreh S. Tolba

Horticulture Dept., Fac. Agric., Fayoum Univ., Egypt.

ABSTRACT

A filed trial was conducted during the two successive seasons of 2007/2008 and 2008/2009 in the Experimental Farm at Demo (reclaimed soil with salinity at 7.84 and 8.01 dSm⁻¹ in the two seasons, respectively), Faculty of Agriculture, Fayoum University to investigate the influence of farmyard manure soil application at the rates of 0, 10, 20 and 30m³ fed.⁻¹, ascorbic acid foliar application at the rates of 0, 100, 200, 300 and 400mg l⁻¹ and their combinations on the possibility of improving growth, yield of onion (*Allium cepa* L.) plants under the above mentioned conditions. In comparison with zero rate of farmyard manure, all other rates significantly increased vegetative growth characters (i. e. plant height, No. of leaves plant⁻¹, leaves fresh weight plant⁻¹, raw bulb fresh weight plant⁻¹, total fresh weight plant⁻¹, leaves dry weight plant⁻¹, bulb dry weight plant⁻¹ and total dry weight plant⁻¹), total yield and some chemical constituents under study (i.e. total chlorophyll, total carotenoids, N, P and K of leaves and/or bulbs). The same results were obtained with all ascorbic acid rates; 100, 200, 300 and 400mg l⁻¹ as compared to the zero rate. Economically, ascorbic acid treatment at the rate of 300mg l⁻¹ combined with soil fertilization treatment at the rate of 20m³ fed.⁻¹ proved to be the best and may counteracted the inhibitory effects of salinity on onion plants.

In view of above mentioned results, it has been concluded that spraying onion plant (cv. Giza 6), producing with the soil fertilized by the farmyard manure at the rate of 20m³ fed.⁻¹, with ascorbic acid at the rate of 300mg l⁻¹ could be counteracted the adverse conditions particularly, salinity up to 5000ppm and consequently, economic yield is obtainable.

Key Words: Onion (*Allium cepa* L.), farmyard manure, ascorbic acid, salinity, vegetative growth, yield, Chemical composition.

INTRODUCTION

In Egypt, onion (*Allium cepa* L.) is considered one of the most important vegetable crops growing for local consumption and exportation.

A general reduction in growth and yield due to salinity is widely documented (El-Saidi, 1997). The drastic influence of salinity on plant growth and metabolism was attributed, principally, to the enhanced Na⁺ uptake which causes ion excess in plant tissues (Greenway and Munns, 1980 and Abbas *et al.*, 1991). One of the primary effects of increasing salinity in the growth medium is the inhibition of K⁺, Ca²⁺ and NO₃⁻ uptake by plant roots (Mass, 1986). In addition, It is well established that salinity stresses damages plant cells through production of reactive oxygen species including superoxide, hydrogen peroxide, hydroxyl anions and singlet oxygen (Scandalios, 1997). On the other hand, some trials have been made to alleviate the disturbances in plant metabolism excreted by salinity stress. It has been suggested that some antioxidants (to which belongs ascorbic acid) may help to overcome some of

these inhibitory effects. Ascorbic acid is an important antioxidant defense in plant cells (Foyer and Halliwell, 1976) to protect them by scavenging the reactive oxygen species. It also stimulates respiration activities, cell division and many enzymes activities (Reda *et al.*, 1977; Innocenti *et al.*, 1990 and Rautenkranz *et al.*, 1994).

Latterly, there are widespread use of natural and safety substances such as antioxidants particularly, ascorbic acid for enhancing growth and productivity of many crops. Since ascorbic acid has synergistic effect on growth, flowering, yield and chemical composition under favourable and unfavourable environmental conditions i.e. salinity (Ali; 2002 on tomato plants and Rady; 2006 on sunflowers plants)

Several recent studies have indicated that addition of organic manure, as a natural or synthetic organic materials, to salinity reclaimed soil is necessary due to their unique ability to improve the chemical, hydrophysical and biological characteristics of soils or growing media. However, their buffering effect helps maintaining an uniform reaction in soil media, beside it can hold up to 20 times its weight in water. This is important particularly for salinity reclaimed soils to improve soil moisture conditions, especially during summer seasons. In this respect, Abo-El-Defan (1990) in his study under saline condition found that addition of organic manure increased fresh and dry yields of tomato shoots, fruits and the concentration of N, P and K in both shoots and fruits.

Organic matter may affect plant growth as a source of growth promoters, auxins, vitamins, amino acids which act on the vegetative growth, yield and quality of the plant product (Melo and De- Oliveira, 1999).

El-Foly (2004) said that the organic manure fertilizers are very important means for providing the plants with their nutritional requirements without having undesirable impact on the environment. For many years, organic fertilizers have been used basically as a mean of alleviation of the problem of chemical residues in the export market commodities.

Accordingly, the present work was planned for studying the effect of different rates of farmyard manure and exogenous application of ascorbic acid on growth parameters, yield and some chemical constituents of onion plants cv. Cleopatra grown under saline reclaimed soils during both 2007/2008 and 2008/2009 seasons.

MATERIALS AND METHODS

A field experiment was conducted using onion (cv. Giza 6) during two successive seasons of 2007/2008 and 2008/2009 in the Experimental Farm at Demo (reclaimed soil with salinity at 7.84 and 8.01 dSm⁻¹ in the two seasons, respectively), Faculty of Agriculture, Fayoum University, Egypt. Prior to the initiation of each season, soil samples to 25cm depth from the experimental site were taken and analyzed according to the published procedures of Wilde *et al.*, (1985) and the results are given in Table (1). Imported onion seeds cv. Giza 6 were sown in the nursery on 20th September, 2007 and 23th September, 2008 seasons. Fifty days after seed sowing, each seedling was transplanted into a plant bed in the field. The used experimental design was a split plot in a randomized complete blocks with four replications. Farmyard manure rates were allocated in the main plots and ascorbic acid rates were assigned randomly to sub-plots. Each experimental unit; 20 m² area consisted of 8 rows; 5 m long and 50 cm width, within row spacing averaged 10 cm apart.

ALLEVIATE THE HARMFUL EFFECTS OF SALINE RECLAIMED... 12

The treatments comprised different farmyard manure rates (0 "as control", 10, 20 and 30m³ fed.⁻¹) were broadcasted and incorporated during the soil preparation. The main chemical characteristics and nutrients status of the applied farmyard manure are presented in Table (2).

Table (1): Physical and chemical characteristics of the experimental site during the seasons of 2007/2008 and 2008/2009.

Property	2007/2008	2008/2009
Physical:		
Clay %	28.30	27.50
Silt %	20.60	21.10
Fine sand %	51.10	51.40
Soil texture	Sandy clay loam	Sandy clay loam
Chemical:		
pH (1:2.5)	7.81	7.78
ECe (ds m ⁻¹)	7.81	8.01
Organic matter (%)	1.26	1.21
Ca CO ₃ (%)	8.48	8.61
N (%)	0.08	0.07
Available elements(mg kg⁻¹ soil):		
K	70.07	65.88
P	16.29	14.96
Zn	0.61	0.56
Fe	5.25	5.03
Mn	4.11	3.78

Table (2): The main chemical analysis and nutrients of the applied farmyard manure.

Character	Value		Character	Value	
	2007/2008	2008/2009		2007/2008	2008/2009
Weight of 1 m ³ (kg)	800	823	C/N ratio	22.02	21.22
pH (1:10 water suspension)	7.41	7.33	Total P %	0.64	0.58
EC (dS/m, 1:10 water extract)	4.1	3.8	Total K %	1.11	1.08
Moisture content %	13.2	19.1	Available Fe (mg kg ⁻¹)	783	775
Organic matter %	48.8	43.7	Available Mn (mg kg ⁻¹)	133	117
Organic carbon %	28.4	26.1	Available Zn (mg kg ⁻¹)	108	93
Total N %	1.29	1.23	Available Cu (mg kg ⁻¹)	35	31

In addition to all different farmyard manure rates, recommended mineral fertilizers (200kg N + 72kg K₂O + 31kg P₂O₅ fed.⁻¹) were added. Nitrogen (Ammonium nitrate 33.5% N) was applied in 4 equal applications; one of them was spread on the rows before transplanting directly and the rest of the amount was added at 30, 50 and 70 days from transplanting. Potassium sulphate (48% K₂O) was used side banded at two equal doses; 50 and 70 days from transplanting. Phosphorus fertilizer, in the form of calcium superphosphate (15.5% P₂O₅) was broadcasted and incorporated during soil preparation. Other

agricultural practices were applied as recommended for commercial onion production.

The plants were received the foliar application of ascorbic acid at 0, 100, 200, 300 and 400 mg l⁻¹ to run off, three times; 30, 45 and 60 days after transplanting. Few drops of Tween-20 were added to the spraying solution as a surfactant.

At maturity stage; 8th and 11rd of April 2008 and 2009, respectively, a random ten plants were collected from the four outer rows from each treatment to estimate the following parameters: plant height (cm), No. of leaves plant⁻¹, fresh and dry weights of leaves plant⁻¹ (g), raw fresh bulb weight (g), dry weight of bulb (g), total fresh and dry weights of bulbs plant⁻¹ (g) were recorded. Also, whole plants were collected from four inner rows from each experimental plot and weighed then used for estimating the total yield (ton fed.⁻¹). Fresh samples of leaves and bulbs from ten plants were dried after their collection randomly from each plot in an electric oven at 70°C for 72h to determine dry weight then, ground and kept for chemical analysis.

Chemical analysis: leaf pigments; ninety days after transplanting, a random sample of fresh mature leaves of four plants was collected for determining leaf pigments; total chlorophyll and total carotenoids contents (mg g⁻¹) were extracted by acetone 80% then, determined using colorimetric method (Welburn and Lichtenthaler, 1984). At harvesting stage, in dried leaves and bulbs, nitrogen (% dry matter) was colorimetrically determined using the Orange-G dye (Hafez and Mikkelsen, 1981). Phosphorus (% dry matter) was colorimetrically determined using chlorostannous molybdophosphoric blue colour method in sulphuric acid system (Jackson, 1967). Potassium (% dry matter) was determined using a Perkin-Elmer Flame Photometer (Page *et al.*, 1982).

Statistical analysis:

All data were subjected to statistical analysis according to (Snedecor and Cochran, 1980). Comparisons of the means were carried out using the least significant difference (LSD) at $p = 0.05$ level.

RESULTS AND DISCUSSION

I- Growth characters:

As for farmyard manure; main treatments, data in **Tables (3 - 5)** show a significant gradual increase in all growth characters under study with increasing farmyard manure rate as compared to the control treatment. The differences among 0, 10 and 20 m³ fed.⁻¹ were significant. On the other hand, the difference between the two rates 20 and 30 m³ fed.⁻¹ was insignificant. Economically, the best farmyard manure rate was 20 m³ fed.⁻¹ which surpassed the control by 8.6, 16.1, 18.2, 14.1, 16.2, 9.5, 11.7 and 10.9% in the first season whereas in the second one surpassed the control by 8.9, 16.8, 19.6, 15.5, 16.9, 10.8, 14.3 and 13.0% for plant height, No. of leaves plant⁻¹, leaves fresh weight plant⁻¹, raw bulb fresh weight plant⁻¹, total plant fresh weight, leaves dry weight plant⁻¹, bulb dry weight plant⁻¹ and total dry weight plant⁻¹ as compared to control treatment; 0 m³ fed.⁻¹ farmyard manure. These enhancing results may be attributed to the improvement in leaf pigments (**Table, 6**) and nutritional status (**Tables, 7 and 8**) which positively reflected on total onion yield (**Table, 6**).

Table 3 Influence of different rates of farmyard manure fertilizer and ascorbic acid foliar application on plant height and leaves number plant⁻¹ of onion plant during the two seasons of 2007/2008 and 2008/2009.

Season	2007 / 2008					2008 / 2009						
	Farmyard manure (m ³ /fed.)					Ascorbic acid (mg l ⁻¹)		Plant height (cm)				
	0	100	200	300	400	Mean	0	100	200	300	400	Mean
Control	55	56	58	61	62	58	52	54	56	59	59	56
10	56	58	60	63	64	60	54	56	59	61	62	58
20	59	60	63	66	67	63	57	59	62	65	65	61
30	60	61	64	67	68	64	58	59	62	65	66	62
Mean	57	59	61	64	65	55	57	60	62	63	63	
L.S.D 5%	F = 1		A = 2		Int = 4		F = 1		A = 2		Int = 4	
	No. leaves plant ⁻¹											
Control	8.58	9.91	11.02	13.12	13.72	11.27	8.51	9.49	10.57	11.58	11.88	10.41
10	9.21	10.32	12.14	14.11	14.41	12.04	8.99	10.00	11.41	12.62	12.63	11.13
20	9.98	11.58	13.24	15.12	15.47	13.08	9.83	11.00	12.52	13.71	13.72	12.16
30	10.18	11.87	13.38	15.19	15.75	13.27	9.88	11.26	12.79	13.95	14.01	12.38
Mean	9.49	10.92	12.45	14.38	14.84	9.30	10.44	11.82	12.97	13.06	13.06	
L.S.D 5%	F = 0.55		A = 0.94		Int = 1.71		F = 0.76		A = 0.77		Int = 1.88	

F = Fertilizer

A = Ascorbic acid

Int = Interaction

Table 4. Influence of different rates of farmyard manure fertilizer and ascorbic acid foliar application on leaves, raw bulb and total fresh weight plant⁻¹ of onion plant during the two seasons of 2007/2008 and 2008/2009.

Season	2007 / 2008						2008 / 2009					
	Farmyard manure (m ³ /fed.)		Ascorbic acid (mg l ⁻¹)				Leaves fresh weight plant ⁻¹ (g)		Raw bulb fresh weight plant ⁻¹ (g)			
	0	100	200	300	400	Mean	0	100	200	300	400	Mean
Control	58.0	63.6	70.8	79.9	81.9	70.8	55.6	61.5	68.0	76.7	78.3	68.0
10	60.8	67.8	76.7	85.3	86.0	75.3	58.2	65.4	74.1	82.5	83.1	72.7
20	66.2	75.8	84.1	95.7	96.8	83.7	64.7	73.5	81.6	92.8	93.9	81.3
30	69.7	78.0	87.2	97.3	98.4	86.1	66.6	75.6	83.9	94.4	95.5	83.2
Mean	63.7	71.3	79.7	89.5	90.8		61.3	69.0	76.9	86.6	87.7	
L.S.D 5%	F = 2.6		A = 5.0		Int = 8.5		F = 2.1		A = 4.7		Int = 7.9	
Control	104	111	121	133	134	121	100	107	117	127	129	116
10	109	118	129	143	145	129	104	114	124	138	140	124
20	115	126	139	154	156	138	112	122	135	149	150	134
30	118	130	143	159	161	142	115	126	139	154	156	138
Mean	112	121	133	147	149		107	117	129	142	144	
L.S.D 5%	F = 7		A = 8		Int = 19		F = 7		A = 7		Int = 19	
Control	162	174	192	212	216	191	155	169	185	204	207	184
10	170	186	206	228	231	204	162	180	198	220	223	197
20	181	202	223	250	253	222	176	196	216	242	244	215
30	188	207	230	256	259	228	182	201	222	248	252	221
Mean	175	192	213	237	240		169	186	205	229	231	
L.S.D 5%	F = 8		A = 10		Int = 20		F = 7		A = 9		Int = 22	

F = Fertilizer

A = Ascorbic acid

Int = Interaction

Table 5. Influence of different rates of farmyard manure fertilizer and ascorbic acid foliar application on leaves, raw bulb and total plant fresh weight plant⁻¹ of onion plant during the two seasons of 2007/2008 and 2008/2009.

Season	2007 / 2008						2008 / 2009					
	Farmyard manure (m ³ /fed.)		Ascorbic acid (mg l ⁻¹)				Leaves dry weight plant ⁻¹ (g)		Bulb dry weight plant ⁻¹ (g)			
	0	100	200	300	400	Mean	0	100	200	300	400	Mean
Control	7.30	7.77	8.61	9.52	9.82	8.61	7.01	7.51	8.27	9.14	9.37	8.26
10	7.54	7.95	8.87	9.78	10.10	8.85	7.32	7.71	8.61	9.48	9.80	8.58
20	7.95	8.52	9.45	10.56	10.67	9.43	7.72	8.26	9.19	10.24	10.35	9.15
30	8.14	8.66	9.65	10.76	10.88	9.62	7.90	8.43	9.37	10.43	10.55	9.34
Mean	7.74	8.22	9.15	10.15	10.37		7.49	7.98	8.86	9.83	10.02	
L.S.D 5%	F = 0.42		A = 0.35		Int = 1.01		F = 0.40		A = 0.35		Int = 0.99	
Control	13.72	14.43	15.36	16.44	16.69	15.33	13.49	13.76	14.44	15.45	15.90	14.61
10	14.24	15.05	16.10	17.27	17.45	16.02	13.60	14.39	15.74	16.77	16.92	15.48
20	15.08	16.14	17.27	18.48	18.65	17.12	14.63	15.65	16.99	18.05	18.17	16.70
30	15.28	16.36	17.62	18.95	19.07	17.45	14.82	15.86	17.09	18.38	18.49	16.93
Mean	14.58	15.49	16.59	17.78	17.96		14.13	14.92	16.07	17.16	17.37	
L.S.D 5%	F = 0.51		A = 0.86		Int = 1.32		F = 0.48		A = 0.80		Int = 1.23	
Control	21.03	22.20	23.97	25.96	26.52	23.94	20.50	21.27	22.71	24.59	25.27	22.87
10	21.78	23.00	24.97	27.05	27.55	24.87	20.92	22.10	24.35	26.26	26.72	24.07
20	23.04	24.65	26.72	29.04	29.32	26.55	22.34	23.91	26.18	28.30	28.52	25.85
30	23.42	25.01	27.28	29.71	29.95	27.07	22.71	24.30	26.46	28.81	29.04	26.27
Mean	22.32	23.72	25.73	27.94	28.33		21.62	22.89	24.92	26.99	27.39	
L.S.D 5%	F = 0.75		A = 1.01		Int = 2.21		F = 0.71		A = 0.97		Int = 1.94	

F = Fertilizer

A = Ascorbic acid

Int = Interaction

The increase in growth characters with the addition of farmyard manure might be due to the improvement of structure of the soil by increasing the soil water holding capacity which positively reflected on the soil aeration and drainage that encourage the root growth and nutrient absorption. Saleh *et al.* (2003) on onion revealed also that organic manure enhanced the availability of certain elements and their supply to the plant during growth period. However, organic manure increased the presence of P, K and Mg in the soil besides the solubility of Ca, Mg and NO_3 as a result of the lowering in pH induced by manure application and the increase in electrical conductivity.

Yung-Yu Shu (2006) studied the effects of applied different kinds of organic composts on some plant characteristics as compared to those treated with the mineral fertilizers alone, and reported that at the most active vegetative growth stages of the plants, the amount of nutrients absorbed from the chemical fertilizers were found to be higher than those treated with organic manure. Whereas, the values of the studied plant parameters and nutrient uptake by plants with the organic manure treatment were the highest at the maturity stage. The later case confirmed by the plants of the next crop, however, the values of the same plant parameters at the organic manure treatment were also recorded the highest among all treatments at maturity stages. It could be interpreted such phenomena on the fact that the chemical fertilizer was a fast-release fertilizer used to supply nutrients at the early stages of growth of the first crop, while the beneficial effect of the composts on plant growth and nutrient uptake was conspicuous in the next crop as compared with that of routine treatment of chemical fertilizer. These results are in accordance with those reported with Ali *et al.* (2001), Salman *et al.* (2002) and Rizk-Fatma (2002).

Regarding ascorbic acid foliar application; sub-main treatment, data illustrated in Tables (3-5) reveal that, as ascorbic acid rate increased up to 300mg l^{-1} vegetative growth traits under study were gradually increased in significant amounts. The rate of 400mg l^{-1} ascorbic acid scored an insignificant difference in favor of it as compared with the rate of 300mg l^{-1} . Economically, the concentration of ascorbic acid at 300mg l^{-1} considers as the favourable treatment which recorded the following increments over the control treatment (0mg l^{-1} ascorbic acid): 12.3, 51.5, 40.5, 31.3, 35.4, 31.5, 22.0 and 25.2% in the first season. Furthermore, the increments in the second season were as follows: 12.7, 39.7, 41.3, 32.7, 35.5, 31.2, 21.4 and 24.8% for plant height, No. of leaves plant^{-1} , leaves fresh weight plant^{-1} , raw bulb fresh weight plant^{-1} , total plant fresh weight, leaves dry weight plant^{-1} , bulb dry weight plant^{-1} and total dry weight plant^{-1} , respectively.

Increased endogenous concentration of ascorbic acid leads to protect plant cells and consequently, protect the photosynthetic apparatus by scavenging reactive oxygen species (Zhang and Schmidt, 2000) thus, vigorous plant growth will be obtained under stress conditions. In this connection, Elade (1992) stated a positive action for antioxidants especially, ascorbic acid on growth and attributed this finding to their effects on counteracting drought, salinity and diseases stresses and protecting plant cells against free radicals that responsible for plant senescence as well as to their auxinic action and consequently, enhancing growth characters. In addition, ascorbic acid might be regulates cell wall expansion, division and elongation through its action on cell vacuolarization (Gonzalez-Reyes *et al.*, 1994; Navas and Gomez-Diaz, 1995 and Cordoba-Pedregosa *et al.*, 1996), improves the nutritional status and

absorbing phenolic compounds which lead to save the growing tissues from toxic effects of the oxidized phenols (Gupta *et al.*, 1980) and / or enhances the biosynthesis of carbohydrates (Ahmed, 2001) and translocation of sugars (Farang, 1996) which could be explain the present results. These findings are in coincidence with those obtained by Rady (2006) and El-Yazal (2007).

Economically, application of farmyard manure at the rate of 20m³ fed.⁻¹ combined with foliar application with ascorbic acid at 300mg l⁻¹ concentration proved to be the best combined treatment among all combined treatments since surpassed the combined control; zero rate ascorbic acid under zero m³ fed.⁻¹ farmyard manure by 20.0, 76.2, 65.0, 48.1, 54.3, 44.7, 34.7 and 38.1% in the first season and 25.0, 61.1, 66.9, 49.0, 56.1, 46.1, 33.8 and 38.1% in the second season for plant height, No. of leaves plant⁻¹, leaves fresh weight plant⁻¹, raw bulb fresh weight plant⁻¹, total plant fresh weight, leaves dry weight plant⁻¹, bulb dry weight plant⁻¹ and total dry weight plant⁻¹, respectively. This distinctive results in relation to vegetative traits under above mentioned best combined control may be attributed to the distinctive results of leaf pigments (Table, 6) and nutritional status of plants (Tables, 7 and 8) under the same combined treatment.

II- Total yield:

It has been showed from data in Table (6) that, all rates of farmyard manure application exhibited significant differences in total yield as compared to the control; zero rate and among of them except for the difference between the highest two rates; 20 and 30m³ fed.⁻¹ which granted an insignificant difference between them. Thus, the best treatment which economically operative is 20m³ fed.⁻¹ farmyard manure application which yielded increases above the control averaged at 19.1, and 20.2% in the first season and the second one, respectively. These distinguished results in total yield may be attributed to the distinctive results obtained for vegetative traits (Table, 3-5), leaf photosynthetic pigments (Table 6) and nutritional status of plants (Tables, 7 and 8).

The positive effect of farmyard fertilizers on yield may be due to increasing the availability of macro and micronutrients in rooting zone and resulting from soil pH reduction by applying organic manure, which could improve physical and chemical properties of the soil. In this respects Negm *et al.* (2003) found that the added organic manure reduced soil pH value and increased the available content of each N, P and K in soil. They also showed that the curve of increase reached its highest value during harvest stage of the grown plants and gradually tended to reduce again.

Singer *et al.* (2007) reported that organic manure amendment increased P and K uptake by plant roots. Organic manure amendment decreased soil pH with the increase in the soil organic matter which reflected on more availability for nutrients in favour of plants and consequently positively reflected on total yield. This trend is in a good accordance with results which outlined by many investigators; Ali *et al.* (2001), Salman *et al.* (2002) and Rizk-Fatma (2002).

As for the influence of ascorbic acid foliar application on total onion yield which illustrated in Table (6), data reveal that there was a gradual increase by significant amounts with increasing the concentration of ascorbic acid used gradually form 0 to 300mg l⁻¹. The concentration of 400mg l⁻¹ ascorbic acid haven't a significant increment above the treatment of 300mg l⁻¹. The latter considered as the best treatment which recorded increases surpassed the control;

zero mg l⁻¹ ascorbic acid recorded at 29.8% and 28.9% for 2007/2008 and 2008/2009 seasons, respectively.

Regarding the distinctive results obtained from the same best treatment of vegetative characters (**Tables, 3-5**), leaf photosynthetic pigments (**Table, 6**) and the nutritional status of plants (**Tables, 7 and 8**), my proved to be the explanation in favour of total onion yield.

In this respect; **Al-Qubaie (2002)** stated that ascorbic acid as an antioxidant has an auxinic action, synergistic effect on the biosyntheses of carbohydrate and controlling the incidence of most fungi on plants which making them in vigorous states and reflects on yields. Furthermore, **Shalaby (2006)** reported an increase in yield bulbs of onion plant as a result of the foliar application with some antioxidants. Their results regarding the beneficial effect of ascorbic acid on yield are confirmed with those reported by **Rady (2006) and El-Yazal (2007)**.

The interactions between the application of farmyard manure and ascorbic acid and their effects on total onion yield are presented in **Table (6)**. The treatment of 20m³ fed⁻¹ farmyard manure combined with foliar application at the rate of 300mg l⁻¹ proved, in this study, to be the best. This treatment gave the increases averaged at 50.5 and 50.3% for the first and second seasons, respectively as compared with the combined control; zero m³ fed⁻¹ farmyard manure and zero mg l⁻¹ ascorbic acid.

The combined treatment of farmyard manure and ascorbic acid at the rates of 20m³ fed⁻¹ and 300mg l⁻¹, respectively proved to be the best, and exhibited the most pronounced counteracted effect on soil salinity under study.

III- Chemical constituents:

Data in **Tables (6-8)** reveal that content of total chlorophyll and total carotenoids, in plant leaves as well as concentration of N, P and K in plant leaves and bulbs were significantly increased as farmyard manure increased till 20m³ fed⁻¹ then, slightly increased in both two seasons of study. Thus, the treatment of 20m³ fed⁻¹ farmyard manure is considered as the best among all farmyard manure treatments. This treatment recorded increments in the order of 33.0 and 30.2% as well as 37.3 and 31.3% for total chlorophyll and total carotenoids in both two seasons, respectively. Furthermore, the best treatment was scored 8.2, 14.3, 6.0, 8.7, 17.4 and 9.9% in the first season, whereas the second one collected 9.4, 14.9, 7.0, 10.0, 18.1 and 10.4% under the same treatment for leaf N, leaf P, leaf k, bulb N, bulb P and bulb K, respectively. These pronounced increments may be due to the best status of leaf photosynthetic pigments consequently more photosynthates producing more energy for more absorption of mentioned elements and others thus the best nutritional status reflecting on the vigourity of vegetative traits and yield under study.

Regarding the influence of ascorbic acids on leaf photosynthetic pigments as well as N, P and K concentration of leaves and bulbs, data shown in **Tables (6-8)** exhibit that all these parameters were gradually increased by significant quantities as a result of foliar application with ascorbic acid up to 300mg l⁻¹ then negligibly increased. The proportion of 48.1, 64.9, 21.4, 22.6, 13.6, 10.5, 24.3 and 46.9% in the first season as well as the proportion of 47.0, 63.9, 21.0, 22.2, 13.2, 10.0, 23.5 and 45.4% in the second season for leaf chlorophyll, leaf carotenoids, leaf N, leaf P, Leaf K, bulb N, bulb p and bulb k, respectively were the increments of the best treatment economically; spraying plant foliage with

ascorbic acid at the rate of 300mg l^{-1} as compared with the treatment free from ascorbic acid (tap water). The promotive effect of ascorbic acid on chlorophyll and carotenoids and the other components under study might be attributed to the enhancing effects of this antioxidant on the nutritional status of onion plants since, N is one of the essential chlorophyll components and consequently more biosynthesis of photosynthates in the face cell elongation.

As regard to the results of the combination between all farmyard manure treatments and different rates of ascorbic acid, data exhibited in **Tables (6–8)** reveal that the combination between $20\text{m}^3 \text{ fed.}^{-1}$ farmyard manure treatment and spraying plant foliage with ascorbic acid at 300mg l^{-1} economically preferable to all other combinations since granted the following increases : 102.8, 120.0, 30.6, 37.7, 19.8, 19.3, 40.5 and 59.7% in the first season while, in the second one granted the following increments : 100.5, 117.0, 33.0, 39.2, 19.8, 20.0, 41.5 and 60.6% for leaf chlorophyll, leaf carotenoids, leaf N, leaf P, Leaf K, bulb N, bulb P and bulb K, respectively as compared to the combined treatment of zero rate farmyard manure and zero rate ascorbic acid.

The beneficial effect of farmyard manure and ascorbic acid on overcoming soil salinity which reflected on improving vegetative growth traits (**Tables, 3, 4 and 5**) and yield (**Table, 6**), surely reflected also on stimulating the nutritional status of plants.

Ali et al. (2001), Rizk-Fatma (2002) and Mohamed and El-Ganaini (2003) They attributed such results to the effect of organic fertilizers in improving soil physical and biological properties as well as chemical characteristic resulting in more release of available nutrient elements to be absorbed by plant roots and the water use efficiency by different plants, which was positively reflected on the biosynthesis processes, particularly, the pigments and carbohydrates.

The promotive effect of ascorbic acid on photosynthetic pigments and chemical constituents might be attributed to the enhancing effect of this antioxidant on the nutritional status of onion plant (**Tables, 6-8**) since, nitrogen is one of essential chlorophyll components. Besides, the presence of iron and manganese is necessary for biosynthesis of chlorophylls (**Rady, 2006**). The role of ascorbic acid as an antioxidant, which directly involved in the regulation and protection of photosynthetic processes (**Farago and Brunold, 1994**) could be led to enhancing the effect of ascorbic acid on photosynthetic pigments under study. This may be explained by the findings of **Foyer et al. (1990)** who stated that, the antioxidant prevented enzyme inactivation, the generation of more dangerous radicals and allowed flexibility in the production of photosynthetic assimilatory power. Moreover, electron transfer to O_2 prevented over reduction of electron transported chain, which reduced the risk of harmful back reactions within the photosystem. In addition, **Elade (1992) and Farag (1996)** proved that most antioxidants were responsible for accelerating the biosynthesis of various pigments. The positive effect of ascorbic acid on photosynthetic pigments obtained by **Ali (2002), Rady (2006) and El-Yazal (2007)** were in agreement with findings of the present study.

Increasing chemical constituents concentrations of plants leaves producing from organic fertilizer fecundated-soil might be due to the increase in plant capacity to absorb nutrients, which increased the roots surface per soil volume unit; **Negm et al. (2003)** showed that the added organic manure reduced soil pH value, increased soil organic matter and increased the available content of N, P

Table 6. Influence of different rates of farmyard manure fertilizer and ascorbic acid foliar application on total yield and total chlorophyll and carotenoids of onion plant during the two seasons of 2007/2008 and 2008/2009.

Season	2007 / 2008					2008 / 2009						
	Farmyard manure (m ³ /fed.)					Ascorbic acid (mg l ⁻¹)						
	0	100	200	300	400	Mean	0	100	200	300	400	Mean
	Total yield (ton fed⁻¹)											
Control	7.07	7.39	8.01	8.77	8.93	8.03	6.86	7.21	7.67	8.36	8.47	7.71
10	7.41	7.89	8.51	9.36	9.56	8.55	7.19	7.61	8.20	8.96	9.09	8.21
20	7.96	8.78	9.57	10.64	10.85	9.56	7.72	8.51	9.28	10.31	10.52	9.27
30	8.19	9.02	9.83	11.01	11.10	9.83	7.94	8.75	9.53	10.69	10.75	9.53
Mean	7.66	8.27	8.98	9.94	10.11		7.43	8.02	8.67	9.58	9.71	
L.S.D 5%	F = 0.45		A = 0.69		Int = 1.12		F = 0.47		A = 0.60		Int = 1.11	
	Total chlorophyll (mg g⁻¹ fresh wt. of leaves)											
Control	0.393	0.455	0.521	0.588	0.599	0.511	0.385	0.439	0.500	0.564	0.575	0.493
10	0.450	0.517	0.587	0.661	0.678	0.578	0.446	0.493	0.569	0.641	0.644	0.559
20	0.530	0.612	0.702	0.797	0.824	0.693	0.513	0.611	0.689	0.772	0.798	0.677
30	0.574	0.647	0.748	0.841	0.855	0.733	0.556	0.634	0.725	0.816	0.829	0.712
Mean	0.487	0.558	0.639	0.721	0.739		0.475	0.544	0.621	0.698	0.712	
L.S.D 5%	F = 0.057		A = 0.057		Int = 0.183		F = 0.051		A = 0.053		Int = 0.180	
	Total carotenoids (mg g⁻¹ fresh wt. of leaves)											
Control	0.060	0.072	0.088	0.104	0.108	0.086	0.059	0.070	0.085	0.099	0.103	0.083
10	0.068	0.081	0.098	0.115	0.119	0.096	0.066	0.079	0.095	0.111	0.115	0.093
20	0.082	0.097	0.115	0.132	0.135	0.112	0.080	0.094	0.112	0.128	0.131	0.109
30	0.087	0.102	0.121	0.137	0.138	0.117	0.085	0.099	0.117	0.133	0.133	0.113
Mean	0.074	0.088	0.106	0.122	0.125		0.072	0.086	0.102	0.118	0.121	
L.S.D 5%	F = 0.007		A = 0.008		Int = 0.019		F = 0.007		A = 0.007		Int = 0.019	

F = Fertilizer

A = Ascorbic acid

Int = Interaction

Table 7. Influence of different rates of farmyard manure fertilizer and ascorbic acid foliar application on concentration of N, P and K on leaves of onion plant during the two seasons of 2007/2008 and 2008/2009.

Season	2007 / 2008					2008 / 2009						
	Farmyard manure (m ³ /fed.)					Ascorbic acid (mg l ⁻¹)						
	0	100	200	300	400	Mean	0	100	200	300	400	Mean
N (% dry matter)												
Control	0.98	1.04	1.11	1.18	1.20	1.10	0.94	1.00	1.06	1.13	1.15	1.06
10	1.02	1.06	1.14	1.23	1.24	1.14	0.97	1.02	1.09	1.18	1.18	1.09
20	1.06	1.12	1.20	1.28	1.31	1.19	1.03	1.09	1.18	1.25	1.27	1.16
30	1.07	1.14	1.24	1.30	1.31	1.21	1.05	1.11	1.20	1.27	1.28	1.18
Mean	1.03	1.09	1.17	1.25	1.26		1.00	1.06	1.13	1.21	1.22	
L.S.D 5%	F = 0.06		A = 0.05		Int = 0.11		F = 0.04		A = 0.05		Int = 0.10	
P (% dry matter)												
Control	1.83	1.90	2.02	2.17	2.22	2.03	1.76	1.84	1.94	2.07	2.12	1.95
10	1.90	2.00	2.13	2.28	2.30	2.12	1.84	1.94	2.07	2.21	2.23	2.06
20	2.02	2.15	2.31	2.52	2.58	2.32	1.96	2.07	2.24	2.45	2.50	2.24
30	2.07	2.22	2.37	2.60	2.63	2.38	2.00	2.13	2.29	2.51	2.54	2.30
Mean	1.95	2.07	2.21	2.39	2.43		1.89	2.00	2.14	2.31	2.35	
L.S.D 5%	F = 0.11		A = 0.11		Int = 0.24		F = 0.10		A = 0.10		Int = 0.22	
K (% dry matter)												
Control	1.30	1.35	1.40	1.46	1.47	1.40	1.27	1.31	1.35	1.40	1.41	1.35
10	1.32	1.37	1.43	1.50	1.51	1.43	1.29	1.33	1.39	1.45	1.46	1.38
20	1.36	1.42	1.49	1.56	1.57	1.48	1.33	1.39	1.45	1.52	1.53	1.44
30	1.39	1.44	1.51	1.59	1.60	1.50	1.34	1.40	1.47	1.54	1.55	1.46
Mean	1.34	1.39	1.46	1.53	1.54		1.31	1.36	1.41	1.48	1.49	
L.S.D 5%	F = 0.03		A = 0.04		Int = 0.07		F = 0.03		A = 0.04		Int = 0.08	

F = Fertilizer

A = Ascorbic acid

Int = Interaction

Table 8. Influence of different rates of farmyard manure fertilizer and ascorbic acid foliar application on concentration of N, P and K on bulb of onion plant during the two seasons of 2007/2008 and 2008/2009.

Season	2007 / 2008					2008 / 2009						
	Farmyard manure (m ³ /fed.)					Ascorbic acid (mg l ⁻¹)						
	0	100	200	300	400	Mean	0	100	200	300	400	Mean
N (% dry wt.)												
Control	1.09	1.11	1.16	1.19	1.20	1.15	1.05	1.08	1.10	1.14	1.15	1.10
10	1.12	1.14	1.18	1.22	1.23	1.18	1.07	1.11	1.14	1.17	1.19	1.14
20	1.17	1.20	1.25	1.30	1.31	1.25	1.13	1.17	1.22	1.26	1.27	1.21
30	1.18	1.22	1.29	1.32	1.33	1.27	1.15	1.18	1.24	1.28	1.29	1.23
Mean	1.14	1.17	1.22	1.26	1.27		1.10	1.13	1.18	1.21	1.22	
L.S.D 5%	F = 0.05		A = 0.05		Int = 0.11		F = 0.05		A = 0.05		Int = 0.11	
P (% dry wt.)												
Control	1.90	1.95	2.07	2.20	2.24	2.07	1.83	1.89	1.98	2.12	2.15	1.99
10	1.98	2.06	2.22	2.39	2.42	2.21	1.92	2.00	2.15	2.31	2.34	2.14
20	2.08	2.24	2.43	2.67	2.71	2.43	2.02	2.17	2.35	2.59	2.63	2.35
30	2.12	2.30	2.49	2.75	2.78	2.49	2.07	2.23	2.42	2.66	2.68	2.41
Mean	2.02	2.14	2.30	2.51	2.54		1.96	2.07	2.23	2.42	2.45	
L.S.D 5%	F = 0.11		A = 0.10		Int = 0.22		F = 0.10		A = 0.10		Int = 0.21	
K (% dry wt.)												
Control	2.04	2.27	2.61	2.96	3.06	2.59	1.97	2.22	2.54	2.88	2.92	2.50
10	2.09	2.36	2.72	3.08	3.11	2.67	2.05	2.30	2.64	2.99	3.01	2.60
20	2.22	2.51	2.91	3.25	3.31	2.84	2.19	2.45	2.82	3.16	3.21	2.76
30	2.24	2.54	2.96	3.32	3.37	2.88	2.23	2.50	2.87	3.22	3.27	2.82
Mean	2.15	2.42	2.80	3.15	3.21		2.11	2.37	2.72	3.06	3.10	
L.S.D 5%	F = 0.03		A = 0.03		Int = 0.07		F = 0.03		A = 0.03		Int = 0.07	

F = Fertilizer

A = Ascorbic acid

Int = Interaction

and K in the soil. These results are in agreement with reported by Ali *et al.* (2001), Rizk-Fatma (2002) Mohamed and El-Ganaini (2003), Saleh *et al.* (2003).

The obtained results were also supported by the results of Ahmed and Abd El-Hameed (2004) who reported that the effect of antioxidants on producing healthy plants leads to enhancing the plants to have a great ability for uptake of elements. Moreover, Gonzalez-Reyes *et al.* (1994) concluded that ascorbate free radical caused hyperpolarization of plasma membranes, and this energization could then facilitate transport processes across such membranes. Most of the previous results are consistent with those of Ali (2002), Rady (2006) and El-Yazal (2007).

CONCLUSION

Within the experimental conditions studied, it has been concluded that this work gave an evidence to the role of farmyard manure especially at the rate of 20m³ fed.⁻¹ by which the soil pH can be reduced consequently more solubility and availability of nutrients for the absorption by plant roots besides, enriching the soil with organic matter. Moreover, the role of ascorbic acid, as an antioxidant by which plant foliage sprayed especially at the concentration 300mg l⁻¹, in inducing salinity tolerance of onion plants cv. Giza 6 cultivated in salt-affected reclaimed soils containing salts concentration at about 500ppm leading to favourable growth and consequently obtain economic yield under such conditions.

REFERENCES

- Abbas, M.A.; Younis, M.E. and Shukry, W.M. (1991). Plant growth, metabolism and adaption in relation to stress conditions. III. Effect of salinity on the internal solute concentrations in *Phaseolus vulgaris*. J. Plant Physiol., 138: 722 – 729.
- Abo-El-Defan T.R. (1990). Effect of organic manure on plant growth and nutrients uptake under saline condition. M. Sc., Fac. Agric., Ain Shams Univ., Cairo, Egypt.
- Ahmed, A.M. (2001). Studies for controlling malformation and improving yield and fruit quality of hindy bisinnara mangoes by using active dry years, ascorbic acid and sulphur. Minia J. of Agric Res. & Develop., 21 (2): 219 – 233.
- Ahmed, F.F. and Abd El-Hameed, H.M. (2004). Influence of some antioxidants on growth, vine nutritional status, yield and quality of berries in banaty grapevines. Assiut J. of Agric. Sci., 35 (4): 131 – 140.
- Ali, A.H.; Abdel-Mouty, M.M. And A.M. Shaheen (2001). Effect of bio-nitrogen, organic and inorganic fertilizer, on the productivity of garlic (*Allium sativum* L.) plants. Egypt J. Appl. Sci. 16 (3): 173 – 188.
- Ali, Z.A. (2002). Effect of foliar application with ascorbic acid on vegetative growth and some biochemical constituents of tomato plants. J. Agric. Sci., Mansoura Univ., 27 (10): 6765 – 6775.
- Al-Qubaie, A.I. (2002). Response of *Ficus nitida* L. seedlings to the application of some antioxidants under soil salinity conditions. Minia J. of Agric. Res. & Develop., 22 (3): 235 – 254.
- Cordoba-Pedregosa, M.C.; Gonzalez-Reyes, J.A.; Sandillas, M.S.; Navas, P. and Cordoba, F. (1996). Role of apoplastic and cell-wall peroxidases

- on the stimulation of root elongation by ascorbate. *Plant Physiol.*, 112: 1119 – 1125.
- Elade, Y. (1992).** The use of antioxidants to control gray mould (*Botrytis cineria*) and white mould (*Sclerotinia sclerotiorum*) in various crops. *Plant Pathol.*, 141: 417 -426.
- El-Foly, H.M.H. (2004).** The productivity of potato plants as affected by different organic and inorganic fertilizers. *Minia J. of Agric Sc.*, (34) 6: 285-305.
- El-Saidi, M.T. (1997).** Salinity and its effect on growth, yield and some physiological processes of crop plants. In: Strategies for improving salt tolerance in higher plants. Jaiwal, P. K.; Singh, R. P. and Anju Gulati (Eds.). Oxford and IBH Publishing Co. Pvt. Ltd. New Delhi, Calcutta and Enfield (USA), pp. 111-127.
- El-Yazal, M.A. (2007).** Effect of some antioxidants on growth, yield and some chemical constituents of onion (*Allium cepa* L.). *Fayoum J. Agric. Res. & Dev.*, 21 (1): 162-176.
- Farag, K.M. (1996).** Use of urea, phenylalanine, thiamine or their combinations to accelerate anthocyanins development and their effect on the storage life of Flame seedless grapes. First Egyptian Hungarian Hort. Vonf., Kafr El-Sheikh, Egypt, 15 – 17 Sept.
- Farago, S. and Brunold, C. (1994).** Regulation of thiol contents maize roots by intermediates and effectors of glutathione synthesis. *J. Plant Physiol.*, 144:433 – 437.
- Foyer, C.H. and Halliwell, B. (1976).** The presence of glutathione and glutathione reductase in chloroplasts: A proposed role in ascorbic acid metabolism *Plants*, 157: 239 – 244.
- Foyer, C.H.; Furbank, R.T.; Harbinson, J. and Horton, P. (1990).** The mechanism contributing to photosynthetic control of electron transport by carbon assimilation in leaves. *Photosynth. Res.*, 25: 83 – 100.
- Gonzalez-Reyes, J.A.; Alcain, F.J.; Serrano, A. Cordoba, F. And Navas, P. (1994).** Relationship between apoplastic ascorbate regeneration and stimulation of root growth in *Allium cepa* L. *Plant Sci.*, 100: 23 – 29.
- Greenway, H. and Munns, R. (1980).** Mechanisms of salt tolerance in nonhalophytes. *Ann. Rev. Plant Physiol.*, 31: 149 – 190.
- Gupta, P.K.; Nadgir, A.L.; Macarentias, A.F. And Jagannathan, V. (1980).** Tissue culture of forest trees: Clonal multiplication *Tecoma grandis* L. (treak) by tissue culture. *Plant Sci. Letters*, 17: 259 – 268.
- Hafez, A.R. and D.S. Mikkelsen (1981).** Colorimetric determination of nitrogen for evaluation the nutritional status of rice. *Commun. Soil Sci. and Plant Analysis*, 12(1): 16-69.
- Innocenti, M.A.; Bitonti, M.; Arrigoni, O. and Liso, R. (1990).** The size of quiescent center in roots of *Allium cepa* L. grown with ascorbic acid. *New Phytol.*, 114: 507 – 509.
- Jackson, M.L. (1967).** *Soil Chemical Analysis*. Prentice-Hall of India, pp.144-197.
- Mass, E.V. (1986).** Crop tolerance to saline soil and water. *Proc. Us. Pak. Biosaline Res. Workshop, Botany Dep., Karachi Univ., Pakistan*, pp. 205 – 219.

- Melo, J.P.L. and A.P.De-Oliveira (1999).** Garlic production as a function of different water levels and bovine manure in soil. *Horticultura, Brasileira*, 17: 11-15.
- Mohamed, S.A. and S.S.S. El-Ganaini (2003).** Effect of organic, mineral and biofertilizers on growth, yield and chemical constituents as well as the anatomy of board bean (*Vicia faba* L.) plants grown in reclaimed soil. *Egypt J. App. Sci.*, 18 (12): 38 – 63.
- Navas, P. and Gomez-Diaz, C. (1995).** Ascorbic free radical and its role in growth control. *Protoplasm*, 184: 8 – 13.
- Negm, M.A.; M.G.M. Rifaat and A.N. Estefanous (2003).** Impact of composted saw-dust and some nitrogenous sources on the production of squash and table beet crops grown on calcareous soil. *Fayoum J. Agric., Res. & Dev.*, 17 (1): 116-137.
- Page, A.I.; R.H. Miller and D.R. Keeney (1982).** *Methods of Soil Analysis. Part II. Chemical and Microbiological Methods.* 2nd ed. Amer. Soc. Agron., Madison, Wisconsin, USA.
- Rady, M.M. (2006).** Efficiency of growth and productivity of sunflower plants as affected by ascorbic acid under saline reclaimed soil conditions. The Second Conference on farm Integrated Pest Management, 16–18 Jan., pp. 186 -200.
- Rautenkranz, A.; machler, F.; Martinoia, E and Oertli, J. (1994).** Transport of ascorbic acid dehydroascorbic acid across protoplast and vacuole membranes isolated from barley (*Hordeum vulgare* L. cv. Gebrel) leaves. *Plant Physiol.*, 106: 187 – 193.
- Reda, F.; Fadl, M.; Abdel-All, R. and El-Moursi, A. (1977).** Physiological studies on *Ammi visnaga* L. The effect of thiamine and ascorbic acid on growth and chromone yield. *Egypt. J. Pharm. Sci.*, 18: 19 -27.
- Rizk-Fatma, A. (2002).** Effect of organic manure with or without inorganic fertilizers, on the productivity of Okra plant (*Hibiscus esculenta* L.). *Egypt J. Appl. Sci.*, 17 (8): 364 – 383.
- Saleh, A. L.; A. A. Abd El-Kader and S. A. M. Hegab (2003).** Response of onion to organic fertilizer under irrigation with saline water. *Egypt. J. Appl. Sci.*, 18(12 B): 707-716.
- Salman, S.R.; Sawan, O.M. and A.M. Eissa (2002).** The effect of different levels and sources of organic fertilizers on NPK status, yield and quality of watermelon in newly reclaimed soil. *Egypt J. App. Sci.*, 17 (10): 665 – 683.
- Scandalios, J.G. (1997).** Molecular genetics of superoxide dismutases in plants. pp. 527–568. In: Scandalios, J. G.(ed.). *Oxidative stress and the molecular biology of antioxidant defenses.* Cold Spring Harbor Lab. Press, Plainview, N. Y.
- Shalaby, O.Y. (2006).** Effect of some antioxidants and ridomil on the incidence of onion downy mildew, purple blotch and plant chemical components. *Egypt., J. of App. Sci.*, 21 (3): 59 – 70.
- Singer, J.W.; Logsdon, S.D. and Meek, D.W. (2007).** Tillage and compost effects on corn growth, nutrient accumulation and grain yield. *Agron. J.*, 99: 80-87.
- Snedecor, G.W. and W.G. Cochran (1980).** *Statistical Methods.* 7th ed., Iowa State Univ. Press, Ames, Iowa, USA.
- Welburn, A.R. and Lichtenthaler, H.(1984).** Formula and program to determine total carotenoids and chlorophylls a and b of leaf extracts in

- different solvents. In "Advances in Photothynthesis Research" (Sybesma C. Ed.), Vol., II, pp. 9-12. Martinus Njihoff Dr. W.Jnk Publishers, The Hague.
- Wilde, S.A.; R.B. Corey; J.G. Lyer and G.K. Voigt (1985).** Soil and Plant Analysis for Tree culture. Oxford and IBM Publishers. New Delhi. India. 3rd ed. pp. 93-106.
- Yung-Yu Shun (2006).** Effect of application of different types of organic composts on rice growth under laboratory conditions. Soil Sci. and Nutrition, 51: 443 – 449.
- Zhang, X. and Schmidt, R.E. (2000).** Hormone-econtaining products impact on antioxidant status of tall fescue and creeping bentgrass subjected to drought. Crop Sci., 40: 1344 – 1349.

تقليل التأثيرات الضارة للأراضي المستصلحة الملحية على نمو ومحصول نباتات البصل باستخدام حمض الأسكوربيك والسماذ العضوى

أشرف شوقى عثمان ومفرح سعداوى طلبية
قسم البساتين - كلية الزراعة - جامعة الفيوم - مصر

أجريت تجربة حقلية خلال موسمى ٢٠٠٧/٢٠٠٨ و ٢٠٠٨/٢٠٠٩ م فى مزرعة كلية الزراعة - جامعة الفيوم بمنطقة دمو (أرض مستصلحة تحتوى على ملوحة قدرت عند ٧,٨٤ و ٨,٠١ ديسي سمنس / م لموسمى النمو على الترتيب)، وذلك لدراسة تأثير الإضافة الأرضية للسماذ العضوى (مخلفات الماشية) بمعدلات صفر (الكنترول)، ١٠، ٢٠، ٣٠ م^٣/فدان، والرش الورقى بحامض الأسكوربيك بمعدلات صفر (كنترول)، ١٠٠، ٢٠٠، ٣٠٠، ٤٠٠ ملجم/لتر ماء وتفاعلاتها معاً على احتمالية تحسين النمو، والمحصول والمكونات الكيماوية لنباتات البصل.

بالمقارنة بالمعدل صفر م^٣/فدان سماذ عضوى، وجد أن جميع المعدلات الأخرى قد أدت إلى زيادة معنوية فى كل من صفات النمو الخضرى (ارتفاع النبات، عدد الأوراق / نبات، الوزن الطازج للأوراق / نبات، الوزن الطازج للنبات، الوزن الجاف للأوراق / نبات، والوزن الجاف للنبات)، والوزن الجاف الكلى للنبات، الوزن الجاف للأوراق / نبات، والوزن الجاف للنبات، والوزن الجاف الكلى وبعض المكونات الكيماوية المختبرة (الكلوروفيل الكلى، الكاروتينويدات الكلية، النيتروجين والفوسفور والبوتاسيوم للأوراق و/ أو الأبصال). وتم الحصول على نفس النتائج مع معدلات حامض الأسكوربيك المستخدم بالمقارنة بالمعدلات صفر.

ومن الناحية الاقتصادية وجد أن الرش بحامض الأسكوربيك بمعدل ٣٠٠ ملجم/لتر مع إضافة السماذ العضوى للتربة بمعدل ٢٠ م^٣/ فدان هى المعاملة الأفضل وادت إلى خفض التأثيرات المثبطة للأملح على نباتات البصل.

وفى ضوء النتائج السابقة، يمكن استنتاج أن الرش بحمض الأسكوربيك بمعدل ٣٠٠ ملجم / لتر على المجموع الخضرى لنباتات البصل (صنف جيزة ٦) النامية فى تربة تم تخصيبها بالسماذ العضوى (سماذ ماشية) بمعدل ٢٠ م^٣/ فدان، يعمل على الحد من الظروف المعاكسة فى التربة، خاصة الملوحة حتى التركيز ٥٠٠٠ جزء فى المليون، وبالتالي إمكانية الحصول على محصول اقتصادى من البصل تحت هذه الظروف.