

PRODUCTIVITY AND QUALITY OF SUGAR BEET AS AFFECTED BY RATES OF POTASSIUM AND SOME MICRONUTRIENTS UNDER TWO LOCATIONS

Alaa I. Badr, N.M.M. Awad and Sahar M. Ibrahim

Sugar Crops Res. Inst., Agric. Res. Center, Giza, Egypt

(Received: Sep. 7, 2009)

ABSTRACT: *Two field experiments were carried out at two locations, (Al-Reyad, Kafr El-Sheikh Governorate (L1) and Kotour, AlGharbia Governorate (L2), Egypt) during 2006/2007 and 2007/2008 seasons, to study the effect of potassium levels (0, 24 and 48 kg K₂O/fed.) and chelated boron at two levels (½, 1 liter/fed.), and micronutrients mixture (Fe + Zn + Mn) at two levels (1.5, 2.25 liter/fed.), as well as water as control treatment.*

Root and sugar yields were significantly greater in L2 than in L1. Sucrose% was higher in L1 than L2. Root yield, sucrose %, white sugar % and sugar yield were significantly increase by adding 24 or 48 K₂O/feddan compared with control. Potassium rate had no significant effect on top yield, K content, α-amino N content, lose sugar % and purity %.

Foliar spraying with liquid chelated boron (1Liter/fed.) or chelated microelements mixture (Fe + Zn + Mn at 2.25 Liter/fed.) produced the highest root yield, top yield, white sugar % and sugar yield.

It can be concluded that application 24 Kg K₂O/fed. along with 1 Liter liquid chelated boron or 2.25 Liter chelated microelements mixture (Fe + Zn + Mn)/fed. could be recommended for optimum root yield at Al Gharbia and Kafr El-Sheikh Governorates.

Key words: *Productivity, Location, micronutrients, microelements, Sugar beet*

INTRODUCTION

Sugar beet ranks the second sugar crop not only in Egypt but also in whole around. Nowadays, fertilization takes a good place in research. Many investigators seriously tried to find out the optimum levels of potassium fertilizer and micronutrients, at different locations. The present work was performed to study the effect of some fertilizers and their levels under different locations on yield and quality of sugar beet. Badr (2009) stated that increasing potassium rate from zero to 48 kg k₂O /fed significantly increased sucrose % and suger yield/ fed . Ibrahim et al. (1988) reported that spraying sugar beet with mixture of Zn, Mn, Fe, Cu and B gave the highest top, root and sugar yields. Martens (1991) mentioned that sugar beet is moderately sensitive to deficiencies of most micronutrients. Safe (1991) found a significant response to micronutrients concentration, he mentioned that the increase in white sugar % may be attributed to decreasing impurities in roots of plants sprayed by B and Fe, and that K content in root not affected by Zn

application. Badr (1999) stated that increasing micronutrients as a foliar spray increased sucrose %, purity %, yields of root and top as well as sugar yield and decreased impurities %. Stevens *et al.*, (2004) clear that foliar applications are recommended sometimes to improve the efficiency of Zn and B assimilation. Nemeat Alla (1997) and Nemeat Alla *et al.* (2005) concluded that application of mixture of (Fe + Zn + Mn) and boron significantly increased root, top, sugar yield. Al Baddrani and Al Rashidi (2007) found that sugar beet response to fertilizer (Zn and/or B) by foliar application at two locations, where sugar percentage and sugar yield were increased by application of (Zn and/or B). Albert (2007) in Minnesota, USA, reported that root yield, root quality, recoverable sugar had no significant effect by application of micronutrients in both locations, while sugar loss to molasses significantly affected in Donaldson site only compared to Dorothy site.

MATERIALS AND METHODS

Two field experiments were carried out on clay soil at two locations (Al-Reyad city, Kafr El-Sheikh Governorate and Kotoor city, Al-Gharbia Governorate, Egypt) during 2006/2007 and 2007/2008 seasons. The preceding crop was cotton at both locations in the two seasons. Chemical analysis of soil samples taken to 30 cm depth in the experimental sites before soil preparation in 2006/07 and 2007/08 seasons are given in Table(1). Potassium sulphate (48 % K₂O) was used at rates of 0, 24 and 48 Kg K₂O/fed. on one dose during land preparation before plant.

Micronutrients solutions were used as foliar spraying as follows:

- 1- Spraying with water (control).
- 2- ½ liter liquid chelated boron (boric acid 9 % B) in 200 liters water / feddan.
- 3- 1 liter liquid chelated boron (boric acid 9 % B) in 200 liters water / feddan.
- 4- 1.5 liters mixture of liquid chelated microelements (0.5 L Fe + 0.5 L Zn + 0.5 L Mn) in 200 liters water
- 5- 2.25 liters mixture of liquid chelated microelements (0.75 L Fe + 0.75 L Zn + 0.75 L Mn) in 200 liters water.

The micronutrients mixture were contained three elements chelated in liquied form as follows:

- 1- Iron (Fe) chelated (7.15% iron oxide and 4% sulphar).
- 2- Zinc (Zn) chelated (7% zinc and 4% sulphar).
- 3- Manganese (Mn) chelated (9.03% manganese oxide and 4% sulphar).

Foliar spraing was applied on one spray after three months from sowing using hand sprayer with 200 liter/fed.

Split- plot design with three replications was used. The main plots were assigned to potassium rates and the sub-plots to micronutrients.

Productivity and quality of sugar beet as affected by

Each plot included six ridges each one 50 cm apart and 7 m long. Sowing took place on 20 of October in both seasons. Seeds of multigerm cultivar (pleno) were sown in hills 20 cm apart on one side ridges. Plants were thinned to one plant per hill at four true leaves stage in both seasons. Nitrogen fertilizer was applied as soil fertilizer in form of urea (46.54% N) with 90 kg/fed. in two equal doses, the first after thinning and the other one month later.

At maturity (210 days after sowing), the central four ridges of each plot were harvested to estimate root and top yields and technological characteristics as follows:

1. Root and top yields per plot transformed to metric ton per feddan
2. Sucrose percentage was determined using sucrometer used lead acetate according to method of [Le-Doct 1927].
3. Potassium and sodium were determined using flame photometer and α -amino nitrogen as determined using minhydrin and hidrindantin method according to (Carruther et al., 1962)
4. Purity % was calculated according to (Devillers, 1988) as follows:

$$\text{Purity \%} = 99.36 - [14.27 (V_1 + V_2 + V_3/V_4)]$$
 Where: V_1 =Sodium, V_2 =potassium, V_3 = α -aminoN, V_4 = sucrose %.
5. Sugar loss to molasses (SM), extractable white sugar % (B) were calculated according to (Devillers, 1988) as follows:
 5. a. Sugar loss to molasses = $(V_1 + V_2) 0.14 + V_1 \times 0.25 + 0.5$
 5. b. Extractable white sugar % (B) = $V_4 - SM - 0.6$ (Deviller et al., 1967).
6. Sugar yield (ton/fed.) was calculated according to the following equation:

$$\text{sugar yield} = \text{root yield} \times \text{sucrose \%}$$

Table (1): Chemical analysis of the soil experimental area at the depth 0-30cm.

| Locations | | pH 1: 2.5 | EC m mhos cm | Available nutrients ppm | | | | | | |
|-------------------|---------|--------------|--------------------|-------------------------|-----|-----|------|------|------|------|
| | | | | N | P | K | B | Fe | Zn | Mn |
| Kafr El-Sheikh | 2006/07 | 7.5 | 3.58 | 23.2 | 8.2 | 390 | 0.20 | 2.01 | 0.48 | 1.85 |
| | 2007/08 | 7.9 | 3.49 | 24.5 | 8.7 | 401 | 0.22 | 2.65 | 0.55 | 1.25 |
| Al-Gharbia | 2006/07 | 8.1 | 2.1 | 18.1 | 6.1 | 400 | 0.41 | 2.62 | 0.73 | 1.1 |
| | 2007/08 | 8.0 | 2.5 | 20.1 | 6.5 | 393 | 0.51 | 3.26 | 0.85 | 1.9 |

The analysis of variance was carried out as a combined analysis for the two locations in each season according to Gomes and Gomes (1984). Treatment means differences were compared by Duncan's Multiple Range test (Duncan, 1955). All statistical analysis was performed using analysis of variance technique by means of "MSTAT" computer software package.

RESULTS AND DISCUSSION

1. Root yield (ton/fed.):

Data in Table (2) show that root yield was greater at Ghariba location than with Kafr El-Sheikh location in the two seasons. Potassium application by 48

kg K₂O/fed. gave the highest root yield in the two seasons. Foliar spraying with chelated boron as a liquid form at the rate of 1 L/fed significantly increased root yield per feddan compared with control treatment and mixture of liquid chelated microelements (Fe + Zn + Mn) at the rate 1.5 L/fed. These results are in harmony with those obtained by Ibrahim, 1988 and Badr, 1999.

Table (2): Root and top yields as affected by locations, potassium rate and microelements spray and their interactions in combined analysis for the two locations in 2006/07 and 2007/08 seasons.

| Factors | Root yield (ton/fed.) | | Top yield (ton/fed.) | |
|--|-----------------------|-----------|----------------------|----------|
| | 2006/07 | 2007/08 | 2006/07 | 2007/08 |
| Locations (L) | * | ** | NS | NS |
| Kafr El-Sheikh | 30.461 b | 28.282 b | 8.310 | 9.394 |
| Al-Gharbia | 33.879 a | 33.611 a | 8.908 | 8.360 |
| Potassium, kg K ₂ O/fed (K) | * | * | NS | Ns |
| 0 Kg | 31.025 b | 32.607 b | 8.759 | 8.663 |
| 24 Kg | 31.938 b | 33.365 ab | 8.666 | 9.003 |
| 48 Kg | 33.547 a | 34.366 a | 9.001 | 8.966 |
| Microelements (M) | ** | * | * | * |
| control | 31.797 b | 32.707 b | 8.202 b | 8.590 b |
| Boron ½ L/Fed | 31.393 b | 33.268 ab | 8.265 b | 8.729 b |
| Boron 1 L/Fed | 34.260 a | 34.642 a | 8.648 ab | 9.572 a |
| (Fe+Zn+Mn) 1.5 L/Fed | 31.437 b | 32.683 b | 8.395 ab | 8.404 b |
| (Fe+Zn+Mn) 2.25 L/Fed | 31.964 b | 33.931 ab | 8.855 a | 9.092 ab |
| Interactions | | | | |
| L x K | * | * | NS | NS |
| L x M | * | * | NS | NS |
| K x M | * | NS | NS | NS |
| L x K x M | NS | NS | NS | NS |

Means of each factor designated by the same letter are not significantly different at 5% level using Duncan's multiple range test.

The interaction between location and potassium rate had a significant effect on root yield per feddan in both seasons (Table 3). Root yield was significantly higher in El-Gharbia site than in Kafr El-Sheikh site at any potassium rate in both seasons. The rate 48 Kg K₂O/fed. significantly increased root yield compared with control at the two locations in both seasons. Application of potassium at 24 or 48 Kg K₂O/fed did not differ in root yield at El-Gharbia site in both seasons. However, no significant difference between 0 and 24 kg K₂O/fed in root yield was recognized in Kafr EL-Sheikh in both seasons.

The effect of interaction between location and microelements was significant on root yield in the two seasons (Table 4). The highest root yield was obtained from foliar spraying with liquid chelated boron at the rate 1 L/fed at Gharbia site in both seasons. Mixture of microelements at the rate 2.25 L/fed did not differ from that at 1 L boron/fed in root yield at Gharbia site. No significant differences among all microelements treatments in root yield were observed at Kafr El-Sheikh site in the second seasons.

Productivity and quality of sugar beet as affected by

Table (3): Root yield (ton/fed.) as affected by the interaction between location and potassium rate in 2006/07 and 2007/08 seasons.

| Potassium K ₂ O/fed | 2006/07 | | 2007/08 | |
|-----------------------------------|----------------|------------|----------------|------------|
| | Kafr El-Sheikh | Al-Gharbia | Kafr El-Sheikh | Al-Gharbia |
| 0 kg | 29.065 c | 32.985 ab | 27.647 c | 37.567 b |
| 24 kg | 29.971 c | 33.905 ab | 28.063 c | 38.667 ab |
| 48 kg | 32.347 b | 34.747 a | 29.135 c | 39.597 a |

Table (4): Root yield (ton/fed.) as affected by the interaction between location and microelements spray in 2006/07 and 2007/08 seasons.

| Microelements | 2006/07 | | 2007/08 | |
|-----------------------|----------------|------------|----------------|------------|
| | Kafr El-Sheikh | Al-Gharbia | Kafr El-Sheikh | Al-Gharbia |
| control | 30.470 def | 33.123 bcd | 28.120 c | 37.293 b |
| Boron ½ L/Fed | 30.873 def | 31.913 cde | 27.750 c | 38.787 ab |
| Boron 1 L/Fed | 32.640 bcd | 35.880 a | 29.360 c | 39.923 a |
| (Fe+Zn+Mn) 1.5 L/Fed | 29.090 f | 33.783 abc | 27.810 c | 37.557 ab |
| (Fe+Zn+Mn) 2.25 L/Fed | 29.232 ef | 34.697 ab | 28.369 c | 39.493 ab |

The interaction between potassium and microelements rate had a significant effect on root yield in the first season, only (Table 5). Application of 48 Kg K₂O/fed and 1 L boron produced the highest root yield. No significant differences in root yield were observed among the mentioned combination and 24 Kg K₂O/fed + 1 L boron or 24 Kg K₂O + 2.25 L microelements mixture. Higher response in the root yield of sugar beet, namely increased root size or early growth, to fertilizer (B and /or Fe,Zn,Mn) application was found in the soil of Al- Gharbia site compared to Kafer El-Sheik site.

Table (5): Root yield (ton/fed.) as affected by the interaction between potassium rate and microelements in 2006/07 season.

| Microelements | potassium rate (K ₂ O/fed) | | |
|-----------------------|--|------------|-------------|
| | Zero | 24 Kg/fed. | 48 Kg /fed. |
| | 2006/07 | | |
| control | 30.128 c | 30.683 c | 34.578 ab |
| Boron ½ L/Fed | 30.708 c | 31.288 bc | 32.183 abc |
| Boron 1 L/Fed | 32.843 abc | 34.863 a | 35.073 a |
| (Fe+Zn+Mn) 1.5 L/Fed | 30.503 c | 31.078 bc | 32.728 abc |
| (Fe+Zn+Mn) 2.25 L/Fed | 30.943 c | 31.778 abc | 33.172a bc |

2. Top yield (ton/fed.):

Data in Table (2) pointed out that locations, and levels of k-fertilizer are not significantly affected on top yield in both seasons. At beet plants sprayed with 2.25 liter/fed. of mixture in the first season and with 1 liter/fed.

of boron in the second season, only, gave the highest top yield compared to other treatments spray. Maximum top yield may be attributed to increase doses of both micronutrients and K₂O fertilizers These results are in a good agreement with those obtained by Ibrahim *et al.* (1988) and Nemeat Alla *et al.* (2005). There was no evidence for significant differences in top yield due to the interaction between all treatments in both seasons.

3. Sucrose percentage:

As for the effect of locations, k-fertilizer and microelements, and its levels on sucrose % Table (6) pointed out that significant differences were observed between mean values of sucrose percentage attributed to mentioned factors. Sucrose percentage had a significant difference due to sites difference in the second season, only. The percentage of sucrose was at kafr Elshikh site higher than EL Gharbia site, This resulted from kafr Elshikh gave smallest root size and lowest root yield than EL Gharbia and the small size of sugar beet have high sugar content than big root size.

Maximum sucrose percentage was obtained when sugar beet plants were fertilized by 24 or 48 kg K₂O/fed in the first, only season compared with zero K. These findings are in agreement with the previous results obtained by Badr (2009). Foliar application of microelements resulted significant differences in sucrose % in both seasons. Foliar spraying with liquid chelated boron(1L/fed) in the first season or chelated microelements mixture (Fe + Zn + Mn at 1.5 or 2.25 L/fed.) in the second season produced the highest sucrose %. Similar results were obtained by Albert (2007), and Al Baddrani and Al Rashidi(2007).

There was no evidence for significant difference in sucrose % due to the interaction between both factors in the two seasons.

4. Soluble non-sugars:

The soluble non- sugars K , Na and α -amino N in the roots are regarded as impurities because they interfere with sugar extraction. Means of these impurities as effected by locations, potassium fertilizer and foliar spraying of microelements in 2006/2007 and 2007/2008 seasons are presented in Table(6).

Data confirmed pointed out that locations had highly significant effect on root content from potassium. Sugar beet planted in Kafr El-Sheikh site gave the highest concentration of K in beet juice in both seasons than Gharbia site which recorded the lowest ones. Levels of k-fertilizer not significantly affected on this trait in both seasons.

All impurities were significantly increased by increasing foliar spray in both seasons. Increasing K levels decreased root content from Na this due to the fact of antagonism between Na and K.

Table (6): Sucrose, concentration of potassium , sodium , α amino N and loss sugar in root juice as affected by locations, potassium rate and microelements spray and their interactions in combined analysis for the two locations in 2006/07 and 2007/08 seasons.

| Factor | Sucrose % | | Potassium meq/100gm | | Sodium meq/100gm | | α -amino N meq/100gm | | Loss sugar % | |
|-----------------------------------|-----------|---------|---------------------|---------|------------------|---------|-----------------------------|---------|--------------|---------|
| | 2006/07 | 2007/08 | 2006/07 | 2007/08 | 2006/07 | 2007/08 | 2006/07 | 2007/08 | 2006/07 | 2007/08 |
| Locations (L) | NS | ** | * | ** | NS | NS | NS | NS | * | * |
| Kafr El-Sheikh | 16.6 | 17.9 a | 6.9 a | 7.2 a | 1.8 | 2.2 | 2.5 | 2.7 | 2.3 a | 2.5 a |
| Aj-Gharbia | 17.0 | 17.0 b | 5.2 b | 5.3 b | 1.4 | 1.8 | 2.4 | 2.7 | 2.0 b | 2.2 b |
| Potassium:K ₂ O/fed(K) | * | NS | NS | NS | ** | ** | NS | NS | NS | NS |
| 0 kg | 16.3 b | 17.0 | 6.0 | 6.2 | 1.9 a | 2.1 a | 2.4 | 2.7 | 2.2 | 2.3 |
| 24 kg. | 17.1 a | 17.5 | 6.0 | 6.2 | 1.8 a | 2.1 a | 2.5 | 2.7 | 2.2 | 2.4 |
| 48 kg. | 17.0 a | 17.6 | 6.2 | 6.4 | 1.4 b | 1.6 b | 2.4 | 2.7 | 2.4 | 2.3 |
| Microelements (M) | * | * | * | * | * | * | * | NS | NS | NS |
| control | 16.6 b | 17.0 b | 6.1 b | 6.1 b | 1.8 a | 2.2 a | 2.6 a | 2.6 | 2.1 | 2.1 |
| Boron ½ L/Fed | 16.5 b | 17.1 b | 5.6 c | 5.7 c | 1.5 b | 1.8 b | 2.2 b | 2.8 | 2.1 | 2.2 |
| Boron 1L/Fed | 17.0 a | 17.2 ab | 5.7 c | 6.1 b | 1.5 b | 2.0 ab | 2.3 b | 2.8 | 2.3 | 2.3 |
| (Fe+Zn+Mn)1.5 L/Fed | 16.9 a | 17.5 a | 6.7 a | 5.8 a | 1.5 b | 2.0 ab | 2.3 b | 2.6 | 2.3 | 2.4 |
| (Fe+Zn+Mn)2.25L/Fed | 16.9 a | 17.6 a | 6.5 a | 6.6 a | 1.6 ab | 2.0 ab | 2.5 ab | 2.6 | 2.4 | 2.4 |
| Interactions | | | | | | | | | | |
| L x K | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| L x M | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| K x M | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| L x K x M | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |

Means of each factor designated by the same letter are not significantly different at 5% level using Duncan's multiple range test.

Foliar spraying with chelated boron as a liquid form at the rate 0.5 L/fed significantly decreased impurities in root juice compared with control treatment or mixture of liquid chelated microelements in both seasons. Nemeat Alla (1997) reported that concentration of K decreased with foliar spraying, Saif (1991) stated that K content in root not affected by Zn application.

It is clear that application of k-fertilizer and microelements spraying recorded highly significant effect on concentration Na in root content. With increasing levels of all fertilizers due to significantly decrease in sodium root content. Badr (1999) concluded that application of (Fe + Zn + Mn) not affected on Na content ratio.

The results showed that the lowest content from α -amino N in root juice was with micronutrients spraying in the first seasons, only.

All interaction effects among location, K-fertilizer and micronutrients spraying on this trait was not significant in both seasons.

5. Loss sugar %:

Data presented in Table (6) appeared that significant differences were observed among values of loss sugar percentage in both seasons due to locations effect. The highest losses in sugar was found in Kafr El-Sheikh site more than in Gharbia site. These high values related with high root content from impurities (K+ Na+ α -amino N) . Albert (2007) stated that loss sugar in molass was increased by increasing B, Fe, Zn and Mn spraying in the two locations. No significant interaction effects were found between all factors under study in both seasons.

6 - Extractable white sugar %

Results in Table (7) showed alternative effect between all factors under study on extractable white sugar % in both seasons. Locations recorded highly significant differences among values of extractable white sugar % in the first season, only and gave maximum value in Gharbia site.

Extractable white sugar % at 24 Kg K₂O/fed did not differ from that at 48 K₂O/fed compared with zero treatment in both seasons. The results not differ than 48 K₂O/fed compared with zero treatment in both seasons.

Application of Microelements had a significant effect on extractable white sugar % in both seasons. The highest of white sugar % was obtained by application mixture of microelements (Fe+Zn+Mn) at the rate 2.25 L/fed, which did not differ from that at 1 L boron/fed in both seasons.

The increase in white sugar % may be attributed to decreasing impurities in root. Similar results were obtained by saif (1991).

No significant interaction effects were found between both factors in this respect in the two seasons. .

Productivity and quality of sugar beet as affected by

Table (7): Extractable white sugar %, purity % and sugar yield ton/fed. as affected by locations, potassium rate and microelements spray and their interactions in combined analysis for the two locations in 2006/07 and 2007/08 seasons.

| Factors | White sugar % | | Purity % | | Sugar yield (ton/fed) | |
|-------------------------------------|---------------|---------|----------|---------|-----------------------|----------|
| | 2006/07 | 2007/08 | 2006/07 | 2007/08 | 2006/07 | 2007/08 |
| Locations (L) | ** | NS | ** | ** | * | ** |
| Kafr El-Sheikh | 13.7 b | 14.5 | 89.5 b | 89.6 b | 4.176 b | 4.106 b |
| Al-Gharbia | 14.9 a | 14.4 | 91.9 a | 91.8 a | 5.001 a | 5.572 a |
| Potassium :K ₂ O/fed (K) | * | * | NS | NS | * | * |
| 0 | 13.6 b | 14.1 b | 90.4 | 90.3 | 4.236 b | 4.609 b |
| 24 kg. | 14.9 a | 14.5 a | 90.6 | 90.3 | 4.748 a | 4.846 ab |
| 48 kg. | 14.2 a | 14.7 a | 90.8 | 90.5 | 4.781 a | 5.061 a |
| Microelements (M) | * | * | * | NS | * | * |
| Control | 13.6 b | 14.3 b | 90.3 b | 90.5 | 3.637 ab | 4.743 b |
| Boron ½ L/Fed | 13.9 b | 14.3 b | 91.1 a | 90.6 | 4.380 b | 4.735 b |
| Boron 1 L/Fed | 14.4 a | 14.4 ab | 91.1 a | 90.4 | 4.943 a | 4.990 a |
| (Fe+Zn+Mn) 1.5 L/Fed | 13.9 b | 14.5 ab | 89.8 c | 90.1 | 4.387 b | 4.747 b |
| (Fe+Zn+Mn) 2.25 L/fed | 14.3 a | 14.6 a | 90.4 b | 90.4 | 4.597 ab | 4.980 a |
| Interactions | | | | | | |
| L x K | NS | NS | NS | NS | * | * |
| L x M | NS | NS | NS | NS | * | * |
| K x M | NS | NS | NS | NS | * | * |
| L x K x M | NS | NS | NS | NS | NS | NS |

Means of each factor designated by the same letter are not significantly different at 5% level using Duncan's multiple range test.

7-Juice purity percentage:

The results obtained in Table (7) showed that locations had highly significant effect on juice purity % and gave the highest values in Gharbia site in both seasons. Potassium fertilization had no significant effect on this trait in both seasons. Microelements spraying recorded significant effect on juice purity % in the first seasons, only. Boron element recorded the highest values of purity % when sugar beet plants were sprayed by it. This may be due to decrease impurities in root juice. While the lowest value of purity % was obtained when mixture of (Fe, Zn and Mn) was applied to sugar beet plants at 1.5 L/fed. in the first season. These findings are in agreement with the previous results obtained by Al-Baddrani 2007.

No significant interaction effects were recognized between both factors in this trait in both seasons.

8-Sugar yield (ton/fed.):

Concerning to sugar yield data in Table (7) revealed that locations exhibited significant differences in the two seasons. Results showed that the highest sugar yield was obtained in Gharbia site in both seasons compared to the another site. The superiority of sugar yield in Gharbia may be attributed to increase in root yield in both seasons. Sugar yield per faddan

was increased by increasing K rate to 48 Kg K₂O/ fed in both seasons. Similar results were obtained by Badr (2009). Beet plants sprayed by boron 1 L/ fed or mixture 2.25 L/ fed produced the same sugar yield in the two seasons. Such increase in sugar yield was obtained from increasing rate of microelements spraying may be attributed to the increase in root yield. Nemeat Alla 1997 and Nemeat Alla 2005 reported that application of boron and micronutrients increased sugar yield.

The effect of interaction between locations and K rate was significant on sugar yield per feddan in both seasons (Table 8). The highest sugar yield resulted from 48 kg K₂O/fed. at Al Gharbia site in both seasons. In (Table 9) there was significant difference in sugar yield due to interaction between locations and microelements, whereas the highest values of sugar yield were at Al Gharbia site with spraying of mixture at 2.25 L /fed in the two seasons.

The interaction between K rate and microelements spraying had a significant effect on sugar yield in both seasons (Table10). Applying 24 or 48 kg K₂O /fed., with born at 1 L/ fed recorded the highest sugar yield per feddan in both seasons. These results indicate a positive association between sucrose % and root yield. Stevens *et al.*, 2004 clear that foliar application are recommended sometimes to improve the efficiency of Zn and B assimilation.

It can be concluded that rate of 48 kg K₂O/fed., foliar spraying of chelated Boron as a liquid form at 1 L/fed and mixture of chelated microelements (Fe+Zn+Mn) at 2.25 L/fed could be recommended for optimum root and extractable white sugar yields feddan area at Kafr El- Sheikh and Al- Gharbia Governorates.

Table (8): Sugar yield (ton/fed.) as affected by the interaction between location and potassium rate in 2006/07 and 2007/08 seasons.

| Potassium rate | 2006/07 | | 2007/08 | |
|-----------------------------|----------------|------------|----------------|------------|
| | Kafr El-Sheikh | Al-Gharbia | Kafr El-Sheikh | Al-Gharbia |
| Zero | 3.681 d | 4.791 b | 3.921 d | 5.297 b |
| 24 kg K ₂ O/fed. | 4.443 c | 5.053 ab | 4.103 cd | 5.589 ab |
| 48 kg K ₂ O/fed. | 4.403 c | 5.159 a | 4.293 c | 5.829 a |

Table (9): Sugar yield (ton/fed.) as affected by the interaction between location and microelements spray in 2006/07 and 2007/08 seasons.

| Microelements | 2006/07 | | 2007/08 | |
|-----------------------|----------------|------------|----------------|------------|
| | Kafr El-Sheikh | Al-Gharbia | Kafr El-Sheikh | Al-Gharbia |
| control | 4.527 cd | 4.747 bcd | 4.067 c | 5.420 b |
| Boron ½ L/Fed | 4.200 de | 4.560 cd | 4.033 c | 5.437 b |
| Boron 1 L/Fed | 4.603 cd | 5.283 ab | 4.273 c | 5.707 ab |
| (Fe+Zn+Mn) 1.5 L/Fed | 3.697 e | 5.077 abc | 4.033 c | 5.460 b |
| (Fe+Zn+Mn) 2.25 L/Fed | 3.853 e | 5.340 a | 4.123 c | 5.837 a |

Productivity and quality of sugar beet as affected by

Table (10): Sugar yield (ton/fed.) as affected by the interaction between potassium rate and microelement spray in 2006/07 and 2007/08 seasons.

| Microelements | 2006/2007 | | | 2007/2008 | | |
|-----------------------|----------------|--------------------------------|--------------------------------|-----------|--------------------------------|--------------------------------|
| | potassium rate | | | | | |
| | zero | 24 kg K ₂ O/fed. | 48 kg K ₂ O/fed. | zero | 24 kg K ₂ O/fed. | 48 kg K ₂ O/fed. |
| control | 4.048 c | 4.803 abc | 5.058 ab | 4.238 e | 4.838 a-d | 5.153 ab |
| Boron ½ L/Fed | 4.103 c | 4.538 abc | 4.498 abc | 4.518 de | 4.748 a-d | 4.938 a-d |
| Boron 1 L/Fed | 4.618abc | 5.178 a | 5.033 ab | 4.793 a-d | 4.998 abc | 5.178 a |
| (Fe+Zn+Mn) 1.5 L/Fed | 4.078 c | 4.508 abc | 4.573 abc | 4.608 cde | 4.718 bcd | 4.913 a-d |
| (Fe+Zn+Mn) 2.25 L/Fed | 4.333 bc | 4.713 abc | 4.743 abc | 4.888 a-d | 4.928 a-d | 5.123 ab |

REFERENCES

Al-Baddrani, W. and S. Al-Rashidi (2007). The effect of zinc on quality of sugar beet (*Beta vulgaris* L.). Department of Soil and Water sciences, College of Agriculture and Forestry, Mosul University, IRAQ. Crops. Org/Zn Crops 2007/PDF/2007, 19 K-adobe PDF-Afficher en html.

Albert, L. Sims (2007). Sugar beet production on sandy soils: The need for non-traditional nutrients WWW. sbreb, org./research/soil 07/sandy soils.pdf.

Badr, A.I. (1999). Effect of ploughing depth and fertilization on yield and quality of sugar beet. M.Sc. Thesis, Fac. of Agric. Al-Azhar Univ. Egypt.

Badr, A.I. (2009). Response of some sugar beet cultivars to planting methods and potassium fertilizer. J. Agric. Res. Kafr El- Sheikh Univ. , 35(1) 2009.

Carruther, A., J.V. Dutton and J.F.T. Oldfield (1962). Chromogenic reducing substances in molasses. International Sugar Journal, 65, 297-301, 300-4.

Devillers, P. (1988). Prevision d sucre melasse sucrerir France 190-200 (C.F. the Sugar Bet Crop. Book).

Duncan, B. D. (1955). Multiple rang and multiple F-test Biometrics 11: 1- 42.

Gomes, K.A. and A.A. Gomes (1984). Statistical procedures for agricultural research. An International Rice Research Institute Book John Willey and Sons. Inc., New York.

Ibrahim, M.H., S. Gh R. Sorour and A.M. Omar (1988). Varietal response of sugar beet to foliar application of some micronutrients. Proc. 3rd Egyptian Conf. Agron. Kafr El-sheikh, 5-7 Sept. 1988, Vol. 11: 324334.

Le-Doct, A. (1927). Commercial determination of sugar in beet root using the aches-Le-Doct Int. Sugar J., 29 = 488-492.

Martens, D.C. and D.T. Westermann (1991). Fertilizer applications for correction micronutrient deficiencies. pp. 549-592. IN: Mortredt, J.J. (ed.) Micronutrients in Agriculture SSSA Book Ser. 5. SSSA, Madison, WI.

Nemeat Alla, S.A. (1997). Agronomic studies on sugar beet. PH.D. Thesis, Fac. of Agric. Kafr El-Sheikh, Tanta Univ. Egypt.

Nemeat Alla, S.A. and G.A. Samia (2005). Agronomical and statistical studies on the response of sugar bet to foliar application with micronutrients

Alaa I. Badr, N.M.M. Awad and Sahar, M. Ibrahim

- under different macronutrients formula. J. Agric. Sci. Mansoura Univ., 30(12): 7265-7284, 2005.
- Saif, L.M. (1991). Yield and quality of sugar beet affected by nitrogen sources and rate of some microelements in Kafr El-Sheikh. Ph.D. Thesis, Fac. of Agric. Ain Shams Univ. Egypt.
- Stevens, W.B., J.G. Davis and J. Blumenthal (2004). Nutrient management in dry bean production and post management. Regional Bull. 562 A. Colo. St. Univ., Univ. Neb. and Univ. Wyo. In Press.

تأثير السماد البوتاسي وبعض العناصر الصغرى على إنتاجية وجودة بنجر السكر في موقعين

علاء ابراهيم بدر ، نبيل مرسى محمد عوض ، سحر سامون ابراهيم
معهد بحوث المحاصيل السكرية - مركز البحوث الزراعية - جيزة - مصر

الملخص العربي

أجريت تجربتان حقليتان في كل موقع من موقعي الدراسة (مركز الرياض - محافظة كفر الشيخ)، (مركز قطور - محافظة الغربية) في موسمي ٢٠٠٦/٢٠٠٧ م، ٢٠٠٧/٢٠٠٨ م لدراسة تأثير معدلات التسميد البوتاسي (صفر - ٢٤ - ٤٨ كجم بوا/فدان) ورش بعض العناصر الصغرى المخليبية وهي البورون بمعدلات ١/٢ - ١ لتر/فدان ومخلوط العناصر الصغرى (حديد - زنك - منجنيز) بمعدلات ١,٥ - ٢,٢٥ لتر/فدان وقد استخدم التصميم الإحصائي قطع منشقة مرة واحدة في ثلاث مكررات. حيث تم توزيع التسميد البوتاسي في القطع الرئيسية ورش العناصر الصغرى في القطع الشقية. وتم استخدام التحليل المشترك بين المواقع في كل موسم. وكانت اهم النتائج ما يلي:

- زاد محصول الجذور والسكر معنويا في موقع محافظة الغربية عما في موقع محافظة كفر الشيخ بينما ارتفعت نسبة السكر في محافظة كفر الشيخ عنها في محافظة الغربية.
- ادى استخدام التسميد بالبوتاسيوم بمعدل ٢٤ او ٤٨ كجم بوا/ف الى زيادة محصول الجذور ونسبة السكر ونسبة السكر الابيض و محصول السكر وذلك مقارنة بمعاملة الكنترول. وادى الرش الورقي بالبورون عند المعدل ١ لتر/ف الى ائصول على قيم لمعظم الصفات إذا ما قورن بباقي العناصر.
- تساوى تأثيرى رش عنصر البورون المخلبي بمعدل ١ لتر للفدان مع رش مخلوط العناصر الصغرى بمعدل ٢,٢٥ لتر/فدان على بعض صفات بنجر السكر، فقد تبين من النتائج ان هناك زيادة معنوية في محصول العرش والجذور ونسبة السكر الابيض و محصول السكر نتيجة اضافة هذه العناصر الغذائية.

- ويستنتج من هذا البحث ان يوصى باضافة ٨ كجم بوز/أف مع الرش الورقى بالبورون المخلبى بمعدل التتر/قدان ومخلوط العناصر الصغرى المخلبية (حديد - زنك - منجنيز) بمعدل ٢,٢٥ لتر/قدان للحصول على اعلى محصول جذور ومحصول سكر من بنجر السكر فى محافظتى كفر الشيخ والغربية.