

EFFECT OF SOME PLANT EXTRACTS AND PARTIAL SUBSTITUTION OF THIOVIT 80 BY PLANT EXTRACT ON POWDERY MILDEW DISEASE OF SUGAR BEET

El-Fahar, Samia A.¹ and Sahar M.E. Moustafa²

1. Sugar Crops Pathology Dept., Sugar Crops Institute, ARC, Egypt

2. Sugar Technology Dept., Sugar Crops Institute, ARC, Egypt

ABSTRACT

Four experiments were conducted during 2004/2005, 2005/2006, 2006/2007 and 2007/2008 seasons at Sakha Agricultural Research Station, Kafr El-Sheikh, Egypt to study the effect of six plant extracts on controlling powdery mildew disease of sugar beet caused by *Erysiphe polygoni*, and its effect on different characters. Six plant extracts were used i.e. *Solanum nigrum*, *Pancratinum maritimum*, *Melia azedarach*, *Anthemis nobilis*, *Ammi visnaga* and *Mentha piperita* in different concentrations, i.e. 1000, 3000 and 5000 ppm in the screen house on two sugar beet cvs. viz. Sultan and Glorious, under artificial inoculation by conidio-spores of *E. polygoni* combinations between *M. azedarach* and Thiovit 80 were used, $\frac{1}{2}$ *M. azedarach* + $\frac{1}{2}$ Thiovit 80, $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80 and $\frac{1}{4}$ *M. azedarach* + $\frac{3}{4}$ Thiovit 80. Those experiments were conducted in the screen house during 2004/2005 and 2005/2006. The combination was applied 3 days before and after inoculation.

Data show that *M. azedarach* at the level of 5000 ppm gave the best disease control before and after inoculation for powdery mildew followed by *Ammi visnaga*. Data showed also that, mixing fungicide with plant extract lead to increasing efficiency of plant extract when $\frac{1}{4}$ plant extract + $\frac{3}{4}$ Thiovit 80, $\frac{1}{2}$ plant extract + $\frac{1}{2}$ Thiovit 80, $\frac{3}{4}$ plant extract + $\frac{1}{4}$ Thiovit 80, efficiency were 93.3, 89.9 and 84.5% before inoculation and 86, 77.5 and 69.9% after inoculation in comparison with the efficiency of Thiovit 80 before inoculation (94.7) and after inoculation (90.1%) for Sultan cv. while efficiency for Glorious the date showed the same trend as 93.5, 90.2 and 84.4% before inoculation, on the other hand, it recorded 87.4, 78.9 and 70.1% after inoculation for the three combinations, respectively. While, Thiovit 80 recorded efficiency of 95.6% before inoculation and 91.2% after inoculation.

In the field experiment, *M. azedarach* (5000 ppm) and the combination $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80, was used because it has high efficiency for controlling powdery mildew disease as well as to reduce pollution and costs.

During 2006/2007, 2007/2008 seasons, disease severity (%) was reduced to more than 50% by utilization of $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80, efficiency reached to 68.1% for Sultan and 70.1 % for Glorious during 2006/2007.

Data also showed that chlorophyll content was high, 67.5 and 77.4 for Sultan and Glorious cvs. All studied characters were affected, like, root weight, TSS%, sucrose (%), purity and increase percentages. Increase % in root weight 61.1 and 58.8% for Sultan and Glorious, respectively. While, increase % in sucrose; was 62.6 and 37.6% for Sultan and Glorious, respectively.

Phenolic compounds showed negative correlation between disease severity and phenolic compounds.

AUDPC affected with disease severity (%), when disease severity increased, AUDPC increased.

So, this research paper pointed out to the possibility of replacing fungicides by plant extracts of for disease control or of foliage diseases of sugar beet, either alone or mixed with Thiovit 80 to reduce the costs and environmental pollution as well as preventing residual effect of the fungicide on produced sugar.

INTRODUCTION

Sugar beet powdery mildew, caused by the fungus *E. ptygoni* now occurs in all sugar beet growing areas and can reduce sugar yield up to 30%. Many crops are infected by similar appearing powdery mildews. Moreover, if the disease is not controlled it can cause a 20 to 35 percent loss in sugar yield. Crop loss is due to reduced root yield and often to a lower concentration of sugar in roots. Both effects apparently are due to a reduced efficiency of diseased leaves and to their premature death when roots are rapidly enlarging (Grimmer *et al.*, 2007). Sugar beet leaves infected with powdery mildew show declining rates of net photosynthesis as the disease develops; relative to healthy controls, reductions of 75% after inoculation (Gordon and Duniway, 1981).

The disease is favored by a warm, dry climate, like Egypt, in addition, failure of the disease to develop on young plants in the field appears to be due to an increased susceptibility of leaves as they grow older, rather than to a microclimate effect within the foliage canopy that might be more favorable for spore germination and growth. It is common to see abundant signs of the mild on the stunted and open leaf canopy of nitrogen deficient plants (Paulus, 2008). In recent study of El-Fahar, 2008, losses in sucrose could be reached to 82.9% for some cvs. due to powdery mildew infection. Controlling powdery mildew disease is the only solution where the resistant variety is not available.

Minimizing chemical application is the main target of the environmental protective people to have an environment free from pollution and chemical hazards, and looking forward to a clean cultures for feed and food. Fungicides were used extensively for controlling most of the susceptible sugar beet cvs. sulfur (Thiovit 80) was applied for controlling powdery mildew (Karaoglanidis and Karadimos, 2005). Use of botanical pesticides (natural plant products) in an agroecosystem is now emerging as one of the prime means to protect crop production and environment from pesticidal pollution. There is practically no risk of developing pest resistance to these products when used in natural forms (Prakash and Rao, 1997). Plant extracts were frequently reported to be fungi toxic to various fungi (Shimon *et al.*, 1993). So this research paper was undertaken to study the efficacy of some plant extracts as well as some combinations between the recommended fungicide and the selected plant extract on controlling powdery mildew disease. In this respect, screen house and field experiments were carried out.

MATERIALS AND METHODS

Plant materials:

a. Sugar beet cultivars:

Two sugar beet cvs. i.e. Sultan and Glorious were selected according to the previous study of El-Fahar (2008) to perform the experiment either in screen house or in the field during the course of this study.

b. Higher plants extract:

Six higher plant samples/parts were used to test their antifungal effects on *E. polygoni* through preparing plant extract solutions. These plants are; *Solanum nigrum* (Black nightshade), *Pancreatinum maritimum* (Soosan bulbs), *Melia azedarach* (Neem), *Anthemis nobilis* (Chamomile), *Ammi visnaga* (Pick tooth or Khella), and *Mentha piperite* (Mente).

Samples of plants shown in Table (1) were collected from different regions of Egypt and were identified by specialists. Extraction procedures were applied according to Scott and Mickibben, 1978 and Ashry *et al.*, 1999.

The plant extractions were diluted to get a series of concentrations i.e., 1000, 3000 and 5000 ppm (mg).

Table (1): Some higher plants screened for their inhibitory effect against powdery mildew disease of sugar beet.

No.	Scientific name	Common name	Arabic name	Part used	Family	Solvent system or extractives
1	<i>Solanum nigrum</i>	Black night shade	عنب الذيب	Leaves	Solanaceae	Methanol/acetone (1: 1) water
2	<i>Pancreatinum maritimum</i>	Soosan bulbs	نرجس بلطيم	Bulbs	Amaryllidaceae	Ethanol/petroleum ether
3	<i>MELia azedarach</i>	Neem	زرنلخت	Fruits	Meliaceae	Petroleum ether (40-60)
4	<i>Anthemis nobilis</i>	Chamomile	شوح	Leaves	Anacardiaceae	Acetone
5	<i>Ammi visnaga</i>	Khella (Pick tooth)	خله	Seeds	Umbelliferaceae	Acetone
6	<i>Mentha piperita</i>	Mente	أوراق النعنع	Leaves	Myrtaceae	Volatile oil

Extraction procedure:

Fresh plant materials were washed, dried in an oven at 40°C and ground to fine powder. The dried powder of each plant material was extracted, almost to exhaustion in host extraction apparatus (Soxhlet) with the specified solvent systems, presented in Table (1). The extracts were filtered, dried over anhydrous sodium sulphate and evaporated to dryness under reduced pressure by a rotary evaporator. The residue was weighted and dissolved in acetone to give the desired concentration. Volatile oils were isolated from plant materials by stem distillation using Clevenger trap. Total alkaloids were isolated from fruits of black pepper by the method described by Scott and McKibben, 1978.

2. Test organism

Artificial inoculation was done by spore suspension of *E. polygoni*, 60 days from planting and was prepared and sprayed at a rate of 40×10^3 conidio-spores/ml on leaves (El-Fahar, 2008), relative humidity was kept high by using plastic sheets. Scoring and disease assessment was done according to Hills *et al.* (1980) scale.

3. Experimental design:

a. Screen house experiments:

Screen house experiment was conducted at Sakha Agricultural Research Station, Kafr El-Sheikh, Egypt during 2004/2005 and 2005/2006 seasons. Microplots of 2 x 10 m² was used. Each cultivar was grown in 3 rows of 8 plants, keeping 60 cm and 20 cm between rows and plants within row, respectively. Split split plot design with three replicates was used. Cultivars were allocated in the main plot, plant extracts were allocated in the sub plots concentrations of plant extract were allocated in sub-sub plots. Plant extracts were evaluated for their efficiency against *E. polygoni* as follow:

The six tested plant extracts were used in three concentrations i.e., 1000, 3000 and 5000 ppm during 2004/2005 and 2005/2006 seasons. Plant extracts were applied 3 days before or after artificial inoculation with *E. polygoni* conidio-spores. The fungicide (Thiovit 80) was applied in recommended dose (2.5 gm/liter) 3 days before or after inoculation. During 2005/2006 season, out of the six plant extracts, only *M. azedarach* was chosen with full dose (5000 ppm), ¾, ½ and ¼ dose mixed with ¼, ½, ¾ dose of chemical fungicide (Thiovit 80), the score of disease severity were recoded after 90 days from sowing, 8 readings were scored with 15 days intervals.

b. Field experiments:

Two field experiments were conducted at Sakha Agricultural Research Station Farm during 2006/2007 and 2007/2008 seasons. A split plot design with three replicates was used. Main plots were devoted to cultivars (Sultan and Glorious), while sub plots were occupied by treatments (*M. azedarach*, *M. azedarach* (3/4) + Thiovit 80 (¼) and Thiovit 80 (2.5 gm/L). Some plots were left for natural infection by fungus without spraying.

Plot size was kept at 3 x 7 m², keeping 50 cm between rows and 20 cm within row. Date of sowing was adjusted on October 15 for both seasons.

All the recommended cultural practices for the experiment were applied according to the recommendation of sugar beet crop in Egypt.

Different measures and traits were measured as follow:

1. Disease severity of *E. polygoni* was recorded when the infection started to appear and 8 readings were recorded at 15 days intervals up to harvest. Disease assessment was done according to Hills *et al.* (1980) scale.
2. Efficiency % = $\frac{\text{Disease severity of control} - \text{disease severity of treatment}}{\text{Disease severity of control}} \times 100$
3. Total soluble solids percentage: was determined in fresh roots for each cultivar using hand refractometer (McGinnis, 1982).
4. Sucrose (%): sucrose percentage was estimated according to A.O.A.C. (1990).
5. Purity (%): It was calculated by dividing percentage of sucrose on total soluble solids (TSS).
6. Root weigh t(kg/plant), the two central ridges of each plot were estimated in Kilograms.
7. The increase (%) due to each treatment over the control = $\frac{\text{Treatment} - \text{control}}{\text{Treatment}} \times 100$.
8. Phenolic compounds:

- a. Free phenols: were determined using folin Ciocalteu reagent described by Bray and Thrope (1954).
 - b. Conjugated phenols:
Conjugated phenols = total phenols-free phenols.
 - c. Total phenols: were determined in the ethanolic extract of sugar beet leaves using the method described by Snell and Snell (1953).
9. Area Under Disease Progress curve (AUDPC):
AUDPC was calculated as according to Pandey *et al.* (1989) as follow:

$$AUDPC = D \left[\frac{1}{2} (Y_1 + Y_k) + (Y_2 + Y_3) \dots Y_{(k-1)} \right]$$

Where:

$Y_1, Y_2 \dots Y_k$ are the K disease score at constant interval of D days.

10. Chlorophyll content: Total chlorophyll content was determined in mg by using chlorophyll meter (SPAD-502) by taking the average of 5 readings in each record for each of the four replicates on the leaves directly.

Statistical analysis was done according to Gomez and Gomez (1983).

RESULTS AND DISCUSSION

a. Screen house:

1. Effect of certain plant extracts on powdery mildew incidence:

The effect of the six plant extracts at three concentrations, i.e. 1000, 3000 and 5000 ppm for each on disease severity of *E. polygoni* was studied in microplots during 2004/2005 season. The fungicide Thiovit 80 at 2.5 gm/liter and untreated plants of Sultan and Glorious cvs. were used for comparison in this study.

Data presented in Table (2) indicated that all concentrations of the tested plant extracts significantly reduced disease severity of *E. polygoni* in comparison with the untreated plants. Application of Thiovit 80 before and after artificial inoculation with *E. polygoni* conidio-spores showed almost completely protection against infection with the causal pathogen.

Table (2): Effect of certain plant extracts as well as chemical fungicide against powdery mildew disease incidence on two sugar beet cvs. during 2004/2005 season (screen house).

Plant extract	Disease severity of Sultan						Disease severity of Glorious					
	1000 pm		3000 ppm		5000 ppm		1000 pm		3000 ppm		5000 ppm	
	Before	After	Before	After	Before	After	Before	After	Before	After	Before	After
<i>S.nigrum</i> P.	48.4	52.1	42.8	47.1	37.8	42.5	29.5	31.2	27.9	29.7	24.5	26.1
<i>maritimum</i> M.	46.5	53.11	42.4	50.0	38.4	46.7	31.9	32.6	29.5	31.7	25.3	26.8
<i>azedarach</i>	27.5	32.7	22.4	27.2	14.5	19.8	21.5	28.3	16.6	22.5	8.7	14.5
<i>An. nobilis</i>	39.7	48.2	32.1	40.3	21.6	28.5	26.4	29.4	20.5	26.9	13.6	18.7
<i>A. nispaga</i>	36.5	42.3	28.4	35.1	18.6	24.5	27.2	27.5	18.9	24.8	11.4	19.8
<i>M. piperita</i>	44.3	51.3	35.7	42.5	26.5	33.2	29.2	30.4	24.1	28.2	19.8	22.9
Thiovit 80	2.7	4.8	2.5	4.1	2.5	4.6	1.1	2.4	1.1	2.6	1.2	2.6
Control	58.4	58.4	58.4	58.4	58.4	58.4	33.7	33.7	33.7	33.7	33.7	33.7

L.S.D. 0.05

2 cv. means at each plant extract x treatments = 2.1

2 plant extracts at each cv. x treatments = 1.9

2 treatments at each plant extract x cv. = 2.0

From the results obtained in Table (2) spraying sugar beet plants with all the tested plant extracts at concentration of 5000 ppm before and after inoculation with the pathogen spore suspension was more effective in reducing disease severity than the other concentrations. The highest effect on powdery mildew disease was obtained when sugar beet plants were sprayed before infection with extracts of *Melia azedarach* and *Ammi visnaga* at concentrations of 5000 ppm.

2. Complementary effect of plant extracts and Thiovit 80:

From the previous data during 2005/2006 season, the effective plant extract of *M. azedarach* (5000 ppm) was selected and mixed with Thiovit 80 as follow; (5000 ppm) *M. azedarach*, $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80, $\frac{1}{2}$ *M. azedarach* + $\frac{1}{2}$ Thiovit 80, $\frac{1}{4}$ *M. azedarach* + $\frac{3}{4}$ Thiovit 80 and Thiovit 80 (2.5 gm/liter) were applied before and after inoculation by *E. polygoni*.

Data illustrated in Table (3) show that, all concentrations of the combinations between plant extract and fungicide significantly reduced disease severity of powdery mildew in comparison with the artificially inoculated plants either before or after inoculation in comparison with the plant extract alone.

The positive effect of plant extracts against powdery mildew disease reflects in turn on the root yield, whereas, it improved the yield potentiality comparable to the untreated control plants. Parameters of plant growth were enhanced due to these treatments, consequently, increasing in total soluble solids (TSS%), sugar purity and its quantity in roots. These results are consistent with those obtained by other investigators who found an antimicrobial activity of some plant extracts against many phytopathogens (Kishore *et al.*, 1982; Gouda, 2001 and El-Fahar, 2003).

The best combinations gave the least disease severity before and after artificial inoculation were $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80 followed by $\frac{1}{2}$ *M. azedarach* + $\frac{1}{2}$ Thiovit 80 and $\frac{1}{4}$ *M. azedarach* + $\frac{3}{4}$ Thiovit 80. On the other hand, efficacy of the three combinations was very high and significantly effective in reducing powdery mildew incidence for the two tested cvs.; Sultan and Glorious in comparison with plant extract alone.

Table (3): Effect of plant extract (alone or mixed with a fungicide) on powdery mildew disease of two sugar beet cvs. during 2005/2006 season under screen house conditions.

Combination plant extract fungicide	Concentration (ppm)	Sultan				Glorious			
		Disease severity %		Efficiency %		Disease severity %		Efficiency %	
		Before	After	Before	After	Before	After	Before	After
<i>M.azedarach</i>	5000	15.5	22.	71.1	57.8	8.8	13.7	70.5	53.5
<i>M.azedarach</i> +Thiovit80	3750+625	8.3	16.1	84.5	69.9	4.6	8.8	84.4	70.1
<i>M.azedarach</i> +Thiovit80	2500+1250	5.4	11.9	89.9	77.5	2.9	6.2	90.2	78.9
<i>M.azedarach</i> +Thiovit80	1250+1875	3.6	7.5	93.3	86.00	1.9	3.7	93.5	87.4
Thiovit80	25g/liter(2500ppm)	2.8	5.3	94.7	90.1	0.9	2.6	95.6	91.2
Control		53.6	53.6	0.0	0.00	29.5	29.5	0.0	0.0
L.S.D. 0.05:									
2 cv. means at each of conf. x Tr.		2.97		4.5		2.1		3.1	
2 con. means at each of cv x Tr.		2.2		5.1		1.98		2.9	

Efficiency of the three combinations as mentioned before were 84.5, 89.9 and 93.3 % before inoculation, while fungicide (Thiovit 80) efficiency was 94.7%. On the other hand, the efficiency of the three combination after inoculation were 69.9, 77.5 and 86.0%, while the fungicide efficiency recorded 90.1% for Sultan cv.

The same trend was obtained for Glorious cv., where the efficiency for the three combination before inoculation were; 84.4, 90.2 and 93.5% in comparison with fungicide which gave 95.6%. On the other side, the data show that the efficiency after inoculation for the three combination were; 70.1, 78.9 and 87.4% while for Thiovit 80 it was 91.2%. Induced resistance could be achieved by applying of plant extracts like *M. azedarach* (Schmitt, 2006).

We selected the dose of $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80, although the other combinations $\frac{1}{4}$ *M. azedarach* + $\frac{3}{4}$ Thiovit 80, $\frac{1}{2}$ *M. azedarach* + $\frac{1}{2}$ Thiovit 80 and Thiovit 80 alone, have a high efficiency than $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80. So, we selected the first combination to minimize both pollution and costs as well as their residual effect on the chemical fungicide on sugar.

b. Field experiments:

The effect of *M. azedarach* at concentration of 5000 ppm and $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80 on the disease incidence of powdery mildew disease using Sultan and Glorious sugar beet cvs. was studied under field conditions during 2006/2007 and 2007/2008 growing season. Thiovit 80 at concentration of 2.5 gm/liter and untreated plants were used for comparison (control). Data presented in Tables 4, 5, 6 and 7 show that plant extract, or Thiovit 80 and the combination significantly reduced disease severity of *E. polygon* compared with untreated plants. However, the combination between plant extract and fungicide came the best in this respect for both Sultan and Glorious cvs. It is obvious that the efficiency of plant extract alone exceeded 50%, while the efficiency of the combination was more effective in reducing disease severity of powdery mildew. It is clear from the results obtained that treated sugar beet plants with Thiovit 80 or plant extract (*M. azedarach*) and the combination between $\frac{3}{4}$ *M. azedarach* + $\frac{1}{4}$ Thiovit 80 tested significantly increased sucrose, purity, TSS content and root weight of the two sugar beet cvs.; Sultan and Glorious.

Table 4 show that increase (%) in root weight of each treatment ranged from 29.1 to 65.4% for Sultan cv. and from 53.1 to 61.2% for Glorious cv.

On the other hand, increase (%) in sucrose ranged between 51.5 to 67.7% for Sultan cv. while it ranged between 30.3 to 41.4% for Glorious cv. during 2006/2007 season.

During 2007/2008 season, root yield increase ranged from 40.5 to 58.8% for Sultan cv., while it ranged from 19.5 to 32.8% for Glorious cv. (Table 6).

Sucrose increase percentage ranged from 31.7 to 45.4% for Sultan cv., while it ranged from 17.8 to 28.1% for Glorious cv. (Table 6).

The data obtained for increase (%) either in sucrose or root weight was confirmed by Wolf and Verreet (2002). Chlorophyll content of leaves

(Tables 4 and 6) affected by disease severity (%) and decreased by increasing of disease severity (%), as mentioned by Gordon and Duniway (1981), when sugar beet leaves infected by powdery mildew show reduction of net photosynthesis as long as disease developing in comparison with healthy plants.

Table (4): Effect of plant extract (alone or mixed with a fungicide) on powdery mildew disease and certain parameters of two sugar beet cvs. under field conditions during 2006/2007 season.

Cultivar	Treatments	Disease severity (%)	Chlorophyll content (mg/gm)	Root weight (kg/plant)	TSS (%)	Sucrose (%)	Purity (%)	Increase (%)	
								Root weight	Sucrose
Sultan	M.az.(5000 pm)	20.2	52.3	0.910	17.9	13.4	74.8	29.1	51.5
	M.az.+Thiovit80*	14.8	67.5	1.659	21.1	17.4	82.5	61.1	62.6
	Thiovit80(2.5g/L)	8.5	74.1	1.866	24.0	20.1	83.7	65.4	67.7
	Control	46.5	29.8	0.645	11.3	6.5	57.5	0	0
Glorious	M.az.(5000ppm)	10.8	70.3	1.731	225	18.1	80.4	53.1	30.4
	M.az.+Thiovit80*	7.2	77.4	1.970	24.5	20.2	82.4	58.8	37.6
	Thiovit80(2.5g/L)	4.2	80.6	2.100	26.3	21.5	81.7	61.2	41.4
	Control	26.9	54.3	0.811	17.1	12.6	73.7	0	0
L.S.D. 0.05: 2 cv. means at each treatment		2.1	2.3	0.46	1.46	1.87	2.6	2.10	1.9

* *M. azedarach* + Thiovit = 3750 + 625 pm (mg/L)

Table (5): Effect of different treatments on disease severity (%), phenols, and AUDPC of two sugar beet cvs. under field conditions during 2006/2007 season.

Treatments	Sultan						Glorious					
	Disease severity (%)	Efficiency (%)	Phenols			AUDPC	Disease severity (%)	Efficiency (%)	Phenols			AUDPC
			Free	Conjugated	Total				Free	Conjugated	Total	
M.az. 5000	20.2	56.5	20.4	59.3	79.7	1811.31	10.8	59.8	28.6	74.5	103.1	915.82
M.az.+ Thiovit 80*	14.8	68.1	27.6	80.2	107.8	1438.5	7.2	73.2	32.8	99.3	132.1	729.75
Thiovit 80	8.5	81.7	32.4	120.6	153.0	934.75	4.2	84.3	51.9	125.4	177.3	366.0
Control	46.5	0	11.9	51.5	63.4	4128.75	26.9	0	17.8	62.2	80.0	1934.25
L.S.D. 0.05 2 treatment means at each cv.	2.8	2.1	1.9	1.8	4.3	10.1	2.9	1.9	2.2	2.7	3.1	5.60

* *M. azedarach* + Thiovit 80 = 3750 + 625 ppm (mg/L)

Regarding phenolic compounds; free, conjugated and total, it decreased by increasing disease severity (%) as shown from Tables (5 and 7), this due to the presence of some enzymes related to the oxidation of phenol compounds. Based on the results of the experiments, we may assert that there is a negative correlation between the resistance of sugar beet varieties to powdery mildew and phenolic activity, these results in accordance with those obtained by Mayer (1987).

Table (6): Effect of plant extract (alone or mixed with a fungicide) on powdery mildew disease and certain parameters of two

sugar beet cvs. under field conditions during 2007/2008 season.

Cultivar	Treatments	Disease severity (%)	Chlorophyll content (mg/gm)	Root weight (kg/plant)	TSS (%)	Sucrose (%)	Purity (%)	Increase (%)	
								Root weight	Sucrose
Sultan	M.az.(5000ppm)	18.4	60.1	1.315	20.8	16.7	80.2	40.5	31.7
	M.az.+Thiovit80*	12.8	71.5	1.730	23.6	19.5	82.6	54.7	41.5
	Thiovit80(2.5g/L)	7.5	82.3	1.887	25.3	20.9	82.6	58.8	45.4
	Control	37.6	47.6	0.782	15.5	11.4	73.5	0	0
Glorious	M.az.(5000 pm)	10.9	75.4	1.845	24.2	19.6	80.9	19.5	17.8
	M.az.+Thiovit 0*	7.4	81.8	1.857	24.6	20.0	81.3	20.0	19.5
	Thiovit80(2.5g/L)	3.2	89.2	2.210	26.1	22.4	85.8	32.8	28.1
	Control	18.7	62.5	1.485	20.5	16.1	78.5	0	0
L.S.D. 0.05: 2 cv. means at each treatment		1.9	2.4	0.39	1.21	1.96	1.12	3.1	2.1

* M. azedarach + Thiovit = 3750 + 625 ppm (mg/L)

Regarding AUDPC as shown from Table 5 and 7, the data show that under natural infection the area recorded the highest, while it recorded the lowest when the plants were treated by Thiovit 80. When disease severity increased, AUDPC increased. On the other side, AUDPC of Glorious cv. were low comparing with Sultan, due to the level of resistance for both cvs.

Table (7): Effect of different treatments on disease severity (%), phenols, and AUDPC of two sugar beet cvs. under field conditions during 2007/2008 season.

Treatments	Sultan						Glorious					
	Disease severity (%)	Efficiency (%)	Phenols			AUDPC	Disease severity (%)	Efficiency (%)	Phenols			AUDPC
			Free	Conjugated	Total				Free	Conjugated	Total	
M.az. 5000	18.4	51.1	24.8	61.5	86.3	1381.5	9.2	54.9	30.8	76.3	107.1	998.23
M. az. + Thiovit 80*	12.8	65.9	34.5	68.9	103.4	1285.5	6.1	70.1	30.1	110.8	140.9	624.25
Thiovit 80	7.5	80.1	40.3	84.7	125.0	768.75	3.2	84.3	54.3	131.5	185.8	285.75
Control	37.6	0	14.6	54.4	69.0	3298.5	20.4	0	21.5	65.2	86.7	1795.41
L.S.D. 0.05 2 treatment means at each cv.	3.1	2.6	2.9	2.3	5.1	9.2	1.5	3.0	2.7	3.2	4.6	6.7

M. azedarach + Thiovit 80=3750 + 625 ppm (mg/L)

It could be concluded that the selected combination gave satisfactory control to powdery mildew disease and could be recommended especially for developing countries, where chemical control would be economically cropping especially if repeated spraying is done. Similar results were reported elsewhere using extracts other than those reported herein i.e. Mori *et al.* (1989) and Chaturvedi *et al.* (1987). The present result would be of a great importance since it would minimize the costs of control process, save the exploited quantity of fungicide and finally will reduce environmental pollution and increase the gross sugar yield, as well as eliminate the fungicidal residual effect in processed sugar.

REFERENCES

- Al-Abed, A.S. (1992). Possible antifungal effects of aqueous extracts and residues of some common wild plant species on certain plant. M.Sc. Thesis, University of Jordan. p. 81.
- A.O.A.C., Association of Official Analytical Chemicals (1990). Official Methods Analysis of the Association of Official Analytical Chemicals. Washington, 25 D.C. USA.
- Ashry, M.A.; Sh. El-El-Hamdy; E.A. Salem; M.A. Abdel-Baki, M.R. Sehly; Z.H. Osman and M.A. Abbasy (1999). *In vitro* and *In vivo* effects of certain plant extracts on rice blast disease caused by *Pyricularia oryzae* Cav. J. Agric. Res. Tanta Univ., 25(2), 1999.
- Bray, H.G. and W.V. Thrope (1954). Analysis of phenolic compounds of interest in metabolism methods of chemical analysis, 27-51.
- Chaturvedi, R.; A. Dikshit and S.N. Dixit (1987). *Adenocalymma ollicea*, a new source of natural fungitoxicant. Trop. Agric. 64: 318-22.
- El-Fahar, Samia, A. (2003). Integrated management of Cercospora leaf spot disease of sugar beet in Delta area. Ph.D. Thesis, Fac. of Agric., Kafr El-Sheikh, Tanta Univ.
- El-Fahar, Samia, A. (2008). Impact of powdery mildew disease on root weight, sugar yield and purity of sugar beet. J. Agric. Res. Kafr El-Sheikh Univ., 34(3): 2008.
- Gomez, K. and A.C. Gomez (1983). Statistical Procedures for Agricultural Research, International Rice Research Institute, Book. John Wiley and Sons, New York, pp. 403.
- Gordon Thomes, R. and John M. Duniway (1981). Effects of powdery mildew infection on the efficiency of CO₂ fixation and light utilization by sugar beet leaves. Plant Physiol. 1982, 69, 139-142.
- Gouda, M.I. (2001). Studies on some causes of sugar beet root rots. Ph.D. Thesis, Fac. of Agric. Tanta Univ. Egypt. p. 146.
- Grimmer, M.K.; K.M.R. Bean and M.J.C. Asher (2007). Mapping of five resistance genes to sugar beet powdery mildew using AFLP and anchored SNP markers. Theoretical and Applied Genetics, 115(1), pp. 67-75.
- Hills, F.J.; L. Chirappa and S. Geng (1980). Powdery mildew on sugar beet. Disease and crop Assessment. Phytopathology. Vol. 70, No. 7.
- Karaoglanidis, G.S. and D.A. Karadimos (2005). Efficacy of strobilurins and mixtures with DMI fungicides in controlling powdery mildew in field-grown sugar beet. Crop Protection. 25(2005): 977-983.
- Kishore, N.; N.K. Dubey; R.D. Tripathi and S.K. Singh (1982). Fungitoxic activity of leaves of some higher plants. National Academy Science Letter, 5(1): 9-10.
- Mayer, M.A. (1987). Polyphenoloxidases in plants recent progress. Phytochemistry, 26(1): 11-20.
- McGinnis, R.A. (1982). Beet sugar technology. 3rd edn. Beet sugar development foundation fort Collins. 855 pp.

- Mori, A.; N. Enaki; K. Shinzuka; C. Nishino and M. Fukushima (1987). Antifungal activity of fatty acids against *Pyricularia oryzae* related to antifungal constitutions of *Miscanthus sinensis*. Agric. Biol., Chem. 51: 3403-05.
- Pandy, H.N.; T.C. Menon and Rao (1989). A simple formula for calculating area under disease progress curve. Rachis, 8(2): 38-39.
- Paulus, A.O. (2008). Sugar beet post management : leaf diseases. U. Bulletin No. 3278. ANR Publications. Oakland, CA 94608-1239.
- Prakash, A. and J. Rao (1997). Botanical pesticides in Agriculture. Lewis Publishers, CRC, Press.
- Schmitt, A. (2006). Induced resistance with extracts of *Reynoutria sachalinensis*: Crucial steps behind the scene induced resistance in plants against insects and diseases, IOBC WPRS Bulletin. Vol. 29(8), 2006.
- Scott, W.P. and G.H. McKibben (1978). Toxicity of black pepper extract to boll weevils. J. Econ. Entomol. 71(2): 343-344.
- Shimon, M.; E. Putievsky; V. Ravid and R. Rcoyeni (1993). Antifungal activity of volatile fractions of essential oils from four aromatic wild plants in Israel. J. of Chem. Ecol., 19(6): 1129-1133.
- Snell, F.D. and C.T. Snell (1953). Calorimetric methods of analysis including some turbidimetric and nephelometric methods. D. Van Nostrand Company Inc. Tornto. New York, London. Vol. III, 606 pp.
- Wolf, P.F.J. and J.A. Verreet (2002). The IPM sugar Beet Model Plant Disease. Vol. 86 No. 4.
- Yoshida, S.; D. Forno; J. Cock and K. Gomez (1976). Laboratory manual for physiological studies of rice. 3rd ed. pp. 83. The International Rice Research Institute, Los Banos, Philippines, Manila, Philippines.

تأثير بعض المستخلصات النباتية والاستبدال الجزئي لمبيد الثيوفيت 80 بواسطة
مستخلص نباتي على مرض البياض الدقيقي في بنجر السكر
سامية عبده الفحار¹ و سحر مأمون إبراهيم مصطفى²
1- قسم بحوث أمراض المحاصيل السكرية - معهد بحوث المحاصيل السكرية
2- قسم تكنولوجيا السكر - معهد بحوث المحاصيل السكرية - مركز البحوث الزراعية -
الجيزة - مصر

تم إقامة أربعة تجارب خلال مواسم 2004/2005 ، 2005/2006 ، 2006/2007 ،
2007/2008 وذلك بمحطة بحوث سخا الزراعية - محافظة كفر الشيخ وذلك لدراسة تأثير بعض
المستخلصات النباتية على مقاومة البياض الدقيقي في بنجر السكر والذي يسببه فطر *Erysiphe*
polygoni وتأثيره على الصفات الأخرى.

تم استخدام ستة مستخلصات نباتية وذلك لاختبار تأثيرها ضد فطر *E. polygoni* وهي عنب
الذيب *Solanum nigrum* والنرجس *Panocratinum maritimum* والزرنكلت *Melia*
azedarach والشايخ *Anthemis nobilis* والخله *Ammi visnaga* وأوراق النعناع
Mentha piperita وذلك بتركيزات مختلفة 1000 ، 3000 ، 5000 جزء في المليون في الصوبة
السلكية وذلك على صنفين من بنجر السكر هما سلطان وجلوريس وتم إجراء التجربة تحت ظروف العدوى

الصناعية باستخدام الجراثيم الكونيدية للفطر. ولخفض الجرعة المستخدمة من المبيد الفطري 80 Thiovit ضد الفطر تم خلط نصف للتركيز الفعال من المستخلص النباتي الزنزلخت بمعدل 2500 جزء في المليون مع نصف التركيز الفعال للمبيد 80 Thiovit (1250 جزء في المليون) وثلاثة أرباع المستخلص النباتي 3750 جزء في المليون مع ربع المبيد (625 جزء في المليون) وأيضا تم خلط ربع المستخلص النباتي (1250 جزء في المليون) مع ثلاثة أرباع المبيد (1875 جزء في المليون) وقد اجريت هاتين التجريبتين في الصوبة لسلكية تحت ظروف لتعوى الصناعية خلال الموسمين 2005/2004م ، 2006/2005م وقد تم الرش بكل المستخلصات النباتية وخليط المستخلص النباتي الزنزلخت مع مبيد الثيوفيت 80 قبل العدوى بثلاثة أيام وبعد العدوى بثلاثة أيام كلا على حده. وقد أظهرت النتائج التي تم الحصول عليها أن المستخلص النباتي الزنزلخت عند تركيز 5000 جزء في المليون كان أفضل المستخلصات (سواء قبل أو بعد العدوى يليه المستخلص النباتي الخله).

وقد أوضحت النتائج أن خلط المبيد مع المستخلص النباتي قد أدى إلى ارتفاع كفاءة المستخلص خاصة عند إضافة 4/3 التركيز الفعال من المبيد مع 4/1 التركيز الفعال من المستخلص يليه 2/1 تركيز المبيد مع 2/1 تركيز المستخلص يليه 4/1 تركيز المبيد مع 4/3 تركيز المستخلص حيث كانت النسب كالاتي: 93.3 ، 89.9 ، 84.54% على التوالي وذلك قبل العدوى ، 86 ، 77.5 ، 69.9% بعد العدوى وذلك مقارنة بكفاءة المستخلص منفردا حيث كانت النسبة 71.1% قبل العدوى ، 57.8% بعد العدوى للصف سلطان. أما الصنف جلورياس كانت لانسبة كالاتي: 93.5 ، 90.2 ، 84.4% وذلك قبل العدوى ، 87.4 ، 78.9 ، 70.1% بعد العدوى مقارنة بالمستخلص منفردا حيث كانت كفاءته بنسبة 70.5% قبل العدوى ، 53.5% بعد العدوى.

وفي التجارب الحقلية تم استخدام المستخلص النباتي الزنزلخت بتركيز 5000 جزء في المليون والخليط المكون من 4/3 المستخلص النباتي الزنزلخت ، 4/1 المبيد الفطري ثيوفيت 80 وذلك لارتفاع كفاءة هذا الخليط في مقاومة مرض البياض الدقيقي وأيضا لتقليل التلوث وخفض التكاليف.

وقد أظهرت النتائج المتحصل عليها من موسمين متتاليين هما 2007/2006م ، 2008/2007م أن شدة الإصابة بالبياض الدقيقي على الأوراق قد أمكن تقليلها بنسبة تفوق 50% باستخدام مستخلص الزنزلخت في حين أظهرت نتائج الخليط المكون من 4/3 المستخلص ، 4/1 المبيد بزيادة كفاءة المقاومة للمرض لتصل إلى 68.1% في الصنف سلطان ، 73.2% للصف جلورياس مقارنة بالمبيد على حده خلال موسم 2007/2006م ، 65.9% للصف سلطان ، 70.1% للصف جلورياس خلال الموسم 2008/2007م.

وأوضحت النتائج أيضا أن محتوى الأوراق من الكلوروفيل كان عاليا 67.5 ، 77.4مجم لكلا من الصنفين سلطان وجلورياس على التوالي عند الرش بالخلوط ولقد تأثرت الصفات الأخرى مثل وزن الجذر والمواد الصلبة للذاتية الكلية والنسبة المئوية للسكريز والنقاوة وذلك تحت ظروف العدوى الطبيعية بالحقل. وكانت الزيادة في وزن الجذر 61.1 ، 58.8% للأصناف سلطان وجلورياس على التوالي بينما كانت الزيادة في السكر فكانت 62.6 ، 37.6% للأصناف سلطان وجلورياس على التوالي بالمقارنة بالمعاملات الأخرى.

وأوضحت النتائج أن المركبات الفينولية تأثرت بالشدة المرضية وأن هناك علاقة عكسية بين الشدة المرضية والمركبات الفينولية. أما عن المساحة تحت منحنى التقدم المرضي فإنها أقل عند المعاملة بالمستخلص ويوضح هذا البحث إلى إمكانية استبدال المبيدات بالمستخلصات النباتية لمقاومة الأمراض في بنجر السكر سواء أكانت منفردة أو بخلطها بنسب بسيطة من المبيد وذلك لخفض التكاليف والتلوث البيئي الناجم عن استخدام المبيدات وأيضا لتلافي خطورة الأثر المتبقى للمبيد في السكر الناتج.