

THE EFFECT OF VITRIFICATION PROCEDURES AND CRYOPROTECTANT TYPES ON POST-THAW SURVIVAL RATE OF RABBIT OOCYTES.

Dorra, Tork M.* ; Kh. EL. Sherif * ; Sh. M. Shamiah** and Sara F. Fouda*

* Poultry production Dept., Fac. of Agric., Man. Univ. Egypt.

**Animal production Dept., Res. Inst., Agric. Rec.Center, Giza, Egypt.

ABSTRACT

Aim of the present study was to determine the effect of some factors affecting oocyte vitrification of rabbit does including, three cryodevices [Straws (S), Open pulled straws (OPS) and Sealed Open pulled straws (SOPS)] using three cryoprotectant types [40% ethylin glycol (EG), 40% dimethyl sulfoxide (DMSO) and their combination, 20% EG + 20% DMSO]. Total of 36 New Zealand white (NZW) rabbit does with (5 mo old and 3.25 ± 0.25 kg LBW) and were used in this study. Ovaries were collected and oocytes were harvested. The oocyte yield was recorded and the number of oocyte/ovary was counted, and oocyte recovery rate was calculated. Results indicated that total number of follicles averaged 36.0/doe and 18.0/ovary. Yield of oocytes averaged 31.3/doe and 15.5/ovary and oocyte recovery rate was 86.2%. Oocyte recovery rate was for 61.96, 11.65, 6.48, and 6.10% for compact, denuded, partial denuded and expanded oocytes, respectively. The corresponding frequency distribution was 71.89, 13.52, 7.52 and 7.07%, respectively. Relative to total number of vitrified oocytes, total survival rate of oocyte was not affected significantly by cryodevice, being 87.30, 83.09 and 85.77% for S, OPS and SOPS, respectively. Post-thaw survival rate of normal oocytes was the highest (68.33%, $P < 0.05$) and abnormal oocytes was the lowest (17.44%, $P < 0.05$) with SOPS and the opposite was obtained for straws (59.43 and 27.87%, respectively, $P < 0.05$). While, OPS showed moderate values (63.31 and 19.78%, respectively), but did not differ significantly from each of straw and SOPS. Total survival rate of oocyte was not affected significantly by cryoprotectant type, being 85.38, 83.45 and 87.22% for EG, DMSO and their combination, respectively. Post-thaw survival rate of normal oocytes was the highest (72.56%, $P < 0.05$) and abnormal oocytes was the lowest (14.66%, $P < 0.05$) with EG+DMSO and the opposite was obtained for EG alone (56.92 and 28.46%, respectively, $P < 0.05$). While, DMSO showed moderate values (61.97 and 21.48%, respectively).

In conclusion, using sealed open pulled straws and combination of 40% ethylin glycol and 40% dimethyl sulfoxide as cryoprotectant solution as vitrification procedures of rabbit oocytes showed the best results of post-vitrification survival rate.

INTRODUCTION

A large number of oocytes are required for various biotechnological techniques including oocyte maturation, cloning and the production of transgenic animals. Thus cryopreservation of mammalian oocytes may be used to establish an ova bank for reproductive biotechnology.

Cryopreservation of oocytes is more difficult than embryos. Many problems are associated with the cooling and freezing of *in vitro* matured or *in vivo* ovulated oocytes, namely spindle disorganization (Magistrini and Szollosi, 1980), loss or clumping of microtubules resulting in scattering of chromosomes (Sathananthan *et al.*, 1988), increased polyploidy during fertilization (Carroll *et al.*, 1989) and a decreased fertilization rate (Wood *et al.*, 1992).

The difficulty, in obtaining acceptable rates of survival and functionality for oocytes after cryopreservation, is due to the size of this cell and its unique morphologic characteristics. Vitrification is a process whereby a solution with a high solute concentration will solidify into a glass-like structure upon cooling, without any ice crystal formation. The process may be thought of as simply increasing viscosity until a solid state is reached (Shaw, *et al.*, 2000a). Vitrification is an alternative that may increase oocyte survival after thawing. Vitrification is the solidification of a solution at low temperatures without ice crystal formation. This phenomenon requires either rapid cooling rates (Rall, 1987) or the use of concentrated cryoprotectant solutions, which depress ice crystal formation and increase viscosity at low temperatures (Vajta, 2000).

The present study was carried out to study the effect of vitrification procedures {straws (S), open pulled straws (OPS) and sealed open pulled straws (SOPS) and cryoprotectant types {ethylin glycol (EG), dimethyl sulfoxide (DMSO) or its combination) on quality of vitrified rabbit oocytes.

MATERIALS AND METHODS

This study was carried out at International Livestock Management Training Center, (ILMTC), Sakha, Kafr El-Sheikh Governorate belonging to the Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture during the period from August 2008 to August 2009.

Animals:

Total of 36 New Zealand white (NZW) rabbit does with average of 5 mo old and 3.25 ± 0.25 kg live body weight (LBW) were used in this study. All rabbit does were subjected as donors of oocytes and 5 rabbit does were used for blood collection for preparing rabbit doe serum (RDS).

All does were kept under the same conditions of feeding and management in the special farm, being individually housed in metal cages (40 x 50 x 60 cm) provided with feeders and water nibble in each cage. Does were fed *ad libitum* on a commercial pelleted concentrate diet.

Ovarian collection:

Immediately after slaughtering, ovaries were removed, washed by saline solution (0.9% NaCl) and dried by cleaning paper. Ovaries were collected and excised, submerged in a flacon plastic tissue culture dishes (60 x 15 mm) containing saline solution supplemented with (Sodium penicillin G, 100 IU/ml and streptomycin, 100 µg/ml) and transported to the laboratory within 5 minute.

Oocyte collection:

All visible follicles on the ovarian surface with ≥ 1 mm in diameter were counted. Oocytes were collected using slicing technique into glass Petri dishes containing 4 ml of harvesting medium (Phosphate buffer saline, PBS) supplemented with sodium penicillin G (100 IU/ml) and streptomycin (100 µg/ml). Ovaries were held by a forceps and by a scalpel blade were made incisions along the whole ovarian surface. Each ovary was washed three times with harvesting medium in dishes containing 2 ml medium for washing and searching oocytes using stereomicroscopy.

Preparation of harvesting medium:

Phosphate buffer saline (PBS) medium was prepared according to Gordon (1994). Two mg/ml of bovine serum albumin (BSA) was added to PBS. The pH value of the medium was adjusted to 7.2-7.4 using pH-meter and to osmolarity of 280-300 mOsmol/kg using osmometer. Then, the medium was filtered by 0.22 µm millipore filter (milieux GV, millipore, Cooperation Bedford MOA.).

Yield and recovery rate of oocytes:

The oocyte yield was recorded and the number of oocyte/ovary was calculated. The recovery rate was determined as percentage of oocytes in proportional to total number of visual follicles presented on the ovarian surface of each doe using the following formula of recovery rate (RR%):

$$RR (\%) = (\text{Number of recovered oocytes/number of follicles}) \times 100$$

Oocyte categories:

After collection, oocytes were washed three times in harvesting medium, counted, evaluated under inverted microscope and classified into six categories, *compact, expanded, denuded, partial denuded and degenerated oocytes*.

Preparation of doe rabbit serum:

Doe rabbit serum was prepared from blood collected from 5 rabbit does. The collected blood was centrifuged two times at 3000 rpm for 15 minutes. Clear sera were aspirated by pasture pipette and placed in another 15 ml-sterile tubes. These tubes were placed into water bath at 56°C for 30 minutes, and then left to cool. Thereafter, sera were placed into 1.5 ml eppendorf tubes and frozen until used.

Vitrification procedures:

Only morphologically high quality immature oocytes (COCs), as determined by uniform granular, homogeneously distributed cytoplasm surrounded by compact layers of cumulus cells were vitrified in this study. Oocytes were vitrified by different vitrification procedures (S, OPS and SOPS) with each of three cryoprotectants (40% EG, 40% DMSO or combination of 20% EG +20% DMSO) using three carrier systems.

The vitrification procedures employed throughout this experiment were based on the methods originally designed by Silvestre et al. (2003). The basal medium used for vitrification (VS) was TCM-199 (Sigma) supplemented with 20% (v:v) of DRS.

Types of cryodevice:

Conventional straws:

Plastic insemination straws (0.25 ml, IVM L' Aigle, France) were used for oocyte vitrification. In this method, ten COCs were transferred in 50 µl of VS then, oocytes with 3 µl loading into a column located between two columns of vitrification solution separated by air bubbles into the center of 0.25 ml plastic insemination straws using a fine glass capillary pipette. After heat-sealing, the straws were plunged immediately into LN₂ (-196°C).

Open pulled straws (OPS):

Conventional plastic insemination straws (0.25 ml, IVM L' Aigle, France) were heat-softened over a Bunsen burner and pulled manually. The straws were cooled in air and then cut at the tapering end with a blade. The

inner diameter of the tip decreased from 1.7 mm to 0.8 mm, and the wall thickness diminished from 0.15 to 0.07 mm according to Vajta et al. (1998).

In this method, ten COCs were transferred in 50 µl VS then, the COCs were transferred to a 3 µl droplet of and loaded into a OPS by capillary action. The OPS were plunged immediately into LN₂ (-196°C).

Sealed open pulled straws (SOPS):

The preparation of SOPS was similar to that described previously for OPS. The straws were cut at the narrowest point with a razor blade. Only half with the cotton plug was used (Lopez-Bejar and Lopez-Gatius, 2002).

In this method, ten COCs were transferred in 50 µl VS then, the COCs were aspirated into a column located between two columns of vitrification solution separated by air bubbles. The other end of the straw is sealed with heat-sealing before plunged immediately into LN₂ (-196°C).

In all methods, three VS containing different tested cryoprotectants were used and oocytes were exposed to VS for 20–25 seconds prior to immersion in LN₂.

Warming method:

After storage for at least 3 weeks in LN₂, vitrified oocytes were warmed by holding the straws for 6 sec in air and then agitating them in water bath at 20 °C for at least 10 sec. The contents of each straw were expelled into Petri dish. To remove of intracellular cryoprotectants, oocytes were transferred to solution of TCM-199 supplemented with 10% DRS and 0.25 M sucrose, then in the same medium with 0.125 M sucrose and finally in the same medium without sucrose, for 5 min per solution,.

Evaluation of post-warming oocyte viability:

Oocyte viability was evaluated morphologically based on the integrity of the oolemma and zona pellucida; loss of membrane integrity (lysis) was obvious upon visual inspection as the sharp demarcation of the membrane disappeared and the appearance of the cytoplasm changed. The criteria used for assessing the post-thaw morphology of vitrified/warmed oocytes were normal and abnormal oocytes.

Statistical analysis:

Data were analyzed using analysis of variance using computer program of SAS (2000). The significant differences among group means were preformed using Duncan Range Test (Duncan, 1955).

RESULTS

Characteristics of follicles and oocytes:

Number of follicles:

Total numbers of follicles and average number of follicles per doe or ovary are shown in Table (1).

Table (1): Total and average number of follicles per doe or ovary observed on the ovarian surface of doe rabbits.

| Number of does | Number of ovaries | Number of follicles | Number of follicles | |
|----------------|-------------------|---------------------|---------------------|-----------|
| | | | Per doe | Per ovary |
| 36 | 72 | 1296 | 36.0 | 18.0 |

Results indicated that total number of follicles including count of vesicular follicles, small follicles < 3 mm, medium follicles 3-10 mm and large follicles

>10 mm, was 1296 follicles, averaging 36.0/doe and 18.0/ovary, regardless the ovarian side.

Yield and recovery rate of oocytes:

Results shown in Table (2) revealed that total of 36 doe rabbits yielded a total number of 1117 oocytes, averaging 31.03 and 15.5 oocytes per doe and ovary, respectively. This reflected in higher oocyte recovery rate, being 86.2%.

Table (2): Total and average number of oocytes per doe or ovary recovered from ovarian follicles of doe rabbits.

| Number of follicles | Number of oocytes | Number of oocytes | | Recovery rate (%) |
|---------------------|-------------------|-------------------|-----------|-------------------|
| | | Per doe | Per ovary | |
| 1296 | 1117 | 31.03 | 15.5 | 86.2 % |

Recovery rate of different oocyte categories:

Results presented in Table (3) show that the total oocyte recovery rate (86.2%) was the highest for compact oocytes (61.96%), followed by denuded and partial denuded (11.65 and 6.48%, respectively), while the lowest rate was obtained for expanded oocytes (6.10%).

Examination of oocytes recovered from the ovaries of doe rabbits in all groups revealed that all oocytes were immature and were evaluated into compact, denuded, partial denuded and expanded oocytes respectively).

Table (3): Number and recovery rate (%) of rabbit oocytes at different categories.

| Number of follicles | Total oocytes | Oocyte category | | | | | | | |
|---------------------|---------------|-----------------|-------|---------|-------|------------|------|----------|-----|
| | | Compact | | Denuded | | P. denuded | | Expanded | |
| | | n | % | n | % | n | % | n | % |
| 1296 | 1117 | 803 | 61.96 | 151 | 11.65 | 84 | 6.48 | 79 | 6.1 |

Frequency distribution of different oocyte categories:

Results shown in Table (4) revealed the highest frequency distribution was obtained for compact oocytes (71.89%), followed by denuded (13.52%) and partial denuded (7.52%), while the lowest distribution was recorded for expanded oocytes (7.07%).

Table (4): Number and frequency distribution (%) of oocytes at different categories.

| Total number of oocytes | Oocyte category | | | | | | | |
|-------------------------|-----------------|-------|---------|-------|------------|------|----------|------|
| | Compact | | Denuded | | P. denuded | | Expanded | |
| | n | % | n | % | n | % | n | % |
| 1117 | 803 | 71.89 | 151 | 13.52 | 84 | 7.52 | 79 | 7.07 |

N= Number

Post-thaw survival rate of vitrified oocytes:

Effect of cryodevice:

Data in Table (5) show that total survival rate of oocyte was not affected significantly by cryodevice, being 87.30, 83.09 and 85.77% for S, OPS and SOPS, respectively. However, post-thaw survival rate of normal and abnormal oocytes relative to total vitrified oocytes was affected significantly ($P<0.01$ and $P<0.001$, respectively) by cryodevice, being significantly ($P<0.05$) the highest (68.33%) with SOPS and the lowest (59.43%), while OPS showed moderate values (63.31%), but did not differ significantly from each of straw and SOPS.

Table (5): Effect of cryodevice on survival rate of different types of oocytes relative to total vitrified rabbit oocytes.

| Cryodevice | N | Post-thawing survival oocytes | | | | | |
|------------|-----|-------------------------------|-------|--------|---------------------|----------|--------------------|
| | | Total | | Normal | | Abnormal | |
| | | n | % | n | % | n | % |
| Straw | 244 | 213 | 87.30 | 145 | 59.43 ^b | 68 | 27.87 ^a |
| OPS | 278 | 231 | 83.09 | 176 | 63.31 ^{ab} | 55 | 19.78 ^b |
| SOPS | 281 | 241 | 85.77 | 192 | 68.33 ^a | 49 | 17.44 ^b |

a and b: Means denoted within the same column with different superscripts are significantly different at $P < 0.05$. N= Total number of vitrified oocytes.

On the other hand, survival rate of oocytes in abnormal type was significantly ($P < 0.05$) higher with S than those with OPS and SOPS (27.87 vs. 19.78 and 17.44%, respectively). These results indicated the beneficial effect of SOPS as cryodevice on survival rate of rabbit oocytes.

Effect of cryoprotectant type:

Results presented in Table (6) revealed that the effect of cryoprotectant type on total survival rate of oocytes relative to vitrified oocytes was not significant, ranging between 87.22% with the combination of DMSO and EG and 83.45% with DMSO. However, post-thaw survival rate of oocytes as normal and abnormal oocytes relative to total number of vitrified oocytes was affected significantly ($P < 0.05$) by cryoprotectant type.

Table (6): Effect of cryoprotectant type on survival rate of different types of oocytes relative to total vitrified rabbit oocytes.

| Type of cryoprotectant | N | Post-thawing survival oocytes | | | | | |
|------------------------|-----|-------------------------------|-------|--------|--------------------|----------|--------------------|
| | | Total | | Normal | | Abnormal | |
| | | n | % | n | % | n | % |
| EG | 253 | 216 | 85.38 | 144 | 56.92 ^b | 72 | 28.46 ^a |
| DMSO | 284 | 237 | 83.45 | 176 | 61.97 ^b | 61 | 21.48 ^b |
| EG+DMSO | 266 | 232 | 87.22 | 193 | 72.56 ^a | 39 | 14.66 ^c |

A and b: Means denoted within the same column with different superscripts are significantly different at $P < 0.05$. N= Total number of vitrified oocytes.

DISCUSSION

Oocyte recovery rate:

Based on the present results of recovery rate and frequency distribution of different oocyte categories, higher yield of oocytes was obtained from rabbit ovaries. Al-Hasani *et al.* (1987) found that the average number of oocytes recovered from ovarian follicles by puncture was 11.7 oocytes/doe versus 31.03/doe as obtained in the present study. Hulot and Matheron (1981) reported that average number of rabbit oocytes that are shed after mating varies according to the strain, and is correlated with live body weight. The wide variation in number oocytes/ovary recovered from rabbits may be attributed to pronounced differences in animal ages, reproductive status, site of the ovary and their interactions (Khalil, 2005). Also, diameter of needle used in oocyte harvesting by aspiration was found to affect recovery rate of oocyte (Abdoon, 2001). Moreover, the differences in number of oocytes/ovary may be related to size of follicles (Shamiah, 1997). In accordance with the present results, data of El-Ratel (2008) clearly revealed that frequency distribution of oocytes at compact category was the

highest in NZW rabbits, being 55.6%. However, frequency distribution of oocytes was 30.5, 8.3 and 5.6% at denuded, partial denude and expanded categories.

Effect of cryodevice:

A variety of new techniques and types of holders/devices have been tested with the aim of improving the overall survival of oocytes post-cryopreservation. Several open-carrier systems and closed carrier methods for oocyte vitrification were developed. Until recently, 0.25 ml standard insemination straws were used almost exclusively for vitrification of oocytes and embryo with a relative cooling speed of 2,500°C/min (Palasz and Mapletoft, 1996) by direct plunging into liquid nitrogen (LN2). Methods using cryodevices such as OPS (Vajta *et al.*, 1998), cryoloops (Lane *et al.*, 1999) and cryotops (Kuwayama and Kato, 2000) allow oocyte vitrification in a volume as little as few microliters. Open pulled straws rendered a high cooling-warming rate of more than 20,000°C/min. Dropping oocytes containing vitrification solution directly into LN2 (Papis *et al.*, 1999) successfully eliminated the insulation effects of the container wall. Yet, plunging a warm object into LN2 resulted in boiling the liquid and creating an isolating layer of nitrogen vapor around the object.

A vitrification device was developed by Arav *et al.* (2000) using LN2 slush by reducing the temperature of LN2 to -205°C. These methods use very high cooling rates coupled with minimal volume of freezing medium containing high concentration of cryoprotectants that permits rapid traversal of temperature zones, corresponding to chilling injury. As applied in this study on rabbit oocytes, Vajta *et al.* (1998) reported vitrification using open pulled straws (OPS) for bovine embryos as well as oocytes. Later on the improved OPS methods using glass capillary (Hochi, *et al.*, 2000 & 2001).

Type of cryoprotectant:

The primary components in successful freezing and vitrification solutions are cryoprotectants (CPAs). Permeating CPAs are essential for dehydrating intracellular water. Additionally, they lower the freezing point, thus giving more time for dehydration. Careful selection of CPAs must be made however, first for toxicity and second for permeability. The CPAs are toxic at high concentrations, but toxicity is minimized at low temperatures, and with short periods of exposure (Agca *et al.*, 1998). The CPAs protect slowly frozen cells by one or more of the mechanisms such as suppressing high salt concentrations, reducing cell shrinkage at a given temperature, reducing the fraction of the solution frozen at a given temperature and minimizing intracellular ice formation. Ice formation can be eliminated entirely, both within the cells and the extra-cellular matrix, when CPAs are used in extremely high concentrations (at least 50% volume/volume) as reported by Fahy (1988). The CPAs used in early experiments are glycerol (Willadsen, 1977). Glycerol and DMSO are the most commonly used CPAs. To achieve high cooling rates requires the use of high concentration of the CPA solution, which depresses ice crystal formation. There is critical concentration required for vitrification. Kanno *et al.* (1975) were able to demonstrate that the temperature at which crystallization begins can be reduced through an increase in the hydrostatic pressure. Finally, reducing the cooling rate can be

achieved by reduction of minimal concentration through the additional use of polymers which are non-permeable and therefore remain in the extra-cellular area (Fahy *et al.*, 1984).

Ethylene glycol is currently the most commonly used permeating CPA for the vitrification of oocytes including human oocytes and embryos, because of its low molecular weight, high permeation ability, and low toxicity. Its low molecular mass permits fast influx and efflux into and out of cells (Kuwayama *et al.*, 2005). O'Neill *et al.* (1997) observed that addition of polyethylene glycol (PEG) resulted in greatly improved viability of oocytes following cryopreservation. Minimizing the toxicity of the CPAs can also be achieved by using a combination of two CPAs, and a stepwise exposure of cells to pre-cooled concentrated solutions. In agreement with the present results, some studies mixed other permeating agents, such as DMSO, to reduce the concentration of single CPAs (Vajta *et al.*, 1998; Yokota *et al.*, 2000). Also, Valdez *et al.* (1992) showed that EG was the least toxic to mouse embryos and successfully used in combination of 20% EG, 20% DMSO and 10% 1, 3-butanediol. Moreover, Valdez *et al.* (1992) found that EG is a very weak glass forming molecule, whereas DMSO is a better glass forming molecule, so often times the two are combined in vitrification solutions.

REFERENCES

- Abdoon, A.S.S. (2001). Factors affecting follicular population, oocyte yield and quality in camels (*Camelus dromedarius*) ovary with special reference to maturation time in vitro. *Animal Reproduction Science*, 66: 71–79.
- Agca, Y., Monson, R.L., Northey, D.L., Mazni, O.A., Schaefer, D.M., Rutledge, J.J., (1998). Transfer of fresh and cryopreserved IVP bovine embryos: Normal calving, birth weight and gestation lengths. *Theriogenology*; 50:147-62
- Al-Hasani, S.K. Diedrich, H. van der Ven, A. Reinecke, M. Hartje and D. Krebs. (1987). Cryopreservation of human oocytes. *Hum. Reprod.* 2:695-700.
- Arav, A., Pearl, M., Zeron, Y. (2000). Does lipid profile explain chill in sensitivity and membrane lipid phase transition of spermatozoa and oocytes. *CryoLett*; 21:179–86.
- Carroll, J., Warnes, G.M., and Matthews, C.D. (1989). Increase in digyny explains polyploidy after in-vitro fertilization of frozen-thawed mouse oocytes. *J. Reprod. Fertil.* 85: 489-494.
- Duncan, D.B. (1955). Multiple range and Multiple F test. *Biometrics*, 11:10.
- El-Ratl, I.T (2008). Physiological studies on in vitro maturation and in vitro fertilization in rabbits. M. Sc. Thesis, Faculty of Agriculture, Mansoura University, Egypt.
- Fahy, G.M., MacFarlane, D.R., Angell, C.A., and Meryman, H.T. (1984). Vitrification as and approach to cryopreservation. *Cryobiology* 21: 407-426.
- Fahy, G.M. (1988). Vitrification in low temperature on Biotechnology: Emerging Applications and Engineering Contributions. (Eds. McGrath, J.J. and Diller, K.R.), New York: ASME, p 133.

- Gordon, I. (1994). Oocyte recovery and maturation in laboratory production of cattle embryos, *Biotechnology in Agriculture* .No II.U.K. CAB International; 20-142.
- Hochi, S., Akiyama, M., Minagawa, G., Kimura, K., and Hanada, A. (2000). Vitrification of in vitro matured bovine oocytes in open-pulled glass capillaries of different diameters. *Theriogenology* 53: 255.
- Hochi, S., Akiyama, M., Minagawa, G., Kimura, K., and Hanada, A. (2001). Effect of cooling and warming rates during vitrification on fertilization of in vitro matured bovine oocytes. *Cryobiology* 42: 69-73.
- Hulot, F. and Matheron, G., (1981). Etude du genotyPe, de l'age et de la saigon sur les composantes de la reproduction chez la lapine. *Ann. Genet. Sel. Anim.*, 13: 131-150.
- Kanno, H., Speedy, R.J., and Angell, C.A. (1975). Supercooling of water to -196°C under pressure. *Science* 189: 880-881.
- Khalil, W.A. (2005). Differences between farm animals in (IVM and IVF) in vitro maturation and in vitro fertilization. M. Sc. Thesis, Faculty of Agriculture, Mansoura University, Egypt.
- Kuwayama, M., and Kato, O. (2000). All-around vitrification method for human oocytes and embryos. *J. Assist. Reprod. Genet.* 17: 477 (abstract).
- Kuwayama, M., Vajta, G. (2005).Improving cryopreservation systems. *Theriogenology* 65:236-44.
- Lane, M., Bavister, B.D. Lyons, E.A. and Forest. K.T. (1999). Containerless vitrification of mammalian oocytes and embryos. *Nature Biotechnol.* 17:1234-1236.
- Lopez-Bejar M, Lopez-Gatius F (2002). Nonequilibrium cryopreservation of rabbit embryos using a modified (sealed) open pulled straw procedure. *Theriogenol* 58, 1541–1552.
- Magistrini, M. and Szöllösi, D. (1980). Effects of cold and isopropyl-N-phenyl carbamate on the second meiotic spindle of mouse oocytes. *Europ. J. Cell. Biol.* 22:699-707.
- O'Neill, L., Paynter, S.J., and Fuller, B.J. (1997). Vitrification of mature mouse oocytes: Improved resulted following addition of polyethylene glycol to a dimethyl sulfoxide solution. *Cryobiology* 34: 295-301.
- Palasz, A. T., Mapletoft R.J., (1996).Cryopreservation of mammalian embryos: recent advances .*Biotechnol Adv* 14, 127–149.
- Papis, K., Shimizu, M., and Izaïke, Y. (2000). Factors affecting the survivability of bovine oocytes vitrified in droplets. *Theriogenology* 15: 651-658.
- SAS, (2000). SAS Institute Inc. SAS User's Guide, Statistics. Cary, NC.
- Sathananthan, A.H., Ng, S.C., Trounson, A.O., Bongso, A., Ratnum, S.S., Ho, J., Mok, H., Lee, M.N. (1998). The effects of ultra rapid freezing on meiotic and mitotic spindles of mouse oocytes and embryos. *Gam. Res.* 21, 385–401.
- Shamiah, S.M. (1997). Some factors affecting in vitro maturation and fertilizing capacity of oocytes in some farm animals. M. Sc. Thesis, Faculty of Agriculture, Minufiya University, Egypt.
- Shaw, J.M., Oranrathachai, A., Trounson, A. (2000a).Cryopreservation of oocytes and embryos. In: Trounson A, Gardner DK, editors. *Handbook of in vitro fertilization*. 2nd ed., Boca Raton, FL: CRC p. 373–412.
- Vajta, G. (2000). Vitrification of the oocytes and embryos of domestic animals. *Anim Reprod Sci*, 60/61:357-364.

- Vajta, G., Holm, P., Kuwayama, M., Booth, P.J., Jacobsen, H. T., and Callesen, H. (1998). Open pulled straw (OPS) vitrification: A new way to reduce cryoinjuries of bovine oocytes and embryos. *Mol. Reprod. Devel.* 51:53-58.
- Valdez, C.A., Abas Mazni, O., Takahashi, Y., Fujikawa, S., Kanagawa, H. (1992). Successful cryopreservation of mouse blastocysts using a new vitrification solution. *J Reprod Fertil*;96:793-802.
- Willadsen, S.M. (1977). Factors affecting the survival of sheep and cattle embryos during deep-freezing and thawing. In: the freezing of Mammalian Embryos. Ciba Foundation Symposium 52 Excerpta Medica, Amsterdam, pp. 175-194.
- Wood, M.J., Whittingham, D.G., Lee, S.H. (1992). Fertilization failure of frozen mouse oocytes is not due to premature corticle granule release. *Biol. Reprod.* 46, 1187-1195.
- Yokota, Y., Sato, S., Yokota, M., Ishikawa, Y., Makita, M., Asada, T., and Araki, Y. (2000). Successful pregnancy following blastocyst vitrification: case report. *Hum. Reprod.* 15: 1802-1803.
- Rall, W.F. (1987). Factors affecting the survival of mouse embryos cryopreserved by vitrification. *Cryobiology*, 24:387-402.

تأثير طرق التجميد بالتزجج وأنواع المواد الواقية على معدل الحيوية أثناء الإسالة لبويضات الأرانبي .

ترك محمد درة*، خليل الشحات شريف*، شريف مغاوري شامية** وسارة فكرى فوده*.

*قسم انتاج الدواجن - كلية الزراعة - جامعة المنصورة.

**معهد بحوث الإنتاج الحيواني - مركز البحوث الزراعية - مصر.

تهدف هذه الدراسة الى تحديد تأثير ٣ أنواع من المواد الواقية ٤٠% إيثيلين جليكول و ٤٠% داي ميثيل سلفوكسيد وخليط منهما معا (٢٠% إيثيلين جليكول + ٢٠% داي ميثيل سلفوكسيد) مع ٣ أنواع من الأدوات المستخدمة في اجراء التجميد بالتزجج (القشاش العادية - القشاش مفتوحة الطرفين - القشاش ذات طرف واحد مفتوح) . تم اجراء هذه الدراسة على عدد ٣٦ أنثى نيوزيلاندى بيضاء متوسط عمرها ٥ شهر تقريبا ومتوسط الوزن الحي ٣,٢٥ كجم تم استخدام الإناث في هذه الدراسة للحصول على البويضاتم جمع البويضات من المبايض وتم حساب معدل استرداد البويضات من المبايض . وقد اشارت النتائج الى ان العدد الكلى للحويصلات المرئية التي تم الحصول عليها من المبايض هي ١٢٩٦ حويصلة بمتوسط ٣٦ حويصلة لكل أم و ١٨ حويصلة لكل مبيضينما كان العدد الكلى للبويضات التي تم الحصول عليها من الحويصلات للمبيضية ١١١٧ بويضة بمتوسط ٣١,٠٣ لكل أم و ١٥.٥ لكل مبيض . وكان المعدل الكلى للبويضات المستردة هي ٨٦,٢% حيث سجلت للبويضات الجيدة اعلى معدل استرداد بنسبة ٦١,٩٦% تعقبها البويضات المعراء كليا ثم البويضات المعراء جزئيا بنسبة (٦,٤٨-١١,٦٥ %) على الترتيب . بينما سجلت البويضات المتوسطة اقل معدل استرداد بنسبة ٦,١٠% وكانت نسبة التوزيع النسبي للبويضات الجيدة أعلى من البويضات المعراء كليا ثم المعراء جزئيا ثم للمتوسطة بنسبة (٧١,٨٩-١٣,٥٢-٧,٥٢-٧,٠٧%) . وقد لوضحت النتائج ان طرق الحفظ بالتزجج باستخدام القشاش ذات طرف واحد مفتوح اظهرت اعلى نتائج على معدل الحيوية للبويضات بعد التجميد وكان العدد الكلى للبويضات الطبيعية وغير الطبيعية هي (٨٥,٧٧-٦٨,٣٣-١٧,٤٤%) على الترتيب مقارنة بالطرق الأخرى .

وكان العدد الكلى للبويضات التي تم حفظها بالتزجج اعلى معنوية عند استخدام خليط من إيثيلين جليكول + داي ميثيل سلفوكسيد (٢٠% إيثيلين جليكول + ٢٠% داي ميثيل سلفوكسيد) عن البويضات التي تم حفظها باستخدام إيثيلين جليكول بمفرده أو داي ميثيل سلفوكسيد بمفرده

قام بتحكيم البحث

أ.د/ عبد الخالق السيد عبد الخالق

أ.د/محمد مصطفى الحباك

كلية الزراعة - جامعة المنصورة

خارجي