

## INTRODUCING AN ENDOPHYTE FOR CONTROLLING TOMATO EARLY BLIGHT DISEASE

Arab, Y. A.\*; Tomader G. Abd-El-Rahman\*\*and Nour-jehan M. M. Eisa\*\*

\* Botany Dept., Fac. Agric., Al-Azhar Univ., Cairo, Egypt.

\*\* Plant Pathol. Inst., Agricultural Research Center , Giza , Egypt.

### ABSTRACT

Diseased tomato samples showed typical symptoms of the early blight disease were collected from four Egyptian governorates , i.e. Sharkia, Kalubia, Gharbia and Ismailia. All *Alternaria solani* isolates affect Peto 86 tomato cultivar. Ismailia was the most aggressive isolet. Three *Bacillus subtilis* isolets (were isolated from planosphere of tomato leaves) and tow *Trichoderma viride* isolates (were isolated from rhizosphere) inhibited *A. solani* growth with different levels. The most aggressive *B. subtilis* isolate was chosen as a tool to induce disease resistance for tomato seedlings. The application of *Bacillus subtilis* on tomato seeds achieved a profuse proliferation of endophytically colonized seedlings, tomato plants were then challenged with *A. solani* suspension. The av. hight of endophytically colonized plants increased from 38 cm for the check plants to 57 cm for the treated ones. After 45 days keeping under greenhouse conditions, the endophytically colonized and challenged tomato plants exhibited a significant reduction in early blight severity (7% compared with 19 % for check plants). The endophytic bacteria was recovered from treated tomato seedlings, indicating that the endophytic *B. subtilis* has the potential to move systemically throughout the plant tissues .

**Keywords:** *Bacillus subtilis*, endophyte, tomato early blight, *Alternaria solani*, disease resistance inducing.

### INTRODUCTION

Tomato (*Lycopersicon esculentum*) is one of the most important vegetable crops in Egypt, and in other countries. The plants are subjected to attack by several diseases that cause great losses for yield production. Early blight caused by *Alternaria solani* (Eil.& Mart.) is one of the most serious diseases of all tomato diseases in Egypt (El-Saman,1986).

Recent researches reported that endophytes (microorganisms inhabit plants without causing visible disease symptoms) may exudate chemicals inhibit the growth of causal organisms, including harmful insects and mammalian herbivorus (Carroll, 1998 and Azevedo, 2000), produce some plant-growth-promoters (Varma *et al.*, 1999 and Adeline *et al.*, 2008) and induce plant resistance against different pathogens (Redman *et al.*,1999, Narisawa *et al.*,2000, Arnold *et al.*, 2003 and Henson *et al.*, 2007). There is always a degree of antagonistic balance between endophytes and their plant hosts (Schulz, Barbara and Christina Boyle, 2005) except when the host is under stress conditions, so, endophytes appear to be associated with initial degradation of the plant tissues followed by senescence (Amico, Margherita *et al.*, 2007).

Many successful trials were carried out on tomato plants including artificially introducing antagonistic endophytes to induce plant disease resistance, i.e. *Acremonium* sp. against Fusarium wilt of tomato (Brgmann and Schonbeck, 1992 and Grunewaldt-Stoecker et al., 1998), a nonpathogenic strain of *Colletotrichum* sp. against tomato anthracnose (Redman et al., 2002), *Bacillus* against tomato wilt (Algam, Soad et al., 2005), Actinomycetes against *Ralstonia solanacearum* (Tan et al., 2006) and nonpathogenic strain of *Fusarium solani* against tomato soil pathogens (Kavroulakis et al., 2007). Grunewaldt-Stoecker and Alten (2003) stated that the injected endophyte must be permanently present within plant tissues before pathogenic infection, to maintain successful systemic disease protection.

In the present study we isolated some organisms from leaves and root tissues of tomato plants to identify the possibility for being biocontrol agents against *Alternaria solani* and to induce disease resistance against tomato early blight disease, as one of the most effective methods in the integrated best management program.

## MATERIALS AND METHODS

Diseased tomato samples showed the typical symptoms of the early blight disease were collected from four Egyptian governorates, i.e. Sharkia, Kalubia, Gharbia and Ismailia. Two fields from each governorate were chosen. The samples were transmitted to the laboratory for isolation and identification. Isolation, purification using single spore technique, and identification was carried out Ellis, 1971.

### Pathogenicity test :

Pathogenicity test was carried out in the greenhouse using the identified isolates as *Alternaria solani*. The inoculum was prepared by growing the pathogenic fungus on Czapek's liquid medium for 10 days. Tomato seeds of the susceptible cultivar Peto 86 were obtained from the Agric. Res. Center, Giza. Seeds were planted in a shallow for 1.5 month, then seedlings were transplanted into green house. Inoculation was carried out by spraying the inoculum of *A. solani* isolets ( $5 \times 10^3$  conidia/ml) on plant leaves. Plants were sprayed with distilled water were used as check. Disease assessment was carried out 10 days after inoculation according to Christ (1991) as follows:

0.1= a few circular lesions ( 1 -2 mm).

0.5= up to 5 % of the leaf area spotted.

0.75= above 5 % , but below 10 % of leaf area spotted.

1= above 10 % , but below 25 % of leaf area spotted.

2= above 25 % , but below 50 % of leaf area spotted.

3= above 50 % , but below 75 % of leaf area spotted.

4= above 75% to 100 % of leaf area spotted.

Mean disease severity for a plant was calculated as the mean of disease severity values of all leaves of that plant.

#### Isolation and identification of antagonistic organisms:

*Bacillus subtilis* was isolated from planosphere of tomato leaves and identified according to Harrigan and McCance (1976), while, *Trichoderma viride* was isolated from rhizosphere of tomato roots and identified according to Rifai (1969).

#### The antagonistic tests:

The antagonistic effect of both organisms above was studied against *A. solani* *in vitro*. Regarding *Bacillus* isolates, The method described by Abo El-Naga, Heidi, 2006 was applied as follows: Plates of PDA medium was inoculated with two equal discs of *A. solani*. The antagonistic bacterium (*B. subtilis*) was streaked at the centre of each plate by a loop loaded with 48 hr old bacterial culture grown at 27°C on nutrient broth. Four plates were used for each combination. Antagonistic effect was determined by measuring zones between the antagonistic bacteria and the tested fungus. Petri dishes with *A. solani* discs only were served as control check.

However, mycelial discs (5 mm in diameter) were cut from 7-day-old PDA cultures of *A. solani* and *T. viride*, two discs (one of each fungus) were transmitted into each experimental dish (containing PDA media), and facing each other with 40 mm distance between them (one disc in each side). Four plates per each combination were incubated at 25±2°C. Dishes containing PDA medium and discs of *A. solani* only were used as control check. Inhibition zone diameters (in mm) were determined after 7 days of incubation period (Sabet and Khan, 1969).

#### Endophytically colonization of tomato seeds :

Pots (50 cm in diameter) were filled with sterilized sandy-clay soil. At the same day of planting, surface sterilized tomato seeds of the susceptible cultivar, Peto 86, were soaked five hours before planting in Petri-dish containing a 3-days-old culture of *Bacillus subtilis* (108-109 cfu/ml) and was grown in liquid tryptic soybean medium (Anonymous., 2003). [One liter of the medium contains: 17g Tryptone, 3g Soytone (enzymatic, digest of soybean meal), 2.5g Dextrose, 5g Sodium Chloride and 2.5g K<sub>2</sub>HPO<sub>4</sub>]. Ten treated seeds were planted in each pot to be artificially inoculated with *Alternaria solani*. Surface sterilized un-soaked seeds were served as a control check. Five pots were used for each treatment, Five pots were cultivated with treated seeds also prepared to study the bacterial effect on tomato growth. Tomato seedlings were sprayed by spore suspension of *A. solani* (5x10<sup>3</sup> conidia/ml) after five rows of true leave stage.

Assessment of disease severity of tomato early blight was estimated after 7 days of inoculation as described by (El-Saman, 1986). The recovery of *Bacillus subtilis* was carried out. All the seedlings were left under greenhouse conditions. Plant height was recorded at 30 and 45 days after sowing. The experiment was implemented for the two successive seasons (2007 and 2008).

#### Statistical analysis:

The obtained results were statistically analyzed according to Gomez and Gomez, 1984.

## RESULTES AND DISCUSSION

### Isolatoin and identification of the causal organism :

Tomato plants showing leathery, dark brown, necrotic lesions on the older leaves, which develop concentric black rings due to the conidia formation, were taken to isolate the causal pathogen from the affected leaves. The obtained cultures were identified according to Ellis ( 1971) as *Alternaria solani* fungus.

### Pathogenicity test :

Data present in table (1) indicate that all the tested isolates affected tomato plants. Ismailia isolate was the most aggressive one followed by Kalubia isolate, while Sharkia and Gharbia isolates were less effective, this may be due to the environmental conditions in Ismailia and Kalubia that was more favorable to the prevalent isolate of *A. solani*, and to the development of early blight disease. Similar result was obtained in the investigation of (Rashed, 1999) on the same tomato cultivar.

Table (1) : Effect of different isolates of *Alternaria solani* on early blight disease severity of tomato plants cv. Peto 86.

Governorate of isolate	Location	Disease severity
Gharbia	Kafr El-Zayat	7.5
	Mahala	7.0
Ismailia	Ismailia	23.2
	Kasaseen	20.4
Kalubia	Banha	16.5
	Kanater El-Khyria	14.3
Sharkia	Zagazig	10.5
	Hehia	9.5
LSD at 5 %		2.9

### Isolation and identification of antagonistic organisms :

Two *Trichoderma viride* isolates were isolated from rhizosphere of tomato roots and three *Bacillus subtilis* isolates were isolated from planosphere of tomato leaves. The antagonistic tests which carried out *in vitro* using discs on PDA media indicated that all the tested isolates of *B. subtilis* or *T. viride* inhibited the mycelial growth of *Alternaria solani* but the bacterium was more antagonistic than Trichoderma.

Table ( 2 ) : Effect of the antagonistic organisms on *A. solani* growth after 7days of incubation.

Treatments	Location	Av. of inhibition zone after 7 days ( in mm.)
<i>Bacillus subtilis</i>	Ismailia	21.0
	Kasaseen	17.5
	Banha	19.0
<i>Trichoderma viride</i>	Kasaseen	11.0
	Mahala	8.5
Check ( <i>A. solani</i> alone )	Ismailia	0.0
LSD at 5 %		5.5

The three *Bacillus subtilis* isolates exhibited inhibition effect with different degrees against *Alternaria solani* growth on PDA medium. The most aggressive *B. subtilis* isolate (Ismailia) was chosen as a tool to induce disease resistance in tissues of tomato seedlings. This effect agreed with the effect of using *B. pumilus* to inhibit mycelial growth of many other fungi ( Bollone and Pelaso, 2003 and Zhang *et al.*, 2007 )

**Endophytically colonization of tomato seedlings :**

The application of *Bacillus subtilis* on tomato seeds achieved many effects:

- No significant damage was noticed on endophytically colonized seedlings. This is in agreement with the meaning of the term "endophyte" of Carroll, 1998.
- There was a profuse proliferation of endophytically colonized plants. The height of endophytically colonized plants increased from 38 cm for the check plants to 57 cm. for the treated plants. So, there was a remarkable enhancement to the growth of the endophytically colonized tomato plants. This result is in harmony with the findings of Verma, 1999 and Redman *et al.*, 2002.
- Seven days after infection with *Alternaria solani*, the endophytically colonized and challenged tomato seedlings - exhibited a significant reduction in early blight disease under greenhouse conditions.

**Table (3): Blighted tomato seedlings % and plant height/seedling (in cm.) after *Bacillus* application inside plant tissues, seasons 2007 and 2008.**

Treatments	season 2007			season 2008		
	Blighted seedlings %	Plant height / seedling ( in cm. ) after		Blighted seedlings %	Plant height / seedling ( in cm. ) after	
		30 days	45 days		30 days	45 days
Protected seedlings	7.0	35.0	57.0	7.9	33.0	55.5
Control I*	19.0	23.0	38.0	20.3	21.5	36.5
Control II**	2.6	30.0	54.0	2.2	28.3	51.2
LSD at 5 %	1.5	5.3	4.2	1.02	3.12	4.0

\* Control I = Free endophyte and challenged seedlings.

\*\* Control II = Endophytically colonized and not challenged seedlings.

This result agreed with those reported by Georgy, 1977, Podile, 1993 and Abo El-Naga, Heidi, 2006. They used *Bacillus subtilis* to induce resistance against the pathogens of onion white-rot, peanut root-rot and sugar beet damping-off , respectively.

To discover the mechanisms behind inducing systemic effects of delayed and reduced tomato early blight symptoms and reduce pathogen spread by using *Bacillus* spp., Katz and Demain, 1977 stated that *Bacillus* species produce 167 biological compounds active against pathogenic bacteria, fungi, protozoa and viruses. Several peptide antifungal compounds synthesized by *Bacillus* spp. are active against filamentous fungi and yeast such as Mycocerein, Pumiucidin, Rhizocticin, Fungicin M<sub>4</sub> and Hcxacne (Wakayama *et al.*, 1984, Karuse *et al.*, 1990, Kugler *et al.*, 1990, Lebbadi *et*

al., 1994, Kudryashava et al., 2005). Also, Fiddaman and Rossak, 1993 reported that *Bacillus subtilis* produces some antifungal volatiles.

The endophytic bacteria was recovered from the imerged treated seeds after tow weeks from application, indicating that the endophytic *B. subtilis* has the potential to move throughout the plant tissues . This result is in agreement with the findings of Grunewaldt-Stoecker et al., 1998 who stated that endophytic organisms are easily detected inside the target plant tissues in the second week after injection.

## REFERENCES

- Abo El-Naga, Heidi, 2006. *Bacillus subtilis* as a biocontrol agent for controlling sugar beet damping-off disease. Egypt. J. Phytopathol., 34(1):51-59.
- Adeline, S.Y. , S. Meon, J. Kadir, S.Radu and G. Singh, 2008. Endophytic microorganisms as potential growth promoteterss of banana. Biocontrol, 53 (3) : 541-553.
- Algarn, Soad A., X.G. Lin and J. Coosemans, 2005. Delivery methods for introducing endophytic *Bacillus subtilis* into tomato and their effect on growth promotion and suppression of tomato wilt. Plant. Pathol. J., 4 (1): 6974.
- Amico, Margherita D., S.W. Frisullo and M. Cirulli, 2007. Endophytic fungi occurring in fennel, lettuce, chicory and celery commercial crops in Southern Italy. -Micological Research, 112 : 100-107.
- Anonymous, 2003. Tryptic soy agar medium. International Harmonisation Pharmeuropa, 15 (3):563-573.
- Arnold, A. Elizabeth, L.C. Mejia, D. Kyllo, E.I. Rojas, Zuleyka Maynard, Nancy Robbins and E.A. Herre, 2003. Fungal endophytes limit pathogen damage in tropical tree. Proc. Nat. Acad. Sc., 100 : 15649-15654.
- Azevedo, T.M., 2000. Endophytic microorganisms :A review on insect control and recent advances to tropical plants. E. J. B. Electronic J. of Biotechnology, 3 (1): 40-60.
- Bargmann, C. and F. Schonbeck, 1992. *Acremonium kiliens* as inducer of resistance to wilt disease on tomatoes. J. Plant Diseases and Protection, 99 : 266-272.
- Bollone, E.J. and R.W. Pelaso, 2003. Production by *Bacillus pumilus* of an antifungal compound that is active against Mucoraceae and Aspergillus species. J. Med. Microbiol., 52:69-74.
- Carroll, G.C., 1998. Fungal endophytes in stems and leaves from latent pathogen to mutualistic Symbiont. Ecol., 69 :2-9.
- Christ , B.J. ,1991. Effect of disease assessment method on Ranking potato cultivars for resistance to early blight. Plant Dis., 4 : 353.
- Ellis, M. B., 1971. Dematiaceous hyphomycetes. Kew, Surrey, Commonwealth Mycological Institute, 608 pp.
- El-Saman, M.G., 1986. Studies on early blight disease of tomato. M. Sc. Thesis, Fac. of Agric., AinShams Univ., 85 pp.
- Fiddaman, P.J. and S. Rossal, 1993. The production of antifungal volatiles by *Bacillus subtilis*. J. Appl. Bacteriol., 74 :119-126.

- Georgy, N.I. 1977. Studies on the fungus *Sclerotium cepivorum* causing white rot of onion and its control. Ph.D.Thesis, Fac. of Agric., AinShams Univ., 90 pp.
- Gomez, K.A. and Gomez, A.A. 1984. Statistical Procedures for Agricultural Research. John Wiley & Sons, Inc. New York, U.S.A. 680 pp.
- Grunewaldt-Stoecker, G. and H.V. Alten, 2003. Plant health of *Acremonium* root endophytes compared to those of arbuscular mycorrhiza. Developments in Plant and Soil Sciences, 101 : 445-454.
- Grunewaldt-Stoecker, G. , T. Kraska and E. Maiss, 1998. PCR- based detection of soil-borne and root-colonizing *Acremonium* endophytes inducing Fusarium-wilt resistance in linseed and tomato plants. Mittlg. Bundesanstalt, 357 : 260.
- Harrigan, W.F. and M. McCance ,1976. Laboratory Methods in Microbiology, Acad. Press, New York, p. 362.
- Henson, J.M., Kathy B. Sheehan , Russell J. Rodriguez and R. S. Redman, 2007. Use of endophytic fungi to treat plants. United States Patent., 7232565.
- Karuse , N., O. Tenyo, S. Kobarus, H. Kamei, T. Miyaki, M. Konishi and T. Oki, 1990. Pumilucidin, a complex of new antiviral antibiotics. Production , isolation , chemical properties, structure and biological activity. J. Antibiot.,43:267-280.
- Katz, E. and A.L.Demain, , 1977. The peptide antibiotics of *Bacillus*. Chemistry, biogenesis and possible functions. Bacteriol. Rev., 40 : 449-474.
- Kavroulakis, N., S. Ntougias, G.I. Zervakis, C.Ehalotis, K. Haralamoidis and K.K. Papadopoluou, 2007. Role of ethylene in the protection of tomato plants against soil -borne fungal pathogens conferred by an endophytic *Fusarium solani* strain . J. Exoermental Botany , 58 : 3853-3864.
- Kudryashava, E., N. Vinokurova and E. Ariskina , 2005. *Bacillus subtilis* and phenotypically similar producing Hcxacne antibiotic . Appl. Biochem. Microbiol., 41 :486-489.
- Kugler, M., W. Loeffler, C. Rapp, A.Kein, A.G.Jung, 1990. Rhizocticin A , an antifungal phosphono-oligopeptide of *Bacillus subtilis*. Arch. Microbiol., 153 :267-281.
- Narisawa, K., T. Ohki and T. Hushiba , 2000. Suppression of clubroot and Verticillium yellows in Chinese cabbage in the field by the root endophytic fungus, *Heteroconium chaetospora*. Plant Pathol., 49 : 141-146.
- Podile,A.R., 1993. Induction of host-plant resistance in peanut against soil borne pathogens by *Bacillus subtilis*. 6<sup>th</sup> International Cong. of Plant Pathol.
- Rashed, A.A., 1999. Pathological and physiological studies on tomato early blight and its control in Egypt. M. Sc.Thesis, Fac. Agric. , Al Azhar Univ., 89 pp.

- Redman, R. S., M. R. Rossinck, S. Maher, Q. C. Andrews, W. L. Schneider and R. J. Rodriguez, 2002. Field performance of cucurbit and tomato plants infected with non-pathogenic mutant of *Colletotrichum magna*. *Symbiosis*, 32 : 55-70.
- Redman, R.S. , S. Freeman, D.DR. Clifton, J. Morrel, G. Brown and R.J. Rodriguez, 1999. Biochemical analysis of plant protection afforded by a nonpathogenic endophytic mutant of *Colletotrichum magna*. *Plant Physiol.*, 119 : 795-804.
- Rifai, M. A. , 1969. A revision of the genus *Trichoderma*. *Mycol.*, 116 : 1-56.
- Sabet, K.A. and I.D. Khan, 1969. Interaction of cotton root- infecting fungi. *Cott. Gr. Rev.*, 46 : 201-222.
- Schulz, Barbara and Christin Boyle, 2005. The endophytic continuum. *Mycological Research of Cambridge Univ.*, 109 ( 9 ) : 661-686.
- Tan, H. M., L. X. Cao, Z. F. He, G. J. Su, B. Lin, S. N. Zhou, 2006. Isolation of endophytic actinomycetes from different cultivars of tomato and their activities against *Ralstonia solanacearum in vitro*. *World J. Microbiol. Biotechnol.*, 22 (2): 1275-1280.
- Verma, A., S. Verma, N. Sahay, B. Butehorn and P. Franken, 1999. *Piriformospora indica*, a cultivable plant- growth- promoting root endophyte. *App. Environ. Microbiol.*, 65 : 2741-2744.
- Wakayama, S. , F.Ishikawa, K.Oishi, 1984. Mycocerein, a novel antifungal peptide antibiotic produced by *Bacillus cereus*, antimicrob. Agents Chemother., 26 : 939-940.
- Zhang, T., Z. Q. Shi, L. B. Hu, L.g.Gheng and F. Wang, 2007. Antifungal compounds from *Bacillus subtilis* inhibiting the growth of *Aspergillus flavus*. *World J. Microbiol. And Biotechnol.*, 783-788.

### إدخال عامل حيوي لمكافحة مرض النفحة المبكرة في الطماطم

- يوسف السعيد عرب \* ، تناصر جمعه عبد الرحمن\*\* و .  
نور جيهان محمد محمود عيسى\*\*  
\* قسم النبات الزراعي ، كلية زراعة الأزهر .  
\*\* معهد أمراض النباتات ، مركز البحوث الزراعية ، الجيزة.

تم جمع عينات طماطم مصابة بأعراض مثالية لمرض النفحة المبكرة من ٤ محافظات مصرية هي الشرقية ، القليوبية ، الغربية و الإسماعيلية ، كانت كل عزلات الفطر المسبب ( الترناريا سولاني ) مؤثرة في إحداث المرض علي صنف الطماطم القابل للإصابة بيتو ٨٦ ، وكانت أشد العزلات شراسة في إحداث الإصابة هي عزلة الإسماعيلية .

تم الحصول على ثلاث عزلات من بكتيريا باسيلس ساتلس من على سطح أوراق النباتات وعزلاتان من فطر ترايكودرما فيردى من المحيط الجذري للنباتات ، وقد أظهرت جميعها قدرة تضادية عالية ضد نمو الفطر المسبب للمرض بدرجات مختلفة ، حيث اختبرت أكثر العزلات البكتيرية شراسة كأداة لاستحداث المقاومة لشتلات الطماطم . تم نفع بذور طماطم صنف بيتو ٨٦ في معلق البكتيريا باسيلس لمدة خمس ساعات ثم زراعة البذور في قصاري تحت ظروف الصوبة موسمين متتاليين ٢٠٠٧ م ، ٢٠٠٨ م وعند وصول الشتلات إلى خمس أوراق حقيقية تم إحداث العدوى بمعلق الفطر الترناريا سولاني .

كانت للمعاملة بالبكتيريا باسيلس ساتلس قبل العدوى أثرها في زيادة متوسط طول النباتات عمر ٤٥ يوما من ٢٨ سم في نباتات المقارنة إلي ٥٧ سم في النباتات المعاملة ، وخفض متوسط نمية النباتات المصابة من ١٩ % بالنسبة لتجربة المقارنة إلي ٧ % في تجربة المعاملة ، كما أمكن عزل البكتيريا من داخل نسيج النباتات مما يدل علي إمكانية تحرك البكتيريا جهازيا داخل نسيج النبات.