

# NUTRITION & HUSBANDRY

## FOR AND AGAINST THE USE OF CONDENSED TANNINS FOR SMALL RUMINANT PRODUCTION – A CRITICAL ANALYSIS

Zafar Iqbal<sup>a</sup>, Muhammad Sarwar<sup>1</sup>, Muhammad Sohail Sajid, Rao Zahid Abbas, Muhammad Kasib Khan, Muhammad Aasif Shahzad<sup>1</sup> and Muhammad Nisar Khan

*Department of Parasitology, University of Agriculture, Faisalabad-38040, Pakistan*

*<sup>1</sup>Institute of Animal Nutrition and Feed Technology, University of Agriculture, Faisalabad-38040, Pakistan*

<sup>a</sup> Corresponding author: Professor Dr. Zafar Iqbal

E-mail: [zafaruaf1@gmail.com](mailto:zafaruaf1@gmail.com)

### ABSTRACT

Indo-Pak subcontinent has a rich history of small ruminant farming. There are many indigenous breeds of sheep and goat contributing to the national economy and food security of the region. Small ruminant farming reared by the pastoralists, settlers and nomadic people mainly relies on the rangelands and temporary community pastures. Historically, the shepherds offer their flocks the leaves and soft stems of many trees and shrubs, now known to have tannin contents. The performance of the animals has always found better with the animals fed on traditional forages containing tannins. Until last couple of decades, there was no scientific evidence of the beneficial and/or adverse effects of tannins. There are, however, many researches published during the last few years, which have reflected upon the multiple benefits or harmful effects of the condensed tannins in small ruminants. More of the published research advocates the use of tannins for enhancing the animal production. Nevertheless, a debate is needed to use, disuse, judicious use and significance of animal feeds supplemented with tannin or the tanniniferous forages. This paper offers a critical analysis of the conclusions based on the research on tannin feeding and potential of tannin containing forages in improving nutrient utilization and availability of proteins for better performance in small ruminants. Suitable recommendations have been made based on logical discussion on “*pros and cons*” of tannin feeding.

*Key Words:* Tannins, Benefits, Adverse effects, Small ruminants

## INTRODUCTION

Tannins are polyphenolic compounds, which may be present in any part of plant, including roots, stems or trunk, bark, leaves, fruit and even hairs. Based on their molecular structure, they are usually classified either as hydrolysable tannins (HT) or CT (proanthocyanidins). Among these, CT are the most common types of tannin found in forage legumes, trees and shrubs (Barry and McNabb, 1999). Structurally, CT are complexes of oligomers and polymers of flavanoid units linked by carbon-carbon bonds (Hagerman and Butler, 1981; Foo *et al.*, 1986) without a monosaccharide nucleus. The CT exist as oligomers of flavan-3-ols (catechin) or flavan-3, 4-diols (epicatechin) and those occurring in temperate forages have a relative molecular mass of 2,000 to 4,000 comprising 10 to 12 oligomers of CT (Foo *et al.*, 1986). These differences can produce an infinite variety of chemical structures, which in turn affect the physical and biological properties of the CT. Condensed tannins form pH reversible bonds with forage proteins, which reduce degradation of protein to ammonia by rumen microorganisms, yet release protein at the low pH (2.5–3.5) in the abomasum ( Jones and Mangan, 1977). Reactivity of CT differs between species of plants that contain these compounds and hence CT from some plants increases the net absorption of essential amino acids more than others (Barry and Mc Nabb, 1999; Min *et al.*, 2003).

Small ruminant production is an important agricultural small and medium enterprise of the farmers. Tremendous increase in cultivated lands and increasing animal population pressure have led to squeezing the rangelands for grazers. Thus, animal nutritionists have been actively engaged in exploiting the nutritional potential of traditional forages to pay off the shortage of good quality food. Indo-Pak subcontinent has a rich history of small ruminant farming with many indigenous breeds of sheep and goat which contribute not only the national economy but also as food source. Rangelands and temporary community pastures are the main source of food for small ruminants reared by the pastoralists, settlers and nomadic people. Historically, the shepherds offer the leaves and soft stems of many trees and shrubs to their flocks which have, now known to contain tannin contents. The performance of the animals has always found better with the animals fed on traditional forages containing tannins. Until last couple of decades, there was no scientific evidence of the beneficial and/or adverse effects of tannins. There are, however, many researches published during the last few years, which have reflected upon the multiple benefits or harmful effects of the condensed tannins in small ruminants (Aerts *et al.*, 1999). More of the published research advocates the use of tannins for enhancing the animal production. Nevertheless, a debate is needed to use, disuse, judicious use and significance of animal feeds supplemented with tannin or the tanniniferous forages.

This paper presents a critical analysis of the conclusions based on the research conducted to enhance the small ruminant production by improving nutrient utilization and availability of proteins through feeding and exploiting the potential of tannin

## NUTRITION & HUSBANDRY

containing forages. Moreover, based on discussion on "pros and cons" of tannin feeding, suitable recommendations have been made to operate the animal performance.

### **Biological features of tannins**

Tannins are functionally known by their competence to bind with proteins which forms the bases of many biological effects of tannins (Hagerman and Butler, 1991). They can have detrimental effect against many microorganisms and fungi (Bernays *et al.*, 1989) which may be one of the major reasons of their evolution (Swain, 1979; Bernays *et al.*, 1989; Ayres *et al.*, 1997; Aerts *et al.*, 1999). Intense deposits of tannin contents occur in the epidermis of leaves and stems of many leguminous forage plants, herbs and grasses with different concentrations including *Onobrychis viciifolia* (sainfoin), *Lotus (L.) corniculatus* (birdsfoot trefoil), *L. pedunculatus* (big trefoil), *Hedysarum coronarium* (sulla), and *Lespedeza cuneata* (sericea lespedeza) (Jones *et al.*, 1976; Terrill *et al.*, 1989, 1992). Interestingly, it has been reported that plants higher in tannins produced fewer leaves as compared to those having low tannins (Coley, 1986).

### **Beneficial and toxic effects of condensed tannins**

The beneficial and toxic effects of condensed tannins on production and performance have extensively been studied in small ruminants. However, it appears that tanniferous feeds can produce beneficial effects in low concentration while high levels are inevitably detrimental in small ruminants (Wiegand *et al.*, 1996; Min *et al.*, 2003). Low to moderate concentrations of tannins (2-5 % dry weight) significantly increased the net absorption of amino acids, sulphur and nitrogen digestibility which ultimately enhanced wool growth, ovulation rate, live weight gain, growth rate and milk production and quality (Waghorn *et al.*, 1987; McNabb *et al.*, 1993; Wang *et al.*, 1994, 1996a; Terrill *et al.*, 1992; Min *et al.*, 1999). Moreover, tannins have also been reported to possess anthelmintic activity against many helminthes including *Teladorsagia circumcincta*, *Haemonchus contortus* and *Trichostrongylus clubriiformis* (Niezen *et al.*, 1998; Paolini *et al.*, 2003a; Molan *et al.*, 1999; Molan *et al.*, 2000; Waghorn and Molan, 2001; Molan *et al.*, 2002; Paolini *et al.*, 2005). In contrast, high concentrations of tannins (>5 % dry weight) in feed of small ruminants may lead to reduced voluntary feed intake, % digestion of amino acids in small intestine, rumen fiber digestibility, growth rate, wool growth and digestion of sulphur and nitrogen (Barry and Duncan, 1984; Barry 1985; Barry *et al.*, 1986; Waghorn *et al.*, 1994; Pritchard *et al.*, 1988, 1992). A summary of beneficial and detrimental effects of tannins in relation with their concentrations is shown in the Table 1.

### **Possible explanations of tannin effects**

#### **a. Nutritional aspect in sheep**

To determine the nutritional effects of tannins in the sheep, all the experiments

were performed on both untreated and tannin inactivated plant materials that have been treated with polyethylene glycol (MW 3350). The beneficial effects of tannins in sheep are associated with the greater outflow and absorption of amino acids especially in sheep fed with the forages containing tannin percentage ranging from 2-4% (Waghorn *et al.*, 1987; Wang *et al.*, 1994, 1996b; Min *et al.*, 1999). Enhanced growth rate and increase in production performance and nutritive value of milk in sheep may be due to increased availability of essential amino acids (Waghorn *et al.*, 1987; Wang *et al.*, 1994). Increase outflow of sulphur containing amino acids which are key precursors of wool production may contribute to increased wool production (Wang *et al.*, 1994; McNabb *et al.*, 1993). Although, microorganisms are essential agents in the rumen for degradation of structural components of plant cells but, on the other hand, they also utilize the ammonia and some essential amino acids for synthesis of their cellular proteins. Tannins make complexes with the proteins in which prevent the degradation of proteins in rumen thus increase flow of proteins to the intestines (Waghorn *et al.*, 1987, 1994; McNabb *et al.*, 1996). This is evident from the experimentations of Barry *et al.* (1986) who found a linear relationship between increased tannin concentration and increased surge of non-ammonia nitrogen. Moreover, the pH value in rumen (6.0-7.0) is very favorable for the formation of stable complexes between tannin and proteins. When the complexes come in intestines, the lower pH (2.5-3.5) separates the bond between tannin contents and proteins resulting in enhanced digestion of essential amino acids in the intestines of sheep (Waghorn *et al.*, 1987, 1994). This greater quantity of proteins is available to be absorbed in the intestine of sheep. In contrast, high tannin concentration reduces percentage digestion of proteins thus leads to reduced growth rate, wool production and milk quality and quantity (Waghorn *et al.*, 1994).

#### **b. Control of parasites**

Although, most of the parasitic control programs are based on chemotherapeutic control (Waller, 1999; FAO, 2002) but, various problems have been evolved with this practice such as increasing problem of development of resistance by the parasites to several families of drenches (McKenna *et al.*, 1995; Vermunt *et al.*, 1995; Chandrathani *et al.*, 1999; Chartier *et al.*, 2001; Leathwick *et al.*, 2001), hazards of chemical residues and toxicity (Kaemmerer & Buitenkotter, 1973), un-economical, non-adaptability and non-availability of drugs in remote areas. Recent reports on the small ruminants suggest that tannin containing fodder decrease the detrimental effect of gastrointestinal parasites by killing larval and adult worms (Athanasiadou *et al.*, 2000). For example, Athanasiadou *et al.* (2001) found significantly lower faecal egg count from the sheep drenched with 16% W/W of Quebracho extract (QE). However, worm burden in the digesta of abomasums was significantly lower in sheep treated with 8% W/W of QE. Nevertheless, enhanced feed intake, live weight gain and feed conversion efficacy was observed in sheep treated with 8% W/W of QE while 90% of sheep drenched with 16% W/W of QE stopped eating by day 30 (Athanasiadou *et al.*, 2001).

These results could be attributable to toxic effects of tannins as higher concentrations of tannins reduce feed intake and absorption of other available nutrients in rumen of sheep (Reed, 1995). Similarly, various scientists observed lower faecal count and worm burden with no parasitic species difference in the sheep drenched with several concentrations of condensed tannins (Niezen *et al.*, 1998; Paolini *et al.*, 2003a, 2005; Molan *et al.*, 1999, 2000, 2002; Waghorn and Molan, 2001). Tannins bind with free available proteins in the gastrointestinal tract and reduced nutrient availability of nutrients would have resulted in larval and worm starvation and death (Athanasiadou *et al.*, 2001). Additionally, tannins would also bind with cuticle of larvae which is high in glycoproteins ensuing in death of larvae (Thompson and Geary, 1995). Moreover, increased availability of proteins due to condensed tannins in the feed of sheep has also been found responsible for enhanced immune response against parasites (Coop and Kyriazakis, 1999).

### **c. Bloat safety**

Bloat is frequent muddle in small ruminants caused by the formation of stable protein broth in the rumen of animals fed with high nutritive value legumes including white clover or Lucerne. These protein foams avert the fermented gases to liberate from the rumen resulting in expansion of rumen. The course of bloat formation is very acute and leads to serious damage of vital organs such as lungs and heart (Mangan, 1959) and ultimately leading to death of animals. While, moderate concentration of tannins in the food of animals destabilizes the protein foams which refers them bloat safe (Tanner *et al.*, 1995).

### **d. Reduced proteolytic enzyme activity and growth of rumen bacteria**

CTs considerably reduce the proteolytic enzyme activity and growth of bacteria in the rumen of sheep (Jones *et al.*, 1993). CTs form complexes with the cell coat polymers of bacteria and proteolytic enzymes secreted by them which enable the protein to sidestep in the rumen. These complexes subsequently release the protein when come in the acidic condition of abomasum. These protein molecules undergo enzymatic hydrolysis in the small intestine leading to availability of enormous number of amino acids to be absorbed from the intestine (Jones and Mangan, 1977; Martin and Martin, 1983; McNabb *et al.*, 1998). Although, studies have not been conducted to determine the effect of CTs on proteolytic activity and growth of all ruminal microbes but data published so far is demonstrating that CTs had not affected the total microbial protein synthesis in the rumen or abomasal microbial protein flow (McSweeney *et al.*, 2001; Min *et al.*, 2002, 2003)

## CONCLUSIONS

Adverse effects of tannin feeding in ruminants is associated with their low palatability and affinity to create linkage with nutrients. Tannin interference with digestion is a function of astringency, concentration and potential sites for binding. However, use of tanniferous diets in proper proportion and concentration can improve animal productivity in addition to alleviating their negative effects.

## RECOMMENDATIONS

1. Relatively low concentration of tannins (0.5% of DM intake) is sufficient to destabilize the bloat proteins while high concentration (2-4% of DM intake) is needed for improvement of protein utilization.
2. High concentration (> 5% of dry weight) should be avoided to prevent its toxic effects like reduced feed intake and feed conversion efficiency.
3. Tannins containing forages should be added in the feed of small ruminants for sustainable control of gastrointestinal parasites.

## REFERENCES

- Ademola, I.O., B.O. Fegbemi and S.O. Idowu, 2005. Anthelmintic activity of Extracts of *Spondias mombin* against gastrointestinal nematodes of sheep: Studies *in vitro* and *vivo*. *Trop. Anim. Hlth. Prod.*, 37: 223–235.
- Ademola, I.O., S.O. Fagbemi and S.O. Idowu, 2004. Evaluation of anthelmintic activity of *Khaya senegalensis* extract against gastrointestinal nematodes of sheep: *in vitro* and *vivo* studies. *Vet. Parasitol.*, 122: 151–164.
- Aerts, R.J., T.N. Barry and W.C. McNabb, 1999. Polyphenols and agriculture: beneficial effects of proanthocyanidins in forages. *Agri. Ecosyst. Environ.*, 75: 1–12.
- Athanasiadou, S., I. Kyriazakis, F. Jackson and R.L. Coop, 2000. Effects of short-term exposure to condensed tannins on an adult *Trichostrongylus colubriformis*. *Vet. Rec.*, 146: 728–732.
- Athanasiadou, S., I. Kyriazakis, F. Jackson and R.L. Coop, 2001. Direct anthelmintic effects of condensed tannins towards different gastrointestinal nematodes of sheep: *in vitro* and *in vivo* studies. *Vet. Parasitol.*, 99: 205–219.
- Ayres, M.P., T.P. Clausen, S. MacLean, A.M. Redman and P.B. Reichardt, 1997. Diversity of structure and antiherbivore activity in condensed tannins. *Ecology*, 78: 1696–1712.

## NUTRITION & HUSBANDRY

- Barry, T.N. and S.J. Duncan, 1984. The role of condensed tannins in the nutritional value of *Lotus pedunculatus* for sheep 1 Voluntary intake. *British J. Nutr.*, 51: 485–491.
- Barry, T.N., 1985. The role of condensed tannins in the nutritional value of *Lotus pedunculatus* for sheep 3. Rates of body and wool growth. *British J. Nutr.*, 54: 211–217.
- Barry, T.N. and W.C. McNabb, 1999. The implications of condensed tannins on the nutritive value of temperate forages fed to ruminants. *Br. J. Nutr.*, 81: 263–272.
- Barry, T.N., T.R. Manley and S.J. Duncan, 1986. The role of condensed tannins in the nutritional value of *Lotus pedunculatus* for sheep. 4. Sites of carbohydrate and protein digestion as influenced by the dietary reactive tannin concentration. *British J. Nutr.*, 55: 123–137.
- Bernays, E.A., G. Cooper Driver and M. Bilgener, 1989. Herbivores and plant tannins. In: Begon, M., Fitter, A.H., Ford, E.D., MacFadyen, A. (Eds.), *Advances in Ecological Research*, vol. 19. Academic Press, London, pp. 263–302.
- Brooker, J.D., L.O. O'Donovan, I. Skene and G. Sellick, 2000. Mechanisms of tannin resistance and detoxification in the rumen. In: Brooker, J.D. (Ed.), *Proc. of the Int. Workshop on Tannins in Livestock and Human Nutrition*. ACIAR Proceedings No. 92. Australian Centre for International Agricultural Research.
- Chandrathani, P., M. Adnan and P.J. Waller, 1999. Anthelmintic resistance in sheep and goat farms in Peninsula Malaysia. *Vet. Parasitol.*, 82: 305–310.
- Chartier, C., F. Soubirac, I. Pors, A. Silvestre, J. Hubert, C. Couquet and J. Cabaret, 2001. Prevalence of anthelmintic resistance in gastrointestinal nematodes of dairy goats under extensive management conditions in southwestern France. *J. Helminthol.*, 75: 325–330.
- Coley, P.D., 1986. Costs and benefits of defense by tannins in a neotropical tree. *Oecologia* 70: 238–241.
- Coop, R.L. and I. Kyriazakis, 1999. Parasite-nutrition interaction. *Vet. Parasitol.*, 84: 187–204.
- FAO, 2002. Biological control in a global perspective –a review on emphasis on *Duddingtonia flagrans*. *Final Proceeding of FAO, Technical Co-operation Project in Malaysia*.
- Foley, W.J. and I.D. Hume, 1987. Digestion and metabolism of high tannin Eucalyptus foliage by the brush tail possum (*Trichosarus vulpecula*) (*Marsupialia; Phalangeridae*). *J. Comp. Physiol.*, 157: 67–76.
- Foo, L.Y., W.T. Jones, L.J. Porter and V.N. Williams, 1986. Proanthocyanidin polymers of fodder legumes. *Phytochemistry*, 21: 933–935.

**3<sup>rd</sup> International Scientific Conference on Small Ruminant Development, Hurghada,  
Egypt, 12-15 April, 2010**

- Hagerman, A.E. and L.G. Butler, 1981. The specificity of proanthocyanidin-protein interactions. *J. Biol. Chem.*, 256: 4494–4497.
- Hagerman, A.E. and L.G. Butler, 1991. Tannins and lignins. In: Rosental, G.A., Berenbaum, M.R. (Eds.), *Herbivores: Their Interaction with Secondary Plant Metabolites*, 2nd Edition. Academic Press, San Diego, pp. 355–376.
- Haslam, E., 1996. Natural polyphenols (vegetable tannins) as drugs: possible modes of action. *J. Nat. Prod.*, 59: 205–215.
- Hoskin, S.O., P.R. Wilson, T.N. Barry, W.A.G. Charleston and G.C. Waghorn, 2000. Effect of forage legumes containing condensed tannins on lungworm (*Dictyocaulus* sp.) and gastrointestinal parasitism in young red deer (*Cervus elaphus*). *Res. Vet. Sci.*, 68: 223–230.
- Jones, W.T., R.B. Broadhurst and J.W. Lyttleton, 1976. The condensed tannins of pasture legume species. *Phytochemistry*, 15: 1407–1409.
- Jones, G.A., K.D. Jakober, H.D. Bae, T.A. McAllister and K.J. Cheng, 1993. Some interactions between condensed tannins of forage legumes, bovine serum albumin, and five strains of proteolytic rumen bacteria. In: *Proceedings of the World Conference on Animal Production*, vol. 36, Edmonton, Canada, pp. 68–69.
- Jones, J.T. and J.L. Mangan, 1977. Complexes of the condensed tannins of sainfoin (*Onobrychis viciifolia* Scop.) with fraction 1 leaf protein and with submaxillary mucoprotein and their reversal by polyethylene glycol and pH. *J. Sci. Food Agric.*, 28: 126–136.
- Kaemmerer, K. and S. Butenkotter, 1973. The problem of residues in meat of edible domestic animals after application or intake of organophosphate esters. *Residue Rev.*, 46: 1.
- Kahiya, C., S. Mukaratirwa and S.M. Thamsborg, 2003. Effects of *Acacia nilotica* and *Acacia karoo* diets on *Haemonchus contortus* infection in goats. *Vet. Parasitol.*, 115: 265–274.
- Leathwick, D.M., W.E. Pomroy and A.C.G. Heath, 2001. Anthelmintic resistance in New Zealand. *New Zealand Vet. J.*, 49: 227–235.
- Mangan, J.L., 1959. Bloat in cattle XI. The foaming properties of proteins, saponins, and rumen liquor. *NZ J. Agric. Res.*, 2: 47–61.
- Martin, J.S. and M.M. Martin, 1983. Tannin assays in ecological studies precipitation of ribulose-1,5-bisphosphate carboxylase/oxygenase by tannic acid, quebracho, and oak foliage extracts. *J. Chem. Ecol.*, 9: 285–294.
- McKenna, P.B., C.M. Allan, M.J. Taylor and K.G. Townsend, 1995: The prevalence of



- anthelmintic resistance in ovine case submissions to animal health laboratories in New Zealand in 1993. *New Zealand Vet. J.*, 43: 96–98.
- McNabb, W.C., J.S. Peters, L.Y. Foo, G.C. Waghorn and F.S. Jackson, 1998. Effect of condensed tannin prepared from several forages on the in vitro precipitation of ribulose-1,5-bisphosphate carboxylase (Rubisco) protein and its digestion by trypsin (EC 2.4.21.4) and chymotrypsin (E.C. 2.4.21.1). *J. Sci. Food Agric.*, 77: 201–212.
- McNabb, W.C., G.C. Waghorn, T.N. Barry and I.D. Shelton, 1993. The effect of condensed tannins in *Lotus pedunculatus* on the digestion and metabolism of methionine, cystine and inorganic sulphur in sheep. *British J. Nutr.*, 70: 647–661.
- McNabb, W.C., Waghorn, G.C., Peters, J.S., Barry, T.N., 1996. The effect of condensed tannins in *Lotus pedunculatus* on the solubilization and degradation of ribulose-1,5-bisphosphate carboxylase (EC 4.1.1.39; Rubisco) protein in the rumen and the sites of Rubisco digestion. *Br. J. Nutr.*, 76: 535–549.
- McSweeney, C.S., B. Palmer, D.M. McNeill and D.O. Krause, 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Anim. Feed Sci. Tech.*, 91: 83–93.
- Min, B.R., T.N. Barry, G.T. Attwood and W.C. McNabb, 2003. The effect of condensed tannins on the nutrition and health of ruminants fed fresh temperate forages: a review. *Anim. Feed Sci. Technol.*, 106: 3–19.
- Min, B.R., D. Miller, S.P. Hart, G. Tomita, E. Loetz and T. Sahlu, 2003a. Direct effects of condensed tannins on gastrointestinal nematodes in grazing Angora goats. *J. Anim. Sci.*, 81(Suppl. 2): 23. (Abstr.)
- Min, B.R., W. Pomroy, S. Hart and T. Sahlu, 2002. The effect of forage condensed tannins on gastrointestinal parasite infection in grazing wether goats. *J. Anim. Sci.*, 80 (Suppl. 1): 31. (Abstr.)
- Min, B.R., W.C. McNabb, T.N. Barry, P.D. Kemp, G.C. Waghorn and M.F. McDonald, 1999. The effect of condensed tannins in *Lotus corniculatus* upon reproductive efficiency and wool production in sheep during late summer and autumn. *J. Agri. Sci.*, 132: 323–334.
- Min, B.R., W.E. Pomroy, S.P. Hart and T. Sahlu, 2004. The effect of short-term consumption of a forage containing condensed tannins on gastro-intestinal nematode parasite infections in grazing wether goats. *Small Rumin. Res.*, 51: 279–283.
- Molan, A.L., G.C. Waghorn and W.C. McNabb, 1999. Condensed tannins and gastro-intestinal parasites in sheep. *Proc. New Zealand Grassland Assoc.*, 61: 57–61.
- Molan, A.L., G.C. Waghorn and W.C. McNabb, 2002. Effect of condensed tannins on egg

**3<sup>rd</sup> International Scientific Conference on Small Ruminant Development, Hurghada,  
Egypt, 12-15 April, 2010**

hatching and larval development of *Trichostrongylus colubriformis* in vitro. *Vet. Rec.*, 19: 65–69.

Molan, A.L., G.C. Waghorn, B.R. Min and W.C. McNabb, 2000. The effect of condensed tannins from seven herbages on *Trichostrongylus colubriformis* larval migration in vitro. *Folia Parasitologica*, 47: 39–44.

Niezen, J.H., G.C. Robertson, G.C. Waghorn and W.A.G. Charleston, 1998. Production, fecal egg counts and worm burdens of ewe lambs which grazed six contrasting forages. *Vet. Parasitol.*, 80: 15–27.

Niezen, J.H., G.C. Waghorn and W.A.G. Charleston, 1998a. Establishment and fecundity of *Ostertagia circumcincta* and *Trichostrongylus colubriformis* in lambs fed lotus (*Lotus pedunculatus*) or perennial ryegrass (*Lolium perenne*). *Vet. Parasitol.*, 78: 13–21.

Niezen, J.H., W.A.G. Charleston, H.A. Robertson, D. Shelton, G.C. Waghorn and R. Green, 2002. The effect of feeding sulla (*Hedysarum coronarium*) or lucerne (*Medicago sativa*) on lamb parasite burdens and development of immunity to gastrointestinal nematodes. *Vet. Parasitol.*, 105: 229–245.

Niezen, J.H., T.S. Waghorn, W.A.G. Charleston and G.C. Waghorn, 1995. Growth and gastrointestinal nematode parasitism in lambs grazing either lucerne (*Medicago sativa*) or sulla (*Hedysarum coronarium*) which contains condensed tannins. *J. Agric. Sci.*, 125: 281–289.

Paolini, A., I. Fouraste and H. Hoste, 2004. *In vitro* effects of three woody plant and sainfoin extracts on 3<sup>rd</sup> stage larvae and adult worms of three gastrointestinal nematodes. *Parasitol.*, 129: 69–77.

Paolini, V., F. De La Farge, F. Prevot, Ph. Dorchies and H. Hoste, 2005. Effects of the repeated distribution of sainfoin hay on the resistance and the resilience of goats naturally infected with gastrointestinal nematodes. *Vet. Parasitol.*, 127: 77–283.

Paolini, V., J.P. Bergeaud, C. Grisez, F. Prevot, Ph. Dorchies and H. Hoste, 2003. Effects of condensed tannins on goats experimentally infected with *Haemonchus contortus*. *Vet. Parasitol.*, 113: 253–261.

Paolini, V., P. Dorchies and H. Hoste, 2003a. Effect of sainfoin hay on gastrointestinal nematode infections in goats. *Vet. Rec.*, 152: 600–601.

Pritchard, D.A., D.C. Stocks, B.M. O'Sullivan, P.R. Martin, I.S. Hurwood and P.K. O'Rourke, 1988. The effect of polyethylene glycol (PEG) on wool growth and liveweight of sheep consuming Mulga (*Acacia aneura*) diet. *Proc. Aust. Soc. Anim.*

## NUTRITION & HUSBANDRY

- Prod.*, 17: 290–293.
- Pritchard, D.A., P.R. Martin and P.K. O'Rourke, 1992. The role of condensed tannins in the nutritional value of Mulga (*Acacia aneura*) for sheep. *Aust. J. Agri. Res.*, 43: 1739–1746.
- Redwane, A., H.B. Lazrek, S. Bouallam, M. Markouk, H. Amarouch and M. Jana, 2002. Larvicidal activity of extracts from *Quercus lusitania* var. *infectoria* galls (Oliv.). *J. Ethnopharmacol.*, 79: 261–263.
- Reed, J.D., 1995. Nutritional toxicology of tannins and related polyphenols in forage legumes. *J. Anim. Sci.*, 73: 1516–1528.
- Swain, T., 1979. Tannins and lignins. In: Rosenthal, G.A. and Janzen, D.H. (Eds.), *Herbivores: Their Interaction with Secondary Plant Metabolites*, Academic Press, New York, pp. 657–682.
- Tanner, G.J., P.J. Moate, L.H. Davis, R.H. Laby, Y.-G. Li, and P.J. Larkin, 1995. Proanthocyanidins (condensed tannin) destabilize plant protein foams in a dose dependent manner. *Aust. J. Agric. Res.*, 46: 1101–1109.
- Terrill, T.H., W.R. Windham, C.S. Hoveland and H.E. Amos, 1989. Forage preservation method influences on tannin concentration, intake, and digestibility of sericea lespedeza by sheep. *Agron.*, J. 81: 435–439.
- Terrill, T.H., G.B. Douglas, A.G. Foote, R.W. Purchas, G.F. Wilson and T.N. Barry, 1992. Effect of condensed tannins upon body growth and rumen metabolism in sheep grazing *Hedysarum coronarium* and perennial pasture. *J. Agri. Sci.*, 119: 265–273.
- Thompson, D. and T. Geary, 1995. The structure and function of helminth surfaces. In: Marr, J., M. Muller, (Eds.), *Biochemistry and Molecular Biology of Parasites*. Academic Press, pp. 203–232.
- Ulyatt, M.J., J.A. Lancashire and W.T. Jones, 1976. The nutritive value of legumes. *Proc. New Zealand Grassl. Assoc.*, 38: 107–118.
- Vermunt, J.J., D.M. West and W.E. Pomroy, 1995. Multiple resistance to ivermectin and oxfendazole in *Cooperia* species in cattle. *Vet. Rec.*, 137: 43–45.
- Waghorn, G.C. and A.L. Molan, 2001. Digestion and effect of condensed tannins in *Dorycnium rectum* on its nutritive value and on Anthelmintic properties. *Proc. New Zealand Grass. Assoc.*, 63: 273–277.
- Waghorn, G.C., A. John, W.T. Jones and I.D. Shelton, 1987a. Nutritive value of *Lotus corniculatus* L. containing low and medium concentrations of condensed tannins for sheep. *Proc. New Zealand Soc. Anim. Prod.*, 47: 25–30.
- Waghorn, G.C. and W.T. Jones, 1989. Bloat in cattle 46. Potential of dock (*Rumex obtusifolius*) as an antibloat agent for cattle. *NZ J. Agric. Res.*, 32: 227–235

- Waghorn, G.C., Shelton, I.D., McNabb, W.C., McCutcheon, S.N., 1994. Effects of condensed tannins in *Lotus pedunculatus* on its nutritive value for sheep 2. Nitrogenous aspects. *J. Agric. Sci. (Cam)*, 123: 109–119.
- Waghorn, G.C., M.J. Ulyatt, A. John and M.T. Fisher, 1987. The effect of condensed tannins on the site of digestion of amino acids and other nutrients in sheep fed on *Lotus corniculatus* L. *Br. J. Nutr.*, 57: 115–126.
- Waller, P.J., 1999. International approaches to the concept of integrated control of nematode parasites of livestock. *Int. J. Parasitol.*, 29: 155–164.
- Wang, Y., G.B. Douglas, G.C. Waghorn, T.N. Barry, A.G. Foote and R.W. Purchas, 1996a. Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*). *J. Agri. Sci.*, 126: 87–98.
- Wang, Y., G.B. Douglas, G.C. Waghorn, T.N. Barry and A.G. Foote, 1996b. Effect of condensed tannins in *Lotus corniculatus* upon lactation performance in ewes. *J. Agric. Sci. (Cam)*, 126: 353–362.
- Wang, Y., G.C. Waghorn, T.N. Barry and I.D. Shelton, 1994. The effect of condensed tannins in *Lotus corniculatus* on plasma metabolism of methionine, cystine, cystine and inorganic sulphate by sheep. *Br. J. Nutr.*, 72: 923–935.
- Wang, Y., G.C. Waghorn, W.C. McNabb, T.N. Barry, M.J. Hedley and I.D. Shelton, 1996c. Effect of condensed tannins in *Lotus corniculatus* upon the digestion of methionine and cysteine in the small intestine of sheep. *J. Agri. Sci.*, 127: 413–421.
- Wiegand, R.O., J.D. Reed, D.K. Combs and A.N. Said, 1996. Leaves from tropical trees as protein supplements in diets for sheep. *Trop. Agric.*, 73: 62–68.
- Windham, W.R., J.C. Peterson and T.H. Terrill, 1990. Tannin as anti-quality factor in forage. pp. 127–135. In: D.E. Akin, L.G. Ljungdahl, J.R. Wilson and P.J. Harris, (Eds.), *Microbial and Plant Opportunities to Improve Lingocellulose Utilization by Ruminant*. Elsevier Science publishing Co., inc., New York.
- Woodward, S.L., M.J. Auld, P.J. Laboyrie and E.B.L. Jansen, 1999. Effect of *Lotus corniculatus* and condensed tannins on milk yield and milk composition of dairy cows. *Proc. New Zealand Soc. Anim. Prod.*, 59: 152–155.

## NUTRITION & HUSBANDRY

**Table 1. Condensed tannin content and/or beneficial/toxic action of different plants**

Plant	Content (% DM)	Toxic/beneficial effect	Reference (s)
<i>Acacia aneura</i>	–	Damage to the structure of intestinal villi for a prolonged period	Brooker <i>et al.</i> , 2000
<i>Acacia aneura</i>	12.00	Reduced dry matter intake, wool growth, and nitrogen and sulphur digestion	Pritchard <i>et al.</i> , 1988, 1992
<i>Acacia aneura</i>	> 5	Antinutritional effects	Pritchard <i>et al.</i> , 1988
<i>Acacia karoo</i>	2.22	Reduction in feed conversion efficiency (FCE)	Kahiya <i>et al.</i> , 2003
<i>Acacia nilotica</i>	0.17		
<i>Festuca arundinacea</i>	0.20	Lesser immune response as compared to <i>Lespedeza cuneata</i>	Min <i>et al.</i> , 2003a
Dietary CT	–	May enhance animals' immune response to the parasite infection	Niezen <i>et al.</i> , 2002; Min <i>et al.</i> , 2003a
<i>Eucalyptus Melliodora</i>	> 5.00	Anti nutritional effects	Foley and Hume, 1987
<i>Hedysarum coronarium</i>	–	Higher antibody titers of secretory-excretory antigens against adult worms	Niezen <i>et al.</i> , 2002
<i>Hedysarum coronarium</i>	10.00	Reduced FCE and parasite burdens	Niezen <i>et al.</i> , 1995; 1998, 1998a
<i>Hedysarum coronarium</i>	5.20	Fed to sheep and goats, had a higher nutritive value	Ulyatt <i>et al.</i> , 1976; Niezen <i>et al.</i> , 1995; Min <i>et al.</i> , 2003a
<i>Lespedeza cuneata</i>			
<i>Hedysarum coronarium</i>	4.00–5.00	Increased wool growth rate	Terrill <i>et al.</i> , 1992
<i>Hedysarum coronarium</i>	–	Diarrhoea associated with a heavy parasite burden could be alleviated by feeding PA-containing	Niezen <i>et al.</i> , 1995
<i>Khaya</i>	–	Anthelmintic activity	Ademola <i>et al.</i> ,

Plant	Content (% DM)	Toxic/beneficial effect	Reference (s)
<i>senegalensis</i>			2004
<i>Lotus corniculatus</i>	–	Higher live weight gain, carcass weight and carcass dressing-out percentage, higher serum total protein and albumin concentration, and lower serum gastrin concentration and faecal lungworm larval count, compared with lucerne-fed deer	Hoskin <i>et al.</i> , 2000
<i>Lotus corniculatus</i>	–	Protect plant proteins against rumen degradation and increase the flow of amino acids to the small intestine	Waghorn <i>et al</i> 1987, 1994
<i>Lotus corniculatus</i>	2.00– 4.00	Increased absorption of essential amino acids from the small intestine by 62%	Waghorn <i>et al.</i> , 1987a
<i>Lotus corniculatus</i>	–	Increased milk and wool production in sheep	Barry and McNabb, 1999; Min <i>et al.</i> , 1999
<i>Lotus corniculatus</i>	–	Increased milk production in dairy cows	Woodward <i>et al.</i> , 1999
<i>Lotus corniculatus</i>	2.00– 4.00	Increases in wool growth (11%), live weight gain (8%), milk yield (21%) and ovulation rate (15–30%) without increasing voluntary feed intake, thus improving production efficiency	Terrill <i>et al.</i> , 1992; Wang <i>et al.</i> , 1996, 1996b; Min <i>et al.</i> , 1999
<i>Lotus. corniculatus</i>	3.00	Increased absorption of plant methionine and cysteine	Wang <i>et al.</i> , 1996b
<i>Lotus corniculatus</i>	4.00– 5.00	Increased production of milk protein lactose	Wang <i>et al.</i> , 1996
<i>Lotus corniculatus</i>	5.00– 6.00	Increased net absorption of methionine and increased supply of cystine for body synthetic reactions	McNabb <i>et al.</i> , 1993
<i>Lespedeza cuneata</i>	5.20	Greater immune response	Min <i>et al.</i> , 2003a
<i>Lespedeza cuneata</i>	4.60	57% reduction in FEC and a 61% reduction in total fecal egg output in	Min <i>et al.</i> , 2002

## NUTRITION & HUSBANDRY

Plant	Content (% DM)	Toxic/beneficial effect	Reference (s)
		goats	
<i>Lotus pedunculatus</i>	–	Absorption of P is greater (22%) throughout the GI tract due to CT	Waghorn <i>et al.</i> , 1994
<i>Lotus pedunculatus</i>	6.00–12.00	Depressed voluntary feed intake, inhibited rumen fibre digestion and reduced animal productivity	Terrill <i>et al.</i> , 1992; Wang <i>et al.</i> , 1996, 1996b; Min <i>et al.</i> , 1999
<i>Lotus pedunculatus</i>	6.30–0.60	Depressed feed intake, digestibility, and animal production in sheep	Barry and Duncan, 1984; Waghorn <i>et al.</i> , 1994
<i>Lespedeza cuneata</i>	> 5.00	Antinutritional effects	Windham <i>et al.</i> , 1990
<i>Lespedeza cuneata</i>	4.60	Reduced fecal egg production	Min <i>et al.</i> , 2004
<i>Lotus pedunculatus</i>	–	Reduced the rate of larval development (91%), eggs hatching (34%) and mobility of L <sub>3</sub> larvae (30%)	Molan <i>et al.</i> , 2000
<i>Lotus corniculatus</i>			
<i>Hedysarum coronarium</i>			
<i>Onobrychis viciifolia</i>			
<i>Lotus corniculatus</i>	5.00	To prevent rumen frothy bloat in cattle	Barry and McNabb, 1999
<i>Lotus pedunculatus</i>	2.20–3.50	Increase the absorption of amino acids from sheep small intestine	Waghorn <i>et al.</i> , 1987a
<i>Lotus pedunculatus</i>	6.00	Reduced voluntary intake and fibre digestibility	Barry and Duncan, 1984
<i>Lotus pedunculatus</i>	8.00–9.00	Reduced rates of body and wool growth	Barry, 1985
<i>Lotus pedunculatus</i>	9.00–10.00	Reduced rumen fibre digestion	Barry <i>et al.</i> , 1986
<i>Lotus</i>	5.00–	Reduced dry dry matter intake and	Waghorn <i>et al.</i> ,

Plant	Content (% DM)	Toxic/beneficial effect	Reference (s)
<i>pedunculatus</i>	6.00	reduced % digestion of amino acids in small intestine	1994
Lotus spp.	–	Reduced establishment of <i>O. circumcincta</i> worms	Niezen <i>et al.</i> , 1998
<i>Onobrychis viciaefolia</i>	–	Decrease in faecal egg output	Paolini <i>et al.</i> , 2003a
<i>Onobrychis viciifolia</i>	–	Inhibit egg hatching and larval development of <i>Trichostrongylus colubriformis</i>	Molan <i>et al.</i> , 1999; Waghorn and Molan, 2001; Molan <i>et al.</i> , 2002
<i>Lotus corniculatus</i>			
<i>Lotus pendunculatus</i>			
<i>Dorycnium rectum</i>			
<i>Dorycnium pentaphyllum</i>			
<i>Onobrychis viciifolia</i>	5.00–8.00	Fed to sheep and goats, had a higher nutritive value	Ulyatt <i>et al.</i> , 1976; Niezen <i>et al.</i> , 1995; Min <i>et al.</i> , 2003a
<i>Onobrychis viciifolia</i> Scop.	–	Lower levels of nematode egg excretion, reduced intestinal worms (50%), better host resilience	Paolini <i>et al.</i> , 2005
<i>Quercus lusitania</i> var. <i>infectoria</i>	–	Larvicidal agents	Redwane <i>et al.</i> , 2002
<i>Rubus fruticosus</i>	7.40	Inhibitory effect on L <sub>3</sub> and adult worms of <i>Teladorsagia circumcincta</i> , <i>Haemonchus contortus</i> and <i>Trichostrongylus clubriiformis</i>	Paolini <i>et al.</i> , 2004
<i>Quercus robur</i>	5.30		
<i>Corylus avellana</i>	14.20		
<i>Rumex obtusifolius</i>	–	Bloat could be eliminated	Waghorn and Jones, 1989
<i>Schinopsis</i>	–	Decreased the viability of L <sub>3</sub> of <i>Haemonchus contortus</i> ,	Athanasiadou <i>et</i>



## NUTRITION & HUSBANDRY

Plant	Content (% DM)	Toxic/beneficial effect	Reference (s)
spp. (Quebracho: CT)		<i>Teladorsagia circumcincta</i> and <i>Trichostrongylus vitrinus</i> and in sheep decreased faecal egg count and worm burden and improve the performance of parasitized sheep	<i>al.</i> , 2001
	–	Reduced FEC	Paolini <i>et al.</i> , 2003
<i>Spondias mombin</i>	–	Anthelmintic activity	Ademola <i>et al.</i> , 2005
<i>Crataegus</i> spp.	–	Used as an astringent for the digestive system in humans	Haslam, 1996