

EFFECTIVENESS OF DIFFERENT PROTOCOLS FOR INDUCING ESTRUS OF ARDY GOATS DURING THE NON-BREEDING SEASON.

Sobayil, K. A.

Dept. of Animal Production and Breeding, Fac. of Agric. and Veterinary Medicine, Univ. of Al-Qassim, P.O. Box 6622, Buriedah 51452, Kingdom of Saudi Arabia

ABSTRACT

The experiment was conducted to determine the reproductive performance (kidding rate and litter size) of estrus synchronized local Ardy goats after different protocols and trans-cervical artificial insemination (TC-AI). This experiment was conducted at the beginning of their non-breeding season. Total of 90 Ardy does were randomly allotted into 6 similar groups of 15 in each and treated as the following: group 1: does were synchronized with a controlled intra-vaginal devices release (CIDR) for 14 days + injecting (i.m.) 600 i.u. of equine Chorionic Gonadotrophins (eCG) at withdrawal time; group 2: does were received two doses (10 mg/dose) of PGF₂ α at 11 days apart; group 3: does were treated with insertion of progesterone sponges for 14 days + injecting 600 i.u. of eCG at withdrawal time; group 4: does were treated with GPG method (day 1: the does were given a dose of 25 μ g of GnRH (Cystorelin) ; each doe was injected with 10 mg of PGF₂ α on day 7 and at the 2nd GnRH dose on day 9; group 5: does were synchronized naturally by using male effect where a teaser ram was introduced to the does and left with them for 20 days; and group 6 (control): does did not receive any hormonal treatment and were left for 20 days with a tested fertile buck. All synchronized does were trans-cervically inseminated (TC-AI) with diluted semen of Ardy bucks. The pregnancy was diagnosed by ultrasonic examination on 40 days post insemination. Results showed that does in G1 and G3 showed clear estrus approximately 36 hours post CIDR and sponge removal and were 8 and 12 hours earlier than those in G4, and G2, respectively. Percentages of does kidded were 53.33, 13.33, 40.00, 20.00, 6.67 and 00.00% in G1, G2, G3, G4, G5 and G6, respectively, and the differences were significant ($P < 0.05$). Multiple birth percentages were 13.33, 00.00, 13.33, 6.66, 00.00 and 00.00% per does kidded in the respective groups. Injecting does with eCG in G1 and G3 at the time of pessary withdrawal was associated with fruitful improvements in the kidding rates. The kidding period in all hormonal-treated does was short (about 7 days) and this is an advantage for synchronizing the estrus in Ardy does.

Keywords: Goats, estrous synchronization, CIDR, eCG, GnRH.

INTRODUCTION

Goats have tremendous potential for improvement and manipulation of reproduction due to their nature as a seasonal breeder and multiple ovulators (Amoah *et al.*, 1996; Powell *et al.*, 1996 and Zarkawi *et al.*, 1999). Their short gestation length, compared to cattle, theoretically allows two kiddings per year. Estrus synchronization is a valuable management tool that has been successfully employed to achieve this goal and enhance reproductive efficiency (Kusina *et al.*, 2000).

Goats are seasonally polyestrous animals and their reproductive season is affected by the change of day length. They reproduce during fall, winter and early spring when the day length is short, so they are called as "short day-breeder animals" (Amoah *et al.*, 1996 and Attwood, 2007).

The seasonality of reproduction exhibited by goats and some other farm animals results in inadequacy of their products supply during some months. This situation has created a strong interest among producers to induce estrus during the anestrus season.

Research has dramatically increased the number of synchronization options (Rahman *et al.*, 2008). The producers have many choices to pick a synchronization protocol for their operation, their production goals and their available labor. One of the oldest ways to synchronize estrus is a luteolytic agent, prostaglandin (PGF₂α) or one of its analogues, which causes the regression of the corpus luteum "CL" (Akusu and Egbunike 1984; Greyling and van Niekerk, 1986 and Kusina *et al.*, 2000).

Prostaglandin (PGF₂α) is only effective if administered after day 5 of the estrous cycle and has been used as a co-treatment in effective progestogen-based synchronizing protocols in goats (Kusina *et al.*, 2000; Pierson *et al.*, 2001, 2003; Medan *et al.*, 2002 and Jackson *et al.*, 2003) sheep (Husein and Kridli, 2003), and cattle (Kojima *et al.*, 2000 and Lucy *et al.*, 2001).

Another way of creating estrus synchrony is using gonadotrophin releasing hormone (GnRH) or its analogue, which causes ovulation of the large follicle. Treatment with GnRH is combined with PGF₂α, as these hormones have different functions. Synchrony of estrus and fertility with a combination of GnRH and PGF₂α (GPG method) are good for cyclic females, and some researches indicate that this combination may induce cyclicity in some females that experiencing postpartum anestrus. However, protocols using only GnRH and PGF₂α are not advised for young females due to extremely variable response. A third method for synchronization of estrus is the use of progestins, which will maintain high levels of progesterone (P4) in the female's system even after the regression of the CL. Synchrony of estrus occurs 2 to 5 days following progestin removal (Wildeus, 1999).

In efforts to extend the lifespan of the CL for synchronizing estrus, various forms of progestogens and different methods of administration have been used in cycling does, as well as in seasonally anestrus does, to induce or synchronize estrus (Amoah and Gelaye, 1990; Wildeus, 1999 and Montielmelo *et al.*, 2002). Progestogen administration is common, especially in seasonally anestrus animals and has been used with or without accompanying treatments such as gonadotrophins or prostaglandin analogs. In the United States, no approved progestogen product is being manufactured for use in goats (Rathbone *et al.*, 1998). However, in other countries such as Australia and New Zealand, commercially available progestogen sponges formulated for goats, including those containing fluorogestone acetate (FGA; Cronogest 45) and methyl acetoxy progesterone (MAP; Repromap and Veramix), can be used. Similarly, a controlled internal drug-releasing device (CIDR) in the form of a silicone intra-vaginal progesterone insert is also available for use in goats outside the United

States (Rathbone *et al.*, 1998). The CIDR is an intra-vaginal pessary containing P4 designed for synchronizing estrus in ruminants (Knights *et al.*, 2001a &b). Previously, norgestomet, an ear implant progestin product, marketed under the name SynchroMate-B (SMB), is available for use in sheep in the United States, and thus held promise for approval for use in goats. The successful use of SMB in goats has been reviewed previously (Wildeus, 1999) and has been documented in other countries such as Mexico and Brazil, (Mellado *et al.*, 2000; Oliveira *et al.*, 2001 and Kusina *et al.*, 2000).

Exposure to male after a period of isolation can be used for induction of estrus during the breeding and non-breeding seasons without additional treatments in goats (Wildeus, 1999; Kusina *et al.*, 2000; Ishwar and Pandey, 1992; Mellado *et al.*, 2000 and Delgadillo *et al.*, 2002). There are other methods of synchronizing estrus that were examined by many researchers (du Preez *et al.*, 2001 and Wuliji *et al.*, 2003).

Therefore, the aims of the present study are to investigate; (1) the benefit of using various estrus synchronization protocols on inducing estrus of goats out of their breeding season, and (2) the accuracy of carrying out early pregnancy, diagnosis on day 40 pos-breeding in goats.

MATERIALS AND METHODS

Animals and experimental outline

This study was conducted at a local private goat farm in Al-Qassim region, Saudi Arabia, during the beginning of the seasonal anestrous period from May till mid-February 2008. Ninety clinically sound pluriparous Ardy goat does (2 – 3 years of age), kidding in February with an average live body weight of 47 kg were used in this experiment. Does were randomly assigned into six similar groups (n= 15 does/group) as the following: Group 1; does were synchronized with CIDR for 14 days + injecting 600 i.u. of equine Chorionic Gonadotrophins (eCG) at withdrawal time; group 2: does received two doses (10 mg/dose) of PGF₂α (Lutalyse: Pharmacia and Upjohn Co. Kalamazoo, MI) at 11 day- interval; group 3: progesterone sponges were intra-vaginally inserted (sponge/doe) for 14 days + injecting 600 i.u. eCG at withdrawal time; group 4: does were given GnRH-PGF₂α-GnRH protocol (GPG). In GPG protocol, each doe was i.m. injected with 25 µg GnRH on day 1 (Cystorelin: Merial Ltd; Duluth, GA), on d 7; each doe was i.m. injected with 10 mg PGF₂α, and on day 9, a second dose GnRH i.m of 25 µg was administered and trans-cervically inseminated, while group 5: does were synchronized naturally using male effect where a teaser buck was introduced to the does and left with them for 20 days. All does exhibiting estrus were exposed to Trans-cervical artificial insemination (TC-AI), while does in group 6 (control), does did not receive any hormonal treatment and were left with a tested fertile buck for 20 continuous days during the treatment period of the other groups. The experimental design is shown in Table (1).

Goats in all groups were managed under the same conditions at the farm. Animals of treatment groups were kept in separate pens. All does were identified by plastic ear tags, kept under semi-shaded pens and fed

approximately 0.7 kg alfalfa hay + 0.5 kg barley grains/ head/day. Water and salt licks were freely available all day.

Table (1): Experimental outline of goat does in different experimental groups.

Group	Synchronization method	Type of insemination
G1	CIDR + eCG	TC-AI
G2	Two doses of PGF _{2α} at 14 day-interval	TC-AI
G3	Sponge + eCG	TC-AI
G4	GPG protocol	TC-AI
G5	Male effect	TC-AI
6 (Control)	No hormonal treatment	Natural mating

TC-AI: Trans-cervical artificial insemination.

Insemination and reproductive performance:

Semen from four tested mature bucks was collected using an electro ejaculator. Semen was entirely tested visually and microscopically and only ejaculates of good sperm motility and morphology were used. The ejaculates from the four bucks were pooled and extended using IMV diluent (IMV-France). Semen was diluted to a final concentration of approximately 100 million motile sperm per ml. For each insemination, a volume of two ml containing approximately 200 million motile sperm was used. Diluted samples were subsequently examined microscopically to ensure that spermatozoa are of good quality. All does were artificially inseminated using trans-cervical AI technique (Wulster-Radcliffe *et al.*, 2004). Reproductive performance was determined by measuring the percentage of does kidding to AI at the synchronized estrus (kidding rate) and by determining the number of kids born per female kidded (litter size).

Pregnancy diagnosis:

Ultrasonic examinations were done using an Aloka-SSD500 (Japan) equipped with a 5 MHz linear array abdominal transducer on day 40 post-breeding. Does were restrained in a regular chute under a shaded area and the ultrasonic examinations were done in the early morning before providing the diet. An obstetrical lubricant was applied just to the right of the area in front of the udder devoid of hair, to insure satisfactory acoustic coupling. The accuracy of pregnancy diagnosis was indicated on the basis of kidding results.

Statistical Analysis:

Data were analyzed by GLM procedure using SAS (2000) program to test the effects of synchronization protocol on studied traits. The fixed effect model was as follow:

$$Y_{ij} = \mu + G_i + e_{ij}$$

Where;

Y_{ij} = An observation taken on individual,

μ = Overall mean,

G_i = a fixed effect of the group (synchronization regime),

e_{ij} = Random error assumed to be independent and normally distributed with mean = 0 and variance = σ^2 .

The significance among means was detected using Duncan's Multiple Range Test (Duncan 1955). The percentage values were subjected to arcsine transformation before performing the analysis of variance. Means were presented after being recalculated from the transformed values to percentages.

RESULTS AND DISCUSSION

Data of reproductive performance traits of the does as influenced by the estrous synchronization method are given in Table (2). Results show that does given CIDR+eCG (G1) significantly ($P<0.05$) exhibited the highest percentages of estrus incidence and pregnancy as compared to the other groups. Does in G3 (given sponges and eCG) ranked the second and did not differ significantly ($P>0.10$) from those of G1. However, the male effect (G5) expressed the lowest ($P<0.05$) percentages among all treated groups. It was anticipated that the control does (G6) did not exhibit any estrous activity during the treatment period (Table 2).

Does given CIDR + eCG (G1) and those given sponge + eCG (G3) showed clear estrous signs approximately 36 hours post CIDR and sponge removal and were 8 hours earlier than those given GPG (G4) and 12 hours earlier than the does given double doses of PGF2 α (G2). Several researchers have reported the onset of estrus to occur within 24-144 hours following progesterone withdrawal. (Billing and Katz, 1999 and Dogan and Nur, 2006). The goats that received either CIDRs or sponges exhibited very clear estrous signs in comparison with the does given other treatments. Among those does that received either CIDRs or sponges, the rate of estrus response was not significantly ($P>0.05$) different, however, it was significantly ($P<0.05$) higher than the does in the other treatment groups.

Additionally, does given CIDR or sponge give higher pregnancy rates (Table 2). Although the addition of eCG to the estrous synchronization program concomitantly with the CIDR or sponge relatively elevated the costs per a head, our data showed a clear advantage of its use.

The insertion and withdrawal of vaginal devices were easy and no intra-vaginal devices were lost during the trial. Some investigators found that using sponges gave better results than using CIDR and this was due to the high retention rate of sponges in comparison to CIDRs (Knight and Hall, 1988). In the present study, the retention rate of sponges and CIDR was 100%, and this is in agreement with the observation of Romano (1998).

None of the does showed estrus while the intra-vaginal devices were in place. Therefore, it can be accepted that the dose of P₄ in the CIDR and Sponge devices absorbed from the vagina during the treatment was sufficient to suppress the preovulatory discharge of pituitary gonadotrophins (Moeini *et al.*, 2007). The source of progestin did not influence the efficiency of estrus synchronization rate in CIDR vs. sponges (80% vs. 73%). Moreover, CIDRs did not cause bad smell in comparison with sponges. However, the source of progestin used for estrous synchronization and AI in small ruminants may still need further investigation.

Table (2): Reproductive performance traits of Ardy does subjected to various estrus synchronization regimes.

Group (Treatment)	Estrus exhibition (%)	Pregnancy rat (%)	Multiple birth per doe (%)	Pregnancy duration (day)
G1 (CIDR + eCG)	80.00±0.106 ^a	53.33±0.13 ^a	13.33±0.09 ^a	149.38±0.53
G2 (Two doses of PGF ₂ α)	33.33±0.125 ^c	13.33±0.091 ^{bc}	0.00±0.00 ^c	152.00±1.00
G3 (Sponges +eCG)	73.33±0.118 ^{ab}	40.00±0.13 ^{ab}	13.33±0.09 ^a	150.00±0.93
G4 (GPG Method)	46.67±0.133 ^{bc}	20.00±0.106 ^{bc}	6.66±0.06 ^b	148.67±0.88
G5 (Male effect)	20.00±0.106 ^{cd}	6.67±0.066 ^c	0.00±0.00 ^c	149.00±0.0
G6 (Control)	0.00±0.00 ^d	0.00±0.00 ^c	0.00±0.00 ^c	0.00±0.00

a, b, c and d: Means in the same column with different superscripts are significantly different at P<0.01.

Using two doses of PGF₂α at 11 day-interval did not give the expected results, since five does (33.33%) displayed estrus and only two (13.3%) of them became pregnant who gave single births. Also, GPG protocol did not give encouraging results where only seven does (46.67%) showed unclear estrus signs from which three (20%) became pregnant. There might be more beneficial effects for PGF₂α in GPG protocol over the use of PGF₂α alone (G2 and G4) during the reproductive season when there are developing follicles and active corpora lutea (CLs). The absence, or by other words, the very low number of CLs reduced the effect of PGF₂α in these two treatments.

Male exposure in G5 did not give good induction of estrus and this does not agree with the results reported by Velize *et al.* (2002) when they introduced active bucks to a group of prepared does. They found that more than 90% of the does displayed estrus behavior in the first 15 days following the introduction of those males. The variation of male effect might be due to the breed of treated goats (Nuti *et al.*, 2003). This also would be attributed to the effect of male pheromone glands. The failure of teasing outside this limited period may be due to the inability of the female to respond during the beginning of the seasonal anestrus period, resulting from refractoriness of the female to the male stimulus. Alternatively, it may be due to inadequate stimulation from the male. Indeed, male behavioral and physiological activity tends to decrease during the period of female seasonal anestrus in both sheep and goats (Chemineau, 1983). In the control group no doe came into estrus during the period of observation (zero estrus and pregnancy).

When ultrasound scanning was used for diagnosing pregnancy at approximately 40 days post breeding, all the positive diagnosed does gave births. The accuracy of diagnosing multiple pregnancies was 100%. Results presented in Table (2) show also that the multiple births percentages were 13.33, 00.00, 13.33, 6.66, 00.00 and 00.00% per doe kidded in the respective treatment groups. Injecting does with eCG in G1 and G3 at the time of pessary withdrawal was associated with fruitful improvements in the kidding rate. This finding is in agreement with those found in beef cattle given

different doses of eCG (Zeitoun, 1990). The kidding period in all hormonal-treated does was short (about 7 days) and this is an advantage for synchronizing the estrus in Ardy does.

Although the fertility results can be affected by many factors such as: type of intra-vaginal device, dose and timing of eCG injection, semen type and dose, time and number of AIs, breed and age of does, body condition, management, season and others. However the fertility results after hormonal treatment are contradictory.

In conclusion, results from this study indicate that goats treated with vaginal CIDRs or sponges accompanied with the injection of eCG showed higher percentage of estrus expression and fertility during the non-breeding season of Ardy goats. There still many studies are need to be conducted as to follow the ovarian follicular dynamics during out of the breeding season in goats.

ACKNOWLEDGEMENTS

The author is grateful to the Research Center of Al-Qassim University for providing the funds and supporting this research. Many thanks to the people of Rokany Goat Farm for giving me the chance to do this experiment at their farm. Thanks go to Prof. Magdy M. Abdulsalam for his assistance and advises in statistical analysis and appreciation to Prof. Moustafa M. Zeitoun for reviewing the manuscript.

REFERENCES

- Akusu, M. O. and G. N. Egbunike (1984). Short Communication. Fertility of West African Dwarf goat in its native environment following prostaglandin F₂ α induced estrus. *Vet. Q.* 6:173-176.
- Amoah, E. A. and S. Gelaye (1990). Superovulation, synchronization and breeding of does. *Small Rumin. Res.*, 3:63-72.
- Amoah, E. A.; S. Gelaye; P. Guthrie and C. E. Rexroad (1996). Breeding season and aspects of reproduction of female goats. *J. Anim. Sci.*, 74:723-728.
- Attwood B. M. (2007). An introduction to reproduction management of fiber and meat goats. 2nd Ed., Department of primary Industries, Melbourne, Australia.
- Billings, H. J. and L. S. Katz (1999). Facilitation of sexual behavior in French-Alpine goats treated with intravaginal progesterone releasing devices and estradiol during the breeding and nonbreeding seasons. *J. Anim. Sci.*, 77:2073-2078.
- Chemineau P. (1983). Effect on oestrus and ovulation of exposing creole goats to the male at three times of the year. *Journal of Reproduction and Fertility*, 67: 65-72.
- Delgadoillo, J. A.; J. A. Flores; F. G. Veliz; H. F. Hernandez; G. Duarte; J. Vielma; P. Poindron; P. Chemineau and B. Malpau (2002). Induction of sexual activity in lactating anovulatory female goats using male goats treated only with artificially long days. *J. Anim. Sci.*, 80:2780-2786.
- Dogan, I. and N, Nur (2006). Different estrus induction methods during the non-breeding season in Kivircik ewes. *Vet. Med.*, 51: 133-138.

- Duncan, D. B. (1955). Multiple Range and Multiple F-test. *Biometrics*, 11:10.
- Du Preez, E. R.; E. F. Donkin; P. A. Boyazoglu; G. H. Rautenbach; D. M. Barry and H. S. Schoeman (2001). Out-of-season breeding of milk goats—the effect of light treatment, melatonin and breed. *J. S. Afr. Vet. Assoc.*, 72:228–231.
- Greyling, J. P. C. and C. H. Van Niekerk (1986). Synchronization of oestrus in the Boer goat doe: Dose effect of prostaglandin in the double injection regime. *South Africa, J. Anim. Sci.*, 16:146–150.
- Husein, M.Q. and R. T. Kridli (2003). Effect of progesterone prior to GnRH-PGF₂ treatment on induction of oestrus and pregnancy in anoestrous Awassi ewes. *Reprod. Domest. Anim.* 38:228–232.
- Ishwar, A. K. and J. N. Pandey (1992). Oestrus synchronization and fertility in Black Bengal goats following administration of progesterone/prostaglandin and gonadotropins. *Res. Vet. Sci.*, 52:141–146.
- Jackson, D. J.; C. M. Fletcher and N. C. Whitley (2003). Attempt to induce estrus in goats during the non-breeding season (Summer). In: *Proc. Assoc. Res. Dir., Inc. 13th Biennial Res. Symp.*, Atlanta, GA. PP 150.
- Knights, M.; T. Hoehn; P. E. Lewis and E. K. Inskeep (2001a). Effectiveness of intravaginal progesterone inserts and FSH for inducing synchronized estrus and increasing lambing rate in aneestrous ewes. *J. Anim. Sci.*, 79:1120–1131.
- Knights, M.; T. D. Maze; P. J. Bridges; P. E. Lewis and E. K. Inskeep (2001b). Short-term treatment with a controlled internal drug releasing (CIDR) device and FSH to induce fertile estrus and increase prolificacy in aneestrous ewes. *Theriogenology* 55:1181–1191.
- Kojima, F. N.; B. E. Salfen; J. F. Bader; W. A. Ricke; M. C. Lucy; M. F. Smith and D. J. Patterson (2000). Development of an estrus synchronization protocol for beef cattle with short-term feeding of melengestrol acetate: 7–11 Synch. *J. Anim. Sci.*, 78:2186–2191.
- Kusina, N. T.; F. Tarwirei; H. Hamudikuwanda; G. Agumba and J. Mukwena (2000). A comparison of the effects of progesterone sponges and ear implants, PGF₂α, and their combination on efficiency of esrus synchronization and fertility of Mashona goat does. *Theriogenology*, 53:1567–1580.
- Lucy, M. C.; H. J. Billings; W. R. Butler; L. R. Ehnis; M. J. Fields; D. J. Kesler; J. E. Kinders; R. C. Mattos; R. E. Short; W. W. Thatcher; R. P. Wettemann; J. V. Yelich; and H. D. Hafs (2001). Efficacy of an intravaginal progesterone insert and an injection of PGF₂α for synchronizing estrus and shortening the interval to pregnancy in postpartum beef cows, peri-pubertal beef heifers, and dairy heifers. *J. Anim. Sci.*, 70:982–995.
- Medan, M.; A. H. Shalaby; S. Sharawy; G. Watanabe; and K. Taya (2002). Induction of estrus during the non-breeding season in Egyptian Baladi goats. *J. Vet. Med. Sci.*; 64:83–85.
- Mellado, M.; R. Olivas and F. Ruiz (2000). Effect of buck stimulus on mature and pre-pubertal norgestomet-treated goats. *Small Rumin. Res.*, 36:269–274.

- Moeini M. M.; Moghaddam A. A.; Bahirale and A. Hajarlan (2007). Effect of breed and progestin source on estrus synchronization and rates of fertility and fecundity in Iranian Sanjabi and Lori ewes. *Pakistan J. of Bio. Sci.*, 10: 3801-3807.
- Montlomelo, K. C.; J. P. C. Greyling and L. M. J. Schwalbach (2002). Synchronization of oestrus in goats: the use of different progestagen treatments. *Small Rumin. Res.*, 45:45-49.
- Nuti, L.; S. Woldesenbet and G. Newton (2003). Effect of breed and progesterone priming on pregnancy rates in anestrus meat goats in response to the buck effect. *J. Anim. Sci.* 81(Suppl. 1):127. (Abstr).
- Oliveira, M. A. L.; S. I. Guido and P. F. Lima (2001). Comparison of different protocols used to induce and synchronize estrus cycle of Saanen goats. *Small Rumin. Res.*, 40:149-153.
- Pierson, J. T.; H. Baldassarre; C. L. Keefer and B. R. Downey (2001). Seasonal variation in preovulatory events associated with synchronization of estrus in dwarf goats. *Theriogenology*, 56:759-769.
- Pierson, J. T.; H. Baldassarre; C. L. Keefer and B. R. Downey (2003). Influence of GnRH administration on timing of the LH surge and ovulation in dwarf goats. *Theriogenology*, 60:397-406.
- Powell, M. R.; M. Kaps; W. R. Lamberson and D. H. Keisler (1996). Use of melengestrol acetate-based treatments to induce and synchronize estrus in seasonally anestrus ewes. *J. Anim. Sci.*, 74:2292-2302.
- Rahman A. N. M. A.; R. B. Abdullah and W. E. Wan-Khadijah (2008). Estrus synchronization and superovulation in goats: A review. *Journal of Biological Science* 8: 1129-1137.
- Rathbone, M. J.; K. L. Macmillan; W. Jo`chle; M. P. Boland and E. K. Inskeep (1998). Controlled-release products for the control of the estrus cycle in cattle, sheep, goats, deer, pigs and horses. *Brit. Rev. Therap. Drug Carrier Syst.*, 15:285-380.
- Romano, J. E. (1998). Effect of two doses of cloprostenol in two schemes for estrous synchronization in Nubian does. *Small Rumin. Res.*, 28: 171-176.
- SAS (2000). *Statistical Analysis System User's Guide (Version 8)*. SAS Institute Inc., NC, USA.
- Veliz F. G.; S. Moreno; G. Duarte; J. Vielma Cheminea P.; P. Poindron; B. Malpoux and J. A. Delgadillo (2002). Male effect in seasonally an ovulatory lactating goats depends on the presence of sexually active bucks, but not estrus females. *Anim. Reprod. Sci.*, 15: 197-207.
- Wildeus, S. (1999). Current concepts in synchronization of estrus: Sheep and goats. *Proceeding of American Society of Animal Science*, pp 1-14.
- Wuliji, T.; A. Litherland; A. L. Goetsch; T. Sahlu; R. Puchala; L. J. Dawson; and T. Gipson (2003). Evaluation of melatonin and bromocryptin administration in Spanish goats. I. Effect on the out of season breeding performance in spring, kidding rate and fleece weight of does. *Small Rumin. Res.*, 49:31-40.
- Wulster-Radcliffe, M. C.; S. Wang and G. S. Lewis (2004). Transcervical artificial insemination in sheep: Effects of a new transcervical artificial insemination instrument and traversing the cervix on pregnancy and lambing rates. *Theriogenology*, 62:990-1002.

- Zarkawi, M.; M. R. Al-Merestani and M. F. Wardeh (1999). Technical Note. Induction of synchronized oestrous in indigenus Damascus goats outside the breeding season. Small Ruminant Res. 33:193-197.
- Zeitoun, M.M. (1990). Effect of PMSG and a monoclonal anti-PMSG on the super-ovulatory response and embryo quality in beef cows. Ph.D Thesis, pp 284. Alexandria University, Egypt.

كفاءة استخدام طرق مختلفة لإحداث الشياح في ماعز العارضي خارج موسم

التناسل

خالد بن عبد الله المسبيل

قسم إنتاج الحيوان وتربيته كلية الزراعة و الطب البيطري - جامعة القصيم المملكة العربية السعودية

أجريت هذه التجربة لدراسة الأداء التناسلي (نسب الحمل ومعدل الولادات وحجم البطن) للماعز العارضي التي تم تنظيم شياحها بواسطة المعاملات الهرمونية المختلفة والتلقيح الاصطناعي. تم استخدام ٩٠ عنزة عارضي خارج موسم التناسل وقسمت للحيوانات إلى ٦ مجاميع كل منها لحتوت على ١٥ حيوان وعوملت الحيوانات كما يلي، للمجموعة الأولى: تم تنظيم الشيق للماعز باستخدام CIDR لمدة ١٤ يوم + الحقن بـ ٦٠٠ وحدة دولية من هرمون eCG عند إزالة الـ CIDR مباشرة، للمجموعة الثانية: أعطيت للماعز جرعتين بالمعضل (١٠ ملجم / جرعة) من هرمون البروستاجلاندين PGF2 α بينهما ١١ يوم، للمجموعة الثالثة: تم استخدام إسفنجة البروجستيرون لمدة ١٤ يوم + الحقن العضلي بـ ٦٠٠ وحدة دولية من هرمون eCG عند إزالة الإسفنجة، للمجموعة الرابعة: تم فيها استخدام نظام الـ GPG (في اليوم الأول يعطى الحيوان ٢٥ ميكروجرام هرمون GnRH ثم اليوم السابع حقنة من البروستاجلاندين PGF2 α وفي اليوم التاسع يعطى الحيوان حقنة أخرى ٢٥ ميكروجرام هرمون GnRH ثم يتم التلقيح اصطناعيا- الحقن كله عضلي)، للمجموعة الخامسة: تم فيها تنظيم الشيق طبيعيا باستخدام تأثير الذكور حيث يتم وضع تيس كشاف للماعز لمدة ٢٠ يوم والإثاث التي ظهرت عليها علامات الشيق تم تلقيحها اصطناعيا، للمجموعة السادسة: استخدمت كمجموعة ضابطة حيث أن الحيوانات لم تعطى أي معاملة هرمونية وتركبت ٢٠ يوم مع ذكور مخصبة مختبرة. وكل الماعز المنظمة تم تلقيحها اصطناعيا، تم جمع السائل المنوي المستخدم في التلقيح من ٤ تيس عارضي مختبرة باستخدام طريقة التثبيبه الكهرببي، وتم تخفيف السائل المنوي بواسطة مخفف IMV وكان حجم التلقيحات المستخدمة ٢ مل تحتوي على ٢٠٠ مليون حيوان منوي حي متحرك، وتم تسجيل معدل الحمل والولادات وتم تشخيص الحمل بواسطة جهاز الموجات فوق صوتية بعد ٤٠ يوم تقريبا من التلقيح. وأوضحت النتائج أن المجموعة الأولى والثالثة أظهرت أعلى معدل استجابة لحدث الشيق (٨٠ و ٧٣) على الترتيب و المجموعة الأولى والثالثة أظهرت شيق واضح بعد ٣٦ ساعة من إزالة الـ CIDR و الإسفنجة وقيل ٨ و ١٢ ساعة في المعاملة الرابعة والثانية على التوالي ، وكانت الاختلافات بين المعاملات في معدل الحمل وعدد المواليد مرتفعة معنويا وكانت النسبة المئوية لمعدل المواليد ٥٣,٣٣ ، ١٣,٣٣ ، ٤٠ ، ٢٠ ، ٦,٦٧ ، صفر% في المعاملات الأولى والثانية والثالثة والرابعة والخامسة والسادسة على التوالي. بينما كانت النسبة المئوية لمعدل النواتم ١٣,٣٣، صفر، ١٣,٣٣ ، ٦,٦٦ ، صفر ، صفر% لكل عنزة على التوالي للمعاملات. الإثاث المحقونة بـ eCG في المعاملات الأولى والثالثة أظهرت تحسن جيد مصاحب لمعدل للولادات ، وكانت فترة الولادة في كل الإثاث المعاملة هرمونيا قصيرة (٧ أيام) وهذا يعتبر ميزة هامة في تنظيم الشيق في الماعز العارضي

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة
مركز البحوث الزراعيه

أ.د / عبد الخالق السيد عبد الخالق
أ.د / علاء الدين فؤاد محرز