

MICROBIOLOGICAL STUDIES ON LUNG INFECTION IN CALVES AT BEHERA ABATTOIR

****S. M. El-Hoshy and *H. A. A. Hamouda**

*Animal Health Research Institute, Dokki, Giza, Egypt.

**Alexandria branch and * Damanhour branch

ABSTRACT

The study was carried for bacteriological and mycological examination of pneumonic lungs of calves from 6-20 month old slaughtered at Behera Abattoir. The examined lungs were collected from a total number of 36 calves. The gross lesions of the affected lungs were recorded followed by collection of tissue specimens.

All collected samples were inoculated directly onto different media for isolation of bacteria and fungi. Results obtained showed that the bacteria isolated as single pathogen were staphylococcus aureus at rate of 22.2% followed by klebsiella pneumonia at rate 19.4%, Esherichia coli at rate 11.9% and streptococcus pneumonia at rate 11.1% Each of pasteurella heamolytica, and salmonella typhimurium isolated at rate 2.8%. Bacteria isolated as mixed pathogens were 6 cases harboured S. aureus and E.coli (16.6%) , 3 cases (8.3%) were due to concurrently infection of strept. Pneumonia and E.coli meanwhile one case (2.8%) showed pasteruella haemolytica and E. coli.

Antimicrobial sensitivity result. Showed that Enrofloxacin was the most effective antibiotic against bacteria followed by amoxycillin and marbofloxacin on the different types of the isolates staph.aureus , klebsiella pneumonia , E.coli and strept pneumonia.

INTRODUCTION

Calf pneumonia can potentially be a significant economic burden to a farm due to the costs of treatment, mortalities, reduced growth rates, additional labour and housing requirements. Pneumonia in young calves may be a chronic disease with very few clinical signs apart from a dry cough and slightly increased respiratory rate the acute form of the disease usually manifests itself in an outbreak involving several calves going down with the disease within a 24 hour period fe-

ver, dullness, in appetite and coughing often combined with nasal discharge are the most common symptoms.

Enzootic pneumonia of calves is not a specific disease attributed a single etiological agent and the variety of microbiological agents that may be associated with condition.

The main causes of pneumonia are bacteria, fungi and viruses whereas poor hygienic measures and climatic disorders are the most

predisposing factors to infection (**Rohman and Iyer 1979. Hafez et al, 1991 and Elyas, 1993**).

Among the most bacterial cause of pneumonia are *Escherichia coli*, *klebsiella pneumoniae*, *pseudomonas aeruginosa*, *streptococcus spp*, *staphylococcus spp*, and *pasteurella*, (**Elyas 1993 Fodor et al., 1999 and Odenoal and Henton 1995**).

The aim of the present study is to determine the different pathogenic bacteria species and in affected lung.

MATERIAL AND METHODS

36 lung samples from 6-20 month old calves showing respiratory disorders and nasal discharge slaughtered at Behlora Abattoir were collected.

The lungs had different forms of affections such as different stages of pneumonia.

The affected parts of the lung tissue were collected directly in sterile disposable containers for bacteriological examination.

Bacterial examination:

Samples were prepared according to **Thatcher and Clark (1978)** the samples were transferred directly into sterile homogenizer flask containing peptone water and homogenized for 2 minutes at 3000 rpm. A loopfull from each sample was directly inoculated into different media including nutrient agar, blood agar, MacConkey's agar, S.S agar and Baird Parker agar plates and incubated aerobically at 37°C for 24-48 hours. All isolates were mor-

phologically studied, then identified biochemically according to **Cruickshank et al. (1975) and Collee et al. (1996)**.

In vitro antimicrobial sensitivity test :

The technique was according to **Quinn et al. (1994)**.

RESULTS & DISCUSSION

Pneumonia is one of the major disease frequently encountered in farm animals causing major economic losses to calves (**Caldow et al. 1993**). The economic losses from death of affected animals (**Grove- White and White (1999)**) and cost of treatment and the predisposition of animals affected by neonatal pathogens to the development of other disease condition (**House 1978**). It was clear from table (1) 36 examined affected lungs were found to harbour various types of microorganisms.

The result in table (1) represented the bacteriological examination of 36 lung tissue samples of pneumonic calves.

The most prevalent member was belonging to *staph. aureus* and *klebsiella pneumoniae* which were more frequently observed as the sole causative agent in an incidence of 22.2% and 19.4% respectively and these results give a focus about the importance as the cause of pneumonia and this observation was in complete agreement with that mentioned by **Glantz et. al. (1972)** who recorded that the main organisms obtained from dead calves after suffering from respiratory syndromes were *C.pyogenes*, *P.aeruginosa*, *k. pneumoniae* and *staph aureus*, However **Samer (2001)** isolated *Staphylococcus aureus* at low percentage

than our results who isolated one isolate from 52 affected lungs.

Streptococcus pneumoniae were isolated from 4 samples at incidence of 11.1% as single pathogen this results agreed with **Hanaa (2007)** while **Samer (2001)** isolated *strept.pneumoniae* from pneumonic lungs at rate of 1.92% .

Escherichia coli is one of most important live agent of early infections in neonatal period and frequently causative agent of pneumonia. Also *Escherichia coli* is a part of normal intestinal flora of both human and worm blooded animal and under certain conditions, it can invade other organs as lung and cause pneumonia. (**Rauprich et al. 2000**).

Table (1) represented the bacteriological examination of 36 lung tissue samples of pneumonic calves it revealed that *E.coli* was isolated from 5 samples at incidence of 13.9 % as single pathogen. This result nearly similar to result **Abd El- Fattah et. al. (1993)**. However **Samah (2004)** could isolated *E.coli* in higher incidence reached to 72.64%.

The least frequent microorganisms secured from calves suffering from pneumonic lesions were *pasteurella heamolytica*, and *salmonella typhimurum* with an incidence of 2.8% and these results were in agreement with that mentioned by **Riad (1989)**.

A total of 10 calves revealed mixed infec-

tion with an incidence of (27.8%) Including *Staph. aureus* and *E.coli* (16.6%), (8.3%) were due to concurrently infection of *Strept. pneumoniae* and *E.coli*, meanwhile (2.8%) showed *Pasteurella heamolytica* and *E.coli*. This gave an idea about the role played by mixed infection and the observation is an agreement with that mentioned by **Vaisaire et al. (1988)** who mentioned that pneumonia was produced more obviously and pneumonic lesions were more extensive in mixed bacterial infection than when a single bacterium was isolated. Results of the vitro sensitivity test table (4) revealed that *Enrofloxacin* was the most effective antibiotic against bacteria followed by *amoxycillin* and *marboceal* on the different types of the isolates *Staph. aureus*, *klebsiella pneumonia*, *E. coli* and *Strept pneumonia* with the percentage of 75%, 66.7%, 60% and 50% respectively for *enrofloxacin*, 75%, 50%, 80% and 25% respectively for *amoxicillin* and 50%, 66.7%, 80% and 50% respectively for *Marbofloxacin*.

The high susceptibility of organisms to *enrofloxacin* and *Marbofloxacin* may be due to the fact that they are rarely used in therapy accordingly few strains get resistance against these antibiotics.

Therefore, periodic studies on resistance levels are needed to enable selection of appropriate antibiotics to which these microorganisms have the lowest resistance **Costa et. al. (1995)**.

Table (1) : Bacteria isolated from lung of Pneumonic calves as a single culture

Types of bacteria	Lung tissues	No. of positive samples	
Staphylococcus aureus	36	8	22.2
Klebsiella Pneumonia	36	7	19.4
Escherichia coli	36	5	13.9
Streptococcus Pneumoniae	36	4	11.1
Pasteurella haemolytica	36	1	2.8
Salmonella typhimurium.	36	1	2.8
Total	36	26	72.2

Table (2) : mixed culture of bacteria isolated from Lung of pneumonic calves.

Bacteria species	Lung tissues (36)	
Staph. aureus + E.coli	6	16.6
Strept. pneumonia + E.coli	3	8.3
Pasteurella haemolytica + E. coli	1	2.8
Total	10	27.8

Table (3) : Antimicrobial susceptibility of bacterial isolates from affected lung.

Antibiotic Disc	Sensitive Strain to different antibiotics							
	Staph. aureus		Klebsiella Pneumoniae		E-coli		Strept. pneumoniae	
	Total no. = 8		Total no. = 6		Total No. 5		Total No. = 4	
	No	%	No	%	No	%	No	%
Enrofloxacin	6	75	4	66.7	3	60	2	50
Amoxycillin	6	75	3	50	4	80	1	25
Gentamycin	5	62.5	3	50	2	40	2	50
Marbofloxacin	4	50	4	66.7	4	80	2	50
Ampicillin	1	12.5	1	16.7	-	-	-	-
Tetracycline	-	-	-	-	1	20	-	-

REFERENCES

Abd - Fattah, A. M.; Sayed, A. M.; Mona, A. M.; Thabet, A. E. and El-Mileegy, I. M. H. (1993) : Microbial respiratory disorders in buffalo calves and the behavior of some serum biochemical constituents. Assiut Vet. Med. J. 28 (56), 246-254.

Caldow, G. L.; Edwards, S.; Deters, A. R.; Nixon, P.; Iyata, G. and Sayers, R. (1993) : Association between viral infections and respiratory disease in artificially reared calves. Vet. Rec, 133 (24) : 85 - 89.

Costa, E. O.; Melville, P. A.; Ribetro, A. R.; Viani, F. C. and White, C. R. (1995) : Dynamics of resistance of aetiological agents of bovine mastitis to antibiotics. Revista Brasileira de Medicina Veterinaria 17 (6) 260- 263.

Collee, J. G.; Marmion, B. P.; Fraser, A. G. and Simmons, A. (1996) : Mackey and Mac Cartney Practical Medical Microbiology 4th ed. Churchill livingstone N. Y London.

Cruickshank, R.; Duguid, J. R.; Marmion, B. P. and Swain, R. H. A. (1975) : medical Microbiology : The practice of Medical Microbiology. 12th ed. Vol. 2, Churchill, livingstone, London.

Elyas, A. H. (1993) : Some studies on sheep pneumonia of bacterial and fungal origin Assuit Vet. Med. J. 29 (58) : 89-95

Fodor, L. Verga, J. Hajtos , I. and Molnar T. (1999) : Serotypes of pasteurilla heamolytica and pasteurilla trechalosi isolated from farm animals in Hungary.Vet. Med. Series (B), 61 : 241 - 247 .

Glantz, P. J.; Simpson, M.; Wilson, L. L. and Straley, E. J. (1972) : Escherichia coli 037 : B 18 in colibacillosis scientific publication: Oxford London, Edinburgh Boston, Melbourne.

Grove-white, D. H. and White, D. G.

(1999): Abdominal distention in collapsed diarrhaeic calves, biochemical finding and treatment. Vet . Rec . 144 (5) : 639 - 642.

Hafez, A. M.; Razig, R. A.; El-Amrousi, S. and Al-Hendi, A. B. (1991) : Respiratory disease occurring in farm animals in the eastern province of saudia Arabia . Assiut Vet. Med. J. 24:84-88 .

Hanaa, A. Eleem. (2007) : Bacteriological and pathological studies on enzootic. Pneumonia of lambs. Mvsc . Thesis presented to Fac. Vet. Med .Alex . University Egypt .

House, J. A. (1978) : Economic impact of rota virus and other neonatal disease agents of animals JAVMA 173 (5) : 573-576 .

Odendoal, M. W. and Henton M. W. (1995) : The distribution of pasteurilla heamolytica serotypes among cattle , sheep and goats in South Africa and their association with disease. Onderstepoort J.Vet. Res 62 (4) : 223 - 226.

Quinn, P.; Carter, M. E.; Markey, B. K. and Carter, G. R. (1994) : Clinical veterinary microbiology. Mosby , Year Book Europe limited.

Rahman, T. and Lyer, P. K. R. (1979) : Studies on pathology of ovine pneumonia, Indian vet. J., 56 : 455 - 461 .

Rauprich, P.; Moller, O.; Walter, G.; Herting and Robertson, B. (2000) : Influence of modified natural or synthetic surfactant preparations on growth of bacteria causing infections in the neonatal period. Clin. Diag. lab Immunol. 7 (5) 817- 822

Riad, E. R. (1989) : Bacteriological observation on the mortality problem in neonatal calves M.V.sc. Thesis (Microbiology) fac. Vet Med Cairo university.

Samah, S. A. (2004) : Pathological affections in the lungs of the slaughered animals

at El-Behera provine M.V.Sc. Thesis in pathology Fac. Vet. Met Alexandria University Egypt.

Samer, K. I. (2001) : Laboratory diagnosis of some bacterial agents causing Pneumonia and / or enteritis in ovines, caprines and bovines. Ph. D Thesis in Microbiolgy and Diagnostic Microbiology, Fac. Vet. Med. Alexandria University, Egypt.

Thatcher, F. S. and Clark, D. S. (1978) :

Microorganisms in food. International, Committee on Microbiological Specification for foods (ICMSF), 2nd Ed, Academic press, New York.

Vaissatre, J.; Martel. J.; Gelstin, P.; Chiral, C.; Brovillet, P.; Brugere, P. and fremanx, A. (1988) : Pneumococal septicaemia in calves Demonstration in france. Bull. Acad. Vet . France 61 (2) : 173 - 180.

الملخص العربى

دراسات ميكروبيولوجية فى الرئة المصابة فى العجول بمجازر البحيرة

ساميه الحوشى حسن عبدالسلام حموده*

معهد بحوث صحة الحيوان - الدقى - الجيزة - فرع الإسكندرية

معهد بحوث صحة الحيوان - الدقى - الجيزة - فرع دمنهور*

أجريت هذه الدراسة للفحص البكتريولوجى لرئات العجول المصابة بالالتهاب الرئوى والتى يتراوح عمرها من ٦ إلى ٢٠ شهر والتى تم ذبحها بمجازر محافظة البحيرة وقد سجلت الإصابات العينية للنسيج الرئوى المتأثر بالالتهاب عند مجمع العينات من ٣٦ عجل.

كشف الفحص البكتريولوجى بأن البكتريا التى عزلت منفردة كسبب مرضى كانت المكورات العنقودية الذهبية بنسبة ٢٢.٢٪ وتليت بالكبسيلات نيومونى بنسبة ١٩.٤٪ والأيشريشيا كولاي القصبات القولونية بنسبة ١٣.٩٪ والاسترىتوكولوس نيومونى المكورات السبحية الرئوية بنسبة ١١.١٪ أما كلاً من البستريلا هيموليتكم والسالمونيليا تيقوميرم فقد عزلا بنسبة ٢.٨٪.

أما البكتريا التى عزلت كأسباب مرضية مختلطة كانت الاستفيلوكوكوس أوربوس مع الأيشريشيا كولاي بنسبة ١٦.٦٪ أما الاسترىتوكولوس نيومونى مع الأيشريشيا كولاي فقد عزلت بنسبة ٨.٣٪ والباستريلا هيموليتكم مع الأيشريشيا كولاي بنسبة ٢.٨٪ أظهرت نتيجة اختيار الحساسية ضد الميكروبات ان الأنروفيلوكسللين هو الأكثر تأثيراً على البكتريا المعزولة يتبعه الأموكسللين والمريوفلوكساسين على البكتريا المختلفة المعزولة الاستفيلوكوكوس أوربوس والكبسيلات نيومونى والأيشريشيا كولاي والاسترىتوكوكوس نيومونى.