

Animal Health Research Institute,  
Assiut Regional Laboratory.

**STUDIES AFFECT CASES OF BACTERIAL  
RESPIRATORY INFECTION ON SOME BIOCHEMICAL  
CHANGES IN CAMELS AT ASSIUT GOVERNORATE**  
(With 5 Tables)

By

**A.M. MANAA and TH.S. ABD EL-ALL \***

\* Dept. of Animal Medicine, Faculty of Vet. Med., Assiut University  
(Received at 1/1/2011)

**دراسة تأثير الإصابات البكتريولوجية للجهاز التنفسي على بعض التغيرات  
البيوكيميائية في الإبل بمحافظة أسيوط**

**أحمد ممدوح مناع ، ثروت سعيد عبد العال**

أجريت هذه الدراسة على عدد ٩٠ جملاً مذبحين من الجنسين ومن مختلف الأعمار في سلخانة بنى عدى بمحافظة أسيوط ، أظهر الفحص الإكلينيكي قبل الذبح بأن ٢٥ من الجمال كانت تعاني من إصابات تنفسية متمثلة في صعوبة في التنفس مصحوبة بكحة وإفرازات مخاطية وبقاى الجمال سليمة من الناحية الإكلينيكية الظاهرية. اشتملت الدراسة على ١٠٠ عينة (٢٥ مسحة من الأنف ، ٢٥ عينة من كل من الرئة والغدد الليمفاوية الرئوية والسائل البلورى) وذلك لفحصها بكتريولوجياً من الحيوانات المريضة إكلينيكياً. أما الحيوانات السليمة ظاهرياً أخذت منها مسحات من الأنف (٦٥) قبل الذبح لفحصها بكتريولوجياً. أعطت ١٠٣ (٦٢,٤٢%) عينة نتائج إيجابية للعزل الميكروبي ومن بينهم ٢٠ (٣٠,٧٦%) من المسحات الأنفية للحيوانات السليمة ظاهرياً. وكانت الميكروبات المعزولة كالاتى: العنقودى الذهبى ٤٧ (٤٥,٦٣%) ، السبجى الصديدي ١٧ (١٦,٥٠%) والكليسيلا ٢ (١,٩٤%) واستافيلوكوكس إيبيديرميس ١٧ (١٦,٥٠%). كما تم تصنيف الميكروب القولونى سيروlogياً إلى ١٥ (١٤,٥٥%) E.Coli (O<sub>126</sub>/B<sub>6</sub>) ، ٥ (٤,٣٥%) لم يصنفاً. وبإجراء الفحوصات البيوكيميائية لعينات مصل الدم المأخوذة من الحيوانات السليمة ظاهرياً والمريضة إكلينيكياً قبل الذبح تبين وجود انخفاض معنوى جداً في نسبة الحديد مع ارتفاع معنوى جداً في تركيز النحاس في الجمال المريضة بالمقارنة مع السليمة. وقد شملت الدراسة أيضاً معرفة مدى تأثير الإصابات البكتيرية الرئوية على غازات الدم والأتزان الحمضى القاعدى وقد أظهرت النتائج وجود نقص معنوى جداً فى قيم الأس الهيدروجينى وانخفاض معنوي في قيم كل من B.E & Po<sub>2</sub> بينما لوحظ ارتفاع معنوى في قيم PCO<sub>2</sub> فى الجمال المريضة عند مقارنتها بالمجموعة السليمة. وتم عمل اختبار حساسية للميكروبات المعزولة ووجد أن الجاراميسين هو أنسب المضادات الحيوية معملياً ضد البكتريا المعزولة وهو الأكثر تأثيراً عليها.

## SUMMARY

This study was carried out on ninety slaughtered camels of different ages and sex in Bani-Adi abattoir at Assiut Governorate. The clinical examination before slaughter proved that 25 camels had respiratory affections (Dyspnoea, cough, hurried respiration and mucoid nasal discharge) and the remaining were apparently healthy clinically served as control group. 100 samples (25 nasal swabs and 25 from each of lung, pulmonary lymph nodes and pleural fluid) were tested for bacteriological examination from diseased camels. Also in clinically healthy camels 65 nasal swabs were taken before slaughter for bacteriological examination. 103 (62.42%) samples were culturally positive for different microorganisms, from which 20 (30.76%) nasal swabs from clinically healthy camel. Bacteriological examination revealed that the main isolates were Staph. aureus 47 (45.63%), Strept. pyogenes 17 (16.50%), Klebsiella species 2 (71.97%) and Staph. epidermidis 17 (16.50%). The isolated *E. Coli* strains were identified serologically into 15 (14.55%) *E. coli* strain (O<sub>126</sub>/B<sub>6</sub>) and 5 (4.35%) untypable. Blood samples from clinically healthy and diseased animals were taken before slaughter for biochemical analysis. It showed that highly significant ( $P < 0.01$ ) decrease in iron level, while there was a highly significant increase in copper levels in case of clinically diseased camels in comparison with healthy ones. Also, the study comprises the effect of bacterial respiratory affection of camels on blood gases and acid-base balance. There were a marked decrease in blood pH value and base excess (B.E), while significant decrease in Po<sub>2</sub> value as well as there is significant increase in PCO<sub>2</sub> value in clinically diseased camels in comparison with healthy ones. Antibiotic sensitivity tests for the isolated bacteria revealed that Garamycin is the best sensitive antibiotic of choice.

*Key works: Bacterial causes, camels, respiratory affection, antimicrobial susceptibility, blood gases and acid-base balance, Assiut.*

## INTRODUCTION

Camel is an animal of considerable importance in Egypt since it is one of the major sources of meat production while its meat represents 66.46% of the total meat obtained from the imported animals for slaughter purposes, Anon, (1986) and Gobrial, *et al.* (1991). Also it has a great value among our farm animals, working in agriculture fields or traveling between the villages and carrying out the farmers crops.

Respiratory diseases of camels continue to be a major cause of economic losses and adverse in animal. Stress of cold weather, rain, bad hygiene and high humidity rate were incriminated to increase the respiratory infection (Bacterial, viral and parasitic). Respiratory affection is the main cause of death among camel calves all over the world (Chowdhary, 1986 and Khanna *et al.*, 1992).

Bekele (2004) recorded that the major clinical signs observed in camels suffering from respiratory disease were cough, loss of appetite and watery nasal discharge which became mucoid.

Bacterial infection of the lung is one of the main causes of pneumonia in camels (Rana *et al.*, 1993; Thabet, 1993; Alhendi, 2000 and Seddek, 2002).

Several species of micro organisms could be isolated from both apparently healthy and affected respiratory tract camel such as Staphylococci, Streptococci, *E.coli*, Pasteurella and Klebsiella (Ghawi, 1978; Rana *et al.*, 1993; Fatma *et al.*, 2001 and Seddek, 2002).

Arora and Kalara (1973) mentioned that Streptococci, Staphylococci, *E.coli* and Klebsiella spp. Were the predominant isolates from respiratory tract infection in camels.

Ismail *et al.* (2008) recorded that the most prevalent bacteria among respiratory affection of camels was, *Staphylococcus aureus*.

Azam and Zaki (2006) reported that the principle causes in pneumonia in camels were *Staphylococcus aureus*, *E. coli*, *Streptococcus pyogenes* and *P. multocida*.

Tarazi (2001) recorded that Streptococci, *Staphylococcus aureus*, *E. coli* and Klebsiella spp. were the most frequent isolates from cases of pneumonia in one humped camels in Jordan.

Vashishta and Singh (1977) stated that Streptococcal pneumonia due to presence of streptococci in lung abscesses of the affected camels.

Vitovec and Vladik (1983) recorded that the Streptococci organism was isolated from bronchial diseases of camels in Somalia.

El-Magawry *et al.* (1986) found that the most bacterial agent causing respiratory affection in camels were *Staph. aureus*.

Respiratory affections affect either directly or indirectly on some serum biochemical parameters. Oxygenation of the blood and gaseous exchange between the blood and lung tissue is the main function of the lung i.e., regulation of the oxygen tension and carbon dioxide concentration.

In pneumonia and bronchitis, there were a marked increase in pressure  $CO_2$  ( $PCO_2$ ) values associated with drop in blood pH values (El-Sebaie *et al.*, 1988). The effect of acidosis are related chiefly to the

respiratory system. Increased carbon dioxide tension of the blood and depletions of bicarbonates caused an increase in the depth and the rate of respiration by stimulation of the respiratory centre (Radostits *et al.*, 2002).

## **MATERIALS and METHODS**

### **Animals:**

A total of 90 camels were used in this study which slaughtered at Assiut governorate abattoir in Bani-Adi. The age of slaughtered camels ranged from 2-5 years, 25 camels showing signs of respiratory disturbances including moist cough, rapid breathing and watery to mucoid nasal discharge, and the rest of camels were apparently healthy (65) used as control.

### **Bacteriological examination:**

65 Nasal swabs from both apparently healthy and diseased camels were collected before slaughtering. The collected swabs were inoculated into nutrient broth and incubated at 37°C for 24 hr. After slaughtering 75 samples from the lung, pulmonary lymph nodes and pleural fluid were taken from clinically diseased camels under a septic conditions and sent without delay to the laboratory, then inoculated on nutrient broth for 24h at 37°C. Loopful was taken from the inoculated broth and recultured on nutrient agar, 5% sheep blood agar, Manitol Salt agar and MacConkeys agar then incubated at 37°C for 24-48 hr.

The isolates were identified according to colonial morphology, pigment production, microscopically by Gram stain and biochemically according to Bailey and Scott (1974) using catalase activity, coagulase as well as novobiocin (30 mcg) and polymixin sulphate (300 µ) sensitivity test for identification of *Staphylococcus* sp.

### **Antimicrobial susceptibility tests of isolates:**

Antibiotic sensitivity tests were done for bacterial isolates using antibiotic disks (Biomerieux) of Ampicillin (10 µg), chloramphenicol (30 µg), Erythromycin (15 µg), Garamycin (30 µg), Oxytetracycline (30 µg), Kanamycin (30 µg), Neomycin (30 µg), Spectinomycin (20 µg), Tetracycline (30 µg), and Nalidixicc acid (30 µg) according the method of Koneman *et al.* (1997), while, zones of inhibition were determined according to National Committee For Clinical Laboratory Standards (NCCLS, 2002).

**Blood samples:**

10 blood samples were taken from both healthy and diseased camels for serum biochemical analysis of copper ( $\mu\text{g/dL}$ ) and Iron ( $\mu\text{g/dL}$ ) levels were estimated colorimetrically using Digital ultraviolet spectrophotometer Model 292 by means of test kits supplied by Boehringer Manuhem GmbH Diagnostica after the methods described by Trinder, (1956) and Zac (1958) respectively.

Another blood samples with anticoagulant 1: 1000 sodium heparin were collected from both clinically healthy and diseased camels for determination of blood pH,  $\text{PCO}_2$ ,  $\text{Po}_2$ ,  $\text{HCO}_3$ ,  $\text{TCO}_2$  and B.E by using coming pH – blood gas analyzer Model 168. The analyzer directly measured at  $37^\circ\text{C}$ .

Statistical analysis of obtained serum biochemical data was performed according to method of Snedecor and Cochran (1980) using test.

**RESULTS**

All results were illustrated in Tables (1-5).

**Table 1:** Incidence of positive bacterial isolates from respiratory tract of slaughtered camel samples.

| Types of samples      | Conditions         |           |              |            |           |           |            |            |              |
|-----------------------|--------------------|-----------|--------------|------------|-----------|-----------|------------|------------|--------------|
|                       | Apparently healthy |           |              | Diseased   |           |           | Total      |            |              |
|                       | No.                | Positive  |              | No.        | Positive  |           | No.        | Positive   |              |
|                       |                    | No.       | %            |            | No.       | %         |            | No.        | %            |
| Nasal swabs           | 65                 | 20        | 30.76        | 25         | 25        | 100       | 90         | 45         | 50           |
| Lung                  | -                  | -         | -            | 25         | 24        | 96        | 25         | 24         | 96           |
| Pulmonary lymph nodes | -                  | -         | -            | 25         | 20        | 80        | 25         | 20         | 80           |
| Pleural fluid         | -                  | -         | -            | 25         | 14        | 56        | 25         | 14         | 56           |
| <b>Total</b>          | <b>65</b>          | <b>20</b> | <b>30.76</b> | <b>100</b> | <b>83</b> | <b>83</b> | <b>165</b> | <b>103</b> | <b>62.42</b> |

**Table 2:** The isolated micro-organisms of both apparently normal and respiratory diseased camels.

| Isolated micro-organisms | Total number of isolates | % from total number of isolates | Nasal swabs        |       |  |       | Diseased camels  |              |  |       |  |       |
|--------------------------|--------------------------|---------------------------------|--------------------|-------|--|-------|--|--------------|--|-------|--|-------|
|                          |                          |                                 | Apparently healthy |       | Diseased camels                          |       | Lung   |              | Pulmonary lymph nodes                    |       | Pleural fluid                            |       |
|                          |                          |                                 | No.                | %     | No.                                      | %     | No.  | %            | No.                                      | %     | No.                                      | %     |
| Staph. aureus            | 47                       | 45.63                           | --                 | --    | 14                                       | 13.59 | 12   | 11.65        | 12                                       | 11.65 | 9  | 8.74  |
| Staph. epidermidis       | 17                       | 16.50                           | 16                 | 15.53 | 1  | 0.97  | --   | --           | --                                       | --    | --                                       | --    |
| Strept. Pyogenes         | 17                       | 16.50                           | --                 | --    | 6  | 5.82  | 4  | 3.88         | 5  | 4.85  | 2  | 1.94  |
| E.coli                   | 20                       | 19.42                           | 4<br>(untypable)   | 3.38  | 4<br>(O <sub>126</sub> /B <sub>6</sub> ) | 3.88  | 5 (O <sub>126</sub> /B <sub>6</sub> ) / 1<br>untypable | 4.85<br>0.97 | 3<br>(O <sub>126</sub> /B <sub>6</sub> ) | 2.91  | 3<br>(O <sub>126</sub> /B <sub>6</sub> ) | 2.91  |
| Klebsiella pneumonia     | 2                        | 1.94                            | --                 | --    | --                                       | --    | 2  | 1.94         | --                                       | --    | --                                       | --    |
|                          | 103                      | 99.99                           | 20                 | 19.42 | 25                                       | 24.27 | 24   | 23.30        | 20                                       | 19.41 | 14                                       | 13.59 |

**Table 3:** Inhibition Zone diameter (IZD) of test antibiotics against isolates of micro organisms

| Antibiotics             | Isolated micro-organisms |                    |                  |                |            |
|-------------------------|--------------------------|--------------------|------------------|----------------|------------|
|                         | Staph. aureus            | Staph. epidermidis | Strept. Pyogenes | Klebsiella sp. | E.coli sp. |
| Tetracycline (30 µg)    | R(<14 mm)                | I(<17 mm)          | R(<14 mm)        | R(<14 mm)      | R(<14 mm)  |
| Erythromycin (13 µg)    | S(>18 mm)                | S(>18 mm)          | S(>18 mm)        | R(<13 mm)      | R(<13 mm)  |
| Garamycin (30 µg)       | S(>15 mm)                | S(>15 mm)          | S(>15 mm)        | S(>15 mm)      | S(>15 mm)  |
| Neomycin (30 µg)        | R(<10 mm)                | R(<10 mm)          | R(<10 mm)        | R(<10 mm)      | R(<10 mm)  |
| Kanamycin (30 µg)       | R(<10 mm)                | R(<10 mm)          | R(<10 mm)        | I(<17 mm)      | I(<17 mm)  |
| Chloramphenicol (30 µg) | R(<12 mm)                | R(<12 mm)          | R(<12 mm)        | R(<12 mm)      | I(<17 mm)  |
| Oxytetracycline (30 µg) | I(<17 mm)                | S(>14 mm)          | I(<17 mm)        | R(<13 mm)      | R(<13 mm)  |
| Ampicillin (10 µg)      | I(<25 mm)                | I(<25 mm)          | I(<25 mm)        | S(>29 mm)      | I(<13 mm)  |
| Spectinomycin (20 µg)   | R(<11 mm)                | R(<11 mm)          | R(<11 mm)        | R(<11 mm)      | I(<14 mm)  |
| Nalidixic acid (30 µg)  | R(<13 mm)                | R(<13 mm)          | I(<17 mm)        | R(<13 mm)      | R(<13 mm)  |

R: Resistant  
I: Intermediate  
S: Sensitive

**Table 4:** Mean values ± standard deviation of blood serum iron and copper levels diseased camels.

| Animals              | No. | Fe. (µg/dL)  | Cu. (µg/dL)    |
|----------------------|-----|--------------|----------------|
| Apparently healthy   | 10  | 120.40±2.13  | 129.60±3.33    |
| Respiratory infected | 15  | 76.50±12.37* | 257.76±40.22** |

\* Significant (p<0.05)

\*\* highly significant (p<0.01)

**Table 5:** Overall values of blood gases and acid-base balance in clinically healthy and diseased camels.

| Groups of animals                 | N  | pH            | Pco <sub>2</sub> mm Hg | Po <sub>2</sub> mm Hg | HCO <sub>3</sub> mmol/L | TCO <sub>2</sub> mmol/L | B.E. mmol/L |
|-----------------------------------|----|---------------|------------------------|-----------------------|-------------------------|-------------------------|-------------|
| Apparently healthy                | 10 | 7.362±0.007   | 59.28±2.79             | 27.27±1.33            | 35.6±1.75               | 36.81±1.83              | 8.176±1.43  |
| Camels with respiratory affection | 15 | 7.231±0.067** | 62.43±3.80*            | 22.01±1.43*           | 32.5±1.56               | 34.6±1.62               | 2.58±1.31** |

N = Number of camels.

± S.D. = Standard deviation

\* = Significant (P < 0.05)

\*\* = Highly significant (P < 0.01)

## DISCUSSION

The most clinical signs of diseased camels with respiratory affection were moist cough, anorexia, depression, loss of appetite, hurried respiration and watery to mucoid nasal discharge. These findings agree with Abd-El-Kader, (1992) and Bekele, (2004).

In tables 2 the nasal bacterial swabs isolated from apparently healthy camels were *Staphylococcus epidermidis* 16 (15.53%) and *E.coli* 4 (3.88%). These results indicated that the respiratory tract of apparently healthy camels acted as a reservoir for many species of pathogenic and potential pathogenic microorganisms, Stress factors such as changes in the hygienic environmental and climatic conditions play a role in the onset of pneumonia (Phillip, 1972; Buxton and Fraser, 1977) Abd El-Kader (1992), was supported in the present study by the fact of the number of bacteria were isolated from apparently healthy camels and agreed with our data because the author was isolated *Staph. epidermidis* and *E.coli* from nasal swabs of clinically healthy camels. Phillip, (1972) recorded that the bacterial isolated from the nose of apparently healthy camels were *Staph. epidermidis* and *E.coli*. these data indicated that the animals were harboring these micro-organisms as members of nasal microflora.

In Table 2 the obtained results revealed that the total number of bacterial isolates from nose of diseased camel 25 (24.27%). The isolates were 14 (13.59%) *Staph. aureus*, 1 (0.97%) *Staph. epidermidis*, 6 (5.82%) *Strept. pyogenes* and 4 (3.88%) *E.coli* (O<sub>126</sub>/B<sub>6</sub>). A variety of pathogenic and potentially pathogenic bacterial isolates from examined samples with variable incidence and frequency percentages.

Also, Ismail *et al.* (2008) agreed with our data. the authors concluded that the major prevalent bacteria among respiratory affected cases were *Staph. aureus*. These data disagree with Bekele, (2004), who said that the major clinical signs of pneumonia in camels was due to *Pasteurella haemolytica*, Also, Azam and Zaki, (2006) recorded that the isolated strains from pneumonia in camels were *Staph. aureus*, *Strept. pyogenes*, *E.Coli*, *P.multocida* and *P.haemolytica*. The obtained results in table 2 revealed that most bacterial isolates from pulmonary lymph nodes and pleural fluid were 12 (11.65%) *Staph. aureus*; 5(4.85%) *Strept. pyogenes*; 3(2.91%) *E.coli* (O<sub>126</sub>/B<sub>6</sub>) and 9 (8.7%) *Staph. aureus*; 2 (1.94%) *Strept. pyogenes*; 3 (2.91%) *E.coli* (O<sub>126</sub>/B<sub>6</sub>) respectively. These data are nearly similar with Moustafa (2004) who isolated *Staph. aureus* 1 (5%); 2 (8.33%); *E.coli* 1 (5%); 1 (4.16%) and *Strept. pyogenes* 3 (15%); 1 (4.16%) from lung and pulmonary lymph nodes respectively from slaughtered pneumonic camels. From these previous data one could



observe that the bacterial isolates of the nose were more than that isolated from the lungs, pulmonary lymph nodes and pleural fluid because of the direct contact of nose with environmental conditions and also the local antibodies, produced by lymphocytes of the lung overcome such microbial agents.

Antibiotic sensitivity tests for the isolated bacteria revealed that the Garamycin and Ampicillin were the most effective antibiotic. For the isolated bacteria which representative in table 3 these findings were in agreement with Thabet, (1993) and Seddek, (2002). While Bekele, (2004) disagree with obtained data, he reported that Oxytetra-cycline was more effective in the treatment of respiratory diseased camels while *Staphylococcus* sp. and *Streptococcus pyogenes* were highly sensitives to Erythromycin.

Cheyne, *et al.* (1977) reported that an apparent improvement of the pneumonic camels following intramuscular injection of Oxytetracline. It had been suggested that resistance of bacterial isolates to some antibiotics may be attributed to wrong dosage, duration of treatment and route of administration (Amstutz *et al.*, 1982). On the other hand Azam and Zaki (2006) mentioned that camel affected with bacterial respiratory affection (*Staph.aureus*, *E.coli*, *Strept. pyogenes*, *Pastruella multocida*) were highly sensitive to Gentamycin followed by Nitro furantion, Amikacin, Enrofloxacin and Choramphenicol, respectively.

Finally respiratory disorders is still serious problem due to its special property that multifactors are responsible and the difficulty to determine the definite cause, so more efforts must be done to overcome that problem, such efforts as periodical, clinical and bacteriological examination of apparently healthy animals to avoid misuse of antibiotics.

In Table 4 the value of blood serum iron ( $120.40 \pm 2.13$  mg/dl) and copper ( $129.60 \pm 3.33$  mg/dl) in clinically healthy camels were in agree with these recorded by Abd-El-Kader, (1992). El-Magawary *et al.* (1986) they mentioned that the mean value of blood serum iron in normal dromedary camel was  $120.4 \pm 2.15$ . While El-Amrousi *et al.* (1984) mentioned that the main value of copper in blood serum of female healthy camel was  $110.9 \pm 7.82$ . On the other hand, there was highly significant ( $p < 0.01$ ) decrease in iron level and highly significant increase in copper level in diseased camel in comparison with healthy ones. These results were similar to that obtained by El-Magawry *et al.* (1986) and Abdel-Kader, (1992). The reduction of blood trace elements mainly to impaired absorption or increased excetion of respective element. Kaneko and Crnelius, (1970) recorded the loss of appetite and loss of blood and the main general causes

of these blood trace elements deficiency which accompany bacterial and parasitic diseases.

Ghargariu and Kadar (1979) mentioned that there was a significant decrease in iron serum level of calves affected with pneumonia, while an increase in serum copper level was recorded. On the other hand, Schulz *et al.* (1987) disagreed with our data, where the author recorded that there were no changes in iron and ketones in calves with pneumonia.

Abdel Kader, (1992) recorded that changes take place in levels of iron and copper associated with pneumonia in camels.

In pneumonia and bronchitis, there were a marked increase in  $PCO_2$  values associated with drop in blood pH values (Rosenberger, 1979 and Sodeman, 1961). The effect of acidosis are related chiefly to the respiratory system. The increased carbon dioxide tension of the blood and depletions of bicarbonate causes an increase in depth and the rate of respiration by stimulation of the respiratory center (Radostits *et al.*, 2002). Respiratory acidosis may occur in animals during pneumonia (Dukes, 1964)

The values of blood gases and acid-base balance in clinically healthy and diseased camels in Table 5.

The values of blood gases acid-base balance (pH,  $PCO_2$ ,  $PO_2$ ,  $HCO_3$ ,  $TCO_2$  and base excess) in clinically healthy camels were in agreement with values recorded by Habashy, (1985). In clinically diseased camels where there were highly significant decrease ( $p < 0.01$ ) in value of pH and base excess (B.E) (7.231 and 2.58 mmol/l) while there was a significant ( $p < 0.05$ ) decrease in value of  $PO_2$  (22.01 mmHg) and significant ( $p < 0.05$ ) increase of  $PCO_2$  (62.43 mmHg).

These obtained results nearly similar with previously results recorded by El-Sebaie *et al.* (1988) mentioned that pneumonia of calves associated with marked drop in blood pH and B.E while there were highly increase in values of  $PCO_2$  and significant decrease in  $PO_2$ .

The drop of pH values and base excess in diseased camels due to respiratory acidosis and this finding were in agreement with data after Coles, (1980) and Haskins, (1983).

Also, a significant increase in power  $CO_2$  and a significant decrease in  $O_2$  tension could be attributed to defect in oxygenation process of the lung during the course of pneumonia which leads to retention of  $CO_2$  in blood (Coles, 1980). Respiratory acidosis (Coles, 1986) occur when  $CO_2$  elimination is decreased, and blood carbonic acid concentration and  $PCO_2$  are increased. Other causes of respiratory acidosis include intrathoracic and pneumonia.

### **Acknowledgment**

I would like to express my deep thanks to the department of biochemistry and microbiology faculty of medicine Assiut EL-Azhar University.

### **REFERENCES**

- Abd El-Kader, H.A. (1992):* Studies on bacterial and parasitic causes of respiratory infection among dairy animals. Ph.D. Thesis, Fac. Vet. Med., Assiut Univ.
- Alhendi, A.A.A. (2000):* Common diseases of camels (*Camelus dromedarius*) in Eastern province of Saudi Arabia. *Pakistan Vet. J.* 20(2): 97-99.
- Amstutz, H.; Morte, R. and Armstrong, C. (1982):* Antimicrobial resistance of strains of *Pasteurella* species isolated from feed lot cattle. *Bovine Practice.* 16: 52-55.
- Anon (1986):* Vet. Service Organization publication, Mist. Agri. A.R.E.
- Arora, R.G. and Kalara, D.S. (1973):* A note on isolation of *Klebsiella pneumoniae* and *Diplococci* from cases of bronchopneumonia in camels. *Indian Journal of Animal Science* 43, 12: 1093-1095.
- Azam, I.M.I.A. and Zaki, M.H. (2006):* The first research laboratory investigation on bacterial pneumonia in Camels. *Egypt J. Comp. Path., Clinic. Path.* Vol. 19 N. 1: 319-344.
- Baily, W.R. and Scott, E.G. (1974):* *Diagnostic Microbiology. A text book for isolation and identification of pathogenic microorganisms* 4<sup>th</sup> Ed., the C.V. Mosby Company Saint Louis.
- Buxton, A. and Fraser, G. (1977):* *Animal Microbiology.* Black Well Scientific Publication, Oxford, London.
- Bekele, T. (2004):* Studies on the respiratory diseases 'Sonbabe' camels in the Eastern Lowlands of Ethiopia. *Tropical Animal Health and Production*, Vol. 31, No. 6.
- Cheyne, L.A.; Pegram, R.G. and Cartwright (1977):* *Tropical animal health and production* 9. 238.
- Chowdhary, B. (1986):* Some important biological and production characters of the Bikaner: Camel. *Ind. J. of Animal Production and Management.* 2(3): 145-151.
- Coles, L.L.E. (1980):* *Veterinary Clinical Pathology.* 3<sup>rd</sup> Ed. W.B. Saunders Company. Philadelphia London Toronto.
- Coles, E.H. (1986):* *A Text Book of Veterinary Clinical Pathology* 4<sup>th</sup> Ed., W.B. Saunders Co., Philadelphia, London.

- Dukes, H.H. (1964):* The Physiology of Domestic Animals. Pp: 476-483. 5<sup>th</sup> Ed., Baillare, Tindall and Lox. London.
- El-Amrousi, S.; Hafez, A.M. and Wasfi, I.A. (1984):* Some biochemical parameters of mature female camels in the Estern Province of Saudi Arabia. 1<sup>st</sup> Scientific Congress, Fac. Vet. Med., Assiut Univ.
- El-Magawry, S.; Okela, M.; Ezzat, M. and El-Attar, H.M. (1986):* A etiological study on respiratory affection in camels and its relation to haematological and biochemical changes. Assiut Vet. Med., Vol. 17, No. 33: 96-103.
- El-Sebaie, A.; Ibrahim, H. and Manaa A. (1988):* Blood gases and acid-base balance in association with bronchopneumonia in calves. Assiut Vet. Med. J. Vol. 19, No. 38 (1988).
- Fatma, M. Darwish; Hammad, A.M. and Hala, S. Ibrahim (2001):* Pathological studies on pneumonia in camels with special reference to mycotic and bacterial infection. J. Egypt. Vet. Med. Ass., 91 (2): 143-172.
- Ghawi, A.M. (1978):* Public health importance of camel lung affection. M.V.Sc. Thesis Fac. Med., Cairo Univ.
- Ghergariu, S. and Kadar, L. (1979):* Behaviour of Cu, Fe and Zn blood levels in diseased cattle. I-Changes in neonatal diarrhea. II-Changes during respiratory diseases in young cattle. Zentl blatt fur Veterinar Medizini 26 A(B0 666-676.
- Gobrial, N.; Ahmed, L.S.; Ali, S.M.; Elyas, A.H.; Nashed, S.M. and Amer, A.A. (1991):* Myco and microflora of the nasal cavity of apparently health camels. Assiut Vet. Med. J. 24 (48): 125-130.
- Habashy, E.S. (1985):* Blood Gases and acid-base balance in camels at Assiut under Various physiological conditions M.V.Sc. Fac. Vet. Med., Assiut University.
- Haskins, S.C. (1983):* Current Vet. Therapy. 6<sup>th</sup> Ed. W.B. Saunders company Philadelphia, London, Toronto and Mexicocity.
- Ismail, A.R. Adwan; Mamdoh Abdelghani; Sabry, A. Hassan and Samy, D. Abd. Alla (2008):* Studies on bacterial causes of preumo-enteritis in camels. (Camels Dromedaries). 13<sup>th</sup> Sci. Cong. 2008, Fac. Vet. Med., Assiut Univ., Egypt.
- Kaneko, J.J. and Cornelius, C.E. (1970):* Clinical biochemistry of domestic animals. 2<sup>nd</sup> Ed. Acad. Press New York and London.
- Khanna, N.D.; Landon, S.N.; Sahani, M.S.; Allen, W.R.; Higgins, A.J.; Mayhew, I.G.; Snow, D.H. and Wade, J.F (1992):* Calf mortality on Indian camel Confernce, Dubai, 2<sup>nd</sup> – 6<sup>th</sup> February 1992, 89-92.

- Koneman, E.W.; Allen, S.D.; Danda, W.M.; Soherchenberger, P.C. and Winn, W.C. (1997):* Colour Atls and Textbook of Diagnostic Microbiology. 4<sup>th</sup> Ed. J.B. Lippincott Co., U.S.A.
- Moustafa, A.H. (2004):* Study of some aerobic bacterial causes of respiratory affection in slaughtered camels in Dakahlia Governorate Assiut. Vet. Med. J. Vol. 50 No. 102 July 2004.
- "NCCLS" National Committee for clinical Laboratory Standards (2002):* Performance standards for Antimicrobial Disk and Dilution susceptibility tests for bacteria isolated from Animals; Approved standard 2<sup>nd</sup> Ed. NCCLS document M31-A2. NCCLS, Wayne, PA.
- Phillip, J.I.H. (1972):* Bovine respiratory disease: Is control possible? Vet. Record 90 No. 20, 552-555.
- Radostits, O.M.; Blood, D.C. and Gay, C.C. (2002):* Veterinary Medicine. 10<sup>th</sup> Ed. Bailliere Tindal.
- Rana, M.Z.; Ahmed, A.; Sindhu, S.T.A.K. and Mohamed, G. (1993):* Bacteriology of Camel Lungs. Camel-News Letter, No. 10, 30-32.
- Rosenberger, G. (1979):*clinical examination of cattle pp:138-141. Roughton, 2<sup>nd</sup> Ed., Berlin and Hamburg.
- Schulz, O.; Walzel, R.; Umlauf, K.; Blohm, H.; Liebig, F. and Claus, D. (1987):* Behaviour of some biological values in calves with pneumonia. Montshefte fur Veterinary Medizin, 42 (7) 248-251.
- Snedecor, G.W. and Cochran, W.G. (1980):* Statistical methods. 7<sup>th</sup> Ed. Iowa State University Press, Ames, U.S.A.
- Seddek, R.S. (2002):* Bacterial causes of lung affections in slaughtered camels in Assiut Governorate. Assiut Vet. Med. J. 46 (92): 169-177.
- Sodeman, T.A. (1961):* Pathologic physiology pp:114-121. 3<sup>rd</sup> Ed., W.B. Saunders company, Philadelphia and London
- Thabet, A.El-R. (1993):* Some microbial studies on lung of clinically healthy and respiratory infected camels. Assiut Vet. Med. J. 30 (59): 188-195.
- Tarazi- Al, Y.H. (2001):* Bacteriological and Pathological Study on Pneumonia in the One-Humped camel (camels dromedaries) in Jordan. Revue Elev. Med. Vet. Pays trop., 2001, 54(2): 97.
- Trinder, P. (1956):* Colorimetric method for the determination of iron. J. Clin. Path. 9: 170-173.
- Vashishta, M.S. and Singh, R.P. (1977):* Camel diseases in India, Scientific book agency.

- Vitovec, I. and Vladik, P. (1983):* Bronchial diseases of camels in Somalia. Bulletin of Animal Health and Production in Africa 31 (3): 291-292.
- Wahbi, A.G.A.; Salah, El-Din, E.A.; Awad El-Seid and Idris, O.F. (1979):* The plasma electrolytes and minerals of normal camels in Sudan. Vet. Med. J., XI No. 19: 215-238.
- Zac, B. (1958):* Colorimetric method for the determination of copper. Clin. Chim. Acta, 3: 328-334.