

Veterinary Serum and Vaccine Research Institute, Abbasia, Cairo, Egypt
Central Veterinary Diagnostic Lab, Ministry of Agriculture, Riyadh. KSA.

EPIDEMIOLOGICAL STUDIES OF *T. EVANSI* IN CAMELS IN SAUDI ARABIA USING REAL TIME POLYMERASE CHAIN REACTION TECHNIQUE (With One Table and Two Figures)

By

R.M. AL-KHATIB; M.R. YOUSEF and H.M. AL NAKHLI*

* National Agriculture and Animal Resources Research Center,
Ministry of Agriculture, Riyadh. KSA.

(Received at 13/6/2011)

**دراسات وبائية على التريبانوسوما ايفانساي في الإبل بالمملكة العربية
السعودية باستخدام تفاعل البلمرة المتسلسل ذو الزمن الحقيقي**

رجب محمد الخطيب ، محمد ربيع يوسف ، حبيب مقبول المناخلي

مرض التريبانوسوما (سارا) في الجمال ما زال يسبب مشاكل على مستوى العالم. ويصاحبه خسائر اقتصادية في المناطق التي تنتشر فيها تربية الجمال. ونظرا لقلّة البيانات المنشورة عالميا باستخدام تفاعل البلمرة المتسلسل ذو الزمن الحقيقي كاختبار سريع للكشف عن طفيل التريبانوسوما ايفانساي في الجمال. فقد استهدفت هذا الدراسة استخدام اختبار البلمرة المتسلسل ذو الزمن الحقيقي للكشف عن الحامض النووي الخاص بطفيل التريبانوسوما ايفانساي وذلك باستخدام بادئ تتابع وراثي خاص بهذا الطفيل باستخدام صبغة السبير الأخضر الفلوريسنت. حيث تم جمع عدد ٢٤٢ عينة دم من جمال يحتمل إصابتها بالتريبانوسوما ايفانساي من محافظات مختلفة من السعودية (الرياض - القصيم - الخرج - الدوادمي) وبإجراء الاختبار كانت نقطة التعاظم للعينات الايجابية للإصابة بطفيل التريبانوسوما ايفانساي تظهر اتحادهما عند درجة حرارة ٨٥ درجة مئوية والذي ينطبق مع عينات ايجابية ضابطة لطفيل التريبانوسوما ايفانساي. وقد وجد أن عدد ١٠٢ عينة ايجابية من عدد ٢٤٢ عينة وذلك بنسبة ٤٢,١٥% ، كما أشارت الدراسة إلى مدى دقة هذا الاختبار للكشف عن التريبانوسوما ايفانساي في الجمال وتعد هذه أول دراسة وبائية للتشخيص الجزيئي لطفيل التريبانوسوما ايفانساي في الجمال في السعودية والتي تعكس مدى انتشار هذا الطفيل بين الجمال في المناطق المختلفة بالمملكة مما يستوجب مزيدا من الدراسات لمحاولة استكشاف مدى احتمال وجود أنواع أخرى من هذا الطفيل وكذلك محاولة السيطرة على الإصابة بهذه الطفيليات بالمملكة.

SUMMARY

Trypanosomiasis (surra) still a serious problem in camel husbandry world widely. It causes considerable economic losses in many camel-rearing regions. A little date for Real Time polymerase chain reaction (PCR) for rapid detection *Trypanosoma evansi* of blood from naturally infected camels were published. In the present study 242 blood samples from clinically suspected camels to surra from four province of Saudia (Al-Riyadh, Al- Qassim, Alharig, and Al-Dawadmi) were collected and used for further examination. Real Time PCR-based assay for blood samples were conducted in direct detection of *T. evansi* DNA from the collected blood samples using specific primer set (TR3/TR4) derived from nuclear repetitive gene of *T.evansi* and SYBER Green1 fluorescent dye have been used. The melting peak chart of the positive samples showed one single peak at an average T_m of 85.0°C. The performed real time PCR technique was found accurate method for diagnosis of trypanosoma infection of camels where; 102 (42.15%) out of 242 were positive. It is concluded that camel trypanosomiasis in Sudia apparently caused by the present single parasite species *T. evansi* and detection of other typanosomes species are recommeded. In addition, the disease is highly prevalent in the country, which strengthens the need to change control policies and institute measures that help prevent the spread of the parasite. To our knowledge, this is the first molecular diagnosis report, which gives a picture of camel *trypanosomiasis* covering large geographical areas in Saudi Arabia.

Key words: Camels, Real time PCR, SYBER Green1, *Trypanosoma evansi*.

INTRODUCTION

Trypanosomiasis ('surra') caused by *Trypanosoma evansi* is considered to be one of most important diseases of camels in KSA. *Trypanosoma evansi* is the most widely geographically distributed pathogenic trypanosome. It can infect several species of animals, including, camels, horses, cattle and buffaloes causing trypanosomiasis commonly known as surra (Luckins, 1988). *Trypanosoma evansi* (*T. evansi*), the cause of trypanosomiasis (Surra), constitutes one of the major veterinary problems worldwide. The disease causes significant morbidity and mortality in camels (Pathak, and Khanna, 1995; Omer *et al.*, 2004). Surra, manifests itself usually as a chronic infection

characterized by weight loss, anemia, infertility and abortion (Lohr *et al.*, 1986; Luckins, 1988). *Trypanosoma evansi* is a blood-borne parasite that is transmitted mechanically by the bites of haematophagous flies. Initially, trypanosomes may be seen readily in the bloodstream, but in chronic infections parasites are difficult or impossible to find (Luckins, 1988) and in chronic infections is likely to be associated with secondary infections due to immunosuppression (Njiru *et al.*, 2004). Clinical signs and pathological lesions caused by *T. evansi* in camels are unreliable for definitive diagnosis (Chaudhary and Iqbal, 2000) in addition, detection of parasites in the blood is difficult because parasitaemia is intermittent (Mahmound and Gray, 1980). Serological tests have been developed and evaluated for diagnosis of trypanosomiasis in camels. They include card agglutination test and enzyme-linked immunosorbent assay (Ab-ELISA) (Nantulya, 1994; Davison *et al.*, 1999). In general serological techniques are useful for detection of a past infection but not for detection of an active infection with *T. evansi* (Viseshakul and Panyim 1990; Hopkins *et al.*, 1998). To address these problems; Real time polymerase chain reaction technique have been developed and evaluated for detection of *T. evansi* directly from blood (Wuyts *et al.*, 1995; Omawa *et al.*, 1999; Imadeldin *et al.*, 2006; Yousef *et al.*, 2010). In Saudia, camels a principal mean of meat and milk production, they are used for transportation of crops and do other farm works; also, their hair, wool and hides are used. Trypanosomosis in camels causes considerable economic losses due to a decrease in milk and meat, premature births and abortions (Boid *et al.*, 1985). In-sight the aim of this study was planed to explore the rate of *T. evansi* among camel population in giving four different provinces in KSA by the most resent molecular technique; real time PCR as a preliminary step for designing rational trypanosomiasis control program.

MATERIALS and METHODS

Animals

This study was Realized from May 2010 to April 2011; a total number of 242 suspected clinical camels (*Camelus dromedarius*), 3-8 years old, were examined clinically: body temperature, mucous membrane, muscles of thigh and hump. These animals were located in four province in Saudia (Al- Riyadh, Al- Qassim, Alharig and Al-Dawadmi). The main complaints of the camel owners were loss of appetite, decrease of productivity, and reduce of the body weight.




Fig. 1: Map of Al Riyadh showing study Provinces

Blood samples

Blood samples were collected in clean sterile vacutainers, containing ethylene diamine tetra acetic acid (EDTA), from naturally suspected infected camels.

Extraction of DNA from blood samples:

Automated extraction of total genomic DNA from the whole blood by BioRobot EZ1 workstation (Qiagen®) using EZ1 DNA blood kit (Qiagen®) according to the manufacturer's instructions.

Primers:

T.evansi repetitive DNA primers was designed and synthesized in BiologioBV®, Belgium to amplify a single band of 257 bp PCR product according to Chansiri *et al.* (2002); Yousef *et al.* (2010) for RT-PCR:

TR3 (5' GCGCGGATTCTTTGCAGACGA 3')

TR4 (5' TGCAGACACTGGAATGTTACT 3')

Real times PCR assay:

Polymerase chain reaction was carried out in LightCycler 2.0 (Roach®). The PCR mixture (20 µl) contained 15 µl of reaction mixture containing Fast start DNA Master^{plus} SYBER Green 1(Roach®) and 10 pmol/µl concentration of each primer and 5 µl of genomic DNA. The PCR profile was performed as following; pre-heated for 1 cycle at 90 °C for 10 min and then denatured at 94 °C for 5 sec, annealing at 55 °C for 10 sec and extension at 72 °C for 10 sec. The PCR amplification was performed for 45 cycles. Fluorescence data were acquired at the end of each cycle in a single step. Once the plateau phase of the PCR had been reached, amplification was stopped and a standard melting curve analysis was performed (95°C for 0 second, 65°C for 10 seconds, and a 0.1°C/second rise to 95°C) with continual fluorescence measuring. PCR data were analyzed using LightCycler2 software version 4.05.

RESULTS

Table 1: The percentages of positives from four provinces in Saudi.

Province	No. of examined animals	No. of positive	% of positive to total examined
Al- Riyadh	80	25	10.33
Al- Qassim	70	41	16.94
Alharig	60	31	12.81
Al-Dawadmi	32	5	2.07
Total examined	242	102	42.15

The melting peak chart of the positive samples showed one single peak at an average T_m of 85.0°C. (Fig.2).

The results of examined camels using real time PCR for detection of DNA of trypanosomes in suspected camels, showing a 102(42.15%). positive camels out of 242. The percentages of positives in different provinces as shown in Table (1) in Al- Riyadh, Al- Qassim, Alharig, and Al-Dawadmi are 10.33%, 16.94%, 12.81% and 2.07% respectively.




Fig. 2: Melting curve analysis of amplicon obtained from real time PCR.

DISCUSSION

The economic importance of *T. evansi* infection is mainly attributed to clinical disease in camels in many parts of the world (Dia *et al.*, 1997; Atarhouch *et al.*, 2003). The fact that PCR assays provided evidence of infection in animals that were classified as uninfected by parasitological techniques suggests that PCR could have an important role to play in the detection of infected animals harbouring a low parasitaemia that would be undetectable by smears (Chansiri *et al.*, 2002; Bashir *et al.*, 2011).

The better PCR sensitivity and the fact that samples can be taken as blood spots on filter papers open the opportunity to carry out epidemiological studies in remote regions in countries where accessibility and conditions to store samples are really difficult, Sample processing in PCR does not have to be done within minimum time after collection but can be delayed at least 180 days after preservation at -20°C (Clausen *et al.*, 1998). The main advantages of different PCR assays are very low false-positives and insensitive to other haemoparasite species (Chansiri *et al.*, 2002). Conventional PCR (Omer *et al.*, 1998; Guterrez *et al.*, 2000; Masiga and Nyang'ao 2001; Bashir

et al., 2011) and real time PCR (Yousef *et al.*, 2010.) has been used successfully for detecting of infection with *T. evansi* in camels.

The real time PCR technique as used in this study was able to detect the suspected acute and chronic infection of *T. evansi* and may be the low parasitaemia consistently a new, sensitive and rapid diagnostic technique.

The obtained result (42.15%) was similar to that recorded in Kenya; 39.8% (Njiru *et al.*, 2004) and 34.40% in India (Reghu *et al.*, 2008) but higher than that recorded in India (17.05%) and Kenya (26.6%) (Singh *et al.*, 2004; Njiru *et al.*, 2004), but lower than that in Sudan (57.1%) (Bashir *et al.*, 2011). The improved detection rates found in the present study, was also reported in studies carried out in the Brazilian pantanal (Ventura *et al.*, 2001; Ventura *et al.*, 2002): these may be attributed to selection of assay conditions and primer sets being used. Our results also revealed that the real time PCR assay was beneficial for diagnosis of *T. evansi* among camels and could be useful for epidemiological, following of drug treatment study and designing rational trypanosomiasis control program in the endemic area.

It is concluded that camel trypanosomiasis in Sudai is apparently caused by a single parasite species *T. evansi* and there were no other typanosomes species detected. In addition, the disease is highly prevalent in the country, which strengthens the need to change control policies and institute measures that help prevent the spread of the parasite. To our knowledge, this is the first molecular diagnosis report, which gives a picture of camel *trypanosomiasis* covering large geographical areas in Sudai Arabia.

REFERENCES

- Atarhouch, T.; Rami, M.; Bendahman, M.N. and Dakkak, A. (2003): Camel trypanosomosis in Morocco 1: results of a first epidemiological survey. *Vet. Parasitol.* 111: 277-286.
- Bashir Salim.; Mohammed A Bakheit; Joseph Kamau; Ichiro Nakamura and Chihiro Sugimoto (2011): Molecular epidemiology of camel trypanosomiasis based on ITS1 rDNA and RoTat 1.2 VSG gene in the Sudan.
- Boid, R.; Jones, T.W. and Luckins, A.G. (1985): Protozoal diseases of camels. *Brit. Vet. J.*, 141: 87-105.

- Chansiri, K.; Khuchareontaworn, S. and Sarataphan, N. (2002):* PCR-ELISA for diagnosis of *T.evansi* in animals and vector. *Molecular and Cellular Probes* 16: 173-177.
- Chaudhary, Z.I and Iqbal, M.J. (2000):* Incidence, biochemical and haematological alterations induced by natural trypanosomosis in racing dromedary camels. *Acta Trop* 77: 209-213.
- Clausen, P.-H.; Wiemann, A.; Patzelt, R.; Kakaire, D.; Poetzsch, C.; Peregrine, A. and Mehlitz, D. (1998):* Use of a PCR assay for the specific and sensitive detection of *Trypanosoma* spp. In naturally infected dairy cattle in peri-urban Kampala, Uganda *Ann. N.Y. Acad. Sci.* 849: 21–31.
- Davison, H.C.; Thrusfield, M.V.; Muharsini, S.; Hussein, A.; Partoutomo, S.; Rae, P.F.; Masake, R. and Luckins, A.G. (1999):* Evaluation of antigen and antibody detection tests for *Trypanosoma evansi* infection of buffaloes in Indonesia. *Epidemiol Infect.* 123: 149-155.
- Dia, M.L.; Diop, C.; Aminetou, M.; Jacquiet, P. and Thiam, A. (1997):* Some factors affecting the prevalence of *Trypanosoma evansi* in camels in Mauritania. *Vet. Parasitol.* 72: 111-120.
- Guterrez, C.; Juste, M.C.; Corbera, J.M.; Magnus, E.; Verloo, D. and Montoya, J.A. (2000):* Camel trypanosomosis in the Canary Island; assessment of seroprevalence and infection rate using the card agglutination test (CATT/ *T. evansi*) and parasite detection. *Vet. Parasitol.* 90: 15-9.
- Hopkins, J.S.; Harrison, C.; Noreen, M.; Luckins, A.G.; Rae, P.E.; Bossche, P. and Eisler, M.P. (1998):* Adaptation and validation of antibody – ELISA using dried blood spots on filter paper for epidemiological surveys of tsetse transmitted trypanosomosis in cattle. *Preventive Veterinary Medicine*, 37: 91-99.
- Imadeldin E. Aradaib and Ali A. Majid (2006):* A simple and rapid method for detection of *Trypanosoma evansi* in the dromedary camel using a nested polymerase chain reaction. *Kinetoplastid Biol. Dis.* 5: 2.
- Lohr, K.F.; Pholpark, S.; Siriwan, P.; Leesirikul, N.; Srikitjakarn, N. and Staak, L.C. (1986):* *Trypanosoma evansi* infection in buffaloes in north east Thailand. 2. Abortions. *Trop. Anim. Hlth. Prod.* 18: 103-108.
- Luckins, A.G. (1988):* *Trypanosoma evansi* in Asia. *Parasitol Today* 4: 137-142.

- Mahmound, M.M. and Gray, A.R. (1980):* Trypanosomosis due to *T. evansi*. A review of recent research. *Trop. Anim. Health Prod.* 12: 35-47.
- Masiga, R.C. and Nyang'ao, J.M. (2001):* Identification of trypanosome species from camel using polymerase chain reaction and procyclic transformation test. *Camel Pract. Res.* 8: 17-22.
- Nantulya, V.M. (1994):* Suratex®: A simple latex agglutination antigen test for diagnosis of *Trypanosoma evansi* (Surra). *Trop. Med. Parasitol.* 45: 9-12.
- Njiru, Z.K.; Constantine, C.C.; Ndung'u, J.M.; Robertson, I.; Okaye, S.; Thompson, R.C.A. and Reid, S.A. (2004):* Detection of *Trypanosoma evansi* in camels using PCR and CATT/T. *evansi* tests in Kenya. *Vet. Parasit.* 124(3-4): 187-99.
- Omuwa, S.; Rao, J.R.; Basagoudanavar, S.H.; Singh, R.H. and Butchaiah, G. (1999):* Direct and sensitive detection of *Trypanosoma evansi* by polymerase chain reaction. *Acta. Vet. Hung.* 47: 351-59.
- Omer, R.A.; Elamin, S.M.M.; El Nahas, A.E. and Aradaib, I.E. (2004):* PCR for detection of *Echinococcus granulosus* hydatid cysts collected from camels (*Camelus dromerarius*) The Sudan. *J. of Vet. Sci. and Anim. Husbandry.* 43:139-143.
- Omer, O.H.; Magzouls, M.; Haroun, E.M.; Mahmoud, O.M. and Abdel Hamid, Y.M. (1998):* Diagnosis of *Trypanosoma evansi* in Saudi Arabian camels (*Camelus dromedarius*) by the passive haemagglutination test and Ag-ELISA. *Zentralbl Veterinarmed.* 45: 627-33. *Parasites and Vectors.* 4:31.
- Pathak, K.M.L. and Khanna, M.D. (1995):* Trypanosomiasis in camel with particular reference to Indian subcontinent: A Review. *International J. Anim. Sci.* 10: 157-162.
- Reghu Ravindran, I.; Jammi Raghavendra Rao; Ashok Kumar Mishra, Satheesh, and Sachivothaman Rahul (2008):* *Trypanosoma evansi* in camels, donkeys and dogs in India: comparison of PCR and light microscopy for detection – short communication. *Vet. Arhiv.* 78 (1): 89-94.
- Singh, N.; Pathak, K.M. and Kumar, R. (2004):* A comparative evaluation of parasitological, serological and DNA amplification methods for diagnosis of natural *Trypanosoma evansi* infection in camels. *Vet. Parasitol.* 30, 126(4): 365-73.

- Ventura, R.M.; Paiva, F.; Silva, R.A.M.S.; Takeda, G.; Buck, G.A. and Teixeira, M. (2001):* Trypanosoma vivax: characterisation of the spliced – leader gene of a Brazilian stock and species – specific detection by PCR amplification of and intergenic spacer sequence. *Experimental Parasitology.* 99: 37-48.
- Ventura, R.M.; Takeda, G.F.; Silva, R.A.M.S.; Nunes, V.L.B.; Buck, G.A. and Teixeira, M. (2002):* Genetic relatedness among Trypanosoma evansi stocks by random amplification of polymorphic DNA and evaluation of a synapomorphic DNA fragment for species – specific diagnosis. *International Journal for Parasitology.* 32: 53-63.
- Viseshakul, N. and Panyim, S. (1990):* Specific DNA probe for the sensitivity detection of Trypanosoma evansi, Southeast Asian. *J Trop. Med. Public. Health,* 21: 21–27.
- Wuyts, N.; Chokesajjawatee, N.; Sarataphan, N. and Panyim, S. (1995):* PCR amplification of crude blood on microscope slides in the diagnosis of Trypanosoma evansi infection in dairy cattle *Ann Soc. Belg. Med. Trop.* 75: 229-37.
- Yousef, M.R.; Al-Khatib, R.M.; Mazloun, K.S.; Al-Sukayran, A. and Al Nakhli, H.M. (2010):* Detection of Trypanosoma evansi in camels using SYBER Green1 real time PCR assay. *Assiut Vet. Med. J.* 56 (127): 279-287.