

Dept. of Artificial Insemination and Embryo Transfer,
Animal Reproduction Research Institute, Al Haram,
(P. O. B. 12556), Giza, Egypt.

**EFFECT OF EPIDERMAL AND INSULIN-LIKE
GROWTH FACTOR-1 ON NUCLEAR MATURATION
AND EMBRYO DEVELOPMENT OF BUFFALO
OOCYTES IN VITRO**
(With 5 Tables)

By
M.R. BADR and F.A. EL ZOHERY
(Received at 4/6/2011)

تأثير إضافة عوامل النمو الشبيهة بالإنسولين وبالإبيدرمل على معدل نضوج
وإخصاب ونمو بويضات الجاموس معمليا

مجدى رمضان بدر ، فوزى عبد الحميد الزهيرى

تهدف الدراسة الحالية إلى دراسة تأثير إضافة عوامل النمو (عامل النمو الشبيهة بالإنسولين-1 أو عامل النمو الشبيهة بالإبيدرمل) بتركيزات مختلفه إلى ميديا النضوج المحدد تركيبها الكيميائى على معدل نضوج وإخصاب ونمو بويضات الجاموس معمليا. تم تجميع البويضات من الحويصلات المتوسطة الحجم (٢ الى ٨ مللى) وبعد تقييم البويضات بناء على شكل الخلايا الوسادية المحيطة بالبويضة وكذلك شكل السيتوبلازم للبويضة. تم إنضاج البويضات لمدة ٢٤ ساعة فى ميديا النضوج المحدد تركيبها الكيميائى بدون أى إضافات كمجموعة ضابطة أو مجموعة مضاف إليها عامل النمو الشبيهة بالإبيدرمل بتركيزات (١٠، ٥، ٢٠ نانوجرام/مللى) أو عامل النمو الشبيهة بالانسولين بتركيزات (١٠٠، ٢٥، ٥٠ نانوجرام/مللى) كلا على حده أو معا. وبعد ذلك تم إخصاب البويضات الناضجة بسائل منوى مجمد بعد معالجته بالهيبارين لمدة ١٨ ساعة وفى نفس ظروف الإنضاج. وبعد تقييم معدل الإخصاب تم زراعة البويضات المخصبة معمليا لمدة سبعة ايام وملاحظة النمو حتى طور البلاستوسيست. ولقد أظهرت النتائج أن إضافة ٥ نانوجرام/مللى عامل النمو الشبيهة بالإنسولين-١ و ٥٠ نانوجرام/مللى عامل النمو الشبيهة بالإبيدرمل معا إلى ميديا النمو قد أدى إلى زيادة معنويه كبيرة فى معدل النضوج والإخصاب ومعدل النمو إلى الطور التوتى وطور البلاستوسيست (٨٣،٤٨ ، ٦٢،٠٠ ، ٢٩،١٢ ، ٢٢،٨٧% على التوالي) بالمقارنة بالمجموعه الضابطه (٢٠،٧٥ ، ٦١،٩٣ ، ٧،٢٥ ، ٢٠،٨٠% على التوالي). من خلال نتائج الدراسه الحاليه يمكن أن نستنتج أن عوامل النمو لها تأثير إيجابي على معدل النضوج والإخصاب والنمو المعملى بناء على الجرعة المؤثرة. كما أن إضافة ٥ نانوجرام/مللى من عامل النمو

الشبية بالإتسولين-1 أو ٥٠ نانوجرام/ملى من عامل النمو الشبيه بالإبيدرمل معا إلى ميديا النمو له تأثير تعضيدي مما يؤدي إلى زيادة معنوية كبيرة لمعدل النضوج والإخصاب المعملى ومعدل النمو إلى الطور التوتى وطور البلاستوسيست.

SUMMARY

The present study was aimed to investigate the possible effects of epidermal growth factor (EGF) and insulin-like growth factor 1 (IGF-1) on nuclear maturation, fertilization and embryo development in vitro of buffalo cumulus oocyte complexes (COC's). Oocytes were matured in vitro for 24 h in a defined SOF media with no supplements (control) or in SOF media supplemented with EGF (5, 10 and 20 ng/ml), IGF-1 (25, 50, 100 ng/ml) or with a combination of EGF and IGF-1. After 24 h of in vitro maturation, oocytes were inseminated with sperm prepared in S-TALP medium supplemented with 10 µg/ml heparin. Embryos were evaluated for cleavage and development to the morula and blastocyst stages. The current results revealed that, addition of a combination of 5 ng/ml EGF plus 50 ng/ml IGF-1 to the maturation medium significantly increased ($P < 0.01$) the in vitro maturation, fertilization rate and embryo development to the morula and blastocyst stages (83.48 ± 2.27 , 62.00 ± 3.19 , 29.12 ± 4.13 and $22.87 \pm 2.26\%$, respectively) as compared with the control (61.93 ± 4.09 , 32.75 ± 2.77 , 7.25 ± 1.45 and $2.80 \pm 1.42\%$, respectively). In conclusion, the current results inferred that EGF and IGF-1 had a positive effect on buffalo oocytes maturation, fertilization and subsequent embryo development in a dose dependent trend. Supplementation of maturation medium with a combination of 5 ng/ml EGF and 50 ng/ml IGF-1 enhanced the in vitro maturation, fertilization and embryo development.

Key words: Buffalo, epidermal growth factor, insulin-like growth factor, in vitro fertilization.

INTRODUCTION

Recent studies have stressed the need for the use of chemically defined serum-free media for in vitro maturation (IVM), in vitro fertilization (IVF) and subsequent development of follicular oocytes in vitro. In general, chemically defined culture

media afford development of lower blastocyst yields (Sirisathien and Brackett, 2003). To improve blastocyst development, several common growth factors have been added to embryo culture media as they are known to be involved in embryo development (Baştan *et al.*, 2010). Amongst the growth factors studied as additives in such media, a combination of epidermal growth factor (EGF) and insulin-like growth factor-1 (IGF-1) appears to be an optimum combination. EGF is a mitogenic factor which has the ability to stimulate the proliferation of ovarian granulosa cells (May *et al.*, 1987). Many reports have reported that EGF contributes to the promotion of oocyte maturation (Sanbuissho *et al.*, 1991), germinal vesicle breakdown (GVBD), polar body formation (Das *et al.*, 1991) and cleavage of the oocytes (Coskun *et al.*, 1991). Moreover, several previous studies have shown that addition of IGF-1 to culture media in vitro promotes maturation of oocytes (Harper and Brackett, 1993; Rieger *et al.*, 1995) and also affects their subsequent development in vitro (Herrler *et al.*, 1992; Palma *et al.*, 1997). EGF and IGF-1 in combination have been shown to act synergistically and to accelerate the cumulus expansion and the progression of meiosis (Lorenzo *et al.*, 1994; Purohit, 2001; Sakaguchi *et al.*, 2002). Despite their actions on early embryonic development, growth factors are not routinely included in embryo culture medium. The lack of exposure to the proper growth factor milieu may be one reason why embryos produced in vitro differ from their counterparts derived in vivo (Block *et al.*, 2007). However, a few reports have shown that addition of IGF-1 had no effect on meiotic maturation, fertilization or embryonic development of oocytes in vitro (Gruppen *et al.*, 1997; Guler *et al.*, 2000). Therefore, the current study was designed to evaluate the effect of EGF and IGF-1, singly or in combination, in serum-free SOF medium on in vitro nuclear maturation, fertilization and embryo development of buffalo COC's.

MATERIALS and METHODS

Oocyte selection and in vitro maturation:

Ovaries were obtained from an abattoir and were transported to the laboratory in PBS at 35°C within 2 h. Oocytes were aspirated from medium follicles (2-8 mm in diameter) using

an 18-gauge needle attached to a 10 ml disposable syringe. Only oocytes having a dense cumulus cell mass and homogeneous cytoplasm were selected. Oocytes were washed 3 times with Dulbecco's phosphate-buffered saline. The selected oocytes were cultured in 35 mm Petri dishes at 39°C under an atmosphere of 5% CO₂ in air, 95% humidity for 24 h for maturation. The oocytes were matured in SOF media with the addition of sodium bicarbonate, penicillin 100 IU ml⁻¹ and streptomycin 50 µg ml⁻¹ (control) or with addition of test substances (growth factors). After 24 hours, nuclear maturational status was assessed by aceto-orcien stain as described by Lorenzo *et al.* (1994).

Sperm preparation and in vitro fertilization (IVF):

Three straws of frozen semen were thawed in a water bath at 38°C for 30 sec. After thawing, the most motile spermatozoa were separated by swim up technique in sperm-TALP medium containing 6 mg/ml bovine serum albumin, for 1h (Parrish *et al.*, 1986). The uppermost layer of the medium containing the most spermatozoa was collected. The selected spermatozoa were washed twice by centrifugation at 2000 rpm for 10 minutes. The sperm pellet was re-suspended in the fertilization TALP (F-TALP) medium containing 10 µg/ml heparin. The prepared sperm was incubated in a CO₂ incubator at 38.5°C, 5% CO₂ for 2 h before further use. The matured oocytes were washed with F-TALP medium and the prepared sperm was added into the droplets containing matured oocytes to give a final concentration of 2 X10⁶ sperm cells/ml. Gametes were co-incubated in the fertilization drops under sterile mineral oil for 18 hour. At the end of gametes co-incubation, some of inseminated oocytes were freed of the attached cumulus cells, fixed in acetic acid- ethanol (1:3), stained with 1% aceto-orcien stain and examined under phase-contrast microscope (X 400) for assessing the in vitro fertilization rate according to Totey *et al.* (1992).

In vitro culture:

Presumptive zygotes were denuded from cumulus cells and the extra spermatozoa by gentle pipetting and transferred, into SOFaa culture medium (SOF with 1 mM glutamine, 1% MEM nonessential amino acids and 1% MEM essential amino acids) and covered with mineral oil for 7- days at 38.5 °C in an

atmosphere of 5% CO₂ in air with maximum humidity. The proportional of cleaved oocytes was recorded 48 hour after insemination and those developed to the morula and blastocyst stages were recorded at 5-7 days post-insemination according to Totey *et al.* (1992).

Experiment 1:

The effect of different growth factor concentrations, individually or in a combination, on nuclear maturation of buffalo oocytes were examined. Maturation medium with no extra supplement served as control. Growth factors were added to maturation medium at different concentrations, IGF-1 (25, 50, 100 ng/ml, respectively) or EGF (5, 10 and 20 ng/ml, respectively). Selected COC's were matured randomly in the media of different treatments. After 24 h the oocytes were examined to evaluate in vitro maturation rate.

Experiment 2:

COC's were matured for 24 h in the best concentrations of the growth factors resulted from experiment 1, singly or in combination. The matured oocytes were then fertilized and cultured in vitro to evaluate fertilization and embryo development rates in relation to each treatment.

Statistical analysis:

All data were analyzed by using *Costat Computer Program, Version 3.03* copyright (1986) *Cottort Software*, and were compared by the least significant difference least (LSD) at 1% and 5% levels of probability.

RESULTS

Data presented in Table 1 revealed that, fortification of the maturation medium with different concentrations of IGF-1 improved the in vitro maturation rate compared to the control in a dose-dependent trend. Addition of 50 ng/ml IGF-1 to the maturation medium improved ($P < 0.01$) significantly the in vitro maturation rate (72.04 ± 3.45 %) as compared with the control (51.48 ± 4.55 %).

Table 1: Effect of different concentrations of insulin-like growth factor 1 on the in vitro maturation rate.

Treatment	No. of oocytes	No. of matured oocytes	Maturation rate
IGF-1 25ng/ml	75	50	66.79± 1.95 ^a
IGF-1 50 ng/ml	54	39	72.04± 3.45 ^a
IGF-1 100 ng/ml	72	39	54.17± 1.17 ^b
Control	78	40	51.48± 4.55 ^b

IGF-1: insulin-like growth factor 1

Values with different superscript letters in the same columns are significantly different at least ($P<0.05$).

Results presented in Table 2 revealed that, supplementation of the maturation medium with different concentrations of EGF improved the in vitro maturation rate compared to the control in a dose-dependent trend. Addition of 5 ng/ml EGF to the maturation medium increased ($P<0.01$) significantly the in vitro maturation rate ($79.35\pm 3.42\%$) as compared with the control ($64.73\pm 2.99\%$).

Table 2: Effect of different concentrations of epidermal growth factor on buffalo oocytes maturation rate in vitro.

Treatment	No. of oocytes	No. of matured oocytes	Maturation rate
EGF 5ng/ml	83	66	79.35± 3.42 ^a
EGF 10 ng/ml	81	60	74.15± 3.77 ^{ab}
EGF 20 ng/ml	86	56	65.15± 1.68 ^b
Control	76	49	64.73± 2.99 ^b

EGF: Epidermal growth factor

Values with different superscript letters in the same columns are significantly different at least ($P<0.05$).

Data regarding the effect of replenishing of the maturation medium with the best concentration of the growth factors resulted from the previous experiments either singly or in combination is presented in table 3. The current results revealed that, when growth factors are combined their effects are additive.

Table 3: Effect of insulin-like growth factor 1 and /or epidermal growth factor on the in vitro maturation rate.

Treatment	No. of oocytes	No. of matured oocytes	Maturation rate
IGF-1 50ng/ml	54	41	76.03± 1.99 ^a
EGF 5 ng/ml	57	44	77.07± 1.14 ^a
IGF-1 0ng/ml+ EGF 5 ng/ml	73	61	83.48± 2.27 ^a
Control	71	44	61.93±4.09 ^b

IGF-1: insulin-like growth factor 1 EGF: Epidermal growth factor
 Values with different superscript letters in the same columns are significantly different at least ($P<0.05$).

Results regarding the effect of augmentation of the maturation medium with 50 ng/ml IGF-1 and/or 5 ng/ml EGF on the in vitro fertilization rate are presented in table 4. The current results revealed that, addition of IGF-1 and EGF in combination to the maturation medium had a positive effect on the in vitro fertilization rate ($P<0.01$) as compared with the control (62.00 ± 3.19 vs. 32.75 ± 2.77).

Table 4: Effect of insulin-like growth factor 1 and /or epidermal growth factor on the in vitro fertilization rate

Treatment	No. of oocytes	Penetration rate	Fertilization rate	Abnormal fertilization
IGF-1 50ng/ml	73	54 (73.97±1.16) ^a	32 (45.16±3.57) ^b	13 (17.91±1.91) ^{ab}
EGF 5 ng/ml	81	55 (67.87±2.54) ^a	36 (44.44±1.86) ^b	13 (16.86±2.80) ^{ab}
IGF-1 50ng/ml+ EGF 5 ng/ml	76	55 (72.59±2.50) ^a	47 (62.00±3.19) ^a	9 (11.77±1.92) ^b
Control	71	50 (70.74±2.73) ^a	23 (32.75±2.77) ^c	16 (22.26±1.73) ^a

IGF-1: insulin-like growth factor 1 EGF: Epidermal growth factor
 Values with different superscript letters in the same columns are significantly different at least ($P<0.05$).

Likewise, data presented in Table 5 revealed that, combination of IGF-1 and EGF resulted in a significant increase ($P<0.01$) in the cleavage rate and embryo development to the morula and blastocyst stages (46.56 ± 3.48 , 29.12 ± 4.13 and $22.87\pm2.26\%$, respectively) as compared with the control (28.83 ± 1.66 , 7.25 ± 1.45 and $2.80\pm1.42\%$, respectively).

Table 5: Effect of insulin-like growth factor 1 and /or epidermal growth factor on the in vitro embryo development rate.

Treatment	No. of oocytes	Cleavage rate	Morula stage	Blastocyst stage
IGF-1 25ng/ml	61	24 (39.81 ± 2.65) ^a	10 (16.59 ± 1.53) ^b	6 (9.96 ± 1.97) ^b
EGF 5 ng/ml	70	29 (41.85 ± 3.09) ^a	13 (18.93 ± 2.63) ^b	8 (11.49 ± 1.51) ^b
IGF-1 50ng/ml+ EGF 5 ng/ml	75	35 (46.56 ± 3.48) ^a	22 (29.12 ± 4.13) ^a	17 (22.87 ± 2.26) ^a
Control	69	20 (28.83 ± 1.66) ^b	5 (7.25 ± 1.45) ^c	2 (2.80 ± 1.42) ^c

IGF-1: insulin-like growth factor 1

EGF: Epidermal growth factor

Values with different superscript letters in the same columns are significantly different at least ($P<0.05$).

DISCUSSION

The role of serum on meiosis resumption and oocyte maturation (Younis *et al.*, 1989) has been demonstrated. However, several studies have recommended the use of serum-free medium for in vitro fertilization (Takagi *et al.*, 1991) because serum quality varies from batch to batch and serum may contain materials toxic to the cell culture (Ogawa *et al.*, 1987). It is difficult to define components already contained in the serum. The present study demonstrated that EGF and IGF-1 enhanced the in vitro maturation, fertilization and embryo development in a dose-dependent trend, but the results were marked when both EGF and IGF-1 were combined. These results were in accordance with previous studies having shown that EGF and IGF-1 have a positive effect on oocyte maturation (Kobayashi *et al.*, 1994). Supplementation of EGF resulted in a higher proportion of nuclear maturation and normal fertilization rates compared to the control in a dose-dependent manner (Im and

Park, 1995). The dose dependent effect of EGF supplementation in increasing the proportion of oocytes reaching nuclear maturation was evident only up to 20 ng/ml. As pointed out by Harper and Brackett (1993), EGF in serum is possibly one of the undetermined components contributing to enhanced oocyte maturation. In the present study, EGF supplementation increased significantly the cleavage, and embryo development rate compared with the control. These results are in consistent with, Lonergan *et al.* (1996); Baştan *et al.* (2010) who reported that blastocyst development was significantly improved for oocytes matured in the presence of EGF.

Moreover, one growth factor that modifies embryonic physiology is insulin-like growth factor-1 (IGF-1). In the present study, IGF-1 treatment tended to increase the proportion of oocytes that matured, fertilized and developed to the blastocyst stage. Similar results were also observed in several studies (Lorenzo *et al.*, 1996; Block *et al.*, 2007). The effect of IGF-1 on embryo development may be partly explained by differences in culture systems since there are reports that the actions of IGF-1 to stimulate embryonic development depend upon culture conditions. IGF-1 has been shown to act on the development of bovine embryos indirectly, via granulosa cells (Palma *et al.*, 1997), the IGF-1 receptors being located in the plasma membrane of granulosa cells (Adashi *et al.*, 1988) and that IGF-1 stimulates the proliferation and differentiation of these cells (Spicer *et al.*, 1993). Moreover, IGF-1 can reduce the proportion of blastomeres that are apoptotic (Sirisathien and Brackett, 2003), alter the abundance of some developmentally important genes (Block *et al.*, 2007), and increase cellular resistance to heat shock (Jousan and Hansen, 2007). Also, IGF-1 can increase development of bovine embryos to the blastocyst stage (Block and Hansen, 2007) and can increase blastocyst cell number (Sirisathien *et al.*, 2003).

Furthermore, the proportion of oocytes that matured, fertilized or developed in vitro were significantly higher in the defined SOF medium supplemented with a combination of EGF and IGF-1 compared to their respective controls. These results were in accordance with previous studies having shown that EGF and IGF-1 have a positive effect on embryo development in various species (Lorenzo *et al.*, 1996; Kumar and Purohit, 2004;

Baştan *et al.*, 2010). This finding indicates a synergistic action between the growth factors used and suggests that their actions are collective under in vitro conditions.

In conclusion, the current results inferred that EGF and IGF-1 had a positive action on buffalo oocytes maturation, fertilization and subsequent embryo development in a dose dependent trend. Supplementation of maturation medium with a combination of 5 ng/ml EGF and 50 ng/ml IGF-1 enhanced the in vitro nuclear maturation, fertilization and embryo development.

REFERENCES

- Adashi, E.Y.; Resnick, C.E.; Hernandez, E.R.; Svoboda, M.E. and Vanwyk, J.S. (1988):* Characterization and regulation of a specific cell membrane receptor for somatomedin C/insulin-like growth factor-I in cultured rat granulosa cells. *Endocrinology*, 122: 194-201.
- Baştan, A.; Polat, B.; Bakiacar, D.; Korkmaz, M. and Olak, A. (2010):* Determination of optimal dose of EGF for bovine oocyte maturation and subsequent in vitro fertilization and culture in two media. *Turk. J. Vet. Anim. Sci.*, 34(1): 33-38.
- Block, J. and Hansen, P.J. (2007):* Interaction between season and culture with insulin-like growth factor-1 on survival of in vitro produced embryos following transfer to lactating dairy cows. *Theriogenology*, 67: 1518-29.
- Block, J.; Fischer-Brown, A.E.; Rodina, T.M.; Ealy, A.D. and Hansen, P.J. (2007):* The effect of in vitro treatment of bovine embryos with IGF-1 on subsequent development in utero to Day 14 of gestation. *Theriogenology*, 68: 153-161
- Coskun, S.; Sonbuissho, A.; Lin, Y.C. and Rikihisa, Y. (1991):* Fertilization and subsequent developmental ability of bovine oocytes matured in medium containing epidermal growth factor (EGF). *Theriogenology*, 36: 485-494.
- Costat Computer Program Copyright (1986):* Version 3.03 copyright Cottort Software.
- Das, K.; Tagatz, G.E.; Stout, L.E.; Phipps, W.R.; Hensleigh, H.C. and Leung, B.S. (1991):* Direct positive effect of epidermal growth factor on the cytoplasmic maturation of mouse and human oocyte. *Fertil. Steril.* 55: 1000-1004.

- Gruppen, C.G.; Nagashima, H. and Nottle, M.B. (1997):** Role of epidermal growth factor and insulin-like growth factor-1 on porcine oocyte maturation and embryonic development *in vitro*. *Reprod. Fertil. Dev.* 9: 571-575.
- Guler, A.; Poulin, N.; Mermillod, P.; Terqui, M. and Cognie, Y. (2000):** Effect of growth factors, EGF and IGF-1 and estradiol on *in vitro* maturation of sheep oocytes. *Theriogenology*, 54: 209-218.
- Harper, K.M. and Brackett, G. (1993):** Bovine blastocysts development after *in vitro* maturation in a defined medium with epidermal growth factor and low concentrations of gonadotropins. *Biol. Reprod.*, 48: 409-416.
- Herrler, A.; Lucas-Ham, A. and Nienann, A. (1992):** Effects of insulin like growth factor-I on *in vitro* production of bovine embryos. *Theriogenology*, 37: 1213-1224.
- Im, K.S. and Park, K.W. (1995):** Effects of epidermal growth factor on maturation, fertilization and development of bovine follicular oocytes. *Theriogenology*, 44: 209-216.
- Jousan, F.D. and Hansen, P.J. (2007):** Insulin-like growth factor-I promotes resistance of bovine preimplantation embryos to heat shock through actions independent of its anti-apoptotic actions requiring PI3K signaling. *Mol. Reprod. Dev.*, 74: 189-96.
- Kobayashi, L.; Yamashita, S. and Hoshi, H. (1994):** Influence of epidermal growth factor's and transforming growth factor 1 on *in vitro* maturation of cumulus cell enclosed bovine oocytes in a defined medium. *J. Reprod. Fertil.*, 100: 439-446.
- Kumar, D. and Purohit, G.N. (2004):** Effect of epidermal and insulin-like growth factor-1 on cumulus expansion, nuclear maturation and fertilization of buffalo cumulus oocyte complexes. *Vet. Arhiv.*, 74: 13-25.
- Lonergan, P.; Carolan, C.; Van Langendonck, A.; Donnay, I.; Khatir, H. and Mermillod, P. (1996):** Role of epidermal growth factor in bovine oocyte maturation and preimplantation embryo development *in vitro*. *Biol. Reprod.*, 54: 1420-1429.
- Lorenzo, P.L.; Illera, M.J.; Illera, J.C. and Illera, M. (1994):** Enhancement of cumulus expansion and nuclear maturation during bovine oocyte maturation *in vitro* by the

- addition of epidermal growth factors and insulin like growth factors. *J. Reprod. Fertil.*, 101 : 697-701.
- Lorenzo, P.L.; Reboller, P.G.; Illera, M.J.; Illera, J.C.; Illera, M. and Alvarino, J.M.R. (1996):* Stimulatory effect of insulin-like growth factor 1 and epidermal growth factor on the maturation of rabbit oocytes *in vitro*. *J. Reprod. Fertil.*, 107: 109-117.
- May, J.V.; Buck, P.A. and Schomberg, D.W. (1987):* Epidermal growth factor enhances (125) iodo-follicle stimulating hormone binding by cultured porcine granulosa cells. *Endocrinology*, 120: 2413-2420.
- Ogawa, T.; Ono, T. and Mars, R.P. (1987):* The effect of serum fractions on single cell mouse embryos *in vitro*. *J. in vitro Fert. Embryo Transfer* 4: 153-158.
- Palma, G.A.; Muller, M. and Brem, G. (1997):* Effect of insulin like growth factor-1 (IGF-1) at high concentrations on blastocyst development of bovine embryos produced *in vitro*. *J. Reprod. Fertil.*, 110: 347-353.
- Parrish, J.J.; Susko-Parrish, J.L.; Leibfried-Rutledge, M.L.; Critser, E.S.; Eyestone, W.H. and First, N.L. (1986):* Bovine *in vitro* fertilization with frozen-thawed semen. *Theriogenology*, 25: 591-600.
- Purohit, G.N. (2001):* Effect of serum free media and growth factors on IVM, IVF and Intravaginal culture of buffalo oocytes. Ph.D. Thesis, Rajasthan Agricultural University, Bikaner.
- Rieger, D.; Luciano, A.M.; Modina, S.; Pocar, P.; Luria, A. and Gandolfi, F. (1995):* The effect of EGF and IGF-1 on metabolism and nuclear maturation of cattle oocytes. *J. Reprod. Fertil.*, Abstract.
- Sakaguchi, M.; Dominko, T.; Yamauchi, N.; Leibfried-Rutledge, M.L.; Nagai, T.A. and First, N.L. (2002):* Possible mechanism for acceleration of meiotic progression of bovine follicular oocytes by growth factors *in vitro*. *Reprod.*, 123: 135-142.
- Sanbuissho, A.; Coskun, S. and Lin, Y.C. (1991):* Stimulatory action of epidermal growth factor (EGF) on *in vitro* bovine oocyte maturation. *Assist. Reprod. Technol. Androl.*, 1: 143-153.

- Sirisathien, S. and Brackett, B.G. (2003): TUNEL analyses of bovine blastocysts after culture with EGF and IGF-I. *Mol. Reprod. Dev.*, 65: 51-6.
- Spicer, I.J.; Alpisar, E. and Echternkamp, S.E. (1993): Effects of insulin-like growth factor-1 and gonadotropins on bovine granulosa cell proliferation, progesterone production, estradiol production and (or) insulin-like growth factor 1 production *in vitro*. *J. Anim. Sci.*, 71: 1232-1241.
- Takagi, Y.; Mori, K.; Tomizawa, M.; Takahashi, T.; Sugawara, S.S. and Masaki, J. (1991): Development of bovine oocytes matured, fertilized and cultured in serum-free chemically defined medium. *Theriogenology*, 35:1197-1207.
- Totey, S.M.; Singh, G.; Taneja, M.; Pawshe, C.H. and Talwar, G.P. (1992): *In vitro* maturation, fertilization and development of follicular oocytes from buffalo (*Bubalus bubalis*). *Theriogenology*, 39: 887-898.
- Younis, A.I.; Brackett, B.G. and Fayer-Hosken, R.A. (1989): Influence of serum and hormones on bovine oocyte maturation and fertilization *in vitro*. *Gamete Res.*, 23: 189-201.