

THE RELATIONSHIP BETWEEN LIGHT INTENSITY AND DRY FORAGE YIELD OF COWPEA AND MAIZE UNDER INTERCROPPING

(Received: 29. 9. 2011)

By

K.I. Abdel-Gawad, A.A. Abdel-Hafeez, W. M. Sharawy* and W. M. El-Nabawy*

Department of Agronomy, Faculty of Agriculture, Cairo University

** Forage Crops Research Section, FCRI, Agriculture Research Center, Giza, Egypt.*

ABSTRACT

Two field experiments were conducted at the Agricultural Experiments and Research Station, Faculty of Agriculture, Cairo University, Giza, Egypt, during 2008 and 2009 summer seasons, to study the intercropping of forage cowpea (*Vigna unguiculata* L., var. Buff) with maize (*Zea mays* L.), stay green and single cross hybrid 122. The experiment was planted at the 1st of July after wheat in both seasons. The experimental design was a split-split plot design arranged in randomized complete blocks with three replicates. The main plots were devoted to three intercropping patterns: 1) Maize on one side of the ridge and forage cowpea on the other side, 2) Solid maize, 3) Solid forage cowpea. Sub-plots were arranged with plant density of forage cowpea, one and two plants hill⁻¹ at 20 cm between hills. Sub-sub plots were four nitrogen levels viz., zero, 60, 90 and 120 kg N fed.⁻¹ N was added in two equal doses after 21 and 60 days from planting (after the 1st cut of forage cowpea).

The results indicated significant differences between intercropping patterns, plant density and nitrogen levels for dry yield. Light intensity at top, middle and bottom of cowpea and maize were obtained at each cut in both seasons. Light intensity under solid cowpea was greater for the three levels of light intensity reading than intercropped cowpea plants. At the top of intercropped cowpea plants, light intensity was greater as compared with readings at the middle and bottom of plants. At the middle, light intensity was greater for solid cowpea compared to intercropped cowpea. One plant hill⁻¹ was superior to two plants hill⁻¹ in light intensity at top, middle and bottom of cowpea plants intercropped with maize at each cut in both seasons. The percentage of increases in light intensity gave lower values between the two plant intensities, it could be recommended with planting two plants hill⁻¹ when intercropped cowpea with maize for obtaining high dry forage yield of cowpea. Nitrogen application was significantly decreased light intensity at top, middle and bottom plant of forage cowpea at each cut in both seasons. Nitrogen from zero to 120 kg N fed⁻¹ caused reduction in light intensity, While total dry forage yield of cowpea was increased. The highest value of light intensity was obtained by solid cowpea, one plant hill⁻¹ and zero nitrogen fed⁻¹, while the lowest value of light intensity was at intercropped cowpea with maize, two plants hill⁻¹ and 120 kg N fed.⁻¹ For maize plants there were significant differences in light intensity as affected by intercropping patterns at top, middle bottom of maize plants in both seasons except between intercropping two plants hill⁻¹ of cowpea with maize and solid maize at top plant before the 1st cut of cowpea in the first season. Nitrogen levels was significantly decreased light intensity at top, middle and bottom plant of maize, with not significant differences between 90 and 120 kg N fed.⁻¹ While, grain yield and dry stover yield were increased. Zero N level had the highest value of light intensity at top, middle and bottom of maize plants compared with the other N levels. The highest value was obtained by solid maize and zero nitrogen fed.⁻¹ while the lowest value was at intercropping two plants hill⁻¹ of cowpea with maize and 120 kg N fed.⁻¹

Key words: forage cowpea, intercropping patterns, light intensity, maize, nitrogen levels.

1. INTRODUCTION

Growing mixed stands of two or more crop species, defined as intercropping, has many advantages over sole cropping. It provides for efficient utilization of environmental resources, reduces the cost of production, provides greater

financial stability for farmers, decreases pest damages, suppresses weed growth, improves soil fertility when legumes are included and improves forage yield and quality (Ofori and Stern, 1987).

Li *et al.*, 2003 indicated that intercropping maximizes the use of the above ground and

environment resources, including space, light and nutrients, and improve crop yield and quality.

The advantages from intercropping was maximized when the intercropped species complement each other and use the environmental resources more efficiently. Also, they improve the efficiency of using both above-ground and below-ground resources compared to growing the crops as pure stands. The total productivity of an intercropping system is often more than sole cropping of the component crops (Marsalis and Angadi, 2009).

The present investigation aimed at studying the effect of intercropping forage cowpea with maize as a main crop, on light intensity under intercropping, plant density and nitrogen levels to improve forage quality and quantity of intercropped crops for late summer season.

2. MATERIALS AND METHODS

Two field trials were carried out during 2008 and 2009 summer seasons at the Agricultural Experiments and Research Station, Faculty of Agriculture, Cairo University, Giza, Egypt. Forage cowpea (*Vigna unguiculata* L., var. Buff) was grown, using the maize hybrid S.C. 122, which has short stature plant that stay green after grains maturity. The experiments were planted in the 1st of July, following wheat in both seasons, and were designed as split-split plots arranged in randomized complete blocks with three replicates. The main plots were devoted to three intercropping patterns, viz., one side of the ridge for maize and the other side for forage cowpea, solid maize and solid cowpea. Sub-plots were devoted to two plant densities of cowpea i.e., one and two plants hill⁻¹ (30000, 60000 plants/fed) with hills space 20 cm apart, maize was planted as recommended density (24000 plants/fed) solid or intercropping with forage cowpea with 25 cm between hills and thinning at one plant hill⁻¹. Sub-sub plots were allotted to four nitrogen levels (Urea 46%) viz., zero, 60, 90 and 120 kg N fed⁻¹. N was added in two equal doses after 21 and 60 days from planting (after the 1st cut of cowpea). The experimental unit (sub-sub plot) consisted of 5 ridges, 4 m long and 70 cm apart, with area of 14.0 m² (1/300 fed). Representing soil samples at a depth of (0-30 cm) were taken before sowing for mechanical and chemical soil analyses in both seasons (Table 1). Forage cowpea was cut twice in both seasons, the 1st cut after 60 days of planting for estimating fresh

forage yield of cowpea, while the 2nd one included maize stover mixed with fresh forage from cowpea after 60 days from the 1st cut. Light intensity (Lux) was measured before cutting at the top, middle and bottom (20 cm from the soil surface) of forage cowpea and maize plants using a Lux-meter at 12 p.m. Dry yield (t fed⁻¹) of cowpea forage, grain and stover yields of maize (t fed⁻¹) were recorded. Data were analyzed by MSTAT-C Computer program 1986-V4 (Freed, 2005).

Table (1): Mechanical and chemical analyses of soil of the experimental sites before planting in 2008 and 2009.

Mechanical analysis (%)	2008	2009
Coarse sand	2.1	2.4
Fine sand	24.14	23.34
Silt	18.44	20.30
Clay	47.82	46.28
Soil type classification	Clay loam	Clay loam
Chemical analysis		
Available N (ppm)	54	51
Available P (ppm)	26	28
% CaCO ₃	3.83	3.51
% Organic matter	1.76	1.85
Soil pH	7.24	7.32

3. RESULTS AND DISCUSSION

3.1. Forage cowpea

3.1.1. Effect of treatments on light intensity

Data of light intensity (lux) (Tables 2 to 7) show the effect of intercropping patterns, plant density and nitrogen levels at top, middle and bottom of cowpea plants at the 1st and 2nd cut in both seasons.

3.1.1.1 Intercropping patterns

Significant differences were observed in light intensity as affected by intercropping patterns at top, middle and bottom of cowpea plants during the two cuts in both seasons. These results are in agreement with Badr (1998) who demonstrated that light interception by soybean plants was significantly affected by intercropping systems.

Light intensity was greater for solid cowpea at all three levels of plant canopy reading than intercropped cowpea.

Light intensity was the greatest at plant tops as compared with the intensity at the middle and bottom levels. At plant tops light intensity for solid cowpea was 3.4% and 2.3% greater than intercropped cowpea at the 1st and 2nd cuts in the first season. Light intensity was 3.2% and 3.3% greater for the same treatments in respective order, in the second season (Tables 2 and 3).

Light intensity at the middle of the plant was

greater for solid compared to intercropped cowpea by 8.3% and 6.8% at the 1st and 2nd cut in the first season, whereas it was greater by 12.1% and 11.2% during the same cuts in the second season (Tables 4 and 5).

Light intensity readings at the bottom of the canopy was increased by 14.6% and 15.1% for solid cowpea compared to intercropped cowpea at the 1st and 2nd cut in the first season, and was greater by 21.2% and 19.9% for the same treatments in the second season (Tables 6 and 7).

These data indicate that the increase in light intensity for solid cowpea culture compared to the intercropped cowpea was due to the competition between forage cowpea and maize for light during vegetative growth. Shading increases in intercropped cowpea than solid cowpea culture. These results are in harmony with those obtained by Abdel-Wahab, (2010) who found that light intensity within soybean canopies is significantly affected by the cropping system. Light intensity was higher for solid soybean as compared with intercropped soybean. Light intensity at the middle and the bottom of soybean canopies decreased under intercropping as compared with recommended solid planting, due to shading effects of adjacent maize plants.

3.1.1.2. Plant density

Light intensity was higher for one plant hill⁻¹ than two plants hill⁻¹ at the top, the middle and the bottom of cowpea canopy intercropped with maize at each cut in both growing seasons.

Light intensity in the first season, increased under one plant hill⁻¹ than two plants hill⁻¹ at the top level of the canopy by 2.3% at the 1st cut and 1.3% at the 2nd cut, whereas in the second season, the percentages of increases were 2.3% and 2.0% at the 1st and 2nd cuts, respectively (Tables 2 and 3).

At the middle level of plant canopy the increases in light intensity under one plant hill⁻¹ compared to two plants hill⁻¹ were 4.4% and 4.8% at the 1st and 2nd cuts in the first season, and 4.9% and 3.8%, respectively at each cut in the second season (Tables 4 and 5).

At the bottom level of the canopy, light intensity was higher under one plant hill⁻¹ than two plants hill⁻¹ by 8.1% and 6.3% at the 1st and 2nd cuts, respectively in the first season and by 8.9% and 8.8%, respectively at each cut in the second season (Tables 6 and 7).

It was clear that the percentages of increases in light intensity were lower between the two plant intensities. Thus, it could be recommended to

plant two cowpea plants hill⁻¹ when cowpea is intercropped with maize in order to obtain high dry forage yield of cowpea.

3.1.1.3. Nitrogen levels

Nitrogen application significantly decreased light intensity at top, middle and bottom levels of cowpea canopy at each cut in both seasons.

Under zero N fertilization light intensity was higher at plant tops compared with N treatments (60, 90 and 120 kg N fed⁻¹), being 1.6, 2.6 and 3.7% higher at the 1st cut, 2.6, 3.5 and 5.8% higher at the 2nd cut in the first season. Corresponding increases were 2.2, 3.4 and 4.3% at the 1st cut, and 1.4, 3.1 and 5.5% at the 2nd cut in the second season (Tables 2 and 3).

At the middle level of the canopy, zero fertilizer treatment showed the highest value of light intensity compared with 60, 90 and 120 kg N fed⁻¹ these increases as percentage were 8.2, 12.4 and 16.4% at the 1st cut, 4.6, 9.9 and 21.9% at the second one in the first season. In the second season, these increases were 9.0, 16.4 and 24.9%, by 9.5, 17.0 and 24.9% at the 1st and 2nd cuts, respectively (Tables 4 and 5).

The check treatment (0 kg N fed⁻¹) showed the highest level of light intensity at the bottom level of the canopy reaching 10.5, 19.4 and 24.0% at the 1st cut, and 10.0, 16.0 and 21.7% at the second one in the first season. Also, increases were 6.2, 20.6 and 28.2% at the 1st cut, 7.6, 19.2 and 29.6% at the 2nd cut in the second season, respectively (Tables 6 and 7).

Applying nitrogen at 120 kg N fed⁻¹ reduced light intensity, while increased total dry forage yield of cowpea. This may be due to shading effect of maize on cowpea plants and consequently light interception. These results are in agreement with those obtained by Bowes *et al.*, (1970) who found that plant height of soybean increased in response to reduced light intensity. Moreover, Abou Keriasha *et al.*, (2009) found that cowpea intercropped with maize could be attributed to shading effect by maize plants and hence a low amount of intercepted light by cowpea plants. This shading effect increased stem elongation resulting in the tallest plants were observed when cowpea was intercropped with maize on all ridges.

3.1.1.4. Effect of the interaction

The interaction of intercropping patterns, plant density and nitrogen levels influenced light intensity on cowpea intercropped with maize, with the highest value obtained by solid cowpea, one plant hill⁻¹ and zero N fed⁻¹, while the lowest

Table (2): Effect of intercropping pattern, plant density and nitrogen application on light intensity (lux) at the top of forage cowpea canopy in 2008.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	1398.0	1382.0	1355.0	1345.0	1370.0	1354.0	1317.0	1314.0	1295.0	1320.0
	Two plants	1368.0	1330.0	1315.0	1287.0	1325.0	1341.0	1310.0	1302.0	1283.0	1309.0
	Mean	1383.0	1356.0	1335.0	1316.0	1347.5	1347.5	1313.5	1308.0	1289.0	1314.5
Maize + Cowpea	One plant	1325.0	1311.0	1304.0	1300.0	1310.0	1334.0	1298.0	1285.0	1263.0	1295.0
	Two plants	1314.0	1295.0	1291.0	1272.0	1293.0	1327.0	1291.0	1270	1204.0	1273.0
	Mean	1319.5	1303.0	1297.5	1286.0	1301.5	1330.5	1294.5	1277.5	1233.5	1284.0
General mean	One plant	1361.5	1346.5	1329.5	1322.5	1340.0	1344.0	1307.5	1299.5	1279.0	1307.5
	Two plants	1341.0	1312.5	1303.0	1279.5	1309.0	1334.0	1300.5	1286.0	1243.5	1291.0
Overall mean		1351.3	1329.5	1316.3	1301.0	1324.5	1339.0	1304.0	1292.8	1261.3	1299.3
L.S.D. at 0.05 level:											
Intercropping pattern (A)		5.1					4.9				
Plant density (B)		---					---				
Nitrogen levels (C)		4.3					3.9				
A x B		4.8					4.7				
A x C		6.0					5.5				
B x C		6.0					5.5				
A x B x C		8.5					7.8				

Table (3): Effect of intercropping pattern, plant density and nitrogen application on light intensity (lux) at the top of forage cowpea canopy in 2009.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	1395.0	1370.0	1351.0	1340.0	1364.0	1373.0	1356.0	1337.0	1314.0	1345.0
	Two plants	1361.0	1337.0	1321.0	1317.0	1334.0	1362.0	1340.0	1319.0	1291.0	1328.0
	Mean	1378.0	1353.5	1336.0	1328.5	1349.0	1367.5	1348.0	1328.0	1302.5	1336.5
Maize + Cowpea	One plant	1362.0	1320.0	1315.0	1291.0	1322.0	1348.0	1327.0	1302.0	1267.0	1311.0
	Two plants	1326.0	1299.0	1270.0	1261.0	1289.0	1311.0	1294.0	1269.0	1226.0	1275.0
	Mean	1344.0	1309.5	1292.5	1276.0	1305.5	1329.5	1310.5	1285.5	1246.5	1293.0
General mean	One plant	1378.5	1345.0	1333.0	1315.5	1343.0	1360.5	1341.5	1319.5	1290.5	1328.0
	Two plants	1343.5	1318.0	1295.5	1289.0	1311.5	1336.5	1317.0	1294.0	1258.5	1301.5
Overall mean		1361.0	1331.5	1314.3	1302.3	1327.3	1348.5	1329.3	1306.8	1274.5	1314.8
L.S.D. at 0.05 level:											
Intercropping pattern (A)		4.7					7.4				
Plant density (B)		4.9					5.3				
Nitrogen levels (C)		4.5					5.9				
A x B		7.0					7.5				
A x C		6.4					8.4				
B x C		6.4					8.4				
A x B x C		9.1					11.9				

Table (4): Effect of intercropping pattern, plant density and nitrogen application on light intensity (lux) at the middle level of forage cowpea canopy in 2008.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	627.0	570.0	599.0	528.0	571.0	616.0	582.0	536.0	458.0	548.0
	Two plants	588.0	545.0	523.0	512.0	542.0	551.0	532.0	520.0	449.0	513.0
	Mean	607.5	557.5	541.0	520.0	556.5	583.5	557.0	528.0	453.5	530.5
Maize + Cowpea	One plant	584.0	531.0	502.0	463.0	520.0	547.0	524.0	496.0	441.0	502.0
	Two plants	552.0	513.3	476.0	462.0	500.8	541.0	513.0	480.0	414.0	487.0
	Mean	568.0	522.2	489.0	462.5	510.4	544.0	518.5	488.0	427.5	494.5
General mean	One plant	605.5	550.5	530.5	495.5	545.5	581.5	553.0	516.0	449.5	525.0
	Two plants	570.0	529.2	499.5	487.0	521.4	546.0	522.5	500.0	431.5	500.0
Overall mean		587.8	539.8	515.0	491.3	533.5	563.8	537.8	508.0	440.5	512.5
L.S.D. at 0.05 level:											
Intercropping pattern (A)							3.8				
Plant density (B)							5.2				
Nitrogen levels (C)							3.8				
A x B							5.5				
A x C							6.4				
B x C							6.4				
A x B x C							9.0				

Table (5): Effect of intercropping pattern, plant density and nitrogen application on light intensity (lux) at the middle level of forage cowpea canopy in 2009.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	662.0	608.0	576.0	534.0	595.0	637.0	581.0	538.0	516.0	568.0
	Two plants	640.0	591.0	536.0	473.0	560.0	633.0	577.0	534.0	468.0	553.0
	Mean	651.0	599.5	556.0	503.0	577.5	635.0	579.0	536.0	492.0	560.5
Maize + Cowpea	One plant	594.0	541.0	487.0	446.0	517.0	582.0	540.0	483.0	439.0	511.0
	Two plants	586.0	519.0	477.0	410.0	498.0	577.0	500.0	461.0	402.0	485.0
	Mean	590.0	530.0	482.0	428.0	507.5	579.5	520.0	472.0	420.5	498.0
General mean	One plant	628.0	574.0	531.5	490.0	556.0	609.5	560.5	510.5	477.5	539.5
	Two plants	613.0	555.0	506.5	441.5	529.0	605.0	538.5	497.5	435.0	519.0
Overall mean		620.5	564.8	519.0	465.8	542.5	607.3	549.5	504.0	456.3	529.3
L.S.D. at 0.05 level:											
Intercropping pattern (A)							4.9				
Plant density (B)							7.8				
Nitrogen levels (C)							4.5				
A x B							9.3				
A x C							6.3				
B x C							6.3				
A x B x C							8.9				

Table (6): Effect of intercropping pattern, plant density and nitrogen application on light intensity (lux) at the bottom level of forage cowpea canopy in 2008.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	125.0	116.0	99.0	92.0	108.0	121.0	104.0	97.0	90.0	103.0
	Two plants	112.0	98.0	91.0	87.0	97.0	108.0	100.0	90.0	86.0	96.0
	Mean	118.5	107.0	95.0	89.5	102.5	114.5	102.0	93.5	88.0	99.5
Maize + Cowpea	One plant	105.0	91.0	84.0	80.0	90.0	96.0	88.0	84.0	80.0	87.0
	Two plants	97.0	88.0	80.3	75.0	85.0	93.0	84.0	80.0	71.0	82.0
	Mean	101.0	89.5	82.0	77.5	87.5	94.5	86.0	82.0	75.5	84.5
General mean	One plant	115.0	103.5	91.5	86.0	99.0	108.5	96.0	90.5	85.0	95.0
	Two plants	104.5	93.0	85.5	81.0	91.0	100.5	92.0	85.0	78.5	89.0
Overall mean		109.8	98.3	88.5	83.5	95.0	104.5	94.0	87.8	81.8	92.0
L.S.D. at 0.05 level:-											
Intercropping pattern (A)		2.9					1.9				
Plant density (B)		—					—				
Nitrogen levels (C)		2.9					2.5				
A x B		5.6					4.9				
A x C		4.2					3.6				
B x C		4.2					3.6				
A x B x C		5.9					5.1				

Table (7): Effect of intercropping pattern, plant density and nitrogen application on light intensity (lux) at the bottom level of forage cowpea canopy in 2009.

Intercropping pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	131.0	123.0	107.0	99.0	115.0	126.0	118.0	100.0	92.0	109.0
	Two plants	122.0	115.0	91.0	80.0	102.0	117.0	106.0	95.0	70.0	97.0
	Mean	126.5	119.0	99.0	89.5	108.5	121.5	112.0	97.5	81.0	103.0
Maize + Cowpea	One plant	101.0	95.0	82.0	74.0	88.0	97.0	90.0	81.0	72.0	85.0
	Two plants	96.0	89.0	77.0	70.0	83.0	92.0	85.0	73.0	70.0	80.0
	Mean	98.5	92.0	79.5	72.0	85.5	94.5	87.5	77.0	71.0	82.5
General mean	One plant	116.0	109.0	94.5	86.5	101.5	111.5	104.0	90.5	82.0	97.0
	Two plants	109.0	102.0	84.0	75.0	92.5	104.5	95.5	84.0	70.0	88.5
Overall mean		112.5	105.5	89.3	80.8	97.0	108.0	99.8	87.3	76.0	92.8
L.S.D. at 0.05 level:											
Intercropping pattern (A)		6.2					3.1				
Plant density (B)		—					—				
Nitrogen levels (C)		2.8					3.3				
A x B		4.2					1.7				
A x C		4.1					4.7				
B x C		4.1					4.7				
A x B x C		5.8					6.7				

value of light intensity was noted for cowpea intercropped with maize at two plants hill⁻¹ and fertilized by 120 kg N fed⁻¹.

3.1.2. The effect of light intensity on cowpea dry forage yield

Intercropping patterns significantly affected the dry forage yield of cowpea. The average dry forage yield (t fed⁻¹) of cowpea intercropped with maize and solid cowpea reached 1.0 and 1.1 t fed⁻¹ 1.4 and 1.6 t fed⁻¹ at the 1st and 2nd cuts, in the first season. Corresponding values for the second season were 0.8 and 1.0 t fed⁻¹ 1.1 and 1.3 t fed⁻¹ at the 1st and 2nd cut, respectively (Tables 8 and 9). It was clear that solid cowpea gave the highest dry forage yield because light intensity at plant tops of solid cowpea was greater by 3.4% and 2.3% compared to intercropped cowpea at the 1st and 2nd cuts in the first season. While, corresponding increases were 3.2% and 3.3% in the second season (Tables 2 and 3). Also, light intensity at the middle of the canopy increase for solid cowpea compared to intercropped cowpea by 8.3% and 6.8% at the 1st and 2nd cut in the first season, and by 12.1% and 11.2% for the same cuts in the second season (Tables 4 and 5). Moreover, bottom reading of light intensity increased by 14.6% and 15.1% for solid cowpea compared to intercropped cowpea at the 1st and 2nd cut, in the first season. Also, corresponding increases in the second season were 21.2% and 19.9% at the 1st and 2nd cut (Tables 6 and 7). Therefore, shading increased in cowpea intercropped with maize than solid cowpea. These results are in agreement with El-Zanaty, (2006) who reported that dry forage yield of cowpea in pure stands significantly surpassed its yield under intercropping in both seasons.

Two plants hill⁻¹ gave the highest dry forage yield compared with one plant hill⁻¹ at each cut in both seasons. However, the light intensity at one plant hill⁻¹ was lower than two plants hill⁻¹ at each cut for top, middle and bottom in both seasons. So it could be recommended with planting two plants hill⁻¹ when intercropped cowpea and maize for obtaining high dry forage yield of cowpea.

Dry forage yield was significantly affected by nitrogen levels below 90 and 120 kg N fed⁻¹ applied at the 2nd cut in both seasons. The average dry forage yield (t fed⁻¹) of cowpea supplied with zero to 120 kg N fed⁻¹ recorded 0.6 and 1.3 t fed⁻¹ 1.0 and 1.8 t fed⁻¹ at the 1st and 2nd cut in the first season. Also, it was 0.5 and 1.1 t fed⁻¹ 0.8 and 1.5 t fed⁻¹ in the second season for the same treatments. On the other hand, the control treatment (zero N) was superior than other

nitrogen levels (60, 90 and 120 kg N fed⁻¹) in light intensity at the bottom level. These increases were 10.5, 19.4 and 24.0% at the 1st cut, and 10.0, 16.0 and 21.7% at the 2nd cut in the first season. Also, corresponding increases in the second season were 6.2, 20.6 and 28.2% at the 1st cut, and 7.6, 19.2 and 29.6% at the 2nd cut. The interaction between intercropped cowpea with maize at two plants hill⁻¹ with the addition of 90 kg N fed⁻¹ gave 3.6 t fed⁻¹ for total dry forage yield, while solid cowpea in the same treatment was 4.1 t fed⁻¹ in the first season without significant effects between the two levels 90 and 120 kg N fed⁻¹ (Tables 8 and 9).

3.2. Intercropped maize

3.2.1. Effect of different treatments on light intensity

Light intensity (Lux) at the top, middle and bottom of intercropped maize plants before the 1st and 2nd cuts of cowpea was significantly affected by intercropping patterns and applied nitrogen levels in the first and second seasons.

3.2.1.1 Intercropping patterns

There were significant differences in light intensity at the top, middle and the bottom level of maize plants due to intercropping patterns before the 1st and 2nd cuts of cowpea in both seasons, except between intercropping two plants of cowpea hill⁻¹ with maize and solid maize at top plant before the 1st cut of cowpea in the first and second seasons and the 2nd cut of cowpea in the first season. These results are in agreement with those obtained by Metwally *et al.*, (2005) who mentioned that maize canopy architecture (spatial distribution of shoot organs) plays an important role in the amount of sunlight radiation intercepted by soybean under intercropping.

On the other hand, light intensity at the top of the plants of intercropped maize before the 2nd cut of cowpea was not significantly affected by intercropping patterns, except for intercropping one plant hill⁻¹ of cowpea with maize and solid maize in the first season, while, it was significant in the second season except between intercropping one plant hill⁻¹ of cowpea with maize and intercropping two plants hill⁻¹ cowpea of with maize in the second season.

In the first season, light intensity at the top of maize plants before the 1st cut of cowpea (one plant hill⁻¹) increased by 0.3% compared with solid maize. However, light intensity of solid maize increased by 0.1% compared to intercropping two plants hill⁻¹ of cowpea with maize (Tables 10 and 11).

In the second season at the 1st cut, solid maize

Table (8): Dry forage yield (t fed⁻¹) for 1st and 2nd cuts of forage cowpea as affected by intercropping pattern, plant density and nitrogen application in 2008.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	0.6	0.8	1.3	1.0	0.9	0.8	1.3	1.5	1.6	1.3
	Two plants	0.8	1.1	1.8	1.8	1.4	1.3	1.7	2.3	2.2	1.9
	Mean	0.7	1.0	1.5	1.4	1.1	1.1	1.5	1.9	1.9	1.6
Maize + Cowpea	One plant	0.5	0.7	1.0	0.9	0.8	0.7	0.9	1.4	1.5	1.1
	Two plants	0.7	1.2	1.6	1.4	1.2	1.1	1.5	2.0	1.9	1.6
	Mean	0.6	0.9	1.3	1.2	1.0	0.9	1.2	1.7	1.7	1.4
General mean	One plant	0.5	0.7	1.1	1.0	0.8	0.8	1.1	1.4	1.6	1.2
	Two plants	0.8	1.1	1.7	1.6	1.3	1.2	1.6	2.1	2.0	1.7
Overall mean		0.6	0.9	1.4	1.3	1.1	1.0	1.3	1.8	1.8	1.5
L.S.D. at 0.05 level:											
Intercropping pattern (A)		0.01					0.05				
Plant density (B)		---					---				
Nitrogen levels (C)		0.05					0.06				
A x B		0.05					0.10				
A x C		0.06					0.08				
B x C		0.06					0.08				
A x B x C		0.09					0.12				

Table (9): Dry forage yield (t fed⁻¹) for 1st and 2nd cuts of forage cowpea as affected by intercrop pattern, plant density and nitrogen application in 2009.

Intercropping Pattern	Plant density (Plant hill ⁻¹)	1 st cut					2 nd cut				
		Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
		Zero	60	90	120		Zero	60	90	120	
Solid Cowpea	One plant	0.5	0.7	1.1	0.9	0.8	0.7	1.0	1.2	1.3	1.1
	Two plants	0.7	0.9	1.5	1.5	1.1	1.1	1.4	1.9	1.8	1.6
	Mean	0.6	0.8	1.3	1.2	1.0	0.9	1.2	1.6	1.6	1.3
Maize + Cowpea	One plant	0.4	0.5	0.9	0.8	0.6	0.6	0.8	1.2	1.3	0.9
	Two plants	0.6	1.0	1.3	1.2	1.0	0.9	1.2	1.7	1.6	1.3
	Mean	0.5	0.8	1.1	1.0	0.8	0.7	1.0	1.4	1.4	1.1
General mean	One plant	0.4	0.6	1.0	0.8	0.7	0.6	0.9	1.2	1.3	1.0
	Two plants	0.6	0.9	1.4	1.3	1.1	1.0	1.3	1.8	1.7	1.5
Overall mean		0.5	0.8	1.2	1.1	0.9	0.8	1.1	1.5	1.5	1.2
L.S.D. at 0.05 level:											
Intercropping pattern (A)		0.01					0.04				
Plant density (B)		---					---				
Nitrogen levels (C)		0.04					0.05				
A x B		0.04					0.09				
A x C		0.05					0.07				
B x C		0.05					0.07				
A x B x C		0.08					0.10				

increased by 0.1% compared with intercropping one plant hill⁻¹ of cowpea with maize, and by 0.4% as compared with intercropping two plants hill⁻¹ of cowpea with maize (Tables 10 and 11).

On the other hand, light intensity of intercropped maize before the 2nd cut of cowpea increased for solid maize by 0.4% as compared with intercropping one plant hill⁻¹ of cowpea with maize, by 0.1% as compared with intercropping two plants hill⁻¹ of cowpea with maize in the first season. Also, it was increased by 0.2% and 0.3% for the same respective treatments in the second season (Tables 10 and 11).

Light intensity at the middle level of maize, before the 1st cut of cowpea in the first season was greater for solid maize than intercropped maize by 3.3% and 5.0% when intercropping one plant hill⁻¹ and two plants hill⁻¹ of cowpea with maize, respectively. Corresponding increases were 3.5% and 5.9% for the same treatments in respective order, in the second season. Light intensity increased by 2.5% and 4.5% for solid maize over maize intercropped at one plant hill⁻¹ and two plants hill⁻¹ of cowpea before the 2nd cut in the first season, and by 4.1% and 6.1% for the same treatments in the same order in the second season (Tables 12 and 13).

On the other hand, bottom reading of light intensity increased by 6.1% and 12.1% for solid maize compared to maize intercropped with one and two plants hill⁻¹ of cowpea before the 1st cut of cowpea in the first season, and was increased by 8.0% and 10.7% for the same treatments in the second season.

Light intensity readings at the bottom of maize canopy before the 2nd cut of cowpea was greater by 6.8% and 12.3% for solid maize compared to intercropping one plant and two plants hill⁻¹ of cowpea with maize in the first season. Corresponding increases were 7.6% and 10.0% in the second season (Tables 14 and 15).

3.2.1.2. Nitrogen levels

The addition of nitrogen significantly decreased light intensity at the top, middle and bottom levels of maize before the 1st and 2nd cuts of cowpea in both seasons, except between the two levels 90 and 120 kg N fed⁻¹ at top and middle reading before the 1st and 2nd cuts of cowpea in the first season, respectively.

Zero fertilizer treatment had the highest value of light intensity at the top of maize plants compared with the applied N levels (60, 90 and 120 kg N fed⁻¹), reaching 0.5, 0.9 and 1.1% before the 1st cut of cowpea, and 0.5, 0.8 and 1.2% before

the 2nd cut of cowpea in the first season. Corresponding values in the second season were 0.3, 0.7 and 1.2% before the 1st cut of cowpea, and 0.4, 0.8 and 1.6% before the 2nd cut of cowpea, respectively (Tables 10 and 11).

On the other hand, at the middle level of maize plants, zero fertilizer treatments showed the highest value of light intensity compared with the 60, 90 and 120 kg N fed⁻¹ treatments, these increases in light intensity were 1.6, 3.0 and 4.1% before the 1st cut, and 1.4, 2.9 and 3.3% before the 2nd cut of cowpea in the first season. Corresponding increases were 1.8, 3.5 and 5.0% before the 1st cut of cowpea, and 1.7, 3.2 and 4.0% before the 2nd cut of cowpea, in the second season respectively (Tables 12 and 13).

Zero N fertilizer treatment was superior in light intensity to nitrogen added levels at the 60, 90 and 120 kg N fed⁻¹ at the bottom of maize plant, showing increases of 7.0, 11.6 and 16.7% before the 1st cut of cowpea, and 4.7, 8.0 and 12.7% before the 2nd cut in the first season. Corresponding increases in the second season were 6.4, 10.2 and 15.5% before the 1st cut, and 5.2, 9.3 and 13.8% before the 2nd cut of cowpea in the second season, respectively (Tables 14 and 15).

The application of nitrogen up to 120 kg N fed⁻¹ caused increased reduction in light intensity. As a result, grain yield and dry stover yield were increased.

3.2.1.3. Effect of the interaction between intercropping patterns and N levels

The interaction effect of intercropping patterns and nitrogen levels on light intensity under maize, indicate that the highest light intensity was obtained by solid maize and zero N fed⁻¹ while the lowest value was obtained under intercropping two plants hill⁻¹ of cowpea with maize fertilized with 120 kg N fed⁻¹.

3.2.2. The relationship between light intensity and maize grain yield

Intercropping patterns significantly affected grain yield of maize except when maize was intercropped with two plants hill⁻¹ of cowpea and solid maize in both seasons. At the same time light intensity at the top of maize plants before the 1st cut of cowpea (one plant hill⁻¹) was higher by 0.3% compared with solid maize. While, light intensity of solid maize increased by 0.1% compared to intercropping two plants hill⁻¹ of cowpea with maize in the first season (Table 10).

These results are in agreement with those obtained by Metwally *et al.* (2009) and disagree with Searle *et al.* (1981).

Table (10): Effect of intercropping pattern and nitrogen levels on light intensity (lux) at the top of maize plants before the first cut of forage cowpea in 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	1593.0	1583.0	1578.0	1574.0	1582.0	1603.0	1596.0	1592.0	1581.0	1593.0
Maize + Cowpea (one plant hill ⁻¹)	1595.0	1590.0	1583.0	1580.0	1587.0	1598.0	1593.0	1590.0	1583.0	1591.0
Maize + Cowpea (two plants hill ⁻¹)	1591.0	1580.0	1576.0	1573.0	1580.0	1596.0	1592.0	1581.0	1575.0	1586.0
Mean	1593.0	1584.3	1579.0	1575.7	1583.0	1599.0	1593.7	1587.7	1579.7	1590.0
L.S.D. at 0.05 level:										
Intercropping pattern (A)	4.1					8.7				
Nitrogen levels (B)	3.6					3.5				
A x B	6.2					6.1				

Table (11): Effect of intercropping patterns and nitrogen levels on light intensity (lux) at the top of maize plants before the second cut of forage cowpea 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	1592.0	1583.0	1578.0	1567.0	1580.0	1600.0	1595.0	1588.0	1569.0	1588.0
Maize + Cowpea (one plant hill ⁻¹)	1581.0	1576.0	1573.0	1566.0	1574.0	1597.0	1590.0	1584.0	1569.0	1585.0
Maize + Cowpea (two plants hill ⁻¹)	1588.0	1580.0	1574.0	1570.0	1578.0	1591.0	1586.0	1580.0	1575.0	1583.0
Mean	1587.0	1579.0	1575.0	1567.7	1577.3	1596.0	1590.3	1584.0	1571.0	1585.3
L.S.D. at 0.05 level:										
Intercropping pattern (A)	4.5					2.5				
Nitrogen levels (B)	4.3					4.7				
A x B	7.4					8.1				

Table (12): Effect of intercropping pattern and nitrogen levels on light intensity (lux) at the middle of maize plants before the 1st cut of forage cowpea 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	985.0	971.0	954.0	946.0	964.0	987.0	968.0	958.0	931.0	961.0
Maize + Cowpea (one plant hill ⁻¹)	953.0	936.0	924.0	915.0	932.0	958.0	936.0	912.0	902.0	927.0
Maize + Cowpea (two plants hill ⁻¹)	937.0	921.0	910.0	896.0	916.0	921.0	910.0	896.0	889.0	904.0
Mean	958.3	942.7	929.3	919.0	937.3	955.3	938.0	922.0	907.3	930.7
L.S.D. at 0.05 level:										
Intercropping pattern (A)	5.1					3.6				
Nitrogen levels (B)	3.5					4.4				
A x B	6.1					7.6				

Table (13): Effect of intercropping pattern and nitrogen levels on light intensity (lux) at the middle of maize plants before the 2nd cut of forage cowpea during 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	968.0	955.0	941.0	936.0	950.0	973.0	950.0	941.0	936.0	950.0
Maize + Cowpea (one plant hill ⁻¹)	942.0	931.0	917.0	914.0	926.0	933.0	921.0	900.0	890.0	911.0
Maize + Cowpea (two plants hill ⁻¹)	927.0	910.0	897.0	894.0	907.0	910.0	896.0	884.0	878.0	892.0
Mean	945.7	932.0	918.3	914.7	927.7	938.7	922.3	908.3	901.3	917.7
L.S.D. at 0.05 level:										
Intercropping pattern (A)					3.4					6.9
Nitrogen levels (B)					3.8					4.1
A x B					6.5					7.2

Table (14): Effect of intercropping pattern and nitrogen levels on light intensity (lux) at the bottom of maize plants before the 1st cut of forage cowpea during 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	260.0	237.0	220.0	207.0	231.0	246.0	231.0	218.0	201.0	224.0
Maize + Cowpea (one plant hill ⁻¹)	235.0	220.0	211.0	202.0	217.0	222.0	206.0	201.0	195.0	206.0
Maize + Cowpea (two plants hill ⁻¹)	219.0	207.0	200.0	186.0	203.0	217.0	204.0	196.0	183.0	200.0
Mean	238.0	221.3	210.3	198.3	217.0	228.3	213.7	205.0	193.0	210.0
L.S.D. at 0.05 level:										
Intercropping pattern (A)					3.5					3.8
Nitrogen levels (B)					5.3					3.1
A x B					9.1					5.4

Table (15): Effect of intercropping pattern and nitrogen levels on light intensity (lux) at the bottom of maize plants before the 2nd cut of forage cowpea during 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	239.0	227.0	219.0	195.0	220.0	221.0	213.0	210.0	196.0	210.0
Maize + Cowpea (one plant hill ⁻¹)	220.0	208.0	198.0	194.0	205.0	209.0	196.0	189.0	182.0	194.0
Maize + Cowpea (two plants hill ⁻¹)	201.0	194.0	190.0	187.0	193.0	208.0	196.0	180.0	172.0	189.0
Mean	220.0	209.7	202.3	192.0	206.0	212.7	201.7	193.0	183.3	197.7
L.S.D. at 0.05 level:										
Intercropping pattern (A)					1.9					4.8
Nitrogen levels (B)					5.5					3.9
A x B					9.6					6.9

Table (16): Grain yield (t fed⁻¹) of maize as affected by intercropping pattern and nitrogen levels during 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	3.0	3.9	4.1	4.2	3.8	2.5	3.2	3.4	3.5	3.2
Maize + Cowpea (one plant hill ⁻¹)	3.3	3.9	4.2	4.3	3.9	2.8	3.3	3.5	3.6	3.3
Maize + Cowpea (two plants hill ⁻¹)	3.2	3.9	4.0	4.2	3.8	2.6	3.2	3.3	3.5	3.2
Mean	3.2	3.9	4.1	4.2	3.8	2.6	3.2	3.4	3.5	3.2
L.S.D. at 0.05 level:										
Intercropping pattern (A)	0.1					0.1				
Nitrogen levels (B)	0.1					0.1				
A x B	0.2					0.2				

Table (17): Effect of intercropping patterns and nitrogen levels on stover dry yield (t fed⁻¹) of maize during 2008 and 2009.

Intercropping Pattern	2008					2009				
	Nitrogen levels (kg fed ⁻¹)				Mean	Nitrogen levels (kg fed ⁻¹)				Mean
	Zero	60	90	120		Zero	60	90	120	
Solid Maize	2.1	2.7	3.4	3.8	3.0	1.8	2.2	2.9	3.1	2.5
Maize + Cowpea (one plant hill ⁻¹)	2.4	3.0	3.7	4.2	3.3	1.9	2.5	3.1	3.5	2.7
Maize + Cowpea (two plants hill ⁻¹)	2.8	3.5	4.6	4.5	3.8	2.3	2.9	3.7	3.8	3.2
Mean	2.4	3.1	3.9	4.2	3.4	2.0	2.5	3.2	3.5	2.8
L.S.D. at 0.05 level:										
Intercropping pattern(A)	0.2					0.2				
Nitrogen levels (B)	0.2					0.2				
A x B	0.3					0.3				

Nitrogen application significantly affected grain yield of maize when applied at levels below 90 and 120 kg N fed⁻¹ in both seasons.

The data indicate that grain yield of maize increased with increasing nitrogen fertilizer levels from zero to 120 kg N fed⁻¹ recording 3.2 to 4.2 t fed⁻¹ in the first season and 2.6 to 3.5 t fed⁻¹ in the second season (Table 16). Nitrogen levels significantly decreased light intensity at the top, middle and bottom levels of maize canopy before the 1st and 2nd cuts of cowpea in both seasons, except between the two levels of 90 and 120 kg N fed⁻¹ for the top and middle readings before the 1st and 2nd cuts of cowpea in the first season. The same trend was obtained by (Searle *et al.*, 1981; Baker and Blamey, 1985; Rana *et al.*, 2001 and Safina, 2007).

The interaction between intercropping patterns and nitrogen levels was most favorable when one plant hill⁻¹ of cowpea was intercropped with maize and 120 kg N fed⁻¹ in both seasons, recording 4.3 t fed⁻¹ in the first season and 3.6 t fed⁻¹ in the second season (Table 16). These results are in agreement with those given by Metwally, (1973). On the other hand, the lowest value was obtained by solid maize and zero nitrogen level in both seasons. It was recorded 3.0 t fed⁻¹ in the first season and 2.5 t fed⁻¹ in the second one (Table 16). These results are in line with Safina, (2007).

3.2.3. The relationship between light intensity and dry stover yield

Significant differences between intercropping patterns for dry stover yield (t fed⁻¹) in both seasons, except between the intercropping of one plant hill⁻¹ of cowpea with maize, as well as solid maize in the second season. These results are in agreement with those given by Metwally *et al.* (2009) on the effect of cropping systems and maize varieties on maize intercropped with soybean. The workers found that the cropping system had significant effects on maize dry weight.

The highest dry weight was obtained by intercropping two cowpea plants hill⁻¹ with maize in both seasons.

Nitrogen fertilization significantly increased dry maize stover yield in the first and second seasons and reduced light intensity.

Dry stover yield of maize supplied with zero to 120 kg N fed⁻¹ ranged from 2.4 to 4.2 t fed⁻¹ in the first season, 2.0 to 3.5 t fed⁻¹ in the second one. The greater dry stover yield average was obtained by intercropping two cowpea plants

hill⁻¹ with maize fertilized with 120 kg N fed⁻¹ in both seasons (Table 17).

4. REFERENCES

- Abdel-Wahab Sh.I. (2010). Growing soybean with maize under different intercropping patterns and high population densities. Ph.D. Thesis, Fac. Agric. Cairo Univ. Egypt, 189 p.
- Abou Keriasha M.A., Abd El-Hady M.A. and Nawar F.R. (2009). Response of some cowpea varieties to intercropping in maize fields under upper Egypt conditions. Egypt. J. of Appl. Sci., 24(2B):495-514.
- Badr M. M. (1998). Biological efficiency of different intercropping patterns of soybean with maize. Ph. D. Thesis, Fac. Agric., Mansoura Univ., Egypt, 102 p.
- Baker C.M. and Blamey F.P.C. (1985). Nitrogen fertilizer effects on yield and nitrogen uptake of sorghum and soybean, grown in sole cropping and intercropping systems. Field Crops Research, 12: 233-240.
- Bowes G., Ogren W.L. and Hageman R.H. (1970). Light saturation photosynthetic rate, RuDP carboxylase: activity and specific leaf weight in soybeans grown under different light intensities. Crop Sci., 12: 77-79.
- El-Zanaty Rafea I.A. (2006). Intercropping maize with cowpea. 1- Effect of intercropping patterns on forage yield, quality and seed yield of cowpea. Egypt. J. of Appl. Sci., 21(6B):523-534.
- Freed R (2005). MSTAT-C Crop and Soil Sciences. Michigan State University, MI, USA.
- Li L., Zhang F.S., Li X.L., Christie P., Sun J.H., Yang S.C. and Tang C. (2003). Inter specific facilitation of nutrient uptake by intercropped maize and faba bean. Nutr. Cycling Agro. Eco., 68:61-71.
- Marsalis M. and Angadi S. (2009). Can intercropping legumes with sorghum improve sustainability? Past - articles & Itemid, November, 2009.
- Metwally A.A. (1973). The study of some companionship between maize and other crops. M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, 151 p.
- Metwally A. A., Mohamed Gmalat O., Sherief M. N. and Abo-Hegazy D. R. (2009). Yield and land equivalent ratio of intercropped maize and soybean under three intercropping patterns. The 4th Conf., Recent Technologies in Agric., 3-5 Nov., Cairo Univ., 2: 284-290.

- Metwally A. A., Shafik M.M., El Morshedy W.A. and Aly H.R. (2005). Yield and land equivalent ratio of intercropped maize and soybean. Proc. 1st Sci. Conf. Cereal Crops, Alex., : 113-120.
- Ofori F. and Stern W.R. (1987). Cereal-legume intercropping systems. Advance in Agronomy. 41: 41-90.
- Rana R.S., Bhupinder S., Negi S.C. and Singh B. (2001). Management of maize/legume intercropping under mid-hill sub-humid conditions. Indian-Journal-of-Agricultural-Research. 35(2): 100-103.
- Safina S.A. (2007). Effect of biofertilization and mineral nitrogen applications on solid and intercropping maize with soybean. Ph.D. Thesis, Fac. Agric., Cairo Univ., Egypt, 151 p.
- Searle P.G.E., Comudom Y., Shedden D.C. and Nance R.A. (1981). Effect of maize + legume intercropping systems and fertilizer nitrogen on crop yielding and residual nitrogen. Field Crops Research, 4: 133-145.

العلاقة بين شدة الاضاعة وحاصل العلف للجاف للوبيا العلف والذرة الشامية تحت ظروف التحميل

قرنى اسماعيل عبد الجواد- عبد الحفيظ عبد الرحمن عبد الحفيظ - وفاء محمد شعراوى * - ولاء محمد النبوى *

قسم المحاصيل - كلية الزراعة - جامعة القاهرة.

* قسم بحوث محاصيل العلف - معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية - الجيزة - مصر

ملخص

أجريت تجربتان حقليتان بمحطة التجارب والبحوث الزراعية بجامعة القاهرة بالجيزة خلال موسمي ٢٠٠٨ ، ٢٠٠٩ بهدف دراسة تأثير نظم التحميل وكثافة النباتات ومستويات التسميد الأزوتى على العلاقة بين شدة الاضاعة وحاصل العلف الجاف للوبيا العلف المحملة على الذرة الشامية، زرعت للوبيا على ريشه وزرعت الذرة على الريشة المقابلة من الخط. استخدم في الدراسة صنف لوبيا العلف Buff وهجين فردى ١٢٢ من الذرة الشامية وهو صنف هجين قصير الطول تظل أوراقه خضراء بعد نضج الحبوب. تمت الزراعة في بداية شهر يوليو في كلا الموسمين وذلك باستخدام كثافتين للوبيا العلف (٣٠ ألف، ٦٠ ألف نبات/فدان) بينما زرعت الذرة بالكثافة الموصى بها. استخدم تصميم القطاعات المنشقة مرتين في ثلاثة مكررات، حيث خصصت القطع الرئيسية لنظم الزراعة (لوبيا العلف محملة على الذرة الشامية ، ذرة شامية منفردة، لوبيا منفردة) والقطع المنشقة الأولى للكثافة النباتية للوبيا العلف (نبات، نباتين/ بالجورة مع ترك ٢٠مسم بين الجور) وتضمنت القطع المنشقة الثانية أربعة مستويات من التسميد الأزوتى هي صفر، ٦٠ ، ٩٠ و ١٢٠ كجم/ن/فدان اضيفت على دفعتين متساويتين، الأولى بعد ٢١ يوم والثانية بعد ٦٠ يوم من الزراعة (بعد الحشة الأولى للوبيا العلف). أظهرت النتائج وجود اختلافات معنوية بين تأثير نظم التحميل ومستويات الكثافة النباتية ومستويات التسميد الأزوتى على شدة الاضاعة في أعلى ومنتصف وأسفل نباتات اللوبيا خلال موسمي الزراعة. كما توقفت الزراعة المنفردة للذرة على الزراعة المحملة في شدة الاضاعة، كما لوحظت زيادة معنوية في شدة الاضاعة في الجزء العلوى من النباتات في الزراعة المحملة مقارنة بشدتها في الجزء الوسطى، والجزء السفلى من النباتات. وكانت شدة الاضاعة في الجزء الوسطى من النباتات في الزراعة المنفردة أكبر منها في الزراعة المحملة، كما توقفت شدة الاضاعة عند زراعة نبات واحد في الجورة مقارنة بزراعة نباتين بالجورة ولكن كانت الفروق بين هذه الزيادة منخفضة بين الكثافتين النباتيتين وعليه يمكن التوصية بزراعة نباتين بالجورة من اللوبيا عند تحميل لوبيا العلف مع الذرة الشامية لتعظيم حاصل العلف الأخضر من اللوبيا. كما أدت اضافة التسميد الأزوتى الى انخفاض شدة الاضاعة وزيادة صفات النمو الخضري وحاصل العلف الأخضر والجاف. وقد أعطت المعاملات غير المسمدة بالأزوت أعلى القيم في شدة الاضاعة مقارنة بالمعاملات المسمدة. أدى للتفاعل بين المعاملات الى تأثير معنوى على شدة الاضاعة وتم الحصول على أعلى شدة اضااعة في حالة الزراعة المنفردة لنبات واحد بالجورة مع عدم اضافة السماد الأزوتى بينما انخفضت هذه القيم مع زيادة معدل التسميد حتى ١٢٠ كجم/ن/فدان مع ترك نباتين بالجورة. كما وجدت اختلافات معنوية بين نظم التحميل في شدة الاضاعة أعلى ومنتصف وأسفل نباتات الذرة الشامية في كلا الموسمين فيما عدا حالة زراعة نباتين في الجورة مع التحميل والزراعة المنفردة للذرة وأدت اضافة السماد الأزوتى الى انخفاض معنوى في شدة الاضاعة في جميع ارتفاعات النباتات ولكن دون فروق معنوية بين مستوى ٩٠ و ١٢٠ كجم/ن/فدان ولكنها أدت الى زيادة حاصل الحبوب ووزن الحطب الجاف، كما أظهرت المعاملات غير المسمدة بالأزوت أعلى شدة اضااعة، و كان تأثير التفاعل بين نظم التحميل ومستويات التسميد الأزوتى على شدة الاضاعة معنويا في كل مستويات القراءة وأعطت الزراعة المنفردة للذرة مع عدم اضافة السماد الأزوتى أعلى القيم بينما أعطت الزراعة المحملة نباتين في الجورة مع اضافة ١٢٠ كجم/ن/فدان أقل القيم.