

EFFECT OF SUPPLEMENTED DIFFERENT LEVELS OF VITAMINS E AND C TO LAYERS HEN DIETS ON: 1- PRODUCTIVE AND REPRODUCTIVE PARAMETERS.

Mona S. Ragab¹; S. F. Youssef² and Kout El-Kloub M. El. Mostafa²

¹*Poultry Production Department, Faculty of Agriculture, Fayoum University, Egypt.*

²*Animal Production Institute, Agriculture Research Center, Ministry of Agriculture, Dokki, Giza, Egypt.*

(Received 25/10/2011, Accepted 23/12/2011)

SUMMARY

This study was carried out at the Poultry Research Station, El-Azab, Fayoum, to study the effects of two dietary levels of vitamin E (Vit. E) (10 or 20 mg/Kg diet), vitamin C (Vit. C) (200 or 400 mg/Kg diet) and their mixtures on egg performance, egg quality, fertility, hatchability and economic efficiency of El-Salam laying hens. A total number of 243 (216 breeder hens and 27 cocks) birds at 25 weeks of age were used in this experiment. Birds were wing banded and randomly distributed into 9 equal treatment groups of 27 birds each (24 breeder hen and 3 cock each). Each group was equally subdivided into three replicates of 11 (eight ♀ and one ♂/replicate) birds each. The experimental treatments were as follows:

- 1- Birds were fed control diet (unsupplemented with Vit. E or Vit. C (D1)).
- 2- Birds were fed D1 supplemented with 10 mg/Kg diet Vit. E (D2).
- 3- Birds were fed D1 supplemented with 20 mg/Kg diet Vit. E (D3).
- 4- Birds were fed D1 supplemented with 200 mg/Kg diet Vit. C (D4).
- 5- Birds were fed D1 supplemented with 400 mg/Kg diet Vit. C (D5).
- 6- Birds were fed D1 supplemented with Vit. E 10 mg/Kg diet + Vit. C 200 mg/Kg diet (D6).
- 7- Birds were fed D1 supplemented with Vit. E 10 mg/Kg diet + Vit. C 400 mg/Kg diet (D7).
- 8- Birds were fed D1 supplemented with Vit. E 20 mg/Kg diet + Vit. C 200 mg/Kg diet (D8).
- 9- Birds were fed D1 supplemented with Vit. E 20 mg/Kg diet + Vit. C 400 mg/Kg diet (D9).

Results obtained could be summarized in the following:

Laying hens fed diet containing mixtures of Vit. E and C supplementation had lower feed intake value, while, laying hens fed control diet had higher feed intake value. No significant effect were observed for average egg weight, total egg mass, egg production% and feed conversion during the experimental period. Laying hens fed diet containing 10 mg/kg diet Vit. E had higher shell thickness and shell% while, those fed diet containing 20 mg Vit. E and 400 mg/kg diet Vit. C had lower shell thickness and shell% during the experimental period. Laying hens fed diet containing 20 mg Vit. E and 400 mg/kg diet Vit. C had higher fertility%. El-Salam laying hens fed diet containing 20 mg Vit. E and 200 mg/kg diet Vit. C gave the best economical and relative efficiency values. In conclusion: Feeding El-Salam laying hens on diets containing 20 mg Vit. E and 200 mg/kg diet Vit. C improved the productive, reproductive performance and relative economic efficiency values.

Keywords: *El-Salam laying hen; productive; reproductive performance.*

INTRODUCTION

Vitamin E (Vit. E) plays important roles in various biochemical and physiological processes, including antioxidation and signaling transduction (Brigelius-Flohe *et al.*, 2002 and Ricciarelli *et al.*, 2002). The addition of α -tocopherol to hen diets increases the content of Vit. E in the egg yolk in a dose dependent manner (Jiang *et al.*, 1994; Surai *et al.*, 1997 and Meluzzi *et al.*, 2000). Tocopherols may also provide health benefits mainly in preventing cancer and coronary diseases (Diplock, 1991 and Knekt *et al.*, 1991) so that the incorporation of Vit. E to the egg may both increase the oxidative stability and provide a source of tocopherols useful for human nutrition and health.

As poultry fowls are able to synthesize vitamin C (Vit. C) and it is not transferred into the egg. The research attention was mainly focused on the effects of the vitamin on improving the egg-shell quality (Pardue and Thaxton, 1986). Even if Vit. C is neither contained nor transferred to the egg, it could play its

antioxidative role in regenerating Vit. E in laying hens. Furthermore it is well known that ascorbic acid (AA) is a water-soluble vitamin) is required for the hydroxylation of proline residues necessary for the synthesis of pro-collagen (Weiser *et al.*, 1990), also it could be involved in the synthesis of egg proteins. Plasma protein concentration was insignificantly increased with Vit. C (El-Badry *et al.*, 2011), the beneficial effect of Vit. C supplementation on plasma protein could be attributed to Vit. C which work as coenzyme playing an important role in the metabolism of amino acid (Kutlu and Forbes, 1993) while, Rice (2000) reported that Vit. C is effective as antioxidant and it play an important role in metabolic activity.

Vitamin C has been demonstrated to enhance antioxidant activity of Vit. E by reducing the tocopheroxyl radicals back to their active form of Vit. E (Packer, 1992 and Jacob, 1995) or by sparing available Vit. E (Retsky and Frei, 1995). Some reports indicated improvements in the egg production and egg shell quality by Vit. C (Bell and Marion, 1990; Balnave and Muheereza, 1997 and Al-Shoquiry, 1999). As well as by adding Vit. E (Bollengier *et al.*, 1999 and Sahin *et al.*, 2002). On the other hand, Vit. E serves as a physiological antioxidant through inactivation free radicals, improves egg production, feed intake, egg yolk and albumen solids (Kirunda *et al.*, 2001), and improves egg quality (Puthongsiriporn, 1998).

Therefore, the objective of this study was to determine the effects of two dietary levels of Vit. E (10 or 20 mg/Kg diet), Vit. C (200 or 400 mg/Kg diet) separately and their mixtures on egg performance, egg quality, fertility, hatchability, embryonic mortality and economic efficiency of El-Salam laying hens.

MATERIALS AND METHODS

This study was carried out at the Poultry Research Station, El-Azab, Fayoum, to study the effects of two dietary levels of vitamin E (Vit. E) (10 or 20 mg/Kg diet), vitamin C (Vit. C) (200 or 400 mg/Kg diet) and their mixtures on egg performance, egg quality, fertility, hatchability, embryonic mortality and economic efficiency of El-Salam laying hens.

A total number of 243 (216 breeder hens and 27 cocks) birds at 25 weeks of age were used in this experiment. Birds were wing banded and randomly distributed into 9 equal treatment groups of 27 birds (having nearly similar body weight) each (24 breeder hen and 3 cock each). Each group was equally subdivided into three replicates of 11 (eight ♀ and one ♂/replicate) birds each. Birds were reared under the same management conditions in egg production batteries (open system). The experimental period was lasted for 14 weeks from 25 to 39 weeks of age. Treatment groups were fed a commercial layer ration (16% CP and 2703.34 Kcal ME/Kg diet, Table 1), (control group) supplemented with 10 or 20 mg/Kg diet α -tocopherol acetate (Vit. E), 200 or 400 mg/Kg diet of L-ascorbic acid (Vit. C) and their mixtures. Artificial light was used beside the normal day light to provide 16-hour day photoperiod. Feed and water were provided *ad libitum*.

The experimental treatments were as follows:

- 1- Birds were fed control diet (unsupplemented with Vit. E or Vit. C (D1)).
- 2- Birds were fed D1 supplemented with 10 mg/Kg diet Vit. E (D2).
- 3- Birds were fed D1 supplemented with 20 mg/Kg diet Vit. E (D3).
- 4- Birds were fed D1 supplemented with 200 mg/Kg diet Vit. C (D4).
- 5- Birds were fed D1 supplemented with 400 mg/Kg diet Vit. C (D5).
- 6- Birds were fed D1 supplemented with Vit. E 10 mg/Kg diet + Vit. C 200 mg/Kg diet (D6).
- 7- Birds were fed D1 supplemented with Vit. E 10 mg/Kg diet + Vit. C 400 mg/Kg diet (D7).
- 8- Birds were fed D1 supplemented with Vit. E 20 mg/Kg diet + Vit. C 200 mg/Kg diet (D8).
- 9- Birds were fed D1 supplemented with Vit. E 20 mg/Kg diet + Vit. C 400 mg/Kg diet (D9).

Egg production (weight and number) was recorded daily and feed intake for each group was calculated weekly and feed conversion calculated as the amount of feed required for producing a unit of egg mass.

Table (1): Composition of the basal diets.

Item	%
Yellow corn, ground	63.50
Soybean meal (44%CP)	24.57
Wheat bran	2.00
Calcium carbonate	7.77
Sodium chloride	0.30
Vit. and Min. premix ¹	0.30
Di-calcium phosphate	1.50
DL-Methionine	0.06
Total	100.0
<i>Calculated analysis %²:</i>	
Crude protein	16.56
Ether extract	2.67
Crude fiber	3.34
Calcium	3.37
Available phosphorus	0.39
Methionine	0.33
Methionine+Cystine	0.61
Lysine	0.84
ME, kcal./Kg	2703
Cost (£.E./ton) ³	2600.0

¹ Each 3.0 Kg of the Vit. and Min. premix contains: Vit. A, 10000000 IU; Vit. D₃ 2000000 IU; Vit. E, 1000 mg; Vit. K₃, 1000 mg; Vit. B1, 1000 mg; Vit. B2, 500 mg; Vit. B6, 1500 mg; Vit. B12, 10 mg; biotin, 50 mg; folic acid, 1 mg; niacin, 3000 mg; Ca pantothenate, 1000 mg; Zn, 50 g; Cu, 4 g; Fe, 30 g; Co, 0.1 g; Se, 0.1 g; I, 0.3 g; Mn, 60 g and anti-oxidant, 10 g. and complete to 3.0 Kg by calcium carbonate.

² According to NRC, 1994.

³ According to the local market price at the experimental time.

Mortality was recorded daily (no mortality of birds were recorded during the study period). Egg quality measurements were determined monthly on eggs of the last three days. Representative egg samples from each treatment were collected monthly throughout the experimental period in order to determine egg and shell quality. Egg shell thickness, including shell membranes, was measured using a micrometer at three locations on the egg (air cell, equator, and sharp end). Haugh unit score was applied from a special chart using egg weight and albumen height which was measured by using a micrometer according to Haugh (1937). Shell surface area (Carter, 1975), shell weight per unit surface area (Hamilton, 1978), egg shape index% (Carter, 1968) and yolk index% (Well, 1968) were calculated. The egg yolk visual color score was determined by matching the yolk with one of the 15 bands of the "1961, Roche Improved Yolk Color Fan". Two batches of eggs (130 egg/treatment) were collected from the 9 treatments at the 38th and 39th weeks of age to study the hatchability and incubated at Chick Master Hatchery. Fertility was determined by candling at 7 days of incubated period. The averages of the fertility and hatchability of the two batches were calculated.

Economical efficiency of egg production was calculated from the input-output analysis which was calculated according to the price of the experimental diets and eggs produced. The values of economical efficiency were calculated as the net revenue per unit of total cost.

An ANOVA with the General Linear Models (GLM) procedure of SPSS software (SPSS, 1999) included the effect of type and treatment means. Treatment means indicating significant differences ($P \leq 0.01$ and $P \leq 0.05$) were tested using Duncan's multiple range test (Duncan, 1955).

RESULTS AND DISCUSSION

Laying hens productive performance:

Effect of supplementing laying hens diets with Vit. E; C and their mixtures on average egg weight (EW), total egg mass (EM), egg production (EP %), daily feed intake (FI) and feed conversion (FC) are shown in Table 2.

As shown in Table 2, type of addition effect was significant only for FI (Table 2), it is clear that laying hens fed diet containing mixtures of Vit. E and C supplementation had lower FI value, while, laying hens fed control diet had higher FI value. No significant effects were observed for EW, EM, EP% and FC during the experimental period. Numerically, laying hens fed diet containing mixtures of Vit. E and C had higher EW value, while, those fed diet containing Vit. C supplementation had higher EM, EP% and the best FC value (the difference is not significant).

Table (2): Effects of supplementing laying hens diets with vitamin E, C and their mixtures on egg production of El-Salam laying hens.

Item	Average egg weight (EW,g)	Total egg mass (EM,g)	Egg production (EP)%	Daily feed intake (FI,g)	Feed conversion (FC)
Type of addition					
Control	44.58	22765.3	68.66	106.94 ^A	3.51
Vitamin (Vit.) E	44.55	22672.6	68.39	104.21 ^B	3.44
Vitamin C	45.13	23572.3	70.06	103.36 ^{BC}	3.29
Mixed (Vit. E & Vit. C)	46.18	23365.3	68.06	102.79 ^C	3.30
±SEM ¹	0.60	858.4	2.33	0.29	0.13
Treatments					
Control	44.58	22765.3	68.66	106.94 ^A	3.51
Vit. E 10 mg/Kg diet	44.68	22367.9	67.25	104.73 ^B	3.51
Vit. E 20 mg/Kg diet	44.42	22977.4	69.53	103.70 ^D	3.38
Vit. C 200 mg/Kg diet	45.03	23187.6	69.20	103.21 ^F	3.31
Vit. C 400 mg/Kg diet	45.23	23957.1	70.92	103.50 ^E	3.26
Vit. E 10 mg/Kg diet + Vit. C 200 mg/Kg diet	45.71	22399.9	65.82	104.23 ^C	3.52
Vit. E 10 mg/Kg diet + Vit. C 400 mg/Kg diet	45.79	22820.2	67.25	101.77 ^H	3.34
Vit. E 20 mg/Kg diet + Vit. C 200 mg/Kg diet	47.00	24301.7	69.53	102.60 ^G	3.15
Vit. E 20 mg/Kg diet + Vit. C 400 mg /Kg diet	46.22	23939.3	69.62	102.55 ^G	3.19
±SEM	0.93	1311.9	3.61	0.06	0.20

¹Pooled SEM

A... G. values in the same column within the same item followed by different superscripts are significantly different (at $P \leq 0.01$ for A to G).

There were insignificant differences among all dietary treatments in productive performance except, FI. It is clear that laying hens fed diet containing 20 mg Vit. E and 400 mg/kg diet Vit. C had lower FI whereas, those fed control diet had higher FI during the experimental period. Numerically, laying hens fed diet containing 20 mg Vit. E and 200 mg Vit. C/kg diet had higher EW, EM and the best FC value while, those fed diet containing 400 mg Vit. C/kg diet had higher EP% (difference is not significant) during the experimental period (Table 2).

These results are in harmony with some experimental results who demonstrated that egg weight was unaffected by a dietary treatment (Gebert *et al.*, 1998; Meluzzi *et al.*, 2000 and Franchini, *et al.*, 2002). Similarly, Puthongsiriporn *et al.* (2001) reported that supplemental Vit. E did not affect egg production of White Leghorn hens. While, these results disagree with those of Ajuyah *et al.* (1993) and Scheideler and Froning (1996) reported that laying hens fed diets with high levels of Vit. E (50 IU/kg diet) greatly improved egg production compared with that of laying hens fed the same diets with low levels of Vit. E (27 IU/kg diet). Also, Whitehead *et al.* (1998) reported that dietary Vit. E at 250 mg/kg provided for optimum egg production compared to 10 mg/kg fed to control hens by Vit. E promoting the release of vitellogenin from liver by protecting cell membranes of hepatocytes from oxidative damage. Bartov *et al.* (1991) reported that Vit. E may minimize the decline in egg production and feed efficiency following the outbreaks of some diseases. El-Mallah *et al.* (2011) reported that dietary Vit. E at either level 0.20 or 0.40mg/kg considerably resulted in positive significant effect on EP values and had no effect on EW compared to the control, also, FI did not differ while, FC values were improved due to Vit. E addition

compared to the control. Vitamin C supplementation of broiler diets did not affect feed consumption or feed conversion rate (Abo Elouun and Al-Huminany, 2011).

As shown in Table 3, type of vitamins (E and C) supplementation had no effect on egg quality except, shell thickness and shell surface area (SSA) throughout the trial. It is clear that laying hens fed control diet had higher shell thickness and SSA, whereas, those fed diet containing mixtures of Vit. E and vitamin C supplementation had lower shell thickness and SSA during the experimental period.

Results presented in Table (3) indicated no significant differences in egg quality among all dietary treatments including the control group except, shell thickness, shell%, SSA and shell weight per unit surface area (SW/SA). Laying hens fed diet containing 10 mg/kg diet Vit. E had higher shell thickness, shell% and SSA while, those fed diet containing 20 mg Vit. E and 400 mg/kg diet Vit. C had lower shell thickness, shell% and SSA during the experimental period. This result agrees with those of Cherian *et al.* (1996) and Franchini *et al.* (2002), who observed no effects of dietary added tocopherols on the Haugh units of eggs.

Moreover, Puthongsiriporn *et al.* (2001) found that egg mass of White Leghorn hens was greater with supplementation of 65 IU of Vit. E /kg. Also, egg yolk was significantly increased when hens were fed 45 and 65 IU/kg compared with the control or Vit. E level (25 IU/kg). Haugh units were higher for hens fed 65 IU of Vit. E /kg compared to 25 and 45 IU/kg. Some reports indicated improvements in the egg production and egg shell quality by Vit. C (Bell and Marion, 1990; Balnave and Muheereza, 1997 and Al-Shoquiry, 1999). As well as by adding Vit. E (Bollengier *et al.*, 1999; Puthongsiriporn 1998; Kirunda *et al.*, 2001 and Sahin *et al.*, 2002). Vit. E serves as a physiological antioxidant through inactivation free radicals, improves egg production, feed intake, egg yolk and albumen solids (Kirunda *et al.*, 2001), and improves egg quality (Puthongsiriporn, 1998). These results confirmed those of El-Mallah *et al.* (2011) who reported that demonstrated that shape index and yolk color which significantly ($P<0.05$) decreased and shell thickness which significantly ($P<0.05$) improved compared to the control. Also, El-Sheikh and Salama (2010) reported that Vit E improved shell thickness and haugh unit score as compared to the control but, did not affect significantly shell weight% and albumen weight% as compared to control, Similar results were reported by Lmann *et al.* (2001), Kirunda *et al.* (2001) and Abdel-Galil and Abdel-Samad (2004). In this connection, the achieved improvement in shell-thickness could be due to enhancement of calcium bioavailability by the action of supplemental Vit. E. These facts confirmed the results of increased serum-ca concentration that has been established in the present study (Abdel-Fattah and Abdel-Azeem, 2007). Moreover, Vit. E addition was stated to influence the oestradiol dependant mechanisms by exerting a direct effect on oestradiol or indirect effect through maintaining more normal function of cellular processes regulating oestradiol and restoration of estrogen secretion (Bollengier *et al.*, 1998).

Fertility and hatchability%:

Fertility, hatchability embryonic mortality at 7 or 18 days and abnormal chicks% as affected by feeding different levels of Vit. E and C to El-Salam laying hens are presented in Table 4. Type of addition effect significantly influenced ($P\leq 0.05$) fertility%, it is clear that laying hens fed diet containing Vit. E supplementation had higher fertility, while, hens fed control diet had lower fertility%. Insignificant ($P\geq 0.05$) effects were observed in hatchability, embryonic mortality at 7 or 18 days and abnormal chicks% during the experimental period (Table 4).

Results presented in Table (4) indicated no significant differences in hatchability embryonic mortality at 7 or 18 d and abnormal chicks% among all dietary treatments including the control group. Dietary treatments effect was significant ($P\geq 0.05$) only for fertility%, hens fed diet containing 20 mg Vit. E and 400 mg/kg diet Vit. C had higher fertility% compared with those fed control diet, while, those fed control diet had lower fertility%. Numerically, there was insignificant increase ($P\geq 0.05$) in hatchability percentages in all groups fed Vit. E and C supplementation compared with the control group.

Similar results were observed by Arscott and Parker (1967) who found that fertility was rapidly restored when Vit. E supplementation was begun at 28 weeks. Comparable results were observed in hens (Machlin *et al.*, 1962) because fertility and hatchability were all drastically restored to normal levels by addition of Vit. E to the diet. Also, some studies have been shown that Vit. E tends to maintain or increase hatchability in heat stressed laying hens (Tengerdy and Nockels, 1973).

Table (3): Effects of supplementing laying hens diets with vitamin E, C and their mixtures on egg quality of El-Salam laying hens.

Item	Yolk color	Shell thickness, mm	Albumen %	Yolk %	Shell%	Yolk index%	Shape index%	Haugh unit	SSA ¹	SW/SA ²
Type of addition										
Control	7.20	0.336 ^a	61.44	29.68	8.67	46.24	75.48	90.54	63.29 ^A	69.30
Vitamin (Vit.) E	7.55	0.333 ^a	61.32	29.55	9.14	45.11	75.13	90.20	62.26 ^{AB}	72.25
Vitamin C	7.15	0.315 ^{ab}	61.23	29.49	9.27	45.76	75.44	89.23	59.91 ^B	74.37
Mixed (Vit. E & Vit. C)	6.78	0.300 ^b	61.52	29.64	8.84	45.34	75.15	87.79	56.06 ^C	70.56
±SEM ³	0.24	0.01	0.67	0.52	0.34	0.57	0.69	1.66	0.94	2.53
Treatments										
Control	7.20	0.336 ^{ab}	61.64	29.68	8.67 ^{ab}	46.24	75.48	90.54	63.29 ^{AB}	69.30 ^{AB}
Vit. E 10 mg/Kg diet	7.70	0.345 ^a	60.82	29.30	9.88 ^a	45.70	75.40	90.75	63.82 ^A	78.67 ^A
Vit. E 20 mg/Kg diet	7.40	0.320 ^{abc}	61.81	29.79	8.40 ^b	44.52	74.86	89.66	60.70 ^{ABC}	65.83 ^B
Vit. C 200 mg/Kg diet	6.90	0.300 ^{bc}	61.64	29.68	8.67 ^{ab}	45.94	75.48	89.10	59.64 ^{BCD}	69.66 ^{AB}
Vit. C 400 mg/Kg diet	7.40	0.330 ^{abc}	60.82	29.30	9.88 ^a	45.57	75.40	89.35	60.19 ^{ABCD}	79.08 ^A
Vit. E 10 mg/Kg diet + Vit. C 200 mg/Kg diet	7.20	0.300 ^{bc}	61.81	29.79	8.40 ^b	44.34	74.86	88.32	57.20 ^{CDE}	66.18 ^B
Vit. E 10 mg/Kg diet + Vit. C 400 mg/Kg diet	6.50	0.290 ^c	61.64	29.68	8.67 ^{ab}	46.13	75.48	87.73	56.27 ^{DE}	70.02 ^{AB}
Vit. E 20 mg/Kg diet + Vit. C 200 mg/Kg diet	6.90	0.320 ^{abc}	60.82	29.30	9.88 ^a	45.81	75.40	88.03	56.82 ^{CDE}	79.50 ^A
Vit. E 20 mg/Kg diet + Vit. C 400 mg /Kg diet	6.50	0.290 ^c	61.81	29.79	8.40 ^b	45.09	74.86	87.06	53.96 ^E	66.52 ^B
±SEM	0.34	0.01	0.97	0.76	0.45	0.81	1.01	2.42	1.32	3.26

¹ Shell surface area² shell weight per unit surface area³ Pooled SEM

a, ..., c, and A, ..., E, values in the same column within the same item followed by different superscripts are significantly different (at $P \leq 0.05$ for a to c; $P \leq 0.01$ for A to E).

Table (4): Effects of supplementing laying hens diets with vitamin E, C and their mixtures on fertility, hatchability, embryonic mortality and abnormal chicks% of El-Salam laying hens.

Item	Fertility %	Hatchability %	Embryonic mortality %			Abnormal chicks%
			at 7 days	at 18 days	Total	
Type of addition						
Control	83.98 ^b	82.59	0.694	0.694	1.389	0.000
Vitamin (Vit.) E	94.40 ^a	91.17	0.385	2.410	2.795	1.282
Vitamin C	90.22 ^{ab}	88.66	1.449	0.725	2.174	0.000
Mixed (Vit. E & Vit. C)	93.83 ^a	91.27	1.649	1.506	2.793	0.669
±SEM ¹	2.21	2.31	0.71	0.73	0.94	0.49
Treatments						
Control	83.98 ^D	82.59	0.694	0.694	1.39	0.000
Vit. E 10 mg/Kg diet	91.67 ^{ABCD}	89.33	0.806	0.645	1.45	0.000
Vit. E 20 mg/Kg diet	96.81 ^{AB}	92.79	0.000	4.020	4.02	2.451
Vit. C 200 mg/Kg diet	86.52 ^{CD}	85.25	1.905	0.476	2.38	0.000
Vit. C 400 mg/Kg diet	93.92 ^{ABC}	91.96	0.980	0.980	1.96	0.000
Vit. E 10 mg/Kg diet + Vit. C 200 mg/Kg diet	88.93 ^{BCD}	86.82	0.571	2.381	2.95	1.667
Vit. E 10 mg/Kg diet + Vit. C 400 mg/Kg diet	93.00 ^{ABCD}	92.43	1.429	0.571	0.57	0.000
Vit. E 20 mg/Kg diet + Vit. C 200 mg/Kg diet	93.62 ^{ABC}	89.78	2.313	1.524	3.84	0.952
Vit. E 20 mg/Kg diet + Vit. C 400 mg /Kg diet	100.00 ^A	96.09	2.344	1.563	3.91	0.000
±SEM	3.01	3.15	1.01	1.03	1.32	0.68

¹Pooled SEM

a, ..., b, and A, ..., D, values in the same column within the same item followed by different superscripts are significantly different (at $P \leq 0.05$ for a to b; $P \leq 0.01$ for A to D).

Table (5): Effects of supplementing laying hens diets with vitamin E, C and their mixtures on economical efficiency.

Item		T1	T2	T3	T4	T5	T6	T7	T8	T9
Price/ k feed (L.E.)	a	2.600	2.602	2.604	2.602	2.604	2.604	2.608	2.604	2.608
Total feed intake (kg)	b	79.564	77.920	77.149	76.786	77.006	77.545	75.718	76.333	76.297
Total feed cost (L.E.)	a x b = c	206.866	202.748	200.897	199.798	200.525	201.927	197.471	198.772	198.982
Total egg mass (Kg)	d	22.7653	22.3679	22.9774	23.1876	23.9571	22.3999	22.8202	24.3017	23.9393
Price/Kg eggs (L.E.)	e	19.819	19.819	19.819	19.819	19.819	19.819	19.819	19.819	19.819
Total price of eggs (L.E.)	d x e = f	451.185	443.31	455.389	459.555	474.81	443.94	452.27	481.64	474.45
Net revenue (L.E.)	f - c = g	244.319	240.561	254.493	259.757	274.281	242.017	254.802	282.863	275.471
Economical efficiency (E.Ef.)	g / c = h	1.181	1.187	1.267	1.300	1.368	1.199	1.290	1.423	1.384
Relative E.Ef.	r	100.00	100.46	107.26	110.08	115.81	101.48	109.25	120.49	117.22

T₁: Control. T₂: Vit. E 10 mg/Kg diet. T₃: Vit. E 20 mg/Kg diet. T₄: Vit. C 200 mg/Kg diet. T₅: Vit. C 400 mg/Kg diet. T₆: Vit. E 10 mg/Kg diet + Vit. C 200 mg/Kg diet. T₇: Vit. E 10 mg/Kg diet + Vit. C 400 mg/Kg diet. T₈: Vit. E 20 mg/Kg diet + Vit. C 200 mg/Kg diet and T₉: Vit. E 20 mg/Kg diet + Vit. C 400 mg/Kg diet.

a..... (based on average price of diets during the experimental time).

e..... (according to the local market price at the experimental time).

g/c..... (net revenue per unit feed cost).

r..... (assuming that economical efficiency of the control groups equals 100).

Economical efficiency(EEf):

Table 5 show the economical efficiency (EEf) and the relative economical efficiency (relative EEf) values. El-Salam laying hens fed diet containing 20 mg Vit. E and 200 mg/kg diet Vit. C gave the best economical and relative efficiency values being 1.423 and 120.49%, respectively followed by hens fed diet containing 20 mg Vit. E and 400 mg/kg diet Vit. C being 1.384 and 117.22%, respectively, all of which are superior compared to the control diet without supplementation, hens fed control diet had the worst corresponding values, being 1.181 and 100%, respectively. The relative efficiency varied between 0.0 to +20.49% which is of minor importance relative to the other factors of production.

CONCLUSION

The results of this study indicated that feeding El-Salam laying hens on diets containing 20 mg Vit. E and 200 mg/kg diet Vit. C improved the productive, reproductive performance and relative economic efficiency values of laying hen.

REFERENCE

- Abdel-Fattah, S. A., and F. Abdel-Azeem (2007). Effect of vitamin E, thyroxine hormone and their combination on humoral immunity, performance and some serum metabolites of laying hens in summer season. *Egypt. Poult. Sci.*, 27 (II): 335-361.
- Abdel Galil, M. A. and M. H. Abdel Samad (2004). Effect of vitamin E, C. Selenium and Zinc supplementation on reproductive performance of two local breeds of chickens under hot climate condition. *Egypt. Poult. Sci.*, 24, 217-229.
- Abo Elouun, S.A. and A. A. Al-Huminany (2011). The prophylactic effect of vitamin C supplementation on sodium chloride induced pulmonary hypertension (PHS) in broiler chickens. *Egypt. Poult. Sci. Vol. (31) (1):121-130.*
- Ajuyah, A. O., R. T. Hardin, and J. M. Sim (1993). Effects of dietary full-flat flax seed with and without antioxidant on the fatty acid composition of major lipid classes of chicken meats. *Poult. Sci.*, 72:125-136.
- Al-Shoquiry, N. E. (1999). Influence of some dietary factors on the performance of chickens. M. Sc. Thesis, Fac. Agric. Univ. Minufiya, Egypt.
- Arscott G.H. and J.E. Parker (1967). In *Physiology and Biochemistry of the domestic fowl*, pp. 380.
- Balnave, D. and S. K. Muheereza (1997). Improving egg shell quality at high temperature with dietary sodium bicarbonate. *Poult. Sci.*, 76: 588-593.
- Bartov, I., Y. Weisman, and E. Wax (1991). Effects of high concentrations of dietary vitamin E and ethoxyanin on the performance of laying hens. *Br. Poult. Sci.* 32: 525-534.
- Bell, D. E. and J. E. Marion (1990). Vitamin C in laying hen diets. *Poult. Sci.*, 69: 1900-1904.
- Bollengier, L. S., P. E. V. Williams, and C. C. Whitehead (1999) Optimal dietary concentration of vitamin E for alleviating the effect of heat stress on egg production in laying hens. *Br. Poult. Sci.*, 40: 102-107.
- Bollengier, L. S., M. A. Mitchell, D. B. Utomo, P. E. V. Williams, and C.C. Whitehead, (1998). Influence of high dietary vitamin E supplementation on egg production and plasma characteristics in hens subjected to heat stress. *Br. Poult. Sci.* 39: 1, 106-112.
- Brigelius-Flohe, R., F. J. Kelly, J. T. Salonen, J. Neuzil, J. M. Zingg, and A. Azzi (2002). The European perspective on vitamin E: Current knowledge and future research. *Am. J. Clin. Nutr.* 76:703-716.
- Carter, T.C. (1968). The hen egg. A mathematical model with three parameters. *Br. Poult. Sci.*, 9: 165-171.

- Carter, T. C. (1975). The hens egg. Estimation of shell superficial area and egg volume using measurements of fresh egg weight and shell length and breadth alone or in combination . Br. Poult. Sci., 16: 541-543.
- Cherian, G., F. H. Wolfe and J. S. Sim (1996). Feeding dietary oils with tocopherols: effects on internal qualities of eggs during storage. J. Food Sci., 61:15-18.
- Diplock, A. T. (1991). Antioxidant nutrients and disease prevention: An overview. Am. J. Clin. Nutr. 53:189-193s.
- Duncan, D.B. (1955). Multiple range and multiple F tests. Biometrics, 11:1-42.
- El-Badry, A. S. O., Kh. A. A. Ali, W. A. H. Ali and M. A. Ahmed (2011). The role of nasal gland and vitamin C in alleviate the adverse effects of osmotic stress on ostrich. Egypt. Poult. Sci., 31: 233-247.
- El-Mallah, G. M., S. A. Yassein, M. M. Abdel-Fattah and A. A. El-Ghamry (2011). Improving performance and some metabolic response by using some antioxidants in laying diets during summer season. J. of American Sci., 7(4):217-224. (ISSN: 1545-1003). <http://www.americanscience.org>
- El-Sheikh, S.E.M. and A.A. Salama (2010). Effect of vitamin C and E as water additives on production performance and egg quality of heat stressed local laying hens in Siwa Aqsis. Egypt. Poult. Sci., 30: 679- 697.
- Franchini, A., F. Sirri, N. Tallarico, G. Minelli, N. Iaffaldano and A. Meluzzi (2002). Oxidative stability and sensory and functional properties of eggs from laying hens fed supranutritional doses of vitamins E and C. Poult. Sci., 81: 1744-17
- Gebert, S., R. Messikommer, H. P. Pfrirter, G. Bee and C. Wenk (1998). Dietary fats and vitamin E in diets for laying hens: effects on laying performance, storage stability and fatty acid composition of eggs. Archiv Geflugelkunde. 62: 214-222.
- Hamilton, R.M.G. (1978). The effect of dietary protein level on productive performance and egg quality of four strains of White Leghorn hens. Poult. Sci., 57 (5): 1355-1364.
- Haugh, R. R. (1937). The Haugh unit for measuring egg quality. US Egg Poult. Mag. 43: 552-555.
- Jacob, R. A. (1995). The integrated antioxidant system. Nutr. Res. 15: 755-766.
- Jiang, Y. H., R. B. McGeachin and C. A. Bailey (1994). α -tocopherol, β -carotene, and retinol enrichment of chicken eggs. Poult. Sci., 73:1137-1143.
- Kirunda, D.F.K., S.E. Scheideler, and S.R. McKee (2001). The Efficacy of vitamin e (dl-tocopheryl acetate) supplementation in hen diets to alleviate egg quality deterioration associated with high temperature exposure. Poult. Sci., 80: 1378-1383.
- Knekt, P. A., A. Aromaa and J. Maatela (1991). Vitamin E and cancer prevention. Am. J. Clin. Nutr. 53: 283-286.
- Kutlu, H. R. and J. M. Forbes (1993). Changes in growth and blood parameters in heat stressed broiler chicks in response to dietary ascorbic acid. Livestock prod. Sci., 36: 335-350.
- Lmann, E. D., J. Halle, H. W. Rauch, H.P. Sallmann and G. Flachowsky (2001). Influence of various vitamin E supplementation on performance of laing hens. Archiv-fur. Geflugelkunde. 65: 182-186.
- Machlin, L.J., R.S. Gordon, J.E. Marr and C.W. Pope (1962). In Physiology and Biochemistry of the domestic fowl, pp. 380.
- Meluzzi, A., N. Tallarico, G. Manfreda, F. Sirri and A. Franchini (2000). Effect of dietary vitamin E on the quality of table eggs enriched with n-3 long chain fatty acids. Poult. Sci., 79:539-545.
- National Research Council, NRC (1994). Nutrient Requirements of Poultry. 9th revised edition. National Academy Press. Washington, D.C., USA.
- Packer, L. (1992). Interactions among antioxidants in health and disease: Vitamin E and its redox cycle. Proc. Soc. Exp. Biol. Med. 200, 271-276.
- Pardue, S. L. and P. J. Thaxton (1986). Ascorbic acid in poultry: A review. World's Poult. Sci. J. 42:107-123.
- Puthongsiriporn, U. (1998). Effects of strain and dietary vitamin E on hen performance, immune and antioxidant status during heat stress. M.S. Thesis. University of Nebraska-Lincoln. Lincoln, NE.

- Puthongsiriporn, U., S.E. Scheideler, J.L. Shell and M.M. Beck (2001). Effect of vitamin E and C supplementation on performance, in vitro lymphocyte proliferation and antioxidant status of laying hens during heat stress. *Poult. Sci.*, 80: 1190-1200.
- Retsky, K. L. and B. Frei (1995). Vitamin C prevents metal iondependent initiation and propagation of lipid peroxidation in human low-density lipoprotein. *Biochem. Biophys. Acta.* 1257:279-287.
- Ricciarelli, R., J. M. Zingg and A. Azzi (2002). The 80th anniversary of vitamin E: Beyond its antioxidant properties. *Biol. Chem.* 383: 457-465.
- Rice, M. E. (2000). Ascorbate regulation and its neuroprotective role in the brain. *Trends in Neuroscience*, 23: 209-216.
- Sahin, K., N. Sahin, and S. Yaralioglu (2002). Effects of vitamin C and vitamin E on lipid peroxidation, blood serum metabolites and mineral concentrations of laying hens reared at high ambient temperature. *Biol. Trace Elem. Res.*85: 35-45.
- Scheideler, S. E., and G. W. Froning (1996). The combined influence of dietary flaxseed variety, level, form, and storage conditions on egg production and composition among vitamin E supplemented hens. *Poult. Sci.*, 75: 1221-1226.
- S.P.S.S. (1999). User's Guide: Statistics. Version 10. SPSS Inc. Chicago, IL, USA.
- Surai, P., I. Ionov, A. Buzhin and N. Buzhina (1997). Vitamin E and egg quality. Pages 387-394 in *Proceedings of the 7th European Symposium on the Quality of Eggs and Egg Products*, Poznan, Poland, Zaklad Poligraficzny "Graf-Com," Poznan, Poland.
- Tengerdy, R. P. and C. F. Nockels (1973). The effect of vitamin E on egg production, hatchability and humoral immune response of chickens. *Poult. Sci.*, 52:778-783.
- Weiser, H., M. Schlachter, H. P. Probst and A. W. Kormann (1990). The relevance of ascorbic acid for bone metabolism. Pages 73-93 in *Ascorbic Acid in Domestic Animals. Proceedings of the 2nd Symposium*. C. Wenk, R. Fenster, and L. Volker (eds.), Kartause Ittingen, Switzerland.
- Well, R. J. (1968). The measurement of certain egg quality: A study of the hens egg. Ed. By T.C. Carter Pub. Oliver and Boyd Edinburgh pp. 220-226 and 235-236.
- Whitehead, C. C., S. Bollenger-Lee, M. A. Mitchell and P. E. V. Williams (1998). The role of vitamin E in alleviating heat stress in laying hens. *Poult. Sci.*, 77 (Suppl. 1):159. (Abstr.).

تأثير إضافة مستويات مختلفة من فيتامين ج، هـ لعلائق الدجاج البيضاء على 1- تحسين بعض الصفات الإنتاجية والتناسلية.

منى سيد رجب¹، صباح فاروق يوسف²، قوت القلوب مصطفى السيد مصطفى²

اكليّة الزراعة - قسم إنتاج الدواجن - جامعة الفيوم- مصر.

²مركز البحوث الزراعية- معهد بحوث الإنتاج الحيواني- القلي- الجيزة- مصر.

اجريت هذه الدراسة بمحطة بحوث الدواجن بالعزب- الفيوم- مصر لدراسة تأثير استخدام مستويين من فيتامين هـ (10 او 20 ملليجرام/كجم عليقة) و فيتامين ج (200 او 400 ملليجرام/كجم عليقة) وخليطهما علي كفاءة البيض، جودة البيض، الخصب، الفقس، والكفاءة الاقتصادية لدجاج السلام البيضاء. استخدم عدد 243 (216 أنثى و 27 ذك) طائر عمر 25 أسبوع قسمت عشوائيا إلي 9 معاملات متساوية 27 طائر / كل معاملة (24 أنثى و 3 ذك) ثم قسمت كل معاملة إلي 3 مكررات 11 طائر/مكرر (8 إناث و 1 ذكور).

وكانت المعاملات التجريبية كما يلي :

1- تغذية الطيور علي عليقة الكنترول (م1).

2- م1 مضاف إليها 10 ملجم/كجم عليقة فيتامين هـ.

3- م1 مضاف إليها 20 ملجم/كجم عليقة فيتامين هـ.

4- م1 مضاف إليها 200 ملجم/كجم عليقة فيتامين ج.

5- م1 مضاف إليها 400 ملجم/كجم عليقة فيتامين ج.

6- م1 مضاف إليها 10 ملجم/كجم عليقة فيتامين هـ+ 200 ملجم/كجم عليقة فيتامين ج.

7- م1 مضاف إليها 10 ملجم/كجم عليقة فيتامين هـ+ 400 ملجم/كجم عليقة فيتامين ج.

8- م1 مضاف إليها 20 ملجم/كجم عليقة فيتامين هـ+ 200 ملجم/كجم عليقة فيتامين ج.

9- م1 مضاف إليها 10 ملجم/كجم عليقة فيتامين هـ+ 400 ملجم/كجم عليقة فيتامين ج.

وتتلخص أهم النتائج المتحصل عليها فيما يلي:-

1- كان للدجاج المغذي علي عليقة مضاف إليها خليط من كل من فيتامين هـ ، ج أقل استهلاك للعليقة، بينما كان للدجاج المغذي علي عليقة المقارنة أعلي استهلاك للعليقة، لم يكن هناك أى تأثير معنوي علي كل من متوسط وزن البيض، كتلة البيض، إنتاج البيض، كفاءة تحويل الغذاء خلال فترة التجربة.

2- كان للدجاج المغذي علي عليقة تحتوي علي 10 ملجم/كجم عليقة فيتامين هـ اعلي سمك للقشرة ونسبة القشرة، بينما كان للدجاج المغذي علي عليقة المقارنة أقل سمك للقشرة ونسبة القشرة خلال فترة التجربة.

3- كان للدجاج المغذي علي عليقة تحتوي علي 20 ملجم/كجم عليقة فيتامين هـ و 400 ملجم/كجم عليقة فيتامين ج اعلي نسبة الخصب.

4- أعطي الدجاج المغذي علي عليقة تحتوي علي 20 ملجم/كجم عليقة فيتامين هـ و 200 ملجم/كجم عليقة فيتامين ج أحسن كفاءة اقتصادية واعلي كفاءة اقتصادية نسبية.

ومن ذلك يمكن استنتاج أن تغذية دجاج السلام علي عليقة تحتوي علي 20 ملجم/كجم عليقة فيتامين هـ و 200 ملجم/كجم عليقة فيتامين ج أدى إلي تحسين الأداء الإنتاجي والتناسلي وأعطى اعلي كفاءة اقتصادية ونسبية.