

TOXICOLOGICAL EFFECTS OF ORGANOPHOSPHORUS INSECTICIDES AND REMEDIATION TECHNOLOGIES OF ITS RESIDUES IN AQUATIC SYSTEM

A. MALATHION

Ahmed H. Massoud; Ismail I. El-Fakhrany; Aly S. Derbalah
and Mostafa S. Saad Allah

Pesticides Department Fac. of Agric. Kafr-EL-Shiekh University,
33516 Egypt

ABSTRACT

This study was carried out to survey the presence of organophosphorus (OP) insecticides in water resources in Kafr-EI-Shiekh governorate, Egypt. Moreover, to evaluate toxicological effects of one of the most frequently detected compound (malathion), with respect to some biochemical parameters (acetyl cholinesterase [AChE], alkaline phasphatase [ALP], glutamic-Pyrovic transaminase [GPT], glutamic-oxaloacetic transaminase [GOT] and glutathion-S-transferase[GST]) in blood and histopathological changes of treated rats organs (kidney and liver). Furthermore, to evaluate efficacy of different remediation techniques (advanced oxidation processes [AOPs] and bioremediation) for removing the tested insecticide in aquatic system. Malathion was detected with high frequency and concentration level relative to other detected organophosphorus in water resources in Kafr-EI-Sheikh Governorate. The tested insecticide at dose level of 10 mg kg⁻¹ induced significant toxicity against the treated rats relative to control with the respect to biochemical parameters and histopathological changes in selected rats organs. Photo-Fenton like reagent was the most effective treatment for malathion removal followed by Fe³⁺/UV, H₂O₂/UV, UV and Fe³⁺/H₂O₂ systems, respectively. Bioremediation of malathion by *Pseudomonas sp.* (EB20) isolate removed 75% of its initial concentration. This study concluded that, malathion at the tested concentration level in water expected to induce side effects on human health. Bioremediation using *Pseudomonas sp.* (EB20) can be regarded as a safe remediation technology for malathion in drinking water and the photo-Fenton like reagent would be more preferable in wastewater.

Key words: malahtion, residues, toxicity, remediation.

INTRODUCTION

Wide spread use and disposal of organophosphorus compounds that have been used as an alternative to organochlorine compounds for pest control (Muller *et al.*, 2001) resulted in the release of their residue into natural water, thus inducing an environmental problem (Amoros *et al.*, 2000). Pesticide pollution of surface waters and wastewaters has increased sharply and it constitutes a major pollutant problem and health hazards due to an extensive use of these substances (Evgenidou *et al.* 2007).

Therefore, monitoring the presence of organophosphorus insecticides in water and evaluation of its side effects on human health considered a source of major concern. Toxicity of

organophosphorus insecticides used compounds against human and animals were always evaluated by assessment of such biochemical parameters alterations and histopathological changes in tissues and organs (Cronelius *et al.*, 1959; Ghanem *et al.*, 2006). However, there is lack of evaluating the toxicity of organophosphorus pesticides at low concentration levels near the environmental level. Since most previous studies were using high doses of the tested pesticides to expect significant effect of these insecticides, however, this is did not reflect real situation.

Malathion is one of the most widely used organophosphate insecticides throughout the world. It is used to control pests affecting agricultural crops, ornamentals, greenhouses, livestock, stored grain, forests, buildings, households and gardens. Contributing to its popularity is its relatively low acute mammalian toxicity (Brenner, 1992; Hazarika *et al.*, 2003).

Due to the great environmental and human risk of pesticide residues in water resources Removal heavily used insecticides and their photoproducts from aquatic environment have become a very important task for human being. Thus, advanced methods are in demand for effective treatment of pesticides-polluted water to achieve complete mineralization of target pesticides and to avoid the formation of toxic end products (Derbalah, 2009). Advanced oxidation processes (AOPs), which are constituted by the combination of several oxidants, are characterized by the generation of very reactive and oxidizing free radicals in aqueous solution such as hydroxyl radicals, which posses a great destruction power (Benitez *et al.*, 2002).

The photo assisted-Fenton reaction process as advanced oxidation process proved to be very powerful in destroying persistent pesticides in the wastewater such as methyl parathion, methamidofos, fenotrothion and alachlor (Penuela and Barcelo 1998; Fallmann *et al.*, 1999; Derbalah *et al.*, 2004). In this respect, it is necessary to apply this method in water with low concentration level of pesticides (near to the environmental level) and by using natural river water to test the efficiency of this method under natural conditions, we would be able to generalize photo-Fenton reaction process for pesticides removal from water (Derbalah *et al.*, 2004). Bioremediation of chemo-pollutants becomes the method of choice because it is economically feasible and safer than chemical remediation technologies (Massoud *et al.*, 2008). *Pseudomonas* sp is known for their versatility in degradatation of xenobiotic compounds such as pesticides in water (Abd El-Razik, 2006; Massoud *et al.*, 2007 b; Derbalah *et al.*, 2008).

Therefore, the present study aimed to monitor the presence of organophosphorus insecticides in different water resources in Kafr El-Sheikh Governorate Egypt, to evaluate the toxic effects of the most frequently detected compound in water resources (malathion), at low

dose near the environmental levels with the respect to some biochemical's parameters (AChE, ALP, GOT, GPT and GST) in blood and histological changes (liver and kidney) in treated rats organs and finally to evaluate the efficacy of different remediation technologies (advanced oxidation processes and bioremediation) for the removal of the tested insecticide residue in the aquatic system.

MATERIALS AND METHODS

Monitoring of the organophosphorus compounds in water resources

Sampling sites and description

Surface water samples were collected in summer 2008 for one time from three sites (Meet Yazid canal, No.7 drain, and Al-Gharbiya main drain) at River Nile, Rosetta branch in Kafrelsheikh Governorate in glass bottles. Water samples transferred directly to the laboratory for filtration, extraction and analysis of pesticides residues.

Meet Yazid canal presents the main source for drinking water in Kafrelsheikh and Seedy Salem cities and adjacent village besides its use as a main source for irrigation water, serve about 150 fadden. Also, fishing activities are carried out. Al-Gharbiya main drain, one of many drainage canals present in the area of study, it receive drainage water from adjacent fields and from industrial area. It serve region about 156 fadden. No.7 drain, this canal is one of many drainage canals present in the area of study, it receive drainage water from adjacent fields. (Drainage system of the irrigation water). It serve region about 64.282 Fadden (**Maasoud et al., 2007a**).

Extraction

In the laboratory, samples were extracted immediately after 24 hr of site collection using disposable solid phase extraction (SPE) with octadecyl-bonded silica cartridges (C₁₈) (500 mg; Phenomenex Scientific) according by **Belden et al., (2000)**.

Gas Chromatography Analysis

Analysis of final extract was conducted using a Hewlett Packed model 6890 equipped with a flame photometric detector (FPD) with phosphorus filter. The analytical column was a 30m×0.32mm HP-5, with a film thickness of 0.25_μm. The 1_L SDME extract was injected into the GC-FPD in splitless mode at 250 °C. The column oven temperature was held at 125 °C for 2 min, then programmed at 75 °C/min to 200 °C for 1 min, 10 °C/min to 220 °C and held for 3 min (**Xiao et al., 2006**). Detection limit and recovery of each detected compound were carried out according to the method described by **Xiao et al., (2006)**.

Toxicity experiment

Adult Wistar male rats *Rattus norvegicus* 8 weeks of age and 80-100 gm in weight, obtained from Faculty of Medicine, Tanta University, Egypt were used. Wister rats were housed in wire cages under standard conditions with free access to drinking water and food. The rats were kept in temperature-controlled room with 14 hrs light and 10 hrs dark cycles and given standard-diet. Before treatment, rats were made adaptation for two weeks during feeding. The animals were randomly divided into four groups each comprising of three animals. Two groups of tested insecticide (24 hrs and 21 days), group for control (without malathion) and group for ethanol control. Rats were treated by active ingredient of the tested insecticide (malathion) with a purity of 99% obtained from Kafr El Zayat for Chemicals and Pesticides Company Limited, Kafr-El-Zayat , Egypt. The tested insecticide was dissolved in ethanol and gave to rats by oral dose at level of 10 mg / kg. Rats were scarified under anesthesia. Then tissue and blood samples were taken after 24 hrs (acute toxicity) and 21 days (sub-chronic toxicity). Blood samples were taken by cardiac puncture in vials containing heparin for enzymes activity determination. For histopathological test, rat organs (liver and kidney) were taken and kept in formalin 10 % for histopathological test (Derbalah, 2009).

Enzymes Assays

The colorimetric methods of The colorimetric methods of **Elman et al., 1961**); **Gornal et al., 1949**); **Belfield and Goldberg, 1971**); **Reitman and Frankel, 1957**); and **Rose and Wallbank, 1986** were used for determining the activity of AChE (acetylcholinesterase), (ALP) alkaline phasphatase, GPT (Glutamic-Pyrovic Transaminase), GOT (Glutamic-Oxaloacetic Transaminase) and GST (glutathion-S-transferase) in blood , respectively.

Histopathological test

The histopathology test was carried out at Dep. of Histopathology, Fac. of Veterinary Medicine, Kafresheikh Univ. Egypt. This experiment was conducted to study the histopathological lesions of the organs in treated rats with the tested insecticide; these organs were removed and prepared for histopathological examination according to **Bancroft and Stevens (1916)**.

Chemical remediation of tested insecticide in aqueous system

A UV mercury lamb (model VL-4 LC (8W) was employed for the irradiation of the tested insecticide (malathion). Ferric chloride was used as a source of iron catalyst because it remains unchanged before and after oxidation and this made the study of the reaction and the future engineering scale-up simpler, because the system remained homogeneous (**Derbalah et al., 2004**). The solution was prepared by addition of desired amounts of malathion technical grade (10 ppm) in distilled water. Then freshly prepared ferric chloride, FeCl_3 , at

concentration level of 1 mM as ferric ion was added followed by addition of H₂O₂ at 20 mg/l and the total volume was reached 100 ml by distilled water. The initial pH of the prepared solution was adjusted at 2.8 using hydrochloric acid 1 Molar for all experiments (**Derbalah et al., 2004; Derbalah, 2009**) using pH meter Jenway (Model 3510, PH/mV/Temperature Meter). All degradation experiments were carried out at room temperature. The solution was transferred from standard flask to glass dish and exposed to irradiation of UV lamp (the distance between the lamp and pesticides solution 15 cm) with a wave length of 265 nm (**Derbalah, 2009**). Illumination times were 10, 20, 40, 80, 160 and 320 min. Samples were removed at these regular intervals for HPLC analysis. Moreover, three experiments were carried out, the first in the absence of hydrogen peroxide (Fe³⁺/UV) to account for the degradation of methomyl under iron, the second in the absence of iron to account for the degradation of methomyl under hydrogen peroxide (H₂O₂/UV) and finally the third in the absence of iron and hydrogen peroxide to account the degradation under UV light only. Moreover, to account the effect of light on Fenton degradation ability, one experiment was carried out in the presence of Fenton components under dark conditions. The irradiated samples were analyzed directly by HPLC system in the central laboratory of pesticides, Agriculture Research Center, El-Dokey, Egypt. A mixture of acetonitrile and distilled water (60:40) was used as mobile phase under the isocratic elution mode. The flow rate of mobile phase was maintained at 1.0 ml /min. The used detector was UV and the wavelength was 254 nm (**Massoud et al., 2007b**).

Bioremediation of tested insecticide in aqueous system

Pseudomonas sp. (EB20) was isolated from El-Hamoul water at Kafr-El-Sheikh Governorate, which polluted by persistent organic pollutants (POPs) (**Ashry et al., 2006**) and identified according to its morphological and physiological parameters as described by **Holt (1984)**. The bioremediation test was carried out at Microbiology Laboratory, Dep. of Agric. Botany, and Fac. of Agric. Kafr El-Sheikh Univ. Egypt. The selected microbial isolate *Pseudomonas sp.* (EB20) was cultured onto Mineral Slat Medium (MSL) spiked with the tested insecticide (malathion) separately for 7 days and then the growing colonies was washed with three ml sterilized MSL medium. The cell suspension of 10⁸cfu/ml (colony forming unit) was used to inoculate 100 ml MSL medium containing 10 ppm of the tested insecticide. The cultures were incubated at 30°C, pH (7) and 150 rpm as optimum conditions for the growth of the tested microbial isolate (**Derbalah et al. 2008**) for 14 days. Samples were collected at 0, 3, 7, 10 and 14 days for monitoring the degradation of the tested insecticide. Control flasks of equal volume of MSL medium and the tested insecticide without the selected microbial isolate were run in parallel at all

intervals to assess a biotic loss. The collected water samples of the tested insecticide were filtered using syringe filtered (**Derbalah et al., 2008**) followed by HPLC analysis as mentioned before.

Statistical analysis Data were subjected to the analysis of variance test and Duncan's multiple range test using a computer program SAS (Version 6.12, SAS Institute Inc., Cary, USA).

RESULTS AND DISCUSSION

Monitoring of organophosphorus in water samples

The analysis of water samples from Kafr El-Sheikh Governorate at the three sampling sites (Meet Yazid canal, No.7 drain and Al-Gharbiya main drain) showed presence of numerous organophosphorus insecticides. The detected organophosphorus compounds were azinophos-methyl, Malathion, dimethoate, phenthoate and fenitrothion in water samples at all sampling sites as shown in Table (1). The concentrations of the detected organophosphorus insecticides ranged from 0.020 to 1.034 ppm. The detection limit and recovery of the detected compounds ranged from 25-56 ng/ml and 90-97%, respectively as shown in Table 1.

Both the detection frequency and concentration level of dimethoate and malathion were higher than the other detected organophosphorus insecticides at all sampling sites. With the concerning the sampling sites, the organophosphorus pesticides were abundant in the following order; Meet Yazid canal > Al-Gharbiya main drain > No.7 drain. These results are in agreement with the findings of **Ashry et al., (2006)** and **Massoud et al., (2007a)**, they recorded that the occurrence of organophosphorus such as dimethoate, fenitrothion, malathion and phenthoate in irrigation and drainage water in Kafr El-Sheikh governorate.

Table (1): Organophosphorus compounds levels (ppm) in irrigation and drainage water of Kafrelsheikh Governorate as well as its detection limit and recovery .

Pesticides	Meet Yazid canal	No.7 drain	Al-Gharbiya main drain	Detection limits	Recovery percentage
Azinophos-methyl	N.D	N.D	0.02	0.27×10^{-3}	97
Malathion	0.59	1.034	0.3	0.23×10^{-3}	92
Dimethoate	0.42	0.83	0.73	0.56×10^{-3}	96
Phenthoate	N.D	N.D	0.017	0.25×10^{-3}	90
Fenitrothion	0.011	N.D	N.D	0.33×10^{-3}	91

*N.D: Not detected.

Toxicity of malathion on some biochemical's parameters in rats

The obtained data in Table (2) showed that, the activity of GOT, GPT and GST enzymes were increased after 24 hrs of treatment with malathion at dose level of 10 mg/kg comparing with

control treatment. On the other hand, the activity of ALP enzyme was decreased after 24 hrs of treatment with malathion at the same dose as shown in Table (2). However, after 21 days of treatment, all liver functions enzymes (GOT, GPT and GST) activity were decreased comparing to control except for GST enzyme that increased as shown in Table 3).

The elevation of liver enzymes in rats treated with malathion might be due to the early damage in the hepatic cells which disturbed normal liver function and this positively correlated with the histopathological changes in the liver tissue mentioned elsewhere in this study. The se results are in agreement with findings of many researches who reported that, when the liver cell membrane was damaged, several enzymes located in the hepatocyte cytosol, including ALP, GPT and GOT were secreted into the blood (**Agrahari et al., 2007; Ncibi et al., 2008**). Moreover, the decrease of hepatic enzymes in rats treated with malathion after 21 days may be due to the presence of ethanol as a solvent which acts as free radical producer, increasing the enzyme inhibition (**Sivapiriya et al., 2006**). Moreover, the detoxification of the tested insecticide with the time after treatment increased by these defense enzymes and subsequently their activity with time gradually decreased.

The obtained data in Tables (2) and (3) showed that, the activity of GST and ALP enzymes were decreased either after 24 hrs or 21days of treatment with ethanol only comparing with control treatment. However, the activity of GPT and GOT enzymes were increased after 24 hrs of treatment with ethanol and decreased after 21days of treatment comparing with control. The decrease of hepatic enzymes in rats treated with ethanol only may be due to ethanol acts as free radical producer, increasing the enzyme inhibition caused by the free radicals (**Sivapiriya et al., 2006**).

Referring to acetylcholinesterase activity, the obtained results revealed that, the activity decreased comparing to the control either after 24 hrs or 21 days of treatment as shown in Tables (2 and 3). These results are in agreement with the findings of many researchers who reported that the most prominent clinical effects of poisoning with OPs were related to their inhibition of the activity of blood acetylcholinesterase (AChE) (**Timur et al., 2003; Hazarika et al., 2003; Massoud et al., 2010**). Moreover, the obtained results indicated that, the activity of AChE enzyme was increased either after 24 hrs or 21days of treatment with ethanol only comparing with control (Tables 2 and 3). The increase in AChE activity in rat treated with ethanol relative to dimethoate may be due to that ethanol known to significantly reduce the inhibition of the AChE inhibitors (**Sivapiriya et al., 2006**)

Table (2): Effect of malathion at dose level of 10 mg/kg on activity of some biochemical's parameters in rats after 24 hrs of treatment.

Treatments	AChE (U/L)	ALP (U/L)	GPT (Units/ml)	GOT (Units/ml)	GST (Units/ml)
Control	$5.51 \times 10^{-1} \pm 0.035^a$	$1.28 \times 10^{-1} \pm 0.0002^a$	$8.49 \times 10^{-1} \pm 0.007^d$	$10.58 \times 10^{-1} \pm 0.007^c$	$11.5 \times 10^{-3} \pm 0.0033^b$
Ethanol	$5.86 \times 10^{-1} \pm 0.023^a$	$4.67 \times 10^{-2} \pm 0.0002^b$	$9.58 \times 10^{-1} \pm 0.0098^c$	$14.47 \times 10^{-1} \pm 0.003^b$	$2 \times 10^{-3} \pm 0.0005^c$
Malathion	$2.22 \times 10^{-1} \pm 0.012^c$	$2.28 \times 10^{-2} \pm 0.00025^d$	$15 \times 10^{-1} \pm 0.0028^b$	$14.97 \times 10^{-1} \pm 0^a$	$5.06 \times 10^{-2} \pm 0.0066^a$

*a, b and c letters shows the significance and non-significance between the means at p value of 0.05 using Duncan's multiple range test.

Table (3): Effect of malathion at dose level of 10 mg/kg on activity of some biochemical's parameters in rats after 21 days of treatment.

Treatments	AChE (U/L)	ALP (U/L)	GPT (Units/ml)	GOT (Units/ml)	GST (Units/ml)
Control	$9.73 \times 10^{-1} \pm 0.012^c$	$1.128 \times 10^{-1} \pm 0.0007^b$	$26.07 \times 10^{-1} \pm 0.009^a$	$21.93 \times 10^{-1} \pm 0.005^a$	$1.7 \times 10^{-2} \pm 0.0017^b$
Ethanol	$14.65 \times 10^{-1} \pm 0.012^a$	$4.77 \times 10^{-2} \pm 0.0003^c$	$10.27 \times 10^{-1} \pm 0.003^c$	$4.19 \times 10^{-1} \pm 0.002^c$	$1.3 \times 10^{-3} \pm 0.0003^c$
Malathion	$6.91 \times 10^{-1} \pm 0.035^d$	$2.1 \times 10^{-2} \pm 0.0001^d$	$8.19 \times 10^{-1} \pm 0.003^d$	$8.39 \times 10^{-1} \pm 0.005^b$	$2.1 \times 10^{-2} \pm 0.006^b$

*a, b and c letters shows the significance and non-significance between the means at p value of 0.05 using Duncan's multiple range test.

Histology test

Histopathological changes in kidney

In normal histological structure of the kidney, the cortex contains glomerular tufts scattered in between proximal and distal convoluted tubules (Fig. 1A). Rats treated with malathion at dose level 10 mg/kg after 24 hrs of treatment showed no lesions (Fig 1B). Rats treated with malathion at dose level 10 mg/kg after 21 days of treatment showed slight tubular degeneration (Fig. 2B) and tubular regeneration (Fig. 2C). The results in this study agree with the findings of Kerem *et al.*, (2007) and Afshar *et al.*, (2008) who reported kidney damage such marked tubular dilation, hydropic degeneration in tubular lining epithelium, moderate congestion and hemorrhage in the cortical male Wistar rats exposed some organophosphate pesticides.




Fig. (1): Sections in kidney of rats treated with malathion at dose level 10 mg/kg after 24 hrs of treatment (B) relative to control (A).




Fig. (2): Sections in kidney of rats treated with malathion at dose level 10 mg/kg after 21 days of treatment (B and C) relative to control (A).

Histopathological changes of liver

There were no changes in the liver of rats treated with malathion at dose level 10 mg/kg after 24 hrs of treatment (Figs. 3B). Rats treated with 10 mg/kg of malathion for 21 days showed more severe changes in the form of moderate perivascular mononuclear cell infiltration (Fig. 4B) and moderate hepatocellular necrosis with mononuclear cell infiltration of the affected area (Fig. 4C). Also, large focal granuloma was observed (Fig. 4D). Severe venous congestion of the hepatic blood vessels and sinusoids were observed (Fig. 4E). The liver is well known target organ of the toxic impact regarding its function in biotransformation and excretion of xenobiotics (Roganovic and Jordanova 1998). Therefore, the liver tissue damage in rats treated with malathion such as inflammatory, cell infiltration, hemorrhage, calcification, vacuolar degeneration, dilation of sinusoids, vascular congestion and necrosis was expected (Sayim, 2007; Yehia et al., 2007). These changes are entirely consistent with the changes in various biochemical parameters that were also observed in this study.




Fig. (3): Sections in liver of rats treated with malathion at dose level of 10 mg/kg after 24.hrs of treatment (B) relative to control (A).





Fig. (4): Sections in liver of rats treated with malathion at dose level of 10 mg/kg after 21.days of treatment (B, C, D and E) relative to control (A).

Chemical remediation of the tested insecticide in aqueous solution


The reduction in malathion initial concentration with the irradiation time under UV, H₂O₂/UV, Fe³⁺/UV, Fe³⁺/H₂O₂ and Fe³⁺/H₂O₂/UV systems was evaluated. The results in Fig.6 showed

that, the degradation rate of the tested insecticide was greatly enhanced by irradiation under $\text{Fe}^{3+}/\text{H}_2\text{O}_2/\text{UV}$ (photo-Fenton like reaction) relative to $\text{H}_2\text{O}_2/\text{UV}$, Fe^{3+}/UV , UV and $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ systems. About 95% of malathion initial concentration was degraded under $\text{Fe}^{3+}/\text{H}_2\text{O}_2/\text{UV}$ system within 320 min of irradiation time compared with 55, 54, 51.2 and 44% of the same insecticide in the presence of, Fe^{3+}/UV , $\text{H}_2\text{O}_2/\text{UV}$, UV and $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ systems, respectively within the same irradiation time (Fig.5).


The photodegradation of malathion under UV light is due to the direct absorbance of UV light (photolysis (**Derbalah et al., 2004**). While the degradation of malathion under $\text{H}_2\text{O}_2/\text{UV}$ system due to firstly, the direct photolysis and secondly due the generation of hydroxyl radicals from hydrogen peroxide eq. (1) (**Benitez et al. 2002; Derbalah et al. 2004b; Derbalah, 2009**)



However, the photodegradation of malathion under Fe^{3+}/UV system is due to the direct photolysis of the tested compounds by absorbance of UV light. Moreover, due to the indirect photolysis of this compound by the hydroxyl radicals generated from in the presence of UV light as shown in eq. (2)






The degradation of malathion under $\text{Fe}^{3+}/\text{H}_2\text{O}_2$ system is due to the generation of hydroxyl radicals under this system in the absence of the light as shown in eq. 3



The degradation rate of malathion under $\text{H}_2\text{O}_2/\text{UV}$ system was slightly slower than its degradation under Fe^{3+}/UV system which may be due to the lower formation rate of hydroxyl radicals under $\text{H}_2\text{O}_2/\text{UV}$ system relative to Fe^{3+}/UV system (**Derbalah et al., 2004**). This low generation rate of hydroxyl radicals under $\text{H}_2\text{O}_2/\text{UV}$ system compared to Fe^{3+}/UV system can attributed to the fact that hydrogen peroxide absorbs weakly above 300 nm and that hydroxyl radicals are only generated through direct photolysis of hydrogen peroxide [**Benitez et al., 2002**]. On the other hand, under Fe^{3+}/UV system the $\text{Fe}(\text{OH})^{2+}$ complex is the predominant species for generating hydroxyl radicals absorbed light at wavelengths up to 410 which may lead to the generation of hydroxyl radicals under this system more than $\text{H}_2\text{O}_2/\text{UV}$ system (**Derbalah et al., 2004**).

The great enhancement in tested insecticide degradation rate under photo-Fenton's like reagent system ($\text{Fe}^{3+}/\text{H}_2\text{O}_2/\text{UV}$) relative to the other advanced oxidation processes is due to the higher generation rate of hydroxyl radicals under this system ($\text{Fe}^{3+}/\text{H}_2\text{O}_2/\text{UV}$) than the other systems (**Derbalah et al., 2004; Derbalah, 2009**). This high generation rate of hydroxyl radicals under photo-Fenton like reagent due to many reasons. Firstly, the photolysis of hydrogen peroxide itself, which leads to the formation of hydroxyl radicals eq. (1) (**Benitez**

et al., 2002). Secondly, the photolysis of $Fe(OH)^{2+}$ complex the predominant specie of Fe^{3+} for generating hydroxyl radicals eq.(2) (*Larson et al., 1991*). Thirdly, the photodecarboxylation of ferric carboxylate complexes generate ferrous ion (eq.3), which can in turn react with hydrogen peroxide to generate additional hydroxyl radicals (eq.4) (*Pignatello and Sun, 1995*).



Results of present study about malathion agree with finding of many researchers who reported that, total disappearance of organophosphorus insecticides were achieved by using photo-Fenton like treatment, after illumination time at pilot-plant scale under natural solar UV light (*Benitez et al., 2002; Oller et al., 2007; Tamimi et al., 2008*).




Fig (5): Degradation of malathion at initial concentration of 10 mg/l under $Fe^{3+}/H_2O_2/UV$, H_2O_2/UV , Fe^{3+}/UV , Fe^{3+}/H_2O_2 and UV systems in distilled water.

Bioremediation of the tested insecticide in aqueous solution

The degradation ability of the selected microbial isolate *Pseudomonas sp.*(EB20) to malathion was illustrated in Fig (6). *Pseudomonas sp.*(EB20) showed high potential in the degradation of the tested insecticide. Since a 75% of malathion of initial concentration level (10 ppm) was degraded within two weeks of incubation with *Pseudomonas sp.* (EB20). On the other hand, the degradation percentage of malathion reached to 4% at the end of incubation time in control or non-inoculated samples. This is implied that the quote of the tested insecticide decay due to temperature

effect, photodecomposition and volatilization is very slight or negligible. The degradation of malathion may be attributed to the secretion of enzymes from either tested bacterial or fungal strains which are capable of degrading pesticides (Bollag and Liu 1990). The genus *Pseudomonas* showed the highest rate of malathion degradation, and has considerable potential for the biotransformation and biodegradation of pesticides with widely differing chemical structures (Massoud et al., 2007b).




Fig (6): Biodegradation of malathion at concentration level 10 ppm by *Pseudomonas* sp. (EB20) isolate in aquatic medium.

CONCLUSIONS

Malathion was detected with high level in different water resources in Kafr-El-Sheikh Governorate relative to other detected insecticides. Malathion induced toxicological effects in treated rats relative to control with the respect to enzymes activity and histological changes in treated organs. However, more extensive studies are needed to evaluate the toxicity of malathion at concentration level near the environmental level which in return helps to evaluate its real toxicity. Photo-Fenton like reagent was the most effective treatment for the removal of malathion residues in wastewater. *Pseudomonas* sp (EB20) could be regarded as a safe removal treatment of malathion in drinking water.

REFERENCES

- Abd El-Razik, M.A.S. (2006). Toxicological studies on some agrochemical Pollutants. M.SC. Thesis, Fac. of Agric. Kafrelsheikh Univ. 1-167.
- Afshar, S. A. Farshid. R. Heidari and M. Ilkhanipour (2008). Histopathological changes in the liver and kidney tissues

- of Wistar albino rat exposed to fenitrothion Toxicol Ind Health. 24. (9):581-586.
- Amoros, I. R. Connon H. Garelick J. L. Alonso and J.M. Carrasco (2000). An assessment of the toxicity of some pesticides and their metabolites affecting a natural aquatic environment using the Microtox TM system. *Water Science Technology* 42: 19-24.
- Ashry, M. A.; O. C. Bayoumi.; I. I. El-Fakharany.; A. S. Derbalah and A. A. Ismail (2006). Monitoring and removal of pesticides residues in drinking water collected from Kafir El-Sheikh governorate, Egypt. *J.Agric.Res.Tanta Univ.*32 (3):691-704.
- Bancroft, J. D. and A. Stevens (1996). *Theory and Practice of Histological Techniques*. Fourth edition.
- Belden, J. B. C. S. Hofelt and M. J. Lydy (2000). Analysis of multiple pesticides: In Urban storm water using Solide-Phase Extrction. *Arch. Environ.Contam.Toxicol.* 38:7-10.
- Belfield, A and D. M. Goldberg (1971). Alkaline phosphates colorimetric method. *Enzyme.* 12:561.
- Benitez, F. J. J. L Acero and F. J. Real (2002). Degradation of carbofuran by using ozone, UV radiation and advanced oxidation processes. *Journal of Hazardous Materials B.*89: 51-65.
- Brenner, L. (1992). Malathion Fact Sheet. *Journal of Pesticide Reform, Northwest Coalition for Alternatives to Pesticides, Eugene, OR.* 12, 1-17.
- Bollag, J.M. and Liu S.Y. (1990) Biological transformation process of pesticides. In: H.H. Chang (ed.), *Pesticides in the Soil Environment*, Soil Science Society of America, Madison, WI, pp. 169-211
- Cronelius, C. E. W. Charles and E. Arhode, (1959). Serum and tissue transaminase activities in domestic animals. *Cornell vet.* 49: 116-121.
- Derbalah, A.S. (2009). Chemical remediation of carbofuran insecticide in aquatic system by advanced oxidation processes. *J.Agric.Res. Kafrelsheikh Univ.* 35(1): 308-327.
- Derbalah, A. S. N. Nakatani and H. Sakugawa (2004). Photocatalytic removal of fenitrothion in pure and natural waters by photo-Fenton reaction. *Chemosphere.*57: 635-644.
- Derbalah. A. S. A. H. Massoud and E. B. Belal (2008). Biodegradability of famoxadone by various microbial isolates in aquatic system. *Land Contamination & Reclamation.* 16(1): 13-23.
- Ellman, G.L. K.D. Courtney V. Andres and R M. Featherstone (1961). A new and rapid calorimetric determination of

- acetylcholinesterase activity. *Biochem. Pharmacol.* 7: 88-95.
- Evgenidou, E. I. Konstantinou K. Fytianos and I. Poulios (2007). Oxidation of tow organophosphorus insecticides by the photo-assisted Fenton reaction. *Water Research.* 41: 2015-2027.
- Fallmann, H. T. Krutzler R. Baue. S. Malato and J. Balanco (1999). Applicability of the photo-Fenton methods for treating water containing pesticides. *Catalysis Today.* 54: 309-319.
- Ghanem N. F. N. A. Hassan A.A. Ismail (2006). biochemical and histological changes induced in rats fed on diets and byproducts of fumigated wheat grains with phostoxin. *Pro. 4th Int. Con. Biol.Sc. (zool)* 259-268.
- Gornal, A. C. C. J. Bardawill and M. M. David (1949). Protein-Biuret colorimetric method. *J. Biol.Chem.* 177:751.
- Hazarika, A S. N. Sarker S. Hajar M. Kataria and J. K. Malik (2003). Influence of malathion pretreatment on the toxicity of anilofos in male rats: a biochemical interaction study. *Toxicology.* 185:1-8.
- Holt, G.H. (1984) *Ordinary gram negative bacteria.* Bergys, Manual of Systematic Bacteriology, (ed. Krieg N.R) Williams and Wilkins, Baltimore.
- Kerem, M.; N.Bedirli.; N. Gurbus.; N. N. Ekinci.; A. Bedirli.; T. Akkaya.; , O .Sakrak. and H. Pasaoglu (2007). "Effects of acute fenthion toxicity on liver and kidney function and histology in rats," *Turk. J. Med. Sci.* 37: 281-288.
- Larson, R. A. M.B. Schlauch and K. A. Marley (1991). Ferric ion promoted photodecomposition of triazines. *Journal of Agriculture and Food Chemistry* .39: 2057-2062.
- Lasram, M. M. A. B. Annabi N. El-elj S. Selmi A. Kamoun S. El-fazaa and N. Gharbi (2009). Metabolic disorders of acute exposure to malathion in adult wistar rats. *Journal of Hazardous Materials.*163:1052-1055.
- Liu, Z. Q. Hong1. J. Xu W. Jun S. Li (2006). Construction of a genetically engineered microorganism for degrading organophosphate and carbamate pesticides. *International Biodeterioration & Biodegradation.*58: 65–69.
- Massoud, A. H.; A. S. Derbalah.; E. B. Belal and I. I. El-Fakhrany (2007b). Bioremediation of malathion in aquatic system by different microbial isolates .*J. Pest Cont & Environ. Sci.* 15(2):13-28.
- Massoud, A. H. I. I. El-Fakhrany and M. A. S. Abd El-Razik (2007a). Monitoring of some agrochemical pollutants in surface water in Kafrelsheikh Governorate. *J. Pest Cont & Environ. Sci.* 15 (2): 21-41.

- Massoud; A.A. Derbalah A.S. Iman A. Abd-Elaziz I.A. and Ahmed M.S. (2010) Oral Toxicity of Malathion at Low Doses in Sprague-Dawley Rats: A Biochemical and Histopathological Study. *Menofia Vet. Journal* 7 :183-196.
- Muller, H and W. Schwack (2001). Photochemistry of organophosphorus insecticides. *Rev. Environ. Contam. Toxicol.* 172: 129-228.
- Oller, I. S. Malato J. A. Sanchez-Perez M. I. Maldanonads and R.Gasso (2007). Detoxification of waste water containing five common pesticides by solar AOPs-biological coupled system. *Catalysis Today* 129: 69-78.
- Penuela, G. A and D. Barcelo (1996). Comparative degradation kinetics of alachlor in water by photocatalysis with FeCl₃, TiO₂ and photolysis, studied by solid phase disk extraction followed by gas chromatographic techniques. *Journal of Chromatography A* . 754: 187-195.
- Penuela, G. A and D Barcelo (1998). Photodegradation and stability of chlorothalonil in water studied by solid phase extraction followed by gas chromatographic techniques. *Journal of Chromatography A* .823: 81-90.
- Pignatello, J. J and Y. Sun (1995). Complete oxidation of metholachlor and methyl parathion in water by the photoassisted Fenton reaction. *Water Research*. 29. (8):1837-1844.
- Reitman, A and S. Frankel (1957). GPT (ALT) Glutamic-Pyruvic Transaminase and GOT (AST) Glutamic -Pyruvic Transaminase colorimetric method. *Amer. J. Clin. Path.* 28:56.
- Rose, H. A and B. E. Wallbank. (1986). Mixed-function oxidase and Glutathion-s-transferase activity in asusceptible and Fenitrothion-resistant strain of *Oryzaephilus surimamensis*. *J. Econ. Entmol.* 79: 896-899.
- Roganovic, Z. D. and M. Jordanova (1998). "Liver lesions in bleak (*Aihurnus alburnus alborella* Filippi) collected from some contaminated sites on lake Ohrid. A histopathological evidence," *Ekol. Zast. Zivot. Sred.* 6: 11-18.
- Sayim, F. (2007). Dimethoate-induced biochemical and histopathological changes in the liver of rats. *Exp. Toxicol. Pathol.* 59: 237-243.
- Singh, R.N. R.K. Pandey N. Nath and V.K. Das (2009). Acute toxicity and behavioral Responses of Common Carp *Cyprinus carpio* (Linn.) to an Organophosphate (Dimethoate). *World Journal of Zoology* 4: 70-75.
- Sivapiriya, V. J. Karan and S. Venkatraman (2006) Effects of dimethoate (O,O-dimethyl S-methyl carbamoyl methyl phosphorodithioate) and Ethanol in antioxidant status of

- liver and kidney of experimental mice. *Pesticide Biochemistry and Physiology* 85:115–121.
- Tamimi, M. S. Qourzal. N.Barka. A. Assabbane and Y. Ait-Ichou (2008). Methomyl degradation in aqueous solution by Fenton's reagent and photo-Fenton system. *Separation and Purification Technology*. 61:103-108.
- Timur, S. S.Onal. N. O. Karabay. F. Sayim and F. Zihioğlu (2003). In vivo effects of malathion on glutathione-s-transferase and acetylcholinesterase in various tissues of neonatal rats. *Turk J. Zool.* 27: 247-252.
- Xiao, Q. Hu, B. Yu, C. Xia, L. Jiang, Z. (2006). Optimization of a single-drop microextraction procedure for the determination of organophosphorus pesticides in water and fruit juice with gas chromatography-flame photometric detection *Talanta* 69: 848–855.
- Yehia, M.A.H. S.G. El-Banna A.B. Okab (2007). Diazinon toxicity affects histophysiological and biochemical parameters in rabbits. *Exp. Toxicol. Pathol.* 59:215-225.

الملخص العربي

التأثيرات السامة لمبيد الملاثيون ومعالجة متبقيات في البيئة المائية

احمد حافظ مسعود – إسماعيل إبراهيم الفخراني - على سليمان درباله - مصطفى سمير
سعد الله

قسم المبيدات – كلية الزراعة - جامعة كفر الشيخ - جمهورية مصر العربية

أجريت هذه الدراسة لعمل حصر سريع واستكشاف وجود المبيدات الفوسفورية في مصادر المياه (شرب و صرف) بمحافظة كفر الشيخ . كذلك لدراسة التأثيرات السامة لحد أكثر المبيدات الفوسفورية اكتشفا في مصادر المياه و هو الملاثيون من خلال تأثيره على بعض الإنزيمات ((AChE و GOT و GPT و GST)) والتغيرات الهستولوجية التي يحدثها في الكبد و الكلى للفئران المعاملة بتركيز ١٠مجم/ كيلوجرام. أجريت هذه الدراسة أيضا لتقييم كفاءة بعض طرق المعالجة (الأكسدة المتقدمة – المعالجة البيولوجية) للتخلص من متبقيات الملاثيون في المياه. أوضحت النتائج أن الملاثيون كان من أكثر المركبات من حيث عدد مرات الاكتشاف و مستوى التركيز في مصادر المياه المختلفة. الملاثيون عند مستوى ١٠مجم/كجم وزن احدث تأثيرات سامة للفئران المعاملة سواء على مستوى التأثيرات البيوكيميائية أو الهستولوجية مقارنة بالكنترول. طريقة مشابهة الفينتون الضوئي (UV/H₂O₂/Fe⁺³) كانت اكفا طرق المعالجة في التخلص من مبيد الملاثيون في المياه. المعالجة البيولوجية لمبيد الملاثيون باستخدام عزلة ال *Pseudomonas sp* أدى الى التخلص من حوالي ٧٥ من متبقيات الملاثيون في المياه. هذه الدراسة توضح ان مبيد الملاثيون بالتركيزات المنخفضة يمكن أن يحدث تأثيرات سامه على صحة الإنسان. كما توضح الدراسة أيضا ان المعالجة البيولوجية باستخدام عزلة ال *Pseudomonas sp* يمكن استخدامها كطريقة معالجة فعالة وأمنة للتخلص من متبقيات الملاثيون في مياه الشرب بينما طريقة مشابهة الفينتون تعتبر مفضله للتخلص من متبقيات الملاثيون في مياه الصرف.