

INHERITANCE OF ADULT PLANT RESISTANCE TO LEAF RUST IN FOUR EGYPTIAN WHEAT CULTIVARS AND THEIR CROSSES WITH FOUR LEAF RUST RESISTANCE MONOGENIC LINES.

Youssef, I. A. M.

Wheat Disease Res. Dept., Plant Pathology Res. Inst., A.R.C. Giza Egypt

ABSTRACT

Leaf rust monogenic lines i.e. *Lr27*, *Lr29* were crossed with four Egyptian wheat cultivars (*Triticum aestivum* L.) also, *Lr46* was crossed with 3 wheat cultivars and *Lr2a* was crossed with one cultivar (Sakha 93). These parents, F1's and F2's were tested at adult plant stage under field conditions against race mixtures of the pathogen (*Puccinia triticina* Eriks.) under the stress of artificial infection. The parents of monogenic lines *Lr2a*, *Lr46*, *Lr29* and *Lr27* showed low disease severity. While, Gemmeiza 7, Sakha 61, Sakha 93 and Sids 1, showed high disease severity against leaf rust disease. The F1's tested plants showed low disease severity with most of crosses except three crosses i.e. *Lr27* + Sakha 61, *Lr27* + Sakha 93 and *Lr27* + Sids 1, which showed high disease severity. The F2 plants populations were segregated into two gene pairs. The dominance tend to the direction of low disease severity (partial leaf rust resistance) with eight crosses. However, the dominance tend to the side of high disease severity with three crosses that were previously mentioned with F1's. But no segregates was recorded with the cross *Lr2a* + Sakha 93 and dominance tend to the aside of low disease severity (partial leaf rust resistance). The cultivar Sakha 93 exhibited the adult plant resistance to the gene *Lr2a* under field conditions. These findings may prove that this gene is an effective under the Egyptian environmental conditions. The partial leaf rust resistance in the tested wheat cultivars was thought to be controlled by digenic pairs in adult plant stage. The selection for partial leaf rust resistant materials in the early generations was possible but delaying it to late ones is more effective, due to the important role of dominance effect in the expression of the character.

Keywords: *Puccinia triticina* Eriks. , Lrs:Leaf rust resistance genes,(L.D.S):Low disease severity,(H.D.S):High disease severity, A.P.R.:Adult plant resistance, P.R.:Partial resistance.

INTRODUCTION

Leaf rust incited by *Puccinia triticina* Eriks, is a worldwide disease of wheat. It causes an important yield loss in temperate regions where wheat is grown. Resistance to leaf rust in wheat cultivars had always been one of the main objectives in breeding programs. The use of resistance genes represents an effective procedure and environment-friendly way to control this disease in wheat.

Meanwhile, this approach demands, a constant effort to identify, characterize and incorporate resistance gene , mainly due to the great capability of rust populations to change host response (McCallum and Seto-Goh, 2005). Two slow-rusting resistance genes i.e. *Lr34* and *Lr46*, condition nonrace-specific adult plant resistant and have provide durable leaf rust resistance (Martines *et al.*, 2001) and Singh *et al.*, 1998). It was reported

about the likelihood that *Lr34* interacts with complementary genes *Lr27* and *Lr31* to produce enhanced resistance under conditions of high disease incidence. The role of *Lr34* in comparing durable resistance to leaf rust, seems to be similar to stem rust resistance gene *Sr2* (Sawhney 1992).

In Egypt, rust diseases are the most common and dangerous of wheat plant. leaf rust, in particular, was the cause of eliminating many cultivars i.e. Giza 139, Mexpake 69, Super x and Chenap 70 because of their susceptibility. It has been widespread on most cultivars of wheat grown to Egypt. After 1980, many cultivars having high level of partial resistance were produced as Giza 155 and Giza 157. Meanwhile, infection might be increased in the late sowing causing considerable losses in grain yield which reached 23% on some cultivars (Nazim *et al.*, 1983). Adult Plant Resistance (APR) was characterized by Zadoks (1961) as a resistance which is not expressed in the seedling stage and which develops with advancing plant stages. Partial Resistance (PR) is assumed to be stable because it is polygenically inherited (Parlevliet, 1978) and is temperature insensitive (Parlevliet, 1975). PR (Partial Resistance) is characterized by a reduced epidemic development despite a susceptible infection type (Parlevliet and Van Ommeren, 1975). The study in this investigation was carried out as an attempt to identify gene (s) for leaf rust resistance in wheat cultivars at adult plant stage.

MATERIALS AND METHODS

This investigation was carried out at the farm of Tag EL-Ezz agricultural research station Temie–Amdid Dakhlyia Egypt, aiming to identify gene(s) for leaf rust resistance in the wheat cultivars i.e. Gemmeiza 7, Sakha 61, Sakha 93 and Sids1 showing high infection type. Crosses were conducted between them and selected monogenic lines i.e. *Lr27*, *Lr29*, *Lr46* and *Lr2a* which displayed low infection type of leaf rust resistance Table (1).

The parental cultivars and monogenic lines were grown in 2008 /2009 growing season, in three successive dates at 15 days intervals to overcome differences in the time of flowering. The monogenic lines under study were used as male parents for crosses with each of commercial cultivars Table (1). Their parents were crossed and resulted in 12 cross. The parents were sown in 1.5 m long rows and 30 cm apart. Each row was sown to 15 seed with a distance of 10 cm. The experimental unit included 4 rows of each parent. The parents were selected according to their susceptibility or resistance on the basis of their reaction to leaf rust in the field during the elapsed growing seasons. Any doubtful of crosses hybrid seeds were discarded and the others were separately harvested.

In 2009/2010 growing season, part of the twelve (*Lr's* x *Cvs.*) cross hybrid seed were sown to produce F1's plants and the other part was kept for the final experiment in the next growing season. In 2010/2011 growing season, the evaluation of parents, F1's, and F2's, plant populations was done against natural and artificial infection with race mixtures of leaf rust pathogen caused by *P. triticina* Eriks. under field conditions. Twelve plots, each one included sixteen rows, one row for each parent and F1's as well as 13 row for

F2 plant populations. Each row with 2 m long, spaced 30 cm apart and seeds were 10 cm apart within row. Each row was planted with 20 seed. The adjacent plots were separated with 1 m wide belt. All plots were surrounded by a spreader area of one meter in width, planted with a mixture of highly susceptible cultivars to the leaf rust pathogen i.e. Giza139, Thatcher, Triticum Spelta Saharensis and Morocco .

For the inoculation , the spreader plants were moistened and dusted with spore – powder mixture of the most prevalent leaf rust races in Egypt (one volume of fresh urediospore mixtures : 20 volume of Talcum powder) . Dusting was carried out in the early evening at (sunset) before dew formation and when air was still in. Seedlings of the susceptible wheat cultivars were placed outside after emergence. The plants were transplanted neighbour of field plots as a race nursery (Zadoks, 1964) with difference that the genotypes were randomized. On both sides and perpendicular to the spreader row, placing about 200 infected plants of the susceptible cultivar in the spreader row, just before they started to sporulate.

The inoculation of the tested plants was carried out at booting stage according to the method suggested by Tervet and Cassel (1951). Data of leaf rust severity were recorded on the adult plant stage of tested plants according to the method of modified cob scale (Peterson *et al.*, 1948).

All regular cultural practices were applied during the growing season. Data were reported according to the technical recommendation as rust severity for each plant. Plants were divided into two classes according to level of rust severity i.e.(0-10 ,11-20,21-30-31-40) and (41-50-51-60,61-70,71-80). Plants grouped in the first four classes were considered as having low disease severity (Phenotypes), while other classes (more than 40%) were considered as having high disease severity (phenotypes) {Negm (2004) and Shahin (2005)}.

Statistical and genetic analysis : Frequency distribution values were computed for parental, F1 and F2 plant populations for leaf rust disease severity under field conditions. In respect to mode of inheritance, goodness of fit of the observed to the expected ratios of phenotypic classes concerning leaf rust infection types and disease severity were determined by X^2 analysis according to Steel and Torrie (1960).

However, the minimal number of effective genes controlling slow – rusting resistance in each cross were estimated by the formula of Wright (1968). Degrees of dominance were calculated according to the method suggested by Romero and Frey (1973). Heritability in broad - sense was estimated according to Lush (1949) as follows:

$$H^2 = \frac{VG}{VP} \times 100$$

Where :

H² = broad – sense heritability

VP = phenotypic variance of F2 individuals

VG = genotypic variance of F2 individuals

VE = environmental variance estimated from variation with the non-segregating populations, i.e. parent and F1 plants.

RESULTS

The present investigation was carried out to study the inheritance of leaf rust resistance in four Egyptian commercial wheat cultivars which were crossed with four wheat monogenic lines, these parents having different levels of high and low leaf rust disease severity.

Data presented in Tables (1 and 2) were qualitatively and quantitatively analyzed as follows : Evaluation of parents, F1 and F2 plant populations against race mixtures of *P. triticina* f.sp. tritici at adult plant stage . Data obtained in Table (1) reported that the monogenic lines parents i.e. *Lr2a*, *Lr46*, *Lr29*, and *Lr27* displayed low disease severity (partial leaf rust resistance). However, the Egyptian commercial wheat cultivars parents showed high disease severity (susceptible) fast-rusting. The twelve crosses could be arranged into two categories i.e. low disease severity (L.D.S.) and high disease severity (H.D.S.). The F1's tested plants exhibited the same trend with one parent and showed low disease severity (partial leaf rust severity) with nine crosses. However, the late three crosses recorded high disease severity i.e. *Lr27*+ Sakha 61, *Lr27* + Sakha 93 and *Lr27* + Sids 1.

Meanwhile, the F2 plant populations showed that one cross out of twelve showed no segregates i.e. (*Lr2a* + Sakha 93), the dominance tend to the side of partial leaf rust resistance. Meanwhile, the rest of crosses displayed segregation with numbers of plant with low and high disease severity as clarified in Table(1) i.e. 113:97 ,170:46, 200:17, 45:164, 167:48, 200:18, 47:167, 162:47, 45:158, 191:17 and 162:45, respectively.

These observed ratios fitted the theoretical expected ratios i.e. 9:7, 13:3, 15:1, 3:13, 13:3,15:1, 3:13, 13:3, 3:13, 15:1, and 13:3 with probable values 0.500. 0.250, 0.500. 0.250, 0.500.0.250, 0.500.250, 0.250.0.100, 0.250.0.100, 0.250.0.100, .0.250.0.100, 0.250.0.100, 0.250.0.100 and 0.500.0.250, respectively, Table (1) .

Quantitative analysis : The genetic behaviour of Egyptian commercial wheat cultivars to leaf rust quantitatively, the parents, F1's and F2's plant population for each of the twelve crosses were tested at adult plant stage under field conditions against race mixtures of the pathogen *P. triticina* is clarified in Table (2). Data obtained in this table explored that the disease severity (%) recorded with eight parents i.e. *Lr27*, *Lr29*, *Lr46*, *Lr2a*, Gemmeiza 7, Sakha 61, Sakha 93 and Sids 1 were : 34.0, 24.5, 6.0, 6.0; 66.0, 64.5, 64.0 and 65.5, respectively. The F1 and F2 means showed values lower than those calculated for their respective mid-parents, revealing the presence of partial dominance for low disease severity (partial leaf rust resistance) which is in accordance with the results obtained from F1's and F2's Table (2).

Expression of gene action measured as the degree of dominance h_1 and h_2 is shown in Table (2). The estimated values of h_1 and h_2 exhibited negative values in these twelve crosses suggesting the manifestation of partial dominance for leaf rust resistance and this is supported the F1's results .

Table (1): Leaf rust severity (%) frequency distributions of the two parents, F1 and F2 plant populations. Phenotype classes, expected ratios, X² and probable values of F2 plant population of 12 (Lr's x Commercial cultivars) crosses as affected by inoculation with race mixtures of wheat leaf rust (*P. triticina*) at adult plant stage under field conditions in 2010/2011 grown seasons.

No.	Cross name	No. of tested plants	Disease severity classes %								Observed ratio		Expected ratio		X ²	Probable values
			Low				High				Low	High	Low	High		
			0-10	11-20	21-30	31-40	41-50	51-60	61-70	71-80						
	Low+High															
1	Lr27	P1	20			2	18									
	Gemmeza 7	P2	20							18	2					
		F1	20			1	19									
		F2	212	5	18	40	50	48	32	19		113	97	9	7	0.482
2	Lr29	P1	20			1	19									
	Gemmeza 7	P2	20							18	2					
		F1	20			3	17									
		F2	219	10	30	50	80	32	9	5		170	46	13	3	0.918
3	Lr46	P1	20	18	2											
	Gemmeza 7	P2	20							18	2					
		F1	20			2	18									
		F2	217	21	39	120	20	14	3			200	17	15	1	0.930
4	Lr27	P1	20			2	18									
	Sakha 61	P2	20							1	19					
		F1	20							19	1					
		F2	209	3	10	15	17	65	55	44		45	164	3	13	1.059
5	Lr29	P1	20			1	19									
	Sakha 61	P2	20							1	19					
		F1	20			2	18									
		F2	215		20	60	87	30	18			167	48	13	3	1.805
6	Lr46	P1	20	18	2											
	Sakha 61	P2	20							1	19					
		F1	20			18	2									
		F2	220	23	26	111	40	16	2			200	18	15	1	1.517
7	Lr2a	P1	20	18	2											
	Sakha 93	P2	20							2	18					
		F1	20			19	1									
		F2	216	35	130	23	20					208	0	1	0	0
8	Lr27	P1	20			2	18									
	Sakha 93	P2	20							2	18					
		F1	20							19	1					
		F2	214		10	37	89	61	17			47	167	3	13	1.448
9	Lr29	P1	20			1	19									
	Sakha 93	P2	20							2	8					
		F1	20			2	18									
		F2	212		22	60	80	30	9	8		162	47	13	3	1.910
10	Lr27	P1	20			2	18									
	Sids 1	P2	20							19	1					
		F1	20							18	2					
		F2	213		6	11	28	79	66	13		45	158	3	13	1.537
11	Lr29	P1	20			1	19									
	Sids 1	P2	20													
		F1	20			3	17									
		F2	208	26	35	57	73	11	6			191	17	15	1	1.389
12	Lr46	P1	20	18	2											
	Sids 1	P2	20							19	1					
		F1	20			3	17									
		F2	204		14	84	64	35	10			162	45	13	3	1.219

Low = low disease severity (resistance)
High = high disease severity (susceptible)

The heritability values for all tested crosses at adult plant stage are considered to be high i.e. 96.64, 94.5, 91.25, 97.02, 94.39, 93.1, 88.25, 91.85, 94.36, 93.62, 94.97 and 90.82, respectively, Table (2).

Table (2):. Mean of P1, P2, F1, F2 and mid-parents, degree of dominance of F1 and F2 as well as broad-sense heritability and number of genes for rust severity (%) of 12 (*Lr*'s x Cvs.) crosses inoculated with race mixtures of *P. triticina* under field conditions in 2010/2011 growing season.

No.	Cross name	Mean of rust severity (%)					Degree of dominance		Heritability	No. of genes
		P1	P2	F1	F2	MP	h1	h2		
	Low + High									
1	<i>Lr27</i> + Gemmeiza 7	34.0	66.0	34.5	38.6	50.0	-1.6	-2.37	96.64	0.578
2	<i>Lr29</i> + Gemmeiza 7	24.5	66.0	33.5	31.53	45.25	-0.566	-1.322	94.5	1.455
3	<i>Lr46</i> + Gemmeiza 7	6.0	66.0	24.0	23.89	36.0	-0.4	0.807	91.25	4.913
4	<i>Lr27</i> + Sakha 61	34.0	64.5	45.5	47.58	49.25	-0.245	0.219	97.02	0.575
5	<i>Lr29</i> + Sakha 61	24.5	64.5	34.0	33.49	44.5	-0.525	-1.1	94.39	1.97
6	<i>Lr46</i> + Sakha 61	6.0	64.5	16.0	25.27	35.25	-0.658	-0.682	93.107	4.234
7	<i>Lr2a</i> + Sakha 93	6.0	64.0	15.0	16.35	35.0	-0.643	-1.286	88.25	7.034
8	<i>Lr27</i> + Sakha 93	34.0	64.0	45.5	46.78	49.0	-0.23	-0.296	91.85	1.273
9	<i>Lr29</i> + Sakha 93	24.5	64.0	34.0	33.5	44.25	-0.52	-1.09	94.36	1.55
10	<i>Lr27</i> + Sids 1	34.0	65.5	46.0	46.2	49.75	-0.238	-0.451	93.62	1.13
11	<i>Lr29</i> + Sids 1	24.5	65.5	33.5	26.25	45.0	-0.561	-1.83	94.97	1.56
12	<i>Lr46</i> + Sids 1	6.0	65.5	23.5	32.25	35.75	-0.412	-0.235	90.82	5.298

Lr's = Leaf rust resistance genes

C.v. = Commercial variety

h1 and h2 = Degree of dominance of F1 and F2 plant populations

MP = Mid-parents

Low = Low disease severity (resistance)

High = High disease severity (susceptible)

Number of genes: The minimum number of effective genes controlling the partial leaf rust resistance or susceptibility was digenic for each of the tested crosses except for the cross number seven (*Lr2a* + Sakha 93) that exhibited no segregation. The estimated numbers were tabulated in Table (2).

DISCUSSION

Wheat rusts are the most important diseases of wheat cultivars in Egypt. Leaf rust, in particular, was the main cause of eliminating many cultivars i.e. Giza 139, Mexpak69, Super x and Chenap70, because of their susceptibility under field conditions. Moreover, some wheat cultivars were discarded very shortly after their release such as Giza 139 (Gomma 1978). The failure of such cultivars was mainly due to the dynamic nature in population of the causal organism which produces new virulence having the ability to breakdown their resistance. However, the use of effective resistance genes is the most economic and desirable method for controlling the disease. Such information is very useful in efficient incorporation of different genes into cultivars for a long-lasting resistance. The Egyptian wheat variety

Giza155 had high level of partial resistance under field conditions (Nazim *et al.*, 1983). They added that , the good method to evaluate a wheat variety to leaf rust, is to test under different epiphytotic levels and different locations. Intercrosses among slow-rusting parents also were made to investigate the genetic diversity of this type of resistance (Herrera Fössel *et al.*, 2007). Singh *et al.*, (2000) mentioned that high yielding cultivars of bread wheat that were nearly immune to leaf rust and stripe rust could be developed by accumulating four or five slow-rusting resistance genes through intercrossing parents that show intermediate disease levels. Also, they pointed to several slow-rusting durum's identified being utilized to develop high yielding cultivars with high levels of slow-rusting resistance to leaf rust.

This investigation represents the evaluation of four monogenic lines i.e. *Lr27*, *Lr29*, *Lr46* and *Lr2a*; and four Egyptian commercial wheat cultivars i.e. Gemmeiza 7, Sakha 61, Sakha 93 and Sids 1 under field conditions during elapsed growing seasons was carried out. All of these parents (*Lr*'s and C.v.) showed susceptibility to leaf rust disease all over the seasons. However, *Lr27*, *Lr29*, *Lr46* and *Lr2a* exhibited different levels of low rust severity (adult plant resistance), while Gemmeiza7, Sakha 61, Sakha 93 and sids1, showed high disease severity to leaf rust disease. The obtained results explored that most of the F1's plants displayed low disease severity (partial leaf rust resistance) slow-rusting except for three crosses i.e. *Lr27* + Sakha 61, *Lr27* + Sakha 93 and *Lr27* + Sids 1 which showed high disease severity. On the other hand, F2's plant populations with all crosses behaved in the same trend with the F1's plants. Results of the present study demonstrated that all of the tested F2 plant populations of the cross between *Lr2a* + Sakha 93 displayed low rust severity with no segregations indecating that this cultivar possess this gene. These findings showed that this gene is effective under the Egyptian environmental conditions. Therefore, it may be transferred to the Egyptian genetic materials as a leaf rust resistance source in breeding programs. Oelke and Kolmar (2005) mentioned that wheat cultivar Alsen proved to have genes *Lr2a*, *Lr10*, *Lr13*, *Lr23* and *Lr34*. Effective leaf rust resistance in this cultivar is conditioned by the combination of *Lr13*, *Lr23* and *Lr34*. Also, they added that virulence to *Lr2a* also very common. McIntosh and Dyck (1972) showed that in some crosses *Lr23* was recessive, while in other crosses it was partially dominant. The dominance effect of *Lr23* increased at higher temperature. Also they mentioned that spring wheat cultivars grown in Minnesota and Dakota may have a fairly complex inheritance of leaf rust resistance based on the genetics of Alsen and Norm. Meanwhile, the rest of crosses were segregated with digenic pairs. Results of the present study demonstrated that the F2 plants of the cross between *Lr27* and wheat cultivar Gemmeiza7 showed complementary gene action with the dominance of resistance over susceptibility. The additive gene action (duplicate dominance) was shown with crossess *Lr46* + Gemmeiza 7, *Lr46* + Sakha 61 and *Lr29* + Sids 1 with dominance of resistance. Also, the inhibitory gene actions were reported with *Lr29* + Gemmeiza 7, *Lr29* + Sakha 61 and *Lr29* + Sakha 93 and *Lr46* + Sids1, the resistance was dominant over susceptibility. However, the susceptibility was dominant and resistance was recessive i.e. *Lr27* + Sakha 61, *Lr27* + Sakha 93 and *Lr27* + Sids 1. Similar

results were recorded by Sawhney (1992) who mentioned that the leaf rust responses of wheat lines carrying the complementary genes *Lr27* and *Lr31* and some genes in a Chinese spring background having *Lr34*, which interacts with the complementary genes gave enhanced levels of field resistance to leaf rust. Likewise, Dyck and Samborski (1982) and Ezzahiri and Roelfs (1985) showed that resistance was controlled by two complementary genes. They added that no segregation was recorded for adult plant resistance in the F₂ families derived from a cross between the wheat cultivars Era and Frontana having adult plant resistance gene *Lr13*. Similar results were reported by Knott and Yadav (1993) who found that in eight lines, the resistance was recessive or partially recessive and was controlled by two or more genes in each line. Hussain *et al.*, (1998) determined the leaf rust resistance genes in four wheat varieties, these varieties exhibited field resistance, probably due to additive effect of adult plant resistance genes. The durable resistance mechanism in wheat is achieved through incorporation of partially resistant minor genes which seems to be more appropriate solution for susceptible wheat production. The obtained results were in accordance with the findings of Kuhn *et al.*, (1980); Bjarko and line (1988); Das *et al.*, (1993); Boulat (1997); Negm (2004) and Shahin (2005) who mentioned that the quantitative analysis demonstrated that the F₂ means of leaf rust severity in the eleven crosses were in general, lower than the estimated means for their respective mid-parents. These results clarified the existence of partial dominance for low disease severity in all crosses. Also, the degree of dominance (h_1 and h_2), were significant and negative in all of the tested crosses. Similar results were confirmed by those obtained with Millus and line (1986) Shehab El Din *et al.*, (1991a, 1991b and 1996); Youssef *et al* (2007).

The heritability in its broad-sense estimated from parents, F₁'s and F₂'s for partial leaf rust resistance is considered to be high in magnitude, since values ranged from 88.25 to 97.02.

Meanwhile, high heritability values are indicative for high rates of success in recovering the desired genes in future generations. These high estimates indicates that selection for this character in early segregating generations could be possible. However, delaying it would be more effective. The obtained results were in accordance with those of Bjarko and Line (1988), Das *et al.*, (1993); Abd-EL-Latif *et al.*, (1995); Boulot (1997); Negm (2004) and Shahin (2005). These findings would be of great importance in wheat breeding programs from the view point of disease resistance.

REFERENCES

- Abd-EL-Latif, A.A.; T.M. Shehab-EL-Din; M.M. EL-Shami and S.A. Abo-EL-Naga (1995). Genetics of *Triticum aestivum* : *Puccinia recondita* tritici interaction "in three Egyptian wheat cultivars ". J. Agric. Res. Tanta Univ. 21:182-188
- Bjarko, M.F. and R.F. line (1988). Quantitative determination of the gene action of leaf rust resistance in four cultivars of wheat (*Triticum aestivum*). Phytopath. 78:451-456

- Boulot, O.A. (1997). "Further studies on leaf rust of wheat" Ph.D. Thesis, Plant Pathology. Minufiya Univ. 290 p.
- Das, M.K. ; S. Rajaram; W.K. Ktonstad; C.C. Mundt and R.P. Singh (1993). Associations and genetics of three components of slow-rusting in leaf rust of wheat. *Euphytica* 68 (112):99-109.
- Dyck, P.L. and D.J. Samborski (1982). The inheritance of resistance to *Puccinia recondita* in a group of common wheat cultivars. *Can. J. Genet. Cytology* 24:273-283 .
- Ezzahiri , B. and A . P . Roelfs (1985). Inheritance of adult plant resistance to leaf rust in Era wheat. *J. Phytopathology* 75:1317.
- Gomma, A.S.A. (1978). Report on the Egyptian wheat varieties and their development. Egyptian Soc. of Genetics Science. Cairo.
- Hussain, M.; M.H. Chaudhry; J.A. Shah and M. Younus (1998). Genetic diversity to *Puccinia recondita* f. sp. tritici in 59 wheat lines. *Pakistan J. Phytopathology*, 10:113-121 .
- Herrera – foessel, S.A.; R.P. Singh; J. Huerta – Espino; J. Cross; A. Djurle and J. Yuen (2007). Evaluation of slow-rusting resistance components to leaf rust in CIMMYT durum wheat. *Euphytica* 155:361-369.
- Knott, D.R. and B. Yadav (1993). The mechanism and inheritance of adult plant leaf rust resistance in 12 wheat lines. *Genome* 1993.36:877-883.
- Kuhn , R.C.; H.W. Ohm and G.E. Shaner (1980). Inheritance of slow leaf rusting in suwon 85 wheat *Crop Sci.* 20:655-659.
- Lush, J.L. (1949). Heritability of quantitative characters in farm animals. *Hereditas*, Supp . Vol . pp. 356-375. (cited after Weber *Agro.j.*44:202-209).
- Martinez, F.; R.E. Nicks; A. Moral; J.M. Urbano; D. Rubiales and P. Hermende (2001). Search for partial resistance to leaf rust in a collection of ancient Spanish wheat. Proceeding of the fourth International Triticale Symposium. Cordoba, Spain. September 2001 *Hereditas – Lund* 2001,135:193-197.
- McCallum, B.D. ; P. Seto – Goh (2005). Physiologic specialization of wheat leaf rust (*Puccinia triticina*) in Canada in 2002. *Can. J. Plant Pathol.* 27:90-99.
- McIntosh, R.A. and P.L. Dyck (1972). Cytogenetical studies in wheat. V11. Gene *Lr23* for reaction to *Puccinia recondita* in Gabo and related cultivars. *Aus. J. Bio. Sci.* 25:765-773.
- Mullus, F.A. and R.F. line (1986). Gene action for inheritance of durable high-temperature, adult plant resistance to stripe rust in wheat. *Phytopathology*, 76:435-441.
- Nazim, M.; A.A. El- Shehidi; Y.A. Abdou and Y.H. El-Daoudi (1983). Yield losses caused by leaf rust on four wheat cultivars under epiphytotic levels. 4th confer. *Microbiol.*, Cairo, 1983, 17-27.
- Negm, S.S.M. (2004). Partial resistance to leaf rust in some Egyptian wheat varieties. Ph.D. Thesis, Faculty of Agric. Minufiya Univ. pp.192.
- Oelke, L.M., and J.A. Kolmer (2005). Genetics of leaf rust resistance in spring wheat cultivars Alsen and Norm. *Phytopathology*. 95:773-778.

Youssef, I. A. M.

- Parlevleit, J.E. (1975). Partial resistance of barley to leaf rust *Puccinia hordei*. I. Effect of cultivars and development stage on latent period. *Euphytica*. 24:21-27.
- Parlevleit, J.E. (1978). Further evidence of polygenic inheritance of partial resistance in barley to leaf rust *Puccinia hordei*, *Euphytica*. 27:369-379
- Parlevleit, J.E. and A. Van Ommeren (1975). Partial resistance of barley to leaf rust, *Puccinia hordei*. II. Relationship between field trials. Microplot tests and latent period. *Euphytica* 24:293-303
- Peterson, R.F.; A.B. Campbell and A.E. Hannah (1948). A diagrammatic scale for estimating rust intensity on leaves and stems of cereals. *Can. J. Res.* 60:496-500.
- Romero, G.E. and K.J. Frey (1973). Inheritance of semi dwarfness in several wheat crosses. *Crop. Sci.* 13:334-337.
- Sawhney, R.N. (1992). The role of *Lr24* in imparting durable resistance to wheat leaf rust through gene interactions. *Euphytica* 61:9-12.
- Shahin, S.I.M. (2005). Resistance of Gemmeiza wheat varieties to leaf rust disease Ph.D. Thesis, faculty of agric. Minufiya Univ. pp.233.
- Shehab El. Din, T.M. and A.H. Abdel-Latiff (1996). Quantitative determination of the gene action of stripe rust resistance in A. parent diallel cross of wheat. *J. Agric. Sci. Mansoura Univ.*, 10:3461-3467.
- Shehab El-Din T.M.; S.A. Abd-Alla; G. El- Fadly and A.H. Abd-El-Latiff (1991a). Genetics of *Triticum aestivum*, *Puccinia graminis* tritici interaction. *J. Agric. Res. Tanta Univ.* 18:426-437.
- Shehab El-Din, T.M.; M.A. Gouda; S.A. Abo El-Naga and M.M. El-Sami (1991b). Quantitative study on wheat resistance to stem rust caused by *Puccinia graminis* tritici. *J. Agric. Sci. Mansoura Univ.* 15:1298-1303.
- Singh. R.P.; A.M. Kazi and J.H. Espino (1998). *Lr46* Agene conferring slow-rusting resistance to leaf rust in wheat. *Phytopath.* 88 -890-894.
- Singh, R.P.; J. Huerta-Espino; S. Rajaram (2000). Achieving near-immunity to leaf and stripe rusts in wheat by combining slow-rusting resistance genes. *Acta Phytopath. Entomol. Hung.* 35:133-139.
- Steel, R.G.D. and T.H. Torrie (1960). Principles and procedures of statistics Mc-Graw Hill, N.Y., USA.
- Tervet, I. and R.C. Cassel(1951). The use of cyclone separation in race identification of cereal rusts. *Phytopath.* 41:282-285.
- Wright, S. (1968). Evolution and genetics of populations. vol .I. genetics of biometric foundations. University of Chicago Press. Chicago and London. 469 pp.
- Youssef, I.A.M.; Gamalat, A. Hermas and M.S. Hamada (2007). Inheritance of resistance to leaf rust in 13 wheat Near – isogenic lines. *J. Agric. Sci. Mansoura, Univ.*, 32(8):6315-6328.
- Zadoks. J.C. (1961). Yellow rust of wheat : studies in epidemiology and physiologic specialization. *Tijdschr. Plantenziekten.* 67:69-256.
- Zadoks. J.C. (1964). The use of race nurseries in cereal resistance breeding. P. 242-249. In : S. Broekhuizen, G. Dantuma, H. Lamberts and W-Lange (Eds). *Barley Genetics I.* Pudoc, Wageningen.

وراثة المقاومة في النبات البالغ ضد مرض صدأ أوراق القمح في الهجن الناتجة عن أربعة أصناف مصرية تجارية وأربع سلالات خاصة بالمقاومة لصدأ الأوراق.

عصام عبد الحميد محمد يوسف

قسم بحوث أمراض القمح - معهد بحوث أمراض النباتات - مركز البحوث الزراعية - الجيزة - مصر.

تم إجراء عدد من التهجينات بين سلالات أحادية الجين مقاومة للصدأ البرتقالي مع أصناف قمح تجارية حيث هجنت السلالتان الأحادية الجين من صدا الورقة *Lr27*, *Lr29* مع أربعة أصناف من القمح (*Triticum aestivum*, L.). وأيضا هجنت السلالة *Lr46* مع ثلاثة أصناف قمح بينما تم تهجين السلالة *Lr2a* مع الصنف سخا ٩٣ فقط. وقد تم اختبار كل من الأباء والجيل الأول والجيل الثاني في مرحلة النبات البالغ وذلك تحت ظروف الحقل ضد مخلوط من سلالات مسبب المرض لصدأ الأوراق (*Puccinia triticina*, Eriks.).

وقد تم تسجيل شدة المرض المنخفضة مع السلالات الأحادية الجين *Lr27*, *Lr29*, *Lr46*

and *Lr2a*,

ومع ذلك فقد أظهرت الأصناف التجارية المصرية مميزة ٧ سخا ٦١ و سخا ٩٣ سيدس ١ استجابة عالية لشدة المرض. اختبار الجيل الأول أوضحت شدة المرض المنخفضة مع معظم الهجن باستثناء ثلاث هجن أوضحت شدة المرض العالية وهى *Lr27 + Sakha 61*, *Lr27 + Sakha 93* and *Lr27 + Sids 1*. اختبار العشيرة في الجيل الثاني والتي تؤكد الاتعزال في زوجين من الجينات مع ثمانية هجن حيث تميل السيادة تجاه شدة المرض المنخفضة (مقاومة صدا الورق الجزئية). بينما تميل السيادة إلى شدة المرض العالية مع ثلاثة هجن والسبق الإشارة إليهم في الجيل الأول ولكن غياب الاتعزال تم تسجيله مع الهجن *Lr2a + Sakha 93*. وسيادة شدة المرض المنخفضة (المقاومة العالية) للصنف سخا ٩٣ ربما يملك *Lr2a* جين مقاومة النبات البالغ وذلك تحت ظروف الحقل. هذا التواجد يؤكد أن هذا الجين مؤثر تحت ظروف الحقل المصرية. وقد تبين أن مقاومة صدا الورقة الجزئية في أصناف القمح المصرية المختبرة في هذا البحث كانت محكومة بزوجين من الجينات وذلك في معظم الهجن في مرحلة النبات البالغ. ومن هنا فإن الانتخاب من أجل المقاومة الجزئية في الأجيال المبكرة من الممكن ولكن التأخير في المراحل المتأخرة يكون أكثر تأثيراً حيث وجد أن الدور الهام في التعبير عن هذه الصفة يرجع إلى تأثير السيادة.

قام بتحكيم البحث

كلية الزراعة بدمياط - جامعة المنصورة

أ.د / محمد سعد إبراهيم حمادة

كلية الزراعة - جامعة كفر الشيخ

أ.د / صلاح الدين عبد الحميد أبو النجا