

## **EFFECTS OF SEAWEED SUPPLEMENTATION TO FATTENING FRIESIAN STEERS RATIONS ON: 2- BLOOD PARAMETERS, PERFORMANCE AND FEED EFFICIENCY**

**Ead, H. M. E.<sup>1</sup> and Eman H. M. Maklad<sup>2</sup>**

**1- Animal Production Res. Inst., Agric. Res. Center, Dokki, Giza, Egypt.**

**2-Animal Production Dept., Fac. Agric., Mansoura University, Egypt.**

### **ABSTRACT**

Twelve male Friesian steers with average body weight  $269 \pm 9$  kg and 15 month of age were used in this study. Steers were distributed into three groups similar in number. The experimental rations were formulated as follows: ration 1 (R1) 81.8% concentrate feed mixture (CFM) + 18.2% clover hay (CH) (control), ration 2 (R2): 81.55% CFM + 18.05% CH + 0.40% seaweed (S) and ration 3 (R3): 81.23% CFM + 17.87% CH + 0.90% S.

The results obtained showed that the mean values of blood parameters were not significantly affected by the feeding on R1 or R2 or R3. The concentration of urea-N in blood was decreased with feeding on R2 and R3 than feeding on R1.

The average daily gains were 0.81, 1.00 and 0.83 kg/d for steers fed on R1, R2 and R3, respectively. The production efficiency was higher when feeding on R2 (24.04 %) than feeding on R1 or R3 (20.97 and 19.17%, respectively), and the economic efficiency, was higher when feeding on R2 (25.39 %) than feeding on R1 or R3 (13.3 and 3.94 %, respectively).

In general, the results indicated that feeding ration at a level 81.55% CFM + 18.05% CH + 0.4% S for fattening Friesian steers was economically improved animal performance compared with the other two rations.

**Keywords:** Friesian steers, seaweed, daily gain and economic efficiency.

### **INTRODUCTION**

Beef production is a large and important segment of animal production and one of the largest industries in the world. Cattle usually weight 270 to 310 kg before they are placed on high grains (high energy) ration. This diet is fed until slaughter weight is achieved. An animal that is gaining weight at a moderate rate needs about 1.5 % of their body weight in concentrates per day.

Rapidly growing cattle, such as calves can be safely fed up to 2.0 to 2.25 % of their weight in concentrates and feeding roughage at least 1.8 to 2.2 kg of hay daily. However, neither should be fed at over 20 % of the diet (Schreder, 2002). Finishing cattle in the terminal stage of fattening are often fed diets with low concentrations of roughage to maintain performance and achieve targeted back fat finish and marbling that desired by beef industry. High energy fattening rations have 70 % TDN and 9 to 15 % CF (Mandell *et al.*, 2001).

Buffer supplementation of high concentrate diets has been shown to improve or stabilize food intake and increase animal performance in some studies (Zinn, 1991), but not in others (Ghorbani *et al.*, 1989).

Aga *et al.* (2000) used a calcified seaweed as a buffer in continuous culture of rumen content. The chemical composition of an ordinary seaweed from *Ascophyllum nodosum*, immediately characterized the material as of low energy content. According to the analytical data, the value of seaweed meal must primarily be sought in its content of vitamins and minerals, among which  $\beta$ -carotene, tocopherols, some B-vitamins, iodine, zinc and potassium are the more important (Scott, 1990).

The most frequently reported benefits associated with inclusion of *Ascophyllum nodosum* in animal diet include increased weight gain and improved carcass quality. Slaughter yield improves with age, for a given carcass weight, animals at high growth rate, receiving more balanced feed and generally have a better carcass yield (Lebas and Colin, 1992).

The objective of this study was to evaluate the effect of seaweed supplementation to fattening steers rations on some blood constituents, growth performance and economic efficiency.

## **MATERIALS AND METHODS**

This study was conducted at El-Karada Animal Production Research Station, Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture in Egypt and Department of Animal Production, Fac. of Agric., Mansoura University.

### **Experimental animals:**

Twelve male Friesian steers with average live body weight  $269 \pm 9$  kg and 15 month of age were distributed into three similar groups (Four for each). The

### **Three experimental rations were formulated as follows:**

R1: ration 1: 81.8% concentrate feed mixture (CFM) + 18.2% clover hay (CH) (as a control ration).

R 2: ration 2 : 81.55% CFM + 18.05% CH + 0.40% seaweed (S)

R 3: ration 3: 81.23% CFM + 17.87% CH + 0.90% S.

The CP concentration of tested rations were ranged from 16 to 17% according to Hunter *et al* (1999). Steers were individually fed the experimental rations to cover the requirements of fattening steers and were adjusted monthly according to their body weight changes. Experimental period extended for five months.

The concentrate feed mixture (CFM) used contained 44% yellow corn , 23% soybean meal (44% protein), 14% wheat bran, 11.5% rice bran , 4.5% molasses, 2%, limestone and 1% salt. The clover hay was made from the 3<sup>rd</sup> cut of Egyptian clover.

The supplement seaweed meal as from *Ascophyllum nodosum* manufactured by Acadian Sea plants Limited, Canada . The approximate label analysis showed that it contains of protein, fiber, carbohydrates, vitamins and minerals.

### **Management of feeding:**

The intake of tested ration by animals was fixed and calculated as the percentage of roughage to concentrate ratio to satisfy their maintenance and production requirements (Ghoneim, 1967).

The CFM fed with or without seaweed was offered to steers at morning. While, clover hay (CH) given after consumption of the concentrate. Drinking fresh and clean water was available at all times.

**Weighing procedure:**

Animals were weighed in the morning before drinking and feeding at the beginning of the trial and monthly thereafter to the nearest kg for each animal.

**Chemical analysis:**

Blood samples were taken from each animal individually during the digestion trials of the tested rations. These samples were taken at 3 hrs post-feeding from jugular vein. Blood samples were immediately separated by centrifugation at 4000 rpm for 10 minutes. The serum samples were stored at (-20°C) until analysis were done. The analysis included total protein (Gornall *et al.*, 1949); albumin (Hill and Wells, 1983); globulin (calculated by differences between the total protein and albumin concentrations); urea (Freidman *et al.*, 1980); creatinine (Ullmann, 1976); Glucose (Teuscher and Richterich, 1971); GOT and GPT (Reitman and Frankel, 1957).

**Production efficiency :**

The ME can be converted to a NEm requirement with an efficiency of 0.576 (NRC, 1996) and NEp will equal (ME-NEm).

The retained energy (RE, Mcal/d) = (live weight<sup>0.2955</sup> × 0.544) × (ADG)<sup>1.262</sup>

Where ADG is in kilograms (Overton, 1999).

Production efficiency = RE/NE<sub>p</sub> × 100

**Economic efficiency :**

Economic efficiency was calculated according to the following formula:

Economic efficiency =  $\frac{(\text{Price of daily gain} - \text{Daily feed cost})}{\text{Daily feed cost}}$

**Statistical analysis:.**

The statistical analysis was performed using the least squares method described by Likelihood programmer of SAS (1994). The obtained data for performance and blood parameters were subjected to one way analysis of variance according to the following model:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where:

Y<sub>ij</sub> = Observation of the tested factor

μ = Overall mean

T<sub>i</sub> = Treatment effect

e<sub>ij</sub> = Error

The differences among means were carried out according to Duncan's New Multiple Range Test (Duncan, 1955).

## **RESULTS AND DISCUSSION**

Concerning blood metabolism, data in Table (1) shows that the values were in the normal range as described by Mohamed and Selim (1999), but without significant effect.

The concentration of urea-N in blood was decreased with feeding on R2 and R3 than feeding on R1. The concentration is affected not only by dietary intake of digestible crude protein in the rumen but also by balance between energy and protein in the diet (Hoffman and Steinhöfel, 1990).

Table (2) shows the effect of feeding tested rations on average body weight (ABW) and average daily gain (ADG). There were no significant effect on the average body weight appeared among tested group. However, the (ADG) was higher ( $p < 0.05$ ) from 17-18 month when animals were fed on R2 or R3 than R1, and then increased ( $p < 0.05$ ) from 19-20 month when fed on R2 than R1 or R3. The average daily gains for whole experimental period were 0.81, 1.00 and 0.83 kg/day for R1, R2 and R3, respectively.

**Table (1): Effect of experimental rations on some blood parameters.**

Item	Experimental rations				
	Ration 1	Ration 2	Ration 3	+SEM	p
Total protein g/100ml	6.49	6.45	5.63	0.300	0.1523
Albumin g/100 ml	4.93	5.42	4.93	0.293	0.4399
Globulin g/100 ml	1.56	1.03	0.70	0.259	0.1363
Creatinine mg/100 ml	0.18	0.31	0.26	0.479	0.2484
Urea-n mg/100 ml	23.76	20.53	20.95	1.873	0.4623
GOT IU/L	61.30	60.30	58.60	5.385	0.9399
GPT IU/L	10.00	12.70	12.60	1.586	0.4410
Glucose (mg/100 ml)	64.64	54.98	53.63	3.95 <sup>a</sup>	0.1805

**Table (2): Effect of feeding tested ration on average body weight (kg) and daily gain (kg).**

Item	ABW					ADG				
	R1	R2	R3	+SEM	p	R1	R2	R3	+SEM	p
Initial BW 15 month	261	271	274	9.33	0.6051	-	-	-	-	-
15-16 month	296	310	310	11.70	0.6405	1.17	1.30	1.19	0.391	0.643
16-17 month	311	322	322	11.28	0.7410	0.50	0.40	0.40	0.066	0.494
17-18 month	331	368	363	12.54	0.1316	0.67 <sup>b</sup>	1.53 <sup>a</sup>	1.38 <sup>b</sup>	0.192	0.024
18-19 month	363	393	382	15.22	0.3910	1.05	0.84	0.63	0.150	0.201
Final BW 19-20 month	383	421	398	15.90	0.2929	0.70 <sup>b</sup>	0.92 <sup>a</sup>	0.53 <sup>b</sup>	0.066	0.008
Mean 15-20 month	-	-	-	-	-	0.81	1.00	0.83	-	-

Table (3) shows that there was no significant effect of DMI between the treatments however, the DMI was increased when fed on R3 than R1 or R2, while the DMI increase ( $p < 0.05$ ) when feeding on R2 than R1, but the production efficiency was higher with feeding on R2 ((24.04%) than feeding on R1 or R3 (20.97 and 19.17%, respectively).

The price of the total fresh feed intake/day was higher when feeding on R3 than feeding on R1 or R2 and the price of R2 was higher than R1 as shown in Table (4). Feeding on R2 was more economic efficiency (25.39 %) than feeding on R1 or R3 (13.35 and 3.94%, respectively).

**Table (3): Production efficiency of growing steers fed the experimental rations.**

Item	R1	R2	R3	+SE	P
DMI (kg/d)	9.99	10.03	10.07	0.000	0.0001
ME (Mcal/kg)	3.02	3.04	3.03	0.5746	0.5504
ME (Mcal/d)	30.20	30.49	31.12	0.5746	0.5504
NEp (Mcal/d)	12.81	12.93	13.19	0.2436	0.5507
Live weight (kg)	393	396	388	12.801	0.8999
ADG (kg/d)	0.87	0.98	0.83	0.0661	0.3363
Retained energy Mcal/d	2.68	3.11	2.54	0.2767	0.3795
Production efficiency	20.97	24.04	19.17	1.969	0.2841

**Table (4): Economic efficiency of steers fed the experimental rations.**

Item	R1	R 2	R 3
Price / kg fresh (LE)	15.35	15.63	15.97
ADG kg/d	0.87	0.98	0.83
Price of daily gain (LE)	17.4	19.6	16.6
Profit (LE)	2.05	3.97	0.63
Economic efficiency %	13.35	25.39	3.94

Market price pt./kg fresh of: concentrate feed mixture= 170; clover hay=80; seaweed= 700; kg body weight gain=2000.

Hersom (2005) reported that, accurate supply of nutrients to cattle can have several positive outcomes. Providing the required nutrients can increase the production potential, reduced feed cost and improve nutrient utilization thereby also reducing nutrient waste and decreasing environmental concerns.

It could be concluded from the presented results that feeding on 82% concentrate feed mixture and 18% clover hay, which contains 0.4 % seaweed of the total ration intake for fattening Friesian steers, successfully and economically improved animal performance, compared with ration without seaweed or with 0.9 % seaweed.

## REFERENCES

- Aga, L. M.; Koski, R. J. and Stern, M. D. (2000). Effect of buffers on pH and microbial metabolism in continuous culture of rumen contents. 25<sup>th</sup> Conference on rumen function, Chicago, Illinois.
- Duncan, D.B. (1955). Multiple Range and Multiple F Test. *Biometrics*, 11:10.
- Freidman, R. B.; Anderson R. E.; Entine S. M. and Hinshberg S. B. (1980). Effects of disease on clinical laboratory tests. *Clin. Chem.*; 26: 26D.
- Ghoneim, A. (1967). *Animal Nutrition, Principles and Feeding-stuffs*. 2<sup>nd</sup> Edt. Anglo. Egyptian Library, University. (Arabic Text Book).
- Ghorbani, G. R.; Jackson, J. A. and Hemken, R. W. (1989). Effect of sodium bicarbonate and sodium sesquicarbonate on animal performance, ruminal metabolism and acid – base status. *J. Dairy Sci.* 72 : 2039.
- Gornall, A. C.; Bard, W. and David M. M. (1949). Determination of total serum protein. *J. Biol. Chem.*, 177: 751.
- Hersom, H. (2005). Phase-feeding the beef herd for improved feed utilization. Florida Ruminant Nutrition Symposium.

- Hill, P. G. and Wells T. N. (1983). Reduction of reaction differences between human mercaptalbumin and human nonmercaptalbumin measured by the bromocresol purple method. *Ann. Clin Biochem.*, 20: 264.
- Hoffman, M. and Steinhofel, O. (1990). Possibilities and restriction in using milk urea concentrations as markers of energy and protein balance. *Mh. Vet. Med.* 45: 223.
- Hunter, R. D.; Drouillard, J. S. and Titgemeyer, E. C. (1999). Effect of level of non-enzymatically browned soybean meal in limited fed, grain sorghum diets for growing fibers. Kansas State University Agricultural Experiment Station and Cooperative Extension Service.
- Lebas, F. and Colin, M. (1992). World rabbit production and research: Situation in 1992. Fifth World Rabbit Congress, Vol. A, p. 29.
- Mandell, I.B.; Campbell, C. P.; Buchanan-Smith, J. G. and Wilton, J. W. (2001). Effects of source and level of dietary roughage on growth performance and carcass quality in yearling calves. Department of Animal and Poultry Science, University of Guelph, Guelph, Ontario.
- Mohamed, H. A. and Selim, H. M. (1999). Hand book of Veterinary Internal Medicine. Department of Animal Medicine, Faculty of Veterinary Medicine. Zagazig Univ.
- NRC (National Research Council) (1996). Nutrient Requirements of beef Cattle (7<sup>th</sup> ed.) National Academy Press, Washington, DC.
- Overton, T. (1999). "Research Feature" Total Dairy Nutrition, Vol, 1, No, 2 April (1999).
- Reitman, A. and Frankel, S. (1957). A colorimetric method of determination of s.GOT and s.GPT. *American J. of Clinical Pathology*, 28: 56.
- SAS (1994). SAS / STAT R User's Guide : Statistics. Ver. 6.04, Fourth Edition SAS Institute Inc, Cary, NC.
- Schreder, P. (2002). Feeding & Riparian the show steer. OUS & Riparian, Linn / Benton / Lan Countries No. L & R.
- Scott, C. (1990). Kelp Help : Livestock thrive on seaweed. Elphin, Lanark Country, Eastern Ontario, Canada.
- Teuscher, A. and Richterich, R. (1971). *Schweiz. Med. Woschr.* 101: 345.
- Ullmann, K. (1976). *Bonitz. Med. Labor.*, 29:137 .
- Zinn, R. A. (1991). Comparative feeding value of steam – flaked corr. and sorghum in finishing diets supplemented with or without sodium bicarbonate. *J. Anim. Sci.* 69 : 905.

## تأثير إضافة الطحالب البحرية لعلائق تسمين عجول الفريزيان على : ٢- الأداء الإنتاجي في المرحلة النهائية

حسين محمد الشافعي عيد<sup>١</sup> و إيمان حنفي محمود مقلد<sup>٢</sup>

- ١- معهد بحوث الانتاج الحيواني - مركز البحوث الزراعية .
- ٢- قسم إنتاج الحيوان - كلية الزراعة - جامعة المنصورة .

أجرى هذا البحث بهدف دراسة تأثير إضافة نسب مختلفة من الطحالب البحرية (صفر ، ٠,٩ ، ٠,٤) من المادة الجافة المأكولة على بعض قياسات الدم ومعدل النمو في تسمين عجول الفريزيان والاستفادة الغذائية والكفاءة الاقتصادية . وتم تكوين ثلاث علائق على النحو التالي: (عليقة أولى) ٨١,٨ % علف مصنع + ١٨,٢ % دريس برسيم (عليقة مقارنة) . (عليقة ثانية) ٨١,٥٥ % علف مصنع + ١٨,٠٥ % دريس برسيم + ٠,٤ % طحالب بحرية . (عليقة ثالثة) ٨١,٢٣ % علف مصنع + ١٧,٨٧ % دريس برسيم + ٠,٩ % طحالب بحرية . وكانت الخلطات الثلاثة متماثلة من حيث نسبة البروتين حيث تروحت بين (١٦,٤٧ - ١٦,٥٥ %) ومستخلص الألياف المتعادلة (٤٢,٦٤ - ٤٢,٦٨ %).

استخدم ١٢ عجل فريزيان متوسط وزن ٢٦٩ كجم، عند متوسط عمر ١٥ شهرا وموزعة في ثلاث مجاميع (أربعة عجول في كل مجموعة) وتم أخذ عينات الدم من كل عجل بعد الأكل بـ ٣ ساعات وتم تسجيل أوزان الحيوانات شهريا لتقدير معدل الزيادة اليومية عند التغذية على العلائق المختبرة وحساب الكفاءة الاقتصادية لكل منها وكانت أهم النتائج المتحصل عليها كما يلي :

- ١- لم تتأثر مكونات الدم معنويا بالتغذية على العلائق المختبرة ولكن لوحظ إنخفاض تركيز الأمونيا لكل من العليقة الثانية والثالثة مقارنة بالعليقة الأولى.
  - ٢- لم تتأثر أوزان الحيوانات معنويا نتيجة المعاملات وكذلك معدل النمو اليومي (كجم / يوم) خلال فترة التجربة ( ٠,٨١ ، ١,٠ ، ٠,٨٣ كجم / يوم) عند التغذية على العليقة الأولى والثانية والثالثة على التوالي.
  - ٣- زادت الكفاءة الانتاجية عند التغذية على العليقة الثانية (٢٤,٠٤ %) مقارنة بالتغذية على العليقة الأولى أو الثالثة (٢٠,٩٧ ، ١٩,١٧ % على الترتيب).
  - ٤- زادت الكفاءة الاقتصادية بالتغذية على العليقة الثانية (٢٥,٣٩ %) مقارنة بالتغذية على العليقة الأولى أو الثالثة (١٣,٣ ، ٣,٩٤ % على الترتيب).
- مما سبق يستنتج أن التغذية على العليقة الثانية ( ٨١,٥٥ % علف مصنع + ١٨,٠٥ % دريس برسيم + ٠,٤ % طحالب بحرية ) هي الأفضل إنتاجيا وإقتصاديا لتسمين العجول الفريزيان مقارنة بالتغذية على العلائق الأخرى.

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة  
اكاديمية البحث العلمي

أ.د / محمد محمد الشناوى  
أ.د / محسن محمود شكرى