

**EFFICACY OF SOME PLANT EXTRACTS AND THE COMMERCIAL BIOCIDES, DOLPHIN ON POTATO TUBER WORM, *Phthorimaea operculella* (ZELLER.) (LEPIDOPTERA: GELECHIIDAE) UNDER FIELD AND STORAGE CONDITIONS**  
Abou Hatab, Eftkhar E. ; Neveen F. Arafat and Naglaa F. Riad  
Plant Protection Research Institute, (ARC). Dokki, Giza.Egypt

**ABSTRACT**

Field and laboratory experiments were conducted at Aga district ,Dakahlia Governorate during two successive seasons 2009 and 2010 to evaluate the efficiency of some plant extracts (demises ) and plant oils(mint oil- fennel oil -cumin oil) in comparison with the biocide (dolphin )against *Phthorimaea operculella*(Zeller.) .

The obtained results showed that the mint oil was the most effective against *P. operculella* larvae (74.2, 75.9%), followed by fennel oil (62.7, 74.2%) ; cumin oil (62.6, 69.1%) and demises plant extracts (61.5, 60.5%). While, dolphin caused a relatively high mortality rate of PTM larvae( 82.6 &81.4 %) during2009 &2010 seasons.

The results indicated that the bioactivity of the tested materials were concentration-dependent, whereas, the inhibition rates of moths resulted from eggs or pupae or moths infestation increased obviously with the rise of concentration. Meanwhile, the bioactivity of all were relatively the most to moths infestation all tested materials than eggs and pupae.

**INTRODUCTION**

Potato (*Solanum tuberosum* L.) is an important vegetable crop in Egypt, which is seriously infested by the potato tuber moth, *Phthorimaea operculella* (Zell.) (Lepidoptera : Gelechiidae) especially, in the field and in stores (Abd El-Salam *et al.*, 1972), Shaheen (1979), Heeder (1983), Doss (1984), Iskander (1985); Khalil *et al.* (1987) and Ahmed *et al* (1991). Infested tubers become completely unmarketable. Chemical control of potato tuber moth is a costly input that contaminated the environment, increase resistance to insecticides and can also cause health hazards for the humans. The present work aims to study the population fluctuation of potato tuber moth *P. operculella* with relation the level infestation with different plant growth stages. Also, field and laboratory studies were carried out evaluate demises plant extract and some plant oils against *P. operculella* larvae.

**MATERIALS AND METHODS**

Field and laboratory experiments were conducted in 2009 and 2010 growing seasons to determine the effects of selected plant extracts and the commercial biocide (Dolphin) on the potato tuber moth (PTM). These materials were subjected in Table (1).

**Field experiments :-**

An area of about 1/4 feddan at Aga district, El-Dakahlia Governorate was selected and divided into 18 plots (each about 55 m<sup>2</sup>). The experimental replicates were arranged in complete randomized block design with three replicates for each treatment. The planting dates were on the 26 and 14 of January for 2009 and 2010 seasons. Sponta variety was cultivated as recommended whereas. All agricultural practices were followed.

To determine the effect of some plant extracts and the commercial biocide (Dolphin) on the potato tuber moth (PTM), *P. operculella* Zeller population, seasonal abundance of PTM larval population was estimated in treated and untreated potato fields throughout two successive years 2009 and 2010.

Inspection of plants was carried out before spray and after 10 days from application to investigate the effect of the different treatments on the reduction rates of PTM larvae populations.

-The efficacy of the previously mentioned plant extract and plant oils were investigated against PTM as follows:

The reducing percentage in infestation was corrected by using Abbott's formula (Abbott, 1925):-

$$\text{Reduction in infestation \%} = \frac{A - B}{A} \times 100$$

Where A = number of infested tubers in control

B = number of infested tubers in treatment

**Estimating the role of some plant extracts and the biocid (Dolphin) on PTM population:-**

Sampling were taken after twenty one days from sowing date. Intervals between each samples were 7 days. Each sample consisted of 10 leaflets picked at random. Collected samples were transferred to the laboratory in paper bags for investigations. The collected samples were examined weekly by using a binocular-microscope to determine the population fluctuation of *P. operculella* larvae in treated and untreated plots. The number of *P. operculella* larvae present in each sample was recorded.

**Laboratory experiments :-**

This laboratory experiment was carried out to compare the efficiency of the previously mentioned natural products against *P. operculella* eggs, larvae and adults.

75 infested potato tubers with PTM eggs approximately equal in size were collected and divided into three replicates (each replicate contain 25 tubers). After that ,potato tubers were treated using various concentrations (5, 2.5, 1.25 and 0.75 %W/V) were prepared from the stock solution for the extract of Demises leaves, fennel oil ;mint oil and cumin oil for conducting the experiments. Three replicates were used for each concentration. Experiments were carried out at 25±2°C and 70±5% R.H. Then the numbers of emerged moths were recorded for one week from the initial moth emergence.

**Table 1: The common and scientific names and rate of application of the tested compounds.**

Compound and scientific names	Family	Rate of application
1-Cumin oil ( <i>Cuminum cyminum</i> L.):	Umbelliferae	200 ml / 100 L water
2- Mint oil ( <i>Mentha piperita</i> L.)	Labiatae	200 ml / 100 L water
3-fennel oil( <i>Foeniculum vulgare</i> L.)	Umbelliferae	200 ml / 100 L water
4-Demises, <i>Ambrosia maritime</i> ,	Compositiae	600 ml / 100 L water
5-Dolphin 85 %, WG, B.T		200 gm/ Fadden

Cumin, mint and fennel oils were bought from the local market, while demises extract was obtained from Horticulture Research Institute (HRI) Egypt,

**Statistical analysis:**

The mortality was corrected using Abbott's formula (1925) with data under the store.

"a Costat" a product of cohort soft ware was used to differentiation between means.

The reduction percentages (% mortality) of different tested materials in the field were calculated according Hinderson and Tilton (1955).

The reduction percentages of the emerged moths from the pupae and eggs were calculated using the following equation:

$$\text{Inhibition} = \frac{\text{No. of emerged moths in control} - \text{No. of emerged moths in treatment}}{\text{No. of emerged moths in control}} \times 100$$

**RESULTS AND DISCUSSION**

The data illustrated in Table (2) showed that infestation by *Phthorimaea operculella* started at 45 days after sowing on March 11<sup>th</sup> 2009 season,(4.8 larvae/10 leaves) on Sponta variety, then increased gradually to reach its maximum (13.9 larva/10 leaves) 108 days after sowing on April 23<sup>rd</sup> , after that the population decreased gradually to reach (3.0 larvae/10 leaves) when plant age 122 days on May 28<sup>th</sup>. Meanwhile, data indicated that the larval stage increased as well as potato plants growth increased, whereas, the level of infestation with potato tuber moth *P. operculella* during the two seasons ,2009 and 2010. PTM larval population started in few numbers, then increased gradually till reached it peak when potato plants age of 108 and 114 days during the two seasons.

In the second season, the obtained results showed that infestation by *P. operculella* stated at 58 days after sowing on March 13<sup>th</sup> 2010 season: (4.5 larva/10 leaves) on Sponta variety, then increased gradually to reach its maximum (12.7 larva/10 leaves) 114 days after sowing on May 15<sup>th</sup> on Sponta variety.

Data show also that incidence of infestation by *P. operculella* as expressed by counts of larvae stage increases sharply with the increase of potato plants age to reach its maximum incidence at 114 days, and infestation dropped to reach minimum rates at ages of more than 128 days on the potato Sponta variety, (Table 2). The obtained data are in agreement with those of Mikhael (1995); Dawood *et al.* (1999); Stein and Vendramin (2000) and Gomma (2002).

**Table (2): Mean larval numbers of the potato tuber moth *P. operculella* in relation to plant age for potato at DaKahlia Governorate during summer plantation of 2009 and 2010 seasons.**

Date of Inspection	Plant age (day)	Mean larval numbers /25 plant leaves	Date of inspection	Plant age (day)	Mean larval numbers /25 plant leaves
		Sponta			Sponta
5/2/2009	10	0.0	6/2/2010	23	0.0
12/2	17	0.0	13/2	30	0.0
19/2	24	0.0	20/2	37	0.0
26/2	31	0.0	27/2	44	0.0
4/3	38	0.0	6/3	51	0.0
11/3	45	4.8	13/3	58	4.5
18/3	52	5.7	20/3	65	5.4
25/3	59	6.6	27/3	72	5.3
2/4	66	6.6	3/4	79	6.4
9/4	73	7.8	10/4	86	8.6
16/4	80	8.9	17/4	93	8.7
23/4	87	9.7	24/4	100	9.4
30/4	94	9.5	1/5	107	11.6
7/5	101	11.8	8/5	114	12.7
14/5	108	13.9	15/5	121	9.4
21/5	115	4.9	22/5	128	2.3
28/5	122	3.0	29/5	135	0.0
4/6	128	0.0	5/6	142	0.0
Mean±S.E. of actual values		5.2±0.73	Mean ±S.E. of actual values		4.7±0.86

The results in Table (3) the mortality in case of after first spray and second spray on treated with Cumin oil, fennel oil, Mint oil, Demises plant extract and Dolphin were (55%& 70.9); (63.5&79.4); (63.5&89.4); (50&73.1%) and (85&80.2%) , with five compounds, respectively, during 2009 season.

The mortality the second season showed nearly the same trend as indicated with after first and second spray. However, all compounds caused a significant mortality in general mean of larvae as compared with control. Also, the Dolphin was relatively the most efficient compound in protecting potato tubers against *P. operculella*

These results agree with the findings Raman *et al.*(1987); Lal (1987) and Doss *et al.* (1994) stated that *L. camara* significant reduced Sponta damage when compared with the untreated control and with those covered with rice straw.

**Table (3):Efficiency of certain treatments against *Phthorimaea operculella* (Zeller) larvae on potato plants during two seasons at Dakahlia Governorate.**

Treatments	Rate / 100L	Pre-spray	Mean larval number											
			Application											
			2009						2010					
			After first spray 10days		After second spray 10days		General mean%	Pre-spray	After first spray 10days		After second spray 10days		General mean%	
No.	%	No.	%	%		No.	%	No.	%	%				
Cumin oil.	200	25	15	55	11	70.9	62.6	13	7	66.1	7	72.0	69.1	
fennel oil	200	32	17	63.5	10	79.4	62.7	18	9	68.5	7	79.9	74.2	
Mint oil.	200	35	17	63.5	8	89.4	74.2	21	9	64.0	5	87.7	75.9	
Demises plant extract	600	27	18	50	11	73.1	61.5	14	11	50.5	8	70.4	60.5	
Dolphin	50	20	4	85.0	6	80.2	82.6	9	2	80.1	3	82.7	81.4	
Control	-	33	44	-	50	-	-	29	46	-	56	-	-	

L.S.D. =8..3

\*= Significant at 5% level.

% Reduction rates are given in brackets.

The effectiveness of plant extract; cumin oil; fennel oil and mint oil against *P. operculella* is demonstrated in Tables (4, 5, 6, and 7).

The results of cumin oil given in Table (4) showed that the average numbers of emerged adults from egg, pupa and moth treatments were reduced with the increase of concentration.

The recorded numbers were, 30.6, 22.6, 16.5 and 10.9 emerged adults from eggs treatments, 29.5, 27.2, 24.0 and 15.5 emerged adults from pupae treatments and 30.5, 29.0, 25.0 and 12.0 emerged adults from moth treatments at the previously mentioned concentrations, respectively.

The obtained results revealed that the efficacy of agent dependent on concentration. The highest mortalities of the treatments were observed at the highest tested concentration.

In this respect, El-wahab *et al.*, (2003) indicated that mortalities of cumin oil of were some what higher at the highest concentration of 10%, to some stored product insects.

Response of *P. operculella* to mint oil is given in Table (4). The results indicated that inhibition rates of moths resulted from eggs, pupae and moth treated dependent on concentration. The highest inhibition rates of moths emerged were 60.3, 69.9% and 71.4% of eggs, pupae and moths treated with the highest concentration (5% W/V) of mint oil for eggs, pupae and moths, respectively, while the lowest reduction rates of moth emergence (12.9,10.0 and 35.7%) were recorded at the lowest concentration (0.75% W/V).

A result of the response of *P. operculella* of fennel oil was given in Table (5). The number of emerged moths was gradually reduced with increasing the concentration at all treatments when compared with control. Consequently, the inhibition rates of moth emergence increased as

increasing the concentration at all treatments, reaching between 12.2-58.8 %, 10.5-56.0 % and 8.5-68.9 % in egg, pupa and moth treatments for fennel oil.

Response of *P. operculella* to demise plant extract is given in Table (6). The results indicated that inhibition rates of moths resulted from eggs; pupae and moths treatment concentrations. The highest inhibition rates of moth's emergence were 55.0, 51.0 and 64.1% of eggs, pupae and moth treatments at the highest Concentration (5%), respectively. While, the lowest reduction rates of moth emergence (8.0 , 7.1 and 18.8%) were recorded at the lowest concentration (0.75%). The obtained results are in complete harmony with that obtained by Kroschel and Koch (1996) when they Studied that the toxicity of three plant extracts, namely chinaberry (*Melia azedarch*), neem (*Azadirachta indica*) on the reduction of *P. operculella*.

Results indicate that the tested plant extract and plan oils were slightly more effective in moths and pupae treatments than eggs treatment and showed promising results at the highest concentrations of the different tested materials. Thus, these plant oils and leaves extracts could be used as tuber Protestants against this species frame of an integrated pest management program (IPM).

Also, the obtained results coincided with the findings of other investigators (Reddy and Urs., 1989; Salem, 1991, Sabbour and Ismail, 2002) and Mikhael,(1995).

**Table (4): Response of *Phthorimaea operculella* (Zeller) to cumin oil (*Cuminum cyminum*).**

Conc. (% w/v)	Average no. of emerged moths from eggs treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from pupae treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from moths treatment	Inhibition rates of moths emergence (%)
5	10.9±3.1	68.8	15.5±2.2	58.0	12.0±2.8	70.0
2.5	16.5±2.9	52.8	24.0±3.4	35.1	25.0±3.6	37.5
1.25	22.6±2.6	35.4	27.2±2.5	26.4	29.9±3.1	27.5
0.75	30.6±3.2	12.6	29.5±2.9	20.2	30.5±2.4	23.8
Control	35.0	-	37.0	-	40.0	-

**Table (5): Response of *Phthorimaea operculella* (Zeller) to mint oil (*Mentha piperita* L).**

Conc. (% w/v)	Average no. of emerged moths from eggs treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from pupae treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from moths treatment	Inhibition rates of moths emergence (%)
5	11.9±2.6	60.3	10.0±2.1	69.9	10.0±2.3	71.4
2.5	13.2±3.4	56.0	17.8±2.5	46.0	14.0±1.9	62.8
1.25	16.4±3.2	45.3	19.5±2.3	40.9	19.0±1.8	45.7
0.75	20.8±1.9	30.6	23.5±2.7	28.7	22.5±2.7	35.7
Control	30	-	33	-	35	-

**Table (6): Response of *Phthorimaea operculella* (Zeller) to fennel oil (*Feoniculum vulgare* L).**

Conc. (% w/v)	Average no. of emerged moths from eggs treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from pupae treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from moths treatment	Inhibition rates of moths emergence (%)
5	18.5±3.1	58.8	20.5±2.2	56.8	13.5±2.8	68.9
2.5	26.5±2.9	41.1	28.5±3.4	40.0	26.5±3.6	39.1
1.25	32.5±2.6	32.2	33.5±2.5	29.4	33.5±3.1	22.9
0.75	39.5±3.2	12.2	42.5±2.9	10.5	40.0±2.4	8.5
Control	45	-	47.5	-	43.5	-

**Table (7): Response of *Phthorimaea operculella* (Zeller) to demises plant extract (*Ambrosia maritime*).**

Conc. (% w/v)	Average no. of emerged moths from eggs treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from pupae treatment	Inhibition rates of moths emergence (%)	Average no. of emerged moths from moths treatment	Inhibition rates of moths emergence (%)
5	22.5±3.1	55.0	24.0±2.2	51.0	19.0±2.8	64.1
2.5	27.0±2.9	46.0	32.6±3.4	33.4	26.9±3.6	49.2
1.25	43.5±2.6	13.0	36.0±2.5	26.5	36.5±3.1	33.3
0.75	46.0±3.2	8.0	45.5±2.9	7.1	43.0±2.4	18.8
Control	50.0	-	49.0	-	53.0	-

## REFERENCES

- Abbott, W. S. (1925): A method of computing the effectiveness of insecticides. *J. Econ. Entomol.*, 18: 265-207.
- AbdEl-Salam, A. M.; Assem, M. A.; Hamad, S. M. and Eid, G. H. (1972): Studies on potato pest in UAR II. Susceptibility of some potato insects' infestation in the field. *Z. Angew. Ent.* 70 (1): 76-82. (c.f. *A.R.E.*, 62 : 1526).
- Abd El-Wahab, H. A.; El-Adl, F. E.; Soad, A. Ibrahim and El-Bouze, M. F. R. (2003): Efficiency of certain materials against potato tuber moth, *Phthorimaea operculella* (Zeller) and its parasitoids with special regard to their residues. 2<sup>nd</sup> International Conference, Plant Prot. 735-741.
- Ahmed, S. A.; Ali, A. M. and Salman, A. M. (1991): Influence of potato varieties, nitrogen fertilization and plant growth regulators on the infestation level of *Myzus persicae* in potato fields. Triennial Conf. of European Assoc. For potato Res. (EAPR), Edinburgh (U.K.), 8-13 July, 56-57.

**Abou Hatab, Eftkhar E. et al.**

- Dawood, M. Z.; El-Rafie, K. K. and Haydar, M. F. (1999): Susceptibility of some potato cultivars to the potato tuber moth, *Phthorimaea operculella* (Zeller) infestation with relation to yield at Giza Governorate. J. Appl. Sci., Egypt, 14 (4).
- Doss, S. A. (1984): Relative susceptibility of seventeen potato varieties to infestation by three insect pest's infestation in stores. Bull. Soc. Ent. Egypt, 65: 157-167.
- Gomaa, A. S. A. (2002): New approaches to control the potato tuber moth, *Phthorimaea operculella* (Zeller) on potato in the A.R of Egypt. Ph. D. thesis, Sci. Tanta Univ., 1-164.
- Heeder, M. F. (1983): Ecological and pest management studies on the potato tuber moth, *Phthorimaea operculella* (Zeller). Ph. D. Thesis, Fac. Agric., Ain Shams Univ.
- Hinderson, C. F. and Tilton, E. W. (1955): Test with acaricides against the brown white mite. J. Econo. Entomo., 48: 157-161.
- Iskander, N. N. (1985): Ecological and biological studies on *Phthorimaea operculella* (Zeller). M. Sc. Thesis, Fac. Agric. Zagazig Univ., Egypt
- Khalil, F. M.; Abdel-Galil, F. A.; Ali. A. M. and Soliman, M. M. (1987): Susceptibility of certain potato varieties to Assuit. J. Agric. Sci., 18 (3): 215-233.
- Kroschel, J. and Koch, W. (1996): Studies on the use of chemicals the potato tuber moth in potato stores. Crop protection, 15-2, 197-203.
- Lal, L. (1987): Studies on natural repellents against potato tuber moth, *Phthorimaea operculella* (Zeller) in country stores. Potato Res., 30 (2): 329-334.
- Mikhael, R. H. (1995): Studies on insect pests infesting potato crop and their control by using non-chemical methods. Ph.D. Thesis, Plant Protection Department Fac. Agric. Moshtohor Benha Branch, Zagazig Univ.
- Raman, K. V.; Both, R. H. and Palacois, M. L. (1987): Control of potato tuber moth, *Phthorimaea operculella* (Zeller) in rustic potato stores. Trop. Sci., 27: 175-194.
- Reddy, G. V. P. and Urs, K. C. D. (1989): Effects of *tribulus terrestris* Linn. On the development of potato tuber moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). Current Science, 58 (4): 212-213.
- Sabbour, M. and Ismail, I. A. (2002): The combined effect of microbial control agents and plant extracts against potato tuber moth, *Phthorimaea operculella* (Zeller). Bulletin of National Research Center, Cairo, 2002, 27: 4, 459-467.
- Salem, S. A. (1991): Evaluation of neem seed oil as tuber protecting against *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). Annals of Agricultural Science Moshtohor, 29:1, 589-595.
- Shaheen, A. H. (1979): Some ecological and biological studies on the potato tuber moth, *Phthorimaea operculella* in Egypt. Bull. Ent. Soc., Egypt, 65: 345-350.
- Stein, C. P. and Vendramin, J. D. (2000): Antibiosis of potato clones to *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). Anais da Sociedade Entomologica do Brasil, 29: 4, 783-788.



فاعلية بعض المستخلصات النباتية والمبيد الحيوي (دولفن) على فراشة درنات البطاطس تحت الظروف الحقلية والمعملية  
أنتخار السيد أبوحطب ، نفين فوزي عرفات و نجلاء رياض  
معهد بحوث وقلية النباتات - مركز البحوث الزراعية

أجريت تجربة حقلية في مركز اجا - محافظة الدقهلية خلال موسمين متتاليين ٢٠٠٩، ٢٠١٠ لدراسة التنذب العددي لفراشة درنات البطاطس *Phthorimaea operculella* وتأثير عمر النبات علي مستوى الإصابة وأوضحت النتائج الإصابة بفراشة درنات البطاطس تبدأ في الاسبوع الثاني من شهر مارس خلال السنتين وقد تزداد تدريجيا لتصل ذروتها في شهر أبريل وشهر مايو أي عند عمر النبات بعد ١٠٨، ١١٤ يوم من الزراعة حيث سجل أعلى تعداد (١٣.٩، ١٢.٧ يرقة/١٠ ورقة) ، في السنة الأولى والثانية علي علي التوالي كما وجد ارتباط معنوي موجب بين عمر النبات وتعداد فراشة درنات البطاطس خلال موسمي الدراسة.  
أجري تقييم لثلاث زيوت نباتية لنبات النعناع والكمون والشمر ومستخلص نبات الدمسيصة بالمقارنة بمبيد حيوي (الدلفن) ضد يرقات فراشة درنات البطاطس في الحقل علي صنف بطاطس سبونتا وأوضحت النتائج أن زيت النعناع أعلى تأثير يليه زيت الشمر وزيت الكمون ومستخلص الدمسيصة علي يرقات فراشة درنات البطاطس بينما بالنسبة لمبيد الدلفن الحيوي كانت النسبة ٨٢.٦ و ٨١.٤% علي التوالي خلال السنتين.

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة  
مركز البحوث الزراعية

أ.د / عبد الستار ابراهيم عبد الكريم  
أ.د / حسن علي طه