

## EFFECT OF DIETARY PROTEIN AND METHIONINE LEVELS ON GROWTH PERFORMANCE AND IMMUNE RESPONSE IN BROILER CHICKS

A. M., Orma; T. I., Mohamed & A. A. M., Abo Elnaga

Dept. of Nutrition & Nutritional Deficiency Diseases, Faculty of Vet. Med.,

Mansoura University

### ABSTRACT

*The present work was carried out to study the effect of decreasing the dietary protein content by 10% (LCP-diet) with supplementation of methionine (0, 0.1, and 0.2%) over the recommended level in broiler diets on growth performance, carcass quality, blood metabolites, immune response. One day old Cobb chicks (120) were used. The chicks were distributed randomly into 4 groups and fed 23% or 20.8% CP for the starter diets, 20 or 18.2% CP for the grower and 18 or 16.2% CP for the finisher diets with different dietary methionine levels (0.5, 0.6 or 0.7%) for starter diets, 0.4, 0.5 or 0.6% for the grower diet and 0.32, 0.42 or 0.52% for the finisher diets, respectively and 3200 Kcal ME/Kg. The chicks were vaccinated against Avian influenza, Newcastle and Gumboro diseases at proper time. The growth performance indices (body weight, feed intake and feed conversion ratio) were measured. Blood samples were collected from the broilers at 14, 21 and 28 days of age without anticoagulant for separation of sera to detect titer of antibodies against Newcastle disease vaccine using haemagglutination inhibition test (HI) as indicative of birds immune response in the different experimental groups and also at 42 days of age, blood samples collected for determination of serum metabolites. At the end of the experiment, 5 chicks from each group were slaughtered to obtain some of the carcass traits.*

*The results revealed that body weight and feed conversion ratio of the broiler fed 10% LCP-diet supplemented with 0.2% methionine above that recommended for the broilers significantly improved on comparison to those fed the LCP diets supplemented with 0, 0.2% methionine. Feeding the broiler chicks the LCP-diet decreased the body weight gain and increased the feed conversion ratio. The abdominal fat % in the carcass of the broiler chicks fed LCP-diet supplemented with 0.2% methionine were lower than those fed on the diets supplemented with 0, 0.2% methionine. Feeding LCP-diet supplemented with 0.2% methionine increased serum total protein, albumin, and globulin and decreased serum total lipids, cholesterol, triglycerides, low density lipoprotein and high density lipoprotein.*

*It could be concluded that feeding the broiler chicks on LCP-diet supplemented with 0.2% methionine diet resulted in better growth performance, lower total lipids,*

*cholesterol, triglycerides, low density lipoprotein and high density lipoprotein and higher total protein, albumin, globulin in comparing to those fed 0 or 0.2% methionine supplemented diet. Supplementing the low protein diet with 0.2% methionine improved the HI titer in the broilers vaccinated against NDV at 14, 21 and 28 days of age.*

### **INTRODUCTION**

Malnutrition has been identified in many countries particularly in the developing countries. Majority of malnourished people live in Asia and Africa (**Iwe and Onadipe, 2001**), thus negatively impacting on the physical and health condition of its people. Poultry production is one major mean of solving part of this problem (**Nworgu et al., 2000**). Feeding poultry presents a great challenge to producers and nutritionists in several tropical countries as meeting the nutritional requirements for growing birds constitutes the majority of costs associated with poultry production (**May et al., 1998**), and certainly is becoming an issue of even greater significance as the prices of feed ingredients continue to rise. A large portion of that cost involves in meeting the amino acid requirements of the birds (**Corzo et al., 2004; Eits et al., 2005**).

The amino acid pattern provided in diets must exactly matches the birds amino acid requirements, excesses can be avoided consequently; protein accretion can occur with maximum efficiency, excess dietary protein increased heat production and water consumption which increased moisture content of litter (**Alleman and Leclercq, 1997**). Lowering crude protein in the diet could reduce the feed costs and decreased nitrogen excretion (**Nahm, 2002 and Namroud et al., 2008**), and allowed for using of a greater variety of feedstuffs (**Kidd et al., 1996**), which

can be valuable in itself as a method to increase flexibility in the choice of locally available feed stuffs and potentially decreasing fluctuation in costs. Dietary protein level therefore, has major effect on growth performance and overall cost of finished product. So using of L-CP-diet supplemented with amino acid for broiler feeding has been received greater interests in the recent years in order to reduced feeding costs and environmental pollution (**Namroud et al., 2008**). The use of synthetic amino acid to meet the amino acid needs of broilers could result in production of cost effective diets.

In chickens as well as mammals, it has been shown that deficiency or excess of dietary protein (**Payne et al., 1990**) or amino acids (**Tsiagbe et al. 1987**) altered immune responses. The magnitude of an immune response and subsequent alterations in nutrient metabolism is dependent upon a nutritionally complete diet (**Webel et al., 1998**). The efficiency of dietary protein is primarily dependent upon an adequacy of lysine and methionine (most limited amino acids). So, appropriate supplementation of ration deficient in these amino acids markedly improved their protein quality (**Ichhponani et al., 1981**). With progress in biotechnology, the cost of production of each amino acid particularly methionine has been significantly reduced, which has been one of the key factors in the expansion of use of amino acids in ani-

mal feed. Therefore, the objective of this study is to further evaluate the probability of lowering the dietary protein content with methionine supplementation and its effect on growth performance, dressed carcass, serum metabolites and immunocompetence in broiler chicks.

### **Materials and Methods**

#### **Birds, diets and management :**

A total of 180 one day old Cobb chicks obtained from a commercial supplier were randomly distributed into 4 equal triplicate groups (15 chicks / replicate) and raised in a floor pen bedded with wheat straw to maintain a good litter. The pen was supplied with feeders, waterers and heaters. The chicks were tagged using wing number tags. The chicks were fed the respective experimental diets according to the suggested design. The diets were formulated to maintain the requirements of energy (3200 kcal ME/kg) with decreasing protein by 10% than that recommended by **NRC (1994)** and supplementation with methionine by (0, 0.1 & 0.2% of the diet) over the recommended. Samples from yellow corn, soybean meal and fish meal were analyzed for dry matter, crude protein, ether extract and ash (**AOAC, 1990**) and used for formulation of the diets. The composition of the experimental diets is shown in table 1.

The broiler chicks were reared during the period from mid of February to first of April, 2010. During the first 2 weeks of chicks' age the brooding temperature was maintained at 35-32°C, and then the temperature was gradually lowered till reach to 24°C by the beginning of the third week of age. The pen was naturally ventilated and electric light was con-

tinuously maintained (24 hours/day). The diets and water were available ad libitum throughout the experimental period (6 weeks). The chicks were vaccinated against avian influenza by subcutaneous injection and Newcastle disease at 8th day of age by Hitchner B1 and at 18th, 28th days of age by lasota vaccine in drinking water and against Gumboro at 10th and 20th days of age in drinking water.

#### **Indices of broiler performance, dressed carcass and abdominal fat**

The chicks were weighed at one day old to obtain the average initial body weight then weighed every week and the average body weights for the chicks in each group were weekly calculated. The average feed intake per chick throughout the experimental period (6w) for each group was recorded and feed conversion ratio was calculated. At the end of the experiment, five birds from each group were randomly taken and fasted for 12 hours, weighed, slaughtered to complete bleeding, dressed and eviscerated for determination of the dressed carcass % and abdominal fat weight and percent in relation to live body weight.

#### **Blood sampling, serum metabolite and immune response**

Blood samples were collected from 5 chicks from each group at 42 days of age via wing vein puncture. Sera were obtained by allowing blood samples to coagulate then centrifuged at 3000 rpm for 15 minutes. The separated sera were frozen at -20°C in a deep freeze until used for determination of glucose (**Trinder and Ann, 1969**), total lipid (**Zollner and Kirsch, 1962**), triglycerides (**Fassati and**

**Precipe, 1982**), cholesterol (**Allain et al., 1974**), high density lipoproteins (**Burstein et al., 1970**), low density lipoproteins (**Friede Wald et al., 1972**) total protein (**Goarnal et al., 1949**) and albumin (**Doumas et al., 1971**) using already prepared analyzing chemical kits and after the instructions of the producer (GOD-PAP, BIODIAGNOSTIC and STANBIO). Globulin concentration was calculated as the difference between total protein and albumin.

Blood samples were collected from 5 chicks of each group at 14, 21 and 28 days of age without anticoagulant for separation of sera which were used to detect titer of antibodies against Newcastle disease vaccine using haemagglutination inhibition test (HI) as indicative of bird's immune response.

#### **Statistical analysis:**

Data obtained in the present work were statistically analyzed for analysis of variance (ANOVA) and least significant difference (LSD) as described by **Snedecor and Cochran (1967)**.

### **RESULTS AND DISCUSSION**

#### **Body weight, body weight gain, feed intake, feed conversion ratio and dressed carcass:**

The data concerned with body weight development (BW), body weight gain (BWG), feed intake (FI) and feed conversion ratio (FCR) showed that decreasing the dietary protein level by 10% than that of the **NRC (1994)** recommendation levels significantly decreased BW, BWG and significantly increased the values of FCR of broiler chicks even when methionine is supplemented by 0.2% above that

recommended by **NRC (1994)**. However, increasing the level of lysine supplementation to 0.4% levels significantly improved the BW, BWG and FCR recorded at 3rd, 4th, 5th and 6th weeks of the experiment (Table 2). **Aletor et al. (2000)** found that supplementing the diet with amino acids improved growth of broilers. Also, **Sterling and Pesti (2003)** demonstrated that different levels of supplemental lysine increased BWG of broiler chicks from 0 to 21 days of age. The improvement in BW of broiler chicks fed LCP-diets supplemented with L-lysine indicated that lysine supplementation improved protein utilization (El-Sherbiny et al., 1997 and Khalifa, 2001). With this concept, Abdallah et al. (2005) found that BWG and FCR of broiler chicks fed L-CP-diets (20 to 18% CP from 0 to 3 wk and 18 to 16% CP from 4 to 7 wk of age) supplemented with methionine and lysine was not significantly different from those fed the 23 to 20% CP-diet. Similarly, **Khan et al. (2010)** found that reduction of protein level from 23% to 19% had no detrimental effects on BWG when supplemented with lysine. The improvement in WG with LCP-diet supplemented with EAA could be due to reduced heat increment, which was associated with the metabolism of excess protein. Reduced heat increment led to reduced heat stress and therefore, improved WG.

The effect of decreasing the dietary crude protein level by 10% in the present study resulted in high feed intake and FCR whatever the diet supplemented with excess lysine or not (Table 3). The recorded FCR are 1.89 for the control diet compared to 2.54, 2.42 and 2.28 for the LCP-diet supplemented with 0, 0.2% or 0.4% lysine diets, respectively. **Nasri-**

**mento, (2003)** found that increasing dietary lysine levels for broilers and balancing the essential amino acids improved FCR without modifying the feed intake because of the increase of muscular growth and body weight gain. Similarly, **El-Hussein et al. (2004)** and **Yamazaki et al. (2006)** found that feed intake of broiler chicks was not significantly affected by LCP amino acid supplemented diets. Also, **Corzo et al. (2004, 2006)** reported that increasing lysine only improved FCR without affecting FI. In addition, **Sterling et al. (2006)** found that increasing dietary lysine improved both FI and FCR and increased BWG. On the other hand, **Khan et al. (2010)** indicated that there was an increase in FI as CP levels were decreased from 23% to 19% but without affecting FCR.

Data of the dressed carcass of broiler chicks fed LCP-diets (10%) and supplemented with lysine (0, 0.2 or 0.4 %) revealed that decreasing dietary protein significantly decreased carcass weight and increased abdominal fat. Lysine supplementation by 0.4% over that of **NRC (1994)** significantly improved dressed carcass and reduced abdominal fat (Table 3). **Kerr and Kidd (1999)** found that a reduction in dietary CP decreased carcass yields, increased abdominal fat, and decreased breast meat yield. Increasing fat accretion in broilers fed the LCP-diet may be attributed to increased feed intake as an attempt to meet their protein and amino acid requirement and the extra energy consumed will be deposited as fat. Similarly, it was found that abdominal fat was significantly increased in association with feeding low-CP diet (18 to 16%) in the finisher period (**Yonemochi et al., 2009; and Yamazaki et**

**al., 2006**). However, it is clear from our results that the LCP-diet supplemented with 0.4% lysine decreased carcass fat deposition near to that of the control diet.

#### **Serum metabolites and HI titer :**

The study of the data related to serum metabolites (Table 4) showed that decreasing DCP level by 10 % than that of **NRC (1994)** significantly ( $p < 0.05$ ) decreased serum total proteins, albumin and globulin and increased total lipids, triglycerides, cholesterol, HDL and LDL. Lysine supplementation by 0.4% above that of **NRC (1994)** to the LCP-diets increased serum total proteins. The increase in plasma proteins was concurrent with an increase ( $P < 0.05$ ) in plasma albumin (non specific immune factor) and an increase ( $P < 0.05$ ) in globulin, the specific immune factor and a decrease ( $P < 0.05$ ) in A/G ratio. The decrease in total lipids, triglycerides, cholesterol, HDL and LDL in broilers were not significantly ( $p < 0.05$ ) different from that of the broilers fed the normal recommended dietary protein and lysine levels. **Thomas and Combs (1967)** indicated that feeding a LCP-diet decreased total plasma protein and vice versa. This may be attributed to the finding of **Harper (1975)** who found that dietary protein served as a precursor for plasma proteins. However, **Corzo et al. (2005)** reported that total plasma protein was not affected by feeding LCP-diet supplemented with EAAs and NEAAs. On the other hand, **Kaamal and Awadalla (2007)** fed the broiler chicks 2% LCP diet with increasing utilizable lysine and methionine levels by 0.25 - 0.1% and 0.2 - 0.1% in starter and finisher diets respectively and found significant increases in serum total protein and albumin.

The effects of feeding the LCP-diet with lysine supplementation on haemagglutination inhibition (HI) titer to NDV are presented in table 4. At 7 day post vaccination (14 days of age), the HI titer was low for broiler chick fed LCP-diet supplemented with 0 or 0.2% lysine. However, lysine supplementation by 0.4% of the diet increased HI titer. By 21 day of age, the HI titer of all treatment start to increase however the highest titer was for the control group followed by broiler chicks group fed LCP-diet supplemented with 0.4% lysine which were not significantly different from the control group. At 28 day of age, the HI titer of different broiler groups reach its highest value and the higher titer was for the broiler chicks fed the recommended CP and lysine levels followed by the broiler chicks fed the LCP-diet supplemented with 0.4% lysine and the lower HI titer was recorded for broiler chick fed the LCP diet supplemented with 0, or 0.2% lysine. **Carlomagno et al. (1980)** found that protein deficiency inhibited antibody production and the development of antibody producing cells in response to T-dependent antigens. **Asibo and Kabasa**

**(1996)** revealed that feeding broiler chicks vaccinated against Newcastle disease diets containing high ME/CP ratios reduced IgM, IgG and HI titers. A deficiency of lysine may reduce immune response of the chicken and resulted in a reduced effect of vaccination. **Chen et al. (2003)** showed that lysine had an influence on chicken immune function and humoral immunity assay and cell mediated immunity function and showed a reduction in antibody response to NDV vaccination in broiler chickens fed a lysine- deficient diet. Also, **Klasing et al. (2000)** found that increasing the dietary concentration of lysine improved the haemagglutination and agglutinin titres, and IgG and IgM levels. Similarly, **Manzoor et al. (2003)** revealed that birds on high protein diets showed higher antibody and cell mediated responses than birds on low protein diets. From the present study it could be concluded that dietary protein level could be reduced by 10% of recommended level with supplementation of lysine at a level of 0.4 % above the recommended level with no adverse effect on growth performance, serum metabolites and immune response.

**Table 2.** Effect of dietary protein level and methionine supplementation on growth performance (mean  $\pm$ SE) of broiler chickens

Methionine supplementation % period	Dietary Protein & lysine levels (%)			
	Control	10% lower than recommended		
		0	0.1	0.2
Initial body weight, g	46.10 $\pm$ 1.33	46.95 $\pm$ 1.73	45.54 $\pm$ 1.65	44.7 $\pm$ 1.60
Body weight, g	139.56 <sup>a</sup> $\pm$ 5.28	124.46 <sup>b</sup> $\pm$ 3.45	130.6 <sup>ab</sup> $\pm$ 3.15	133.82 <sup>ab</sup> $\pm$ 4.55
Body gain, g	93.45 <sup>a</sup> $\pm$ 4.48	77.49 <sup>c</sup> $\pm$ 3.58	85.27 <sup>bc</sup> $\pm$ 3.39	89.28 <sup>abc</sup> $\pm$ 3.92
Feed consumption, g	124.6 <sup>a</sup> $\pm$ 5.77	126.72 <sup>a</sup> $\pm$ 7.57	107.5 <sup>ab</sup> $\pm$ 9.81	94.1 <sup>b</sup> $\pm$ 8.08
Feed conversion ratio	1.08 <sup>b</sup> $\pm$ 0.05	1.43 <sup>a</sup> $\pm$ 0.11	1.34 <sup>ab</sup> $\pm$ 0.08	1.12 <sup>b</sup> $\pm$ 0.1
Body weight, g	331.7 <sup>a</sup> $\pm$ 9.40	280.7 <sup>b</sup> $\pm$ 6.45	285.2 <sup>b</sup> $\pm$ 9.01	299.2 <sup>ab</sup> $\pm$ 7.82
Body gain, g	192.20 <sup>a</sup> $\pm$ 5.94	156.59 <sup>b</sup> $\pm$ 6.51	180.27 <sup>a</sup> $\pm$ 9.18	181.47 <sup>a</sup> $\pm$ 9.71
Feed consumption, g	235.3 <sup>ab</sup> $\pm$ 11.54	263.0 <sup>a</sup> $\pm$ 13.69	256.2 <sup>ab</sup> $\pm$ 15.62	220.9 <sup>b</sup> $\pm$ 16.6
Feed conversion ratio	1.22 <sup>b</sup> $\pm$ 0.12	1.78 <sup>a</sup> $\pm$ 0.13	1.58 <sup>a</sup> $\pm$ 0.16	1.28 <sup>b</sup> $\pm$ 0.11
Body weight, g	625.3 <sup>a</sup> $\pm$ 17.69	547.5 <sup>c</sup> $\pm$ 16.55	570.8 <sup>bc</sup> $\pm$ 20.46	597.2 <sup>ab</sup> $\pm$ 18.39
Body gain, g	291.57 <sup>a</sup> $\pm$ 11.30	266.80 <sup>ab</sup> $\pm$ 11.3	259.74 <sup>b</sup> $\pm$ 13.34	281 <sup>ab</sup> $\pm$ 9.80
Feed consumption, g	453.9 <sup>b</sup> $\pm$ 23.09	518.6 <sup>ab</sup> $\pm$ 27.71	487.2 <sup>ab</sup> $\pm$ 27.03	471.0 <sup>ab</sup> $\pm$ 23.67
Feed conversion ratio	1.58 <sup>b</sup> $\pm$ 0.17	1.95 <sup>a</sup> $\pm$ 0.19	1.82 <sup>ab</sup> $\pm$ 0.20	1.62 <sup>b</sup> $\pm$ 0.14
Body weight, g	1027.5 <sup>a</sup> $\pm$ 32.54	927.2 <sup>b</sup> $\pm$ 23.53	948.87 <sup>b</sup> $\pm$ 23.15	988.02 <sup>ab</sup> $\pm$ 26.2
Body gain, g	403.80 $\pm$ 17.19	379.36 $\pm$ 12.76	378.03 $\pm$ 14.48	390.86 $\pm$ 16.05
Feed consumption, g	691.7 <sup>b</sup> $\pm$ 31.79	824.8 <sup>a</sup> $\pm$ 32.33	758.6 <sup>ab</sup> $\pm$ 34.03	706.8 <sup>b</sup> $\pm$ 32.33
Feed conversion ratio	1.74 <sup>b</sup> $\pm$ 0.20	2.18 <sup>a</sup> $\pm$ 0.26	1.95 <sup>ab</sup> $\pm$ 0.24	1.80 <sup>b</sup> $\pm$ 0.16
Body weight, g	1448.7 <sup>a</sup> $\pm$ 39.10	1286.3 <sup>c</sup> $\pm$ 30.1	1360.9 <sup>bc</sup> $\pm$ 29.2	1393.4 <sup>ab</sup> $\pm$ 33.1
Body gain, g	411.7 <sup>a</sup> $\pm$ 17.82	359.14 <sup>b</sup> $\pm$ 13.26	422.36 <sup>a</sup> $\pm$ 19.30	405.65 <sup>a</sup> $\pm$ 15.64
Feed consumption, g	896.5 <sup>b</sup> $\pm$ 34.64	908.3 <sup>ab</sup> $\pm$ 44.81	1031.5 <sup>a</sup> $\pm$ 40.99	884.3 <sup>b</sup> $\pm$ 40.10
Feed conversion ratio	2.06 <sup>c</sup> $\pm$ 0.24	2.52 <sup>a</sup> $\pm$ 0.32	2.38 <sup>ab</sup> $\pm$ 0.30	2.18 <sup>bc</sup> $\pm$ 0.20
Body weight, g	1943.8 <sup>a</sup> $\pm$ 52.73	1706.66 <sup>c</sup> $\pm$ 39.5	1824.9 <sup>bc</sup> $\pm$ 40.3	1874.3 <sup>ab</sup> $\pm$ 44.6
Body gain, g	504.1 <sup>a</sup> $\pm$ 18.27	420.66 <sup>b</sup> $\pm$ 16.7	456.63 <sup>ab</sup> $\pm$ 25.5	480.24 <sup>a</sup> $\pm$ 19.02
Feed consumption, g	1181.0 $\pm$ 55.72	1237.0 $\pm$ 57.73	1256.7 $\pm$ 61.19	1215 $\pm$ 66.39
Feed conversion ratio	2.39 <sup>c</sup> $\pm$ 0.27	2.92 <sup>a</sup> $\pm$ 0.34	2.70 <sup>ab</sup> $\pm$ 0.35	2.51 <sup>bc</sup> $\pm$ 0.24

Means in the each row followed by a different superscripts are significantly differ ( $p < 0.05$ ).

**Table 3.** Effect of dietary protein level and methionine supplementation on allover growth performance and carcass quality (mean  $\pm$ SE) of broiler chickens

Methionine supplementation %	Dietary protein & methionine levels (%)			
	Control	10% lower than recommended		
		0	0.1	0.2
Items				
Live body weight (g)	1943.8 <sup>a</sup> $\pm$ 52.7	1706.7 <sup>c</sup> $\pm$ 39.5	1824.9 <sup>bc</sup> $\pm$ 40.3	1874.3 <sup>ab</sup> $\pm$ 44.6
Body weight gain (g)	1897.4 <sup>a</sup> $\pm$ 51.9	1659.6 <sup>c</sup> $\pm$ 39.5	1779.31 <sup>b</sup> $\pm$ 40.1	1830.9 <sup>ab</sup> $\pm$ 44.3
Feed intake (g)	3459.5 <sup>c</sup> $\pm$ 69.35	3878.2 <sup>a</sup> $\pm$ 69.0	3907.5 <sup>a</sup> $\pm$ 87.75	3592 <sup>bc</sup> $\pm$ 70.2
Feed conversion ratio	1.82 <sup>c</sup> $\pm$ 0.08	2.33 <sup>a</sup> $\pm$ 0.15	2.17 <sup>ab</sup> $\pm$ 0.17	1.96 <sup>bc</sup> $\pm$ 0.09
Protein efficiency ratio	2.77 <sup>a</sup> $\pm$ 0.17	2.12 <sup>b</sup> $\pm$ 0.19	2.25 <sup>b</sup> $\pm$ 0.24	2.28 <sup>b</sup> $\pm$ 0.30
Cost of Kg BWG	3.60	4.39	4.24	3.84
Dressed carcass %	73.40 <sup>a</sup> $\pm$ 2.75	63.91 <sup>b</sup> $\pm$ 4.45	67.54 <sup>ab</sup> $\pm$ 2.44	69.58 <sup>ab</sup> $\pm$ 1.65
Abdominal fat%	1.10 <sup>c</sup> $\pm$ 0.11	1.88 <sup>a</sup> $\pm$ 0.14	1.50 <sup>b</sup> $\pm$ 0.08	1.35 <sup>bc</sup> $\pm$ 0.06

<sup>bc</sup> means in the each row followed by a different superscripts are significantly differ (P < 0.05).

**Table 4.** Effect of dietary protein levels and methionine supplementation on serum metabolites and HIT of broiler chickens

methionine suppl. %	Diet. protein & methionine levels (%)			
	Control	10% lower than recommended		
		0	0.1	0.2
Items				
Total protein (mg/dl)	4.14 <sup>a</sup> $\pm$ 0.14	2.79 <sup>c</sup> $\pm$ 0.10	3.28 <sup>bc</sup> $\pm$ 0.18	3.75 <sup>ab</sup> $\pm$ 0.11
Albumin (g/dl)	1.85 <sup>a</sup> $\pm$ 0.05	1.45 <sup>c</sup> $\pm$ 0.07	1.53 <sup>bc</sup> $\pm$ 0.02	1.68 <sup>ab</sup> $\pm$ 0.06
Globulin (g/dl)	2.28 <sup>a</sup> $\pm$ 0.09	1.34 <sup>c</sup> $\pm$ 0.07	1.74 <sup>bc</sup> $\pm$ 0.14	2.07 <sup>ab</sup> $\pm$ 0.08
A/G ratio	0.76 <sup>c</sup> $\pm$ 0.11	1.18 <sup>a</sup> $\pm$ 0.11	0.90 <sup>ab</sup> $\pm$ 0.13	0.81 <sup>ab</sup> $\pm$ 0.07
Glucose (mg/dl)	102.86 $\pm$ 2.66	111.09 $\pm$ 4.23	109.49 $\pm$ 4.98	106.66 $\pm$ 3.32
Total lipid (g/dl)	398.6 <sup>c</sup> $\pm$ 7.05	509.34 <sup>a</sup> $\pm$ 5.9	448.24 <sup>b</sup> $\pm$ 7.26	420.82 <sup>bc</sup> $\pm$ 8.95
Total cholesterol (mg/dl)	140.1 <sup>c</sup> $\pm$ 6.25	197.59 <sup>a</sup> $\pm$ 3.69	181.47 <sup>ab</sup> $\pm$ 6.39	157.23 <sup>bc</sup> $\pm$ 6.54
Triglycerides (mg/dl)	53.2 <sup>c</sup> $\pm$ 2.32	90.81 <sup>a</sup> $\pm$ 3.95	74.08 <sup>ab</sup> $\pm$ 2.57	65.37 <sup>bc</sup> $\pm$ 3.02
HDLP (mg/dl)	86.1 <sup>c</sup> $\pm$ 3.21	110.65 <sup>a</sup> $\pm$ 4.40	102.65 <sup>ab</sup> $\pm$ 5.60	98.37 <sup>abc</sup> $\pm$ 3.75
LDL (mg/dl)	43.5 <sup>c</sup> $\pm$ 2.88	70.16 <sup>a</sup> $\pm$ 4.80	62.28 <sup>ab</sup> $\pm$ 2.25	52.3 <sup>bc</sup> $\pm$ 3.10
HI titer (log2) at 14 days	6.24 <sup>a</sup> $\pm$ 0.22	4.03 <sup>b</sup> $\pm$ 0.20	4.28 <sup>b</sup> $\pm$ 0.31	5.55 <sup>a</sup> $\pm$ 0.34
HI titer (log2) at 21 days	6.62 <sup>a</sup> $\pm$ 0.25	5.05 <sup>c</sup> $\pm$ 0.22	5.26 <sup>bc</sup> $\pm$ 0.38	6.20 <sup>ab</sup> $\pm$ 0.40
HI titer (log2) at 28 days	9.92 <sup>a</sup> $\pm$ 0.30	7.01 <sup>c</sup> $\pm$ 0.28	7.69 <sup>bc</sup> $\pm$ 0.40	8.94 <sup>ab</sup> $\pm$ 0.47

<sup>abc</sup> means in the each row followed by a different superscripts are significantly differ (P < 0.05).



**REFERENCES**

**Abdallah, A. G.; Soliman, A. Z. and Saudi, M. (2005)** : Response of broiler chicks to low protein diets supplemented with amino acids administered through feed or drinking water under heat stress conditions. Animal Production Research Institute, Second Conference and the Regional Symposium on Buffalo Production and Future prospects, Sakha, Kafr El-Sheikh, Egypt.

**Aletor, V. A.; Hamid, I. I.; Niebß, E. and Pfeffer, E. (2000)** : Low-protein amino acid-supplemented diets in broiler chickens: Effects on performance, carcass characteristics, whole-body composition and efficiencies of nutrient utilization. *J. Sci. Food Agric.* 80 : 547 - 554.

**Allain, C.; Poon, L. and Chan, C. (1974):** Enzymatic determination of total serum cholesterol. *Clin. Chem* 20, 470.

**Alleman, F. and Leclercq, B. (1997)** : Effect of dietary protein and environmental temperature on growth performance and water consumption of male broiler chickens. *Br. Poult. Sci.* 38: 607-610.

**Asibo, J. O. and Kabasa, J. D. (1996)** : Serum immunoglobulin responses among broiler chickens fed ugandan commercial poultry feed and vaccinated against Newcastle disease. *Trop - Anima. Science Prod.* 28 : 163-173.

**AOAC (1990)** : Association of official analytical chemist. Official methods of Analysis. 13th Ed. Washington DC.USA.

**Attia, Y. A.; Nawar, M. E.; Osman, M. and Abd-El-hady, S. B. (1998)** : optimum levels of metabolizable energy and crude protein in the rations for Avian 34 broiler chicks. Proceeding. 10<sup>th</sup> Conference of Egyptian Society of Animal Production, Assiut Univ- ersity, Egypt, pp. 223–241.

**Burstein, M.; Scholnick, H. and Morfin, R. (1970)** : Rapid method for isolation of lipoproteins from human serum by precipitation with polyanions. *Journal of lipid research* 11., 583-595.

**Carlomagno, M. A.; Alito, A. E.; Rife, S. U. and Glimeno, A. L. (1980)** : B cell immune response during totalprotein deprivation. *Acta Physiol. Lat. Am.*, 30:187-192.

**Chen, C.; Sander, J. E. and Dale, N. M. (2003)** : The effect of dietary lysine deficiency on the immune response to Newcastle disease vaccination in chickens, *Avian. Dis.* 47: 1346-1351.

**Corzo, A.; Dotzer, W. A. and Kidd, M. T. (2006)** : Dietary lysine needs of late- developing heavy broilers. *Poult. Sci.*, 85: 457-461.

**Corzo, A.; Fritts, C. A.; Kidd, M. T. and Kerr, B. J. (2005)** : Response of broiler chicks to essential and non-essential amino acid supplementation of low crude protein diets. *Anim. Feed Sci. Technol.* 118 : 319-327.

**Corzo, A.; Kidd, M. T.; Burnham, D. J. and Kerr, B. J. (2004)** : Dietary glycine needs of broiler chicks. *Poult. Sci.* 83 : 1382-1384.

**Doumas, B.; Waston, W. and Biggs, H., (1971)** : Albumin Standards and the measurement of serum albumin with bromocresol green. *Clin. Chim. Acta.* 31, 87-96.

**Eits, R. M.; Kwakkel, R. P.; Versteegen, M. W. A. and DenHartog, L. A. (2005)** : Dietary balanced protein in broiler chickens. 1. A flexible and practical tool to predict dose-response curves. *Br. Poultry Sci.* 46(3): 300-309.

**El-Hussein, O. M.; Abou -El-Wafa, S. and El-Komy H. M. A. (2004)** : Nutrients allowance for broiler performance. *Egypt. Poult. Sci.*, 24: 575-595.

**El-Sherbiny, A. E., Moahmed. M. A., Hamza, A. S. and El-Affi. T. M. (1997)** : Response of broiler chicks to low protein diets supplemented with lysine and methionine. *Egypt Poult. Sci.* 17: 23-28-5-595.

**Fassati, P. and Precipe, L. (1982)** : Serum triglycerides determined calorimetrically with an enzyme that produce hydrogen peroxide. *Principle, J. Clinic. Chemistry* 28, 20-77.

**Firman, J. D. and S. D. Boling (1998)** : Ideal protein in turkeys. *Poult. Sci.*, 77: 105-110.

**Friedewald, W.; Levy, R. and Frederickson, D. (1972)** : Estimation of the concentration of low density lipoprotein cholesterol in plasma with out use of ultra centrifuge. *Clin. Chem.* 18, 499-502.

**Goarnal, A.; Bardawill, C. and David, M.**

**(1949)** : Determination of serum protein by means of biuret reaction. *J. Biol chem.* 177-751.

**Harper, H. A. (1975)** : The blood, lymph and cerebrospinal fluid. Chapter 10 in *Review of Physiological Chemistry*. 15 th ED. By Harper, H.H, LANG Medical Publications.

**Ichhponani, N. S.; Lodhi, G. N. and Ichhponani, J. S. (1981)** : Effect of supplementing lysine and methionine in the vegetable protein diets on the performance of broilers. *Indian Journal of Poultry Science.* 16: 28-43.

**Iwe, M. O. and Onadipe, O. O. (2001)** : Effect of addition of extruded full-fat soy flour into sweet potato flour on functional properties of the mixture, *J. Sustain. Agri. Environ.*, 3: 109-117.

**Kaamal, A. and Awadalla, S. A. (2007)** : Effect of levels of protein and amino acids (methionine and lysine) on some biochemical parameter and performance of broilers. *J. Egypt. Vet. med. Asse.* 67, no 2: 123-134.

**Kerr, B. J. and Kidd, M. T. (1999)** : Amino acid supplementation of low-protein broiler diets: 1. Glutamic acid and indispensable amino acid supplementation. *J. Appl. Poultry Res.* 8:298-309.

**Khalifa, M. M. (2001)** : Effect of protein/energy feeding systems on the performance of two local chicken strains. Ph. D. Thesis, Faculty of Agriculture- Kfr- El-Sheikh, Tanta University.

**Khan, S.; Jabbar, M. A.; Mehmud, A.; Abbas, M. M. and Mahmood, A. (2010) :** Effect of lysine supplementation in low protein diets on the performance of growing broilers. *Pakistan Vet J*, 30 (1): 17-20.

**Kidd, M. T.; Kerr, B. J.; Firman, J. D. and Boling, S. D. (1996) :** Growth and carcass characteristics of broilers fed low protein-threonine supplemented diets. *J. Appl. Poult. Res.* 5: 180-190.

**Klasing, K. C. and Calvert, C. C. (2000) :** The care and feeding of an immune system: an analysis of lysine needs. *Proceedings of the VIIIth International Symposium on Protein Metabolism and Nutrition.* G. E. Loble, A. White and J. C. MacRae. Wageningen, Wageningen Press: 253-264.

**Manzoor, A.; Cheema, M.; Qureshi, A. and Havenstein, G. B. (2003) :** A comparison of the immune profile of commercial broiler strains when raised on marginal and high protein diets. *Int. J. Poult. Sci.* 2 (5): 300.

**May, J. D.; Lott, B. D. and Simmons, J. D. (1998) :** The effect of environmental temperature and body weight on growth rate and feed:gain of male broilers. *Poult. Sci.* 77: 499-501.

**Nahm, K. H. (2002) :** Efficient feed nutrient utilization to reduce pollutants in poultry and swine manure. *Crit. Rev. Environ. Sci. Technol.* 32 (1): 1-16.

**Namroud, N. F.; Shrivazad, M. and Zahari, M. (2008) :** Effects of fortifying low crude protein diet with crystalline amino ac-

ids on performance, blood ammonia level, and excreta characteristics of broiler chicks. *Poultry Sci.* 87:2250-2258.

**Nascimento, A. A. (2003) :** Lisina otimiza a converso alimentar. *Revista Ave World.* Ano1, n.5, outubro/novembro, p.48-50.

**NRC (1994) :** National Research Council .Nutrient Requirement of Poultry. 9th Rev. ed. National Academy Press, Washington, DC.

**Nworgu, F. C.; Egbunike, G. N.; Fapho-hunda, J. B.; Ogundola, F. O.; Omela, A. G.; Ogbosuka, G. E. and Okpeze, C. N. (2000) :** Performance of broiler chicks fed full-fat extruded soybean meal and full-fat soybean meal *Trop. Anim. Prod. Invest.*, 3: 47-54.

**Payne, C. J.; Scott, T. R.; Dick, J. W. and Glick, B. (1990) :** Immunity to *Pasteurella multocida* in proteindeficient chickens. *Poult. Sci.*, 69: 2134-2142.

**Snedecor, G. W. and Cochran, W. G. (1967) :** *Statistical Methods.* 6th ed., Oxford and IBN Publishing CO., Calcutta. PP. 344-346.

**Sterling, K. G. and Pesti, G. M. (2003) :** Performance of broiler chicks fed various levels of dietary lysine and crude protein. *Poult. Sci.* 82: 1939-1947.

**Sterling, K. G.; Pesti, G. M. and Bakalli, R. I. (2006) :** Performance of different broiler genotypes fed diets with varying levels of dietary crude protein and lysine. *Poult. Sci.* 85:1045-1054.

**Thomas, O. P. and combs, G. F. (1967) :** Relationship between serum protein level and body composition in the chick. *J. Nutr.*, 91: 468-478.

**Trinder, P. and Ann, D. (1969) :** Glucose: Enzymatic Colorimetric Test (GOD-PAP). *Clin. Biochem.* 6:24.

**Tsiagbe, V. K.; Cook, M. E.; Harper, A. E. and Sunde, M. L. (1987) :** Enhanced immune responses in broiler chickens fed methionine-supplemented diets. *Poultry Science* 66, 1147-1154.

**Webel, D. M.; Johnson, R. W. and Baker, D. H. (1998) :** Lipopolysaccharide-induced reductions in food intake do not decrease the efficiency of lysine and threonine utilization for protein accretion in chickens. *Journal of Nutrition* 128, 1760-1766.

*Poult. Sci.*, 55: 243-253.

**Yamazaki, M.; Murakami, H.; Nakashima, K.; Abe, H. and Takemasa, M. (2006) :** Effects of excess amino acids in low protein diet on abdominal fat deposition and nitrogen excretion of the broiler chicks. *J. of Poult. Sci.* 43: 150-155.

**Yonemochi, C.; Fujisaki, H. and Takagi, H. (2003) :** Effects of amino acid, enzyme mixture and phytase added to low protein and low phosphorus diet on performance and excretion of nitrogen and phosphorus in broilers. *J. Poult. Sci.* 40 : 114-120.

**Zollner, N. and Kirsch, K. (1962) :** Determination of the total lipid concentration in serum. *Zentralblatt Fur Gesamte Experimentale Medizin* 135,545-549.

## الملخص العربى

### تأثير مستويات البروتين والميثيونين فى علائق دجاج التسمين على معدلات النمو والاستجابة المناعية

أ.د/ عبد الهادى محمد عرمه      أ.د/ طارق ابراهيم محمد  
ط.ب/ احمد احمد محمد ابوالنجا

قسم التغذية وأمراض سوء التغذية - كلية الطب البيطري - جامعة المنصورة

أجريت التجربة على 120 كتكوت عمر يوم من سلالة كب ثم تقسيمهم الى 4 مجموعات غذيت على علائق بادية ونامى وناهى تحتوى على 3200 كيلو كالورى طاقة تمثيل غذائى مع خفض نسبة البروتين بنسبة 10% من مستوى البروتين فى العليقة وزيادة نسبة الميثيونين بمعدل 0.1 و 0.2 % فوق مستوى الميثيونين فى العليقة ولذا احتوت العلائق على نسبة 20.8% بروتين و 1.3؛ 1.5 و 1.7% ليسين و 3200 كيلو كالورى طاقة فى عليقة البادية و 18.2% بروتين و 1؛ 1.22 و 1.4% ميثيونين و 3200 كيلو كالورى طاقة فى عليقة النامى و 16.2% بروتين و 3200 كيلو و كالورى طاقة 0.93؛ 1.15 و 1.33 ميثيونين فى عليقة الناهى مع استخدام علائق ضابطة تقليدية تحتوى على 3200 كيلو كالورى و 20؛ 23 و 18% بروتين و 1.2؛ 1.4 و 1.6% ميثيونين فى عليقة البادية والنامى والناهى بالترتيب. وتم تحصين كل الكتاكيت ضد أمراض الأنفلونزا والنيوكاسل والجمبورو. وتم وزن الكتاكيت عند نهاية كل اسبوع من العمر لمتابعة النمو وتم تسجيل العلف المستهلك اسبوعيا. وكذلك تم حساب معدل التحويل الغذائى اسبوعيا. كما أخذت عينات من الدم من 5 كتاكيت عند اليوم 18؛ 21 و 28 و تم فصل مصل الدم لقياس مستويات الأجسام المناعية لمصل النيوكاسيل وعند نهاية التجربة لقياس بعض مكونات مصل الدم ( البروتينات الكلية-الجلوبولين-الالبومين-الجلوكوز-الدهون الكلية-الكولسترول-الجلسليريدات). وعند نهاية التجربة ( 6 أسابيع) تم اخذ 5 دجاجات من كل مجموعة وذبحها وقياس الوزن قبل وبعد الذبح وبعد ازالة الاحشاء الداخلية وقياس وزن دهن البطن. وقد خلصت نتائج هذه الدراسة إلى:

- خفض مستوى البروتين بمعدل 10% من مستوى البروتين فى العليقة ادى الى نقص فى وزن الجسم وزيادة فى كمية العلف المستهلك. اضافة الميثيونين بمعدل 0.4% من مستوى الميثيونين فى العليقة ادى الى تحسين فى وزن الجسم؛ خفض معدل استهلاك العلف؛ خفض معدل التحويل الغذائى؛ تحسين نسبة صافى الذبيحة وخفض نسبة الدهن فى الجسم مقارنة بالمجموعة التى اضيف اليها

الميثيونين يعدل 0.2% الى الليسين فى العليقة.  
- اضافة الميثيونين يعدل 0.2% من مستوى الليسين فى العليقة الى العلائق التى تم خفض نسبة البروتين بمعدل 10% من مستوى البروتين فى العليقة ادى الى زيادة معنوية فى مستوى البروتينات الكلية-الجلوبيولين-الالبومين ونقص فى مستوى الدهون الكلية - الكولسيترول- الجلسيريدات مقارنة بالمجموعة التى اضيف اليها الميثيونين يعدل 0.2% من مستوى الميثيونين فى العليقة.