

Effect of road transportation on body weight loss, respiration rate, pulse rate, and rectal temperature in fed *ad libitum*, fasted or electrolytes drenched buffalo.

Raghda A. S. Taghian, F. M. Allam, Abd El-Ati, M. N., and S. F. Abbas.

Fac. Agric., Dept Animal and Poultry Production, Assiut Univ.

Abstract:

This study was carried out at the Animal Experimental Farm, Animal and Poultry Production Department, Faculty of Agriculture, Assiut University. This work was conducted to investigate the effects of truck transportation on live body weight with and without fasting and some physiological parameters in buffalo. Also the effect of drenched electrolyte fluid on transportation stress was investigated. Fifteen mixed sexes buffalo calves separated equally to three groups according to their body weights were used in the present study. Group A Calves ranged in body weight from 126 to 163.5 kg, group B from 176 to 252 kg, and group C from 275 to 368 kg. Insignificant differences among the three animal groups were found in all of the studied parameters. Transportation induced increment percentage of body weight loss, rectal temperature, and increment followed by decrement in respiration rate and pulse rate. Buffaloes Fed *ad libitum* lost more body weight than fasted ones, while drenched electrolyte fluids was less affected

by transportation stress. Transport caused rise in rectal temperature and increase in respiration rate and pulse rate in treated buffaloes. Meanwhile, transported and fed electrolyte drenched buffaloes were less affected by transportation stress. Therefore, drenching the electrolyte fluids treatment had ameliorated the detrimental effects of transport stress throughout the studied parameters such as body weight loss, rectal temperature, respiration rate and pulse rate.

Key words: Buffalo, Transportation stress, Body weight, Electrolyte.

Introduction:

Stress is defined as conduction in an animal that results from the action of one or more stressors that may be of either external or internal origin, whether a stressor can be considered as a harmful depends on the way an organism is able to cope with a threatening situation as it regains homeostasis (Von Borell, 2001). Transportation is an inevitable husbandry practice, which livestock are subjected to as a major stressor (Adenkola *et al.*, 2009a and b Bisalla *et al.*, 2011).

Referees: Prof. Dr. Soliman.M.Mousa,

Prof. Dr. Mohamed.A.Ebrahim.

Received on: 13/6/2012

Accepted for publication on: 21/6/2012

In Egypt, livestock are transported often without interest of any welfare order and in vehicles designed for transportation of goods and not specifically for transportation of animals such as open trailers, trucks, pick-ups.

There is scarcity of researches dealing with the effects of transportation on buffalo. However, earlier work has shown that transport leads to loss of body weight. After a stress response, negative effect on growth rate has been observed (Broucek *et al.*, 1983). Long transportation of food animals without water and feeds, which occurs very often, is accompanied by drastic loss in body weight (Ayo *et al.*, 1996). Adenkola *et al.*, (2009a) showed that pigs transported for 4 h lose 19.50% while those administered with ascorbic acid lose 7.29% indicating that ascorbic acid was able minimize loss often encountered with road transportation of livestock. Atkinson (1992) reported that dehydration occurs in transported calves and that keeping them in lairage may help in their recovery. Baldock and Sibly (1990) observed that placing ewes in a stationary trailer had no effect on heart rates, but transportation of ewes for 20 min in the trailer produced an increase of 12 beats per minute. Adenkola *et al.*, (2008) reported that rectal temperature recorded during 8-h and 4-h journey. Whythens *et al.*, (1985) added that body fluid shifts could be seen to occur

quite rapidly. Fluid supplementation has been shown to be effective in reducing the detrimental effects of transport stress on cattle (Sranmek and Pozdosek, 1987 and Schaefer *et al.*, 1990). The economic ramifications of these findings are clear, however, the biological basis of the effects of fluid treatment is less well defined. There was substantial research work on the effects of handling and transportation of cattle, pigs and poultry (Rollin, 1995). Little work was carried out to assess the effects of stress in transported buffaloes, sheep and goats especially under hot and subtropical conditions. This study was carried out to investigate the effects of *ad libitum*, fasted and drenched with electrolytes on truck transported buffaloes of different weights.

Materials and methods

Fifteen mixed sexes buffalo calves were divided equally into three groups according to there body weights were used in the present study. Group A calves ranged in body weight from 126 to 163.5 kg, group B from 176 to 252 kg, and group C from 275 to 368 kg.

All groups were subjected to the following three treatments:

- 1) Transportation after *ad libitum* feeding.
- 2) Transportation after Fasting for 16 h.
- 3) Transportation after *ad libitum* feeding and drenched with electrolytes fluid (two liters/head) which contains sodium chloride

8.6g, Potassium chloride 0.3g and Calcium chloride $2H_2O$ 0.33 g.

All Animals were fasted during the trip and four h after arrival (time needed to finish all measurements). The trip lasted for about 3 h using truck for 250 km distance at speed ranged from 60-80 km/h. The ambient temperature during the trip and while taking the measurements was around 28 to 34 °C. The trip commenced at 7:00 am. At the day of transport, the measurements were taken just before loading the animals (Time 1), immediately after uploading (Time 2), two hours after arrival (Time 3), and four hours after arrival (Time 4). The following measurements were taken:

- 1- Body weight (kg).
- 2- Rectal temperature (°C).
- 3- Pulse rate (beat/min.) estimated by counting the beat of Jugular vein for one minute.
- 4- Respiration rate (breath/min.) determined by counting the flank movements for one minute.

Statistical analysis: Data were statistically analyzed using three factors (group, treatment and time) analysis of variance, Duncan's multiple range test and multiple linear correlation as described by Gomez and Gomez (1984). All calculations were performed using SPSS/PC.

Results and discussion

No significant differences among the three animal groups were detected in all the studied parameters.

1. Effect of transportation on body weight loss.

Data in Table (1) and Figure (1) showed that road transport caused highly significant loss in buffalo live body weight. Buffaloes in T1 lost 4.2% of their original transport body weight, while those fasted (T2) or electrolytes supplemented (T3) were less affected by transportation (2.55 % and 2.09 %, respectively). Differences between T1 mean and the other two means were highly significant ($P < 0.01$). While, the differences between T2 and T3 means were significant ($P < 0.05$). These results are in agreement with those obtained by Broucek *et al.* (1983) that negative effect on body weight has been observed after animal had been stressed. They added that release of thyroid stimulating hormone (TSH), known to affect growth rates, can be inhibited by negative feed back from adrenocortical hormone after a stress response.

With respect to the effects at different times of taking the measurements, insignificant differences in body weight loss between Time 3 and Time 4 (2 hours and 4 hours after transport) were averaged over the three treatments (4.07 and 4.82 %, respectively). Both values were significantly higher ($P < 0.05$) than the percentage of body weight loss (2.89 %) in Time 1 (immediately after transport).

Both fasted (T2) and electrolytes supplemented (T3) transported buffaloes lost significantly lesser percentages of body weight than those fed *ad libitum* buffaloes (T1). These findings are in concomitance with Schaefer *et al.*, (1997) who reported that the application of oral electrolyte therapy, especially if similar in constituents to interstitial fluid, resulting improvements in both live and carcass weights (less shrink) of up to several percent in treated animals as well as a reduction in meat quality degradation (reduced dark cutting). Similar results were found by Daghash, (2008) that lambs treated with 5% glucose solution were less affected by transportation stress. But, in contrary to our results he found that fasted lambs were more affected by transportation stress than those fed *ad libitum* and transported ones. The same observation was reported by Kannan *et al.*, (2000 & 2002) that transportation and prolonged feed deprivation in

goats may increase stress and live weight losses.

The use of electrolyte solutions for minimizing the effects of stressors on animals in the marketing process has been advocated in the sheep and beef industries without a full understanding of the effects of transport stress on the acid-base physiology of ruminants (Schaefer *et al.*, 1997). Furthermore, Adenkola *et al.* (2009^b) showed that pigs transported for 4 h lost 19.50% while those administered with ascorbic acid lost only 7.29% of live weight indicating that ascorbic acid was able to minimize loss often encountered with road transportation of livestock. Much of this loss is actually from carcass components and not simply gastrointestinal tract fill (Jones *et al.*, 1988; Warriss, 1990 and Schaefer *et al.*, 1997). Indeed, it is well established that different animal species and even animals of the same species but different genetic backgrounds respond differently to the same stressor (Hall *et al.*, 1998).

Table 1: Effect of transportation on body weight loss and some physiological parameters (Mean \pm SD) in fed *ad libitum*, fasted or electrolytes drenched buffalo.

		% Body weight loss	Respiration rate (breath/min.)	Pulse rate (beat/min.)	Rectal temperature (°C)	
<u>TREATMENT</u>		**	**	**	**	
Transp.fed <i>ad libitum</i> (T1)		4.20 ^A ±0.613	23.83 ^A ±1.96	60.45 ^A ±7.65	38.13 ^B ±3.11	
Transp and fasting (T2)		2.55 ^B ±0.412	23.45 ^B ±2.11	55.14 ^C ±4.13	37.66 ^C ±2.981	
Transp and electrolytes (T3)		2.09 ^C ±0.311	21.31 ^C ±1.76	58.14 ^B ±5.16	38.71 ^A ±1.971	
<u>TIME</u>		**	**	**	**	
Before transport (Time 0)		0.00 ^D ±0.00	19.70 ^D ±1.98	51.99 ^D ±4.61	37.53 ^B ±3.25	
Imm. after transport (Time 1)		2.89 ^C ±0.193	25.25 ^A ±2.61	64.42 ^A ±6.11	38.36 ^A ±4.11	
2h after transport (Time 2)		4.07 ^A ±0.246	24.60 ^B ±2.19	59.60 ^B ±6.91	38.33 ^A ±3.87	
4h after transport (Time 3)		4.82 ^A ±0.278	22.84 ^C ±1.96	55.63 ^C ±4.75	38.44 ^A ±4.15	
<u>TIME*TREATMENT</u>		**	**	**	**	
Before transport	Before <i>ad libitum</i>	0.00 ⁱ ±0.00	20.23 ^f ±3.93	52.20 ^j ±3.96	37.62 ^e ±4.13	
	Before fasting	0.00 ⁱ ±0.00	19.67 ^g ±4.76	49.36 ^k ±4.15	37.14 ^f ±4.09	
	Before electrolytes	0.00 ⁱ ±0.00	19.21 ^h ±3.71	54.41 ^h ±6.11	37.82 ^{de} ±3.92	
Time after transportation	0 hour	<i>Ad libitum</i>	3.93 ^d ±0.313	27.28 ^a ±4.61	67.75 ^a ±5.13	38.51 ^b ±3.87
		Fasting	2.74 ^h ±0.251	25.29 ^c ±3.79	62.23 ^c ±4.17	38.02 ^{cd} ±3.11
		Electrolytes	2.00 ⁱ ±0.119	23.17 ^d ±3.81	63.27 ^b ±3.96	38.55 ^b ±4.151
	2 hour	<i>Ad libitum</i>	5.72 ^b ±0.678	25.63 ^b ±4.76	62.22 ^c ±8.71	38.13 ^{cd} ±3.35
		Fasting	3.47 ^c ±0.371	25.64 ^b ±5.81	56.30 ^f ±7.66	37.61 ^e ±4.71
		Electrolytes	3.02 ^g ±0.314	23.19 ^d ±3.71	60.29 ^d ±4.95	39.26 ^a ±4.45
	4 hour	<i>Ad libitum</i>	7.14 ^a ±0.612	22.20 ^e ±2.15	59.64 ^a ±6.12	38.27 ^{bc} ±5.55
		Fasting	3.97 ^c ±0.111	23.22 ^d ±3.97	52.68 ⁱ ±5.42	37.86 ^{de} ±4.76
		Electrolytes	3.35 ^f ±0.105	19.66 ^g ±2.99	54.58 ^g ±4.16	39.20 ^a ±4.04

- Different letter in the same column indicate statically significance (P<0.05) between values according to Duncan's multiple range tests (small superscript letters to compare the interactions). * significant at 5% and ** significant at 1%




Fig 1 : Effect of transportation times on %body weight loss, respiration rate, pulse rate and rectal temperature in experimental buffaloes.

Treatments:

- Fed *ad libitum* and transported.
- Fasted and transported.
- Drenched electrolytes and transported.

Time1: Before transport.

Time2: Immediately after transport.

Time3: 2h after transport.

Time4: 4h after transport.

2. Effect of transportation on respiration rate.

As shown in Table (1) and figure (1), the respiration rate (breath/min) was highly significantly ($P < 0.01$) lower in transported and drenched electrolytes buffaloes (21.31 breath/min) than those fasted (23.45 breath/min) or *ad libitum* fed (23.83 breath/min) transported buffaloes. These results are coinciding with Daghash, (2008) who reported that transported lambs which treated with 5% glucose solution were less affected by transportation stress and showed lower respiration rate. But, in contrary to our results on buffaloes He found that fasted lambs

were more affected by transportation stress than fed *ad libitum* and transported ones. Averaged over the three treatments, the respiration rate was 19.7

(breath/min) before transport (Time 0) and got higher to 25.25 immediately after transport (Time 1) then began to decrease to reach 24.6 then 22.84 at 2h and 4h after transport, respectively. These values highly significantly ($P < 0.01$) differed. Similarly, in study using donkeys during road transportation, Plyaschenko and Sidorov (1987) observed that respiration rate increased from 22.3 to 40.1 breaths per minute within 15 min of loading and it remained high dur-

ing the journey. Transported and given electrolytes buffaloes were significantly less affected by transportation and reached the base line 4h after transport. Indicating that giving the buffaloes electrolyte solutions had alleviated the stressful effects of road transportation. Electrolyte and fluid supplementation has been shown to be effective in reducing the detrimental effects of transport stress on animals (Sranmek and Pozosek, 1987 and Schaefer *et al.*, 1990). Similarly, during transportation of ostriches, Minka and Ayo (2007b; 2008) reported a significant increase in respiration rate of the birds subjected to 6 h road transportation. Daghash (2008) reported that in transported lambs, immediately after transportation, respiration rate was significantly higher than (before transportation). After that, these values were decreased and the lowest values were observed after 4 hours following transportation

3. Effect of transportation on pulse rate.

Table (1) and figure (1) indicate that buffaloes in T1 showed the highest significantly pulse rate (60.45 beat/min) in comparison with buffaloes in T2 which exhibited the lowest pulse rate 55.14 beat/min while intermediate value (58.14 beat/min) was obtained with buffaloes in T3. Furthermore, buffalo road transport resulted in highly significant ($P<0.01$) increase in pulse rate from 51.99 beat/min before transport (Time 0) to 64.42

beat/min immediately after arrival (Time 1). Two hours after arrival (Time 2), pulse rate decreased to 59.6 beat/min and reached 55.63 beat/min at 4h after transport (Time 4). Several investigators have reported the immediate increment in pulse rate after transportation (Bianca, 1976; Vihan and Sahni, 1981; Ayo *et al.*, 1998, Ayo *et al.*, 2005,

Bouwknicht *et al.*, 2007 and Zulkifli *et al.*, 2010).

Obviously, fasted buffaloes exhibited significantly ($P<0.05$) lower pulse rate in the three times after arrival followed by buffaloes that given electrolytes (Table 1). This was inconsistent with the results of Daghash (2008) who reported that transported and fasted lambs exhibited the highest pulse rate followed by fed *ad libitum* and then lambs which treated with 5% glucose solution.

4. Effect of transportation on rectal temperature.

The differences in rectal temperature were around 1 °C. Averaged over the three treatments, the transported animals showed significant higher rectal temperature at Time 1 (38.36 °C) than Time 0 (before transport, 37.53 °C). No significant differences in rectal temperature were found between Time 1, Time 2 and Time3 (38.36, 38.33 and 38.44, respectively) as shown in Table (1) and figure (1).

Transported and electrolytes treated buffaloes (T3)

showed significantly higher rectal temperature (38.71 °C) followed by T1 (38.13 °C) while T2 showed significantly lower rectal temperature (37.66 °C).

This was consistent with the results of Daghash, (2008) who reported that rectal temperature in lambs transported and fed *ad libitum* exhibited the highest values, while the lowest value was found in transported and fasted lambs compared to other treated lambs. Also, Horton *et al.*, (1996) found that rectal temperature was lower in fasted lambs and this supported by the result of Naqui and Rai (1991) who found lower temperature in fasted lambs.

On the topic of body temperature, Bouwknecht *et al.*, (2001) and Zulkifli *et al.*, (2010) found that transportation at ambient temperatures of 30–32°C resulted in hyperthermia (unusually high body temperature) among the goats. The increase in body temperature following transportation could also be attributed to stress-induced hyperthermia. The phenomenon of stress induced hyperthermia has been reported in mammalian species when subjected to mild disturbance (Bouwknicht *et al.*, 2007) and handling (Moe & Bakken, 1997). Stress induced hyperthermia has been closely associated with an activation of the hypothalamic-pituitary-adrenal axis and the sympathetic-adrenal-medullary system.

Rectal temperature (RT), respiratory rate (RR) and heart rate

(HR) are important physiological parameters most relevant for on-the-spot evaluation of the health status and adaptability of animals, including poultry species (Bianca, 1976; Ayo *et al.*, 1998). The parameters are easily measured and are of value in the determination of state of stress especially during the process of transportation in rural areas where laboratory facilities may be lacking (Minka and Ayo, 2007b). These parameters are of importance in evaluating the adaptability of domestic animals to various environmental stress factors (Bianca, 1976; Vihan and Sahni, 1981; Ayo *et al.*, 1998), including transportation stress (Ayo *et al.*, 2005). In contrary, Ali *et al.*, (2006) recorded in desert sheep and goats that road transportation for 2 hours resulted in variable and statistically insignificant increases in heart, pulse and respiratory rates in both control and experimental animals. Therefore, the application of electrolyte solutions to minimize transport stress in animals, while fluid supplementation has been showed to be effective in reducing the detrimental effects of transport stress on cattle, sheep and buffalo.

References:

- Adenkola AY, J.O. Ayo, A.K.B. Sackey, and A.B. Adelaiye (2009a). Modulatory role of ascorbic acid on rectal temperature of pigs transported by road during the harmattan season. *Anim. Prod. Res. Adv.* 5(2): 120- 127.

- Adenkola AY, J.O. Ayo, A.K.B. Sackey, and N.S. Minka (2008). Ameliorative effect of ascorbic acid on rectal temperature of pigs transported by road for eight hours during the harmattan season. Proceedings of the 13th Annual Conference of Animal Science Association of Nigeria. Held in Zaria Kaduna State, Nigeria. pp. 177-181.
- Adenkola, A.Y., J.O. Ayo, A.K.B. Sackey, and A. B. Adelaiye (2009^b) Haematological and serum biochemical changes in pigs administered with ascorbic acid and transported by road for four hours during the harmattan season. *Journal of Cell and Animal Biology* Vol. 3 (2), pp. 021-028.
- Ali, B.H., A.A. Al-Qarawi, and H. M. Mousa (2006). Stress associated with road transport of desert sheep and goats and the effect of pretreatment with xylazine or sedbetaine. *Research Veterinary Science*, vol.80, (3): 343-348.
- Atkinson, P.J. (1992). Investigation of the effects of transport and lairage state on hydration state and resting behaviour of calves for export. *Vet. Rec.* 130: 413-416.
- Ayo JO, S.B. Oladele, and A. Fayomi (1996). Heat stress and its adverse effects on livestock production: A review. *Nig. Vet. J.* 1: 58-68.
- Ayo, J. O., S.B. Oladele, A. Fayomi, S. D. Jumbo, and J. O. Hambolu (1998). Rectal temperature, respiration and heart rates in the Red Sokoto goat during the harmattan season. *Bull. Anim. Hlth. Prod. Afri.* 46, 161-166.
- Ayo, J.O., N.S. Minka, and A. Fayomi, (2005). Effects of ascorbic acid on rectal temperature of pullets transported by road during the hot-dry season in Northern Nigeria. *Proc. 10th Ann. Conf. Anim. Sci. Ass. Nig. Ado-Eketi, Nigeria* 10: 58-60.
- Baldock N.M., and R.M. Sibly (1990). Effects of handling and transportation on the heart rate and behaviour of sheep. *Appl. Anim. Behav. Sci.* 28: 15-39.
- Bianca, W. K. (1976): The significance of meteorology in animal production. *Int. J. Biometeorol.* 20, 139-156.
- Bisalla, M., M.B. Bi-Allah, , and F.O. Nnadiwa, (2011) Some haemato-biochemical parameters of sheep pre-treated with levamisole and transported by road. *Agriculture Journal* 6(4): 131-133.
- Bouwknicht, J.A., B. Olivier and R.E. Paylor, (2007). The stress-induced hyperthermia paradigm as a physiological animal model for anxiety: A review of pharmacological and genetic studies in the mouse. *Neur. Biobehav. Rev.*, 31: 47-59.

- Bouwknicht, J.A., T.H. Hijzen, J. Van Der Gugten, R.A. Maes, R. Hen and B. Olivier. (2001). Absence of 5-HT1B receptors is associated with impaired impulse control in male 5-HT1B knockout mice. *Biol. Psychiatry*, 49: 557-568
- Broucek, J., M. Kovalcikova, and K. Kovalcik, 1983. The effect of noise on the biochemical characteristics of blood in daring cows. *Ziroc. Vyr.* 28 (4): 261 – 267.
- Daghash M. W. H. A. H. (2008) Performance of lambs after transportation and effect of fenugreek seeds supplementation on growth and carcass characteristics. Msc. Thesis, Fac. of Agri. Assiut Uni. Assiut, Egypt.
- Hall, S.J.G., D.M. Broom, and G.N.S. Kiddy. (1998). Effect of transportation on plasma cortisol and packed cell volume in different genotypes of sheep. *Anim.Sci.* 29:233-237.
- Horton, J. J., J. A. Baldwin, S. M. Emanuele, J. E. Wohit and L. R. Mc Douell. (1996). Performance of blood chemistry in lambs flowing fasting and transport. *Animal Science* 62: 49 – 56.
- Jones, S. D. M., A. L. Schaeyer, , A. K. W. Tong, and B. C. Vincent (1988). The effect of fasting and transportation of beef cattle. 2- Body components changes, carcass composition and meat quality. *livest. Prod Sci*: 20: 25-35.
- Kannan,G.,T.H.Terrill.,B.Kouakom.,O.S. Gazal.,S.Gelaya., E. A. Amoah, and S. Samake. (2000).Transportation of goats:Effects on physiological stress responses and live weight loss. *J. Anim. Sci.* 78: 1450 – 1457.
- Kannan,G.,T.H.Terrill.,B. Kouakou, , S. Gelaye, and E.A Amoah. (2002). Stimulated preslaughter holding and isolation effects on stress responses and live weight shrinkage in meat goats. *Journal Animal Science* 80:1771-1780.
- Minka N. S. and J. O. Ayo (2010).Physiological rsponses of food animals to road transportation stress. *African Journal of Biotechnology* Vol. 9(40), pp. 6601-6613,
- Minka N.S., and J.O.Ayo (2007c).Physiological responses of transported goats treated with ascorbic acid during the hot-dry season. *Anim.Sci.J.*78: 164-172.
- Minka N.S., and J.O. Ayo (2008). Assessment of the stress imposed on adult ostrich (*Struthio camelus*) during handling, loading, transportation and unloading. *Vet.Rec.* 162: 846-851.
- Minka S.N., and J.O. Ayo (2007a). Effect of loading behaviour and road transport stress on traumatic injuries in cattle transported by road during the hot-dry season. *Livestock Sci.* 107: 91-95.
- Minka, S.N., and J.O. Ayo (2007b). Road transportation effect on rectal temperature,

- respiration and heart rate of ostrich(*Struthocamelus*) chicks. *Vet. Arhiv.* 77(1):3946.
- Moe, R.O. and M. Bakken, (1997) Effects of handling and physical restraint on rectal temperature, cortisol, glucose and leucocyte counts in the silver fox (*Vulpes vulpes*). *Acta Vet. Scandinavica*, 38: 2939.
- Naqui, S. M. K. and A. K. Rai (1991) Effects of fasting on some physiological responses and blood constituents in native and crossbred sheep. *Indian J. of A. Science* 61:985.
- Plyaschenko S.I., and V.T. Sidorov (1987). Stresses in Farm Animals. *Agropromizdat*, Moscow (in Russian). (cited after Minka and Ayo, 2010. Physiological responses of food animals to road transportation stress. *African Journal of Biotechnology* Vol. 9(40), pp. 6601-6613.)
- Rollin, B.E. (1995). Farm animal welfare. Iowa State University Press, Ames Iowa, U.S.A.
- Schaefer, A. L., S. D. M. Jones, and R. W. Stanley (1997). The use of electrolyte solution for reducing transport stress. *J. Anim. Sci.* 75: 258 – 265.
- Schaefer, A. L., S. D. M., Jones, A. K. W. Tong, and B. A. young. (1990). Effects of transport and electrolyte supplementation on ion concentration, carcass yield and quality in bulls. *Can. J. Anim. Sci.* 70: 107 – 119.
- Sranmek, J. and J. pozdosek. (1987). The effect of sodium – chloride supplement on the contents of sodium and potassium in blood plasma and on sodium and potassium retention in heifers. *Zivoc. Vyr.* 32: 515 – 225.
- Vihan V.S., and K.L. Sahni (1981). Seasonal changes in body temperature, pulse and respiration rate in different genetic groups of sheep. *Indian Vet. J.* 58: 617-621.
- Von Borell, E. H. (2001). The biology of stress and its application to livestock housing and transportation assessment. *J. Anim. Sci.* 79 (Suppl.) 260-267.
- Warriss, P.D. (1990). The handling of cattle pre-slaughter and its effects on carcass and meat quality. *Appl. Anim. Behav. Sci.* 28:171.
- Whythens, J.R., G.N. Johnston, N. Beaman, and P. K. O'Rourke. (1985) Preslaughter handling of cattle. The availability of water during the lairage period. *Aust. Vet. J.* 62:163 – 165.
- Zulkifli I., B. Y. W. Norbaiyah, A.F. Cheah, A. Q. Soleimani, M. A. Sazili, rajion And Y. M. Goh, (2010). Physiological Responses in Goats Subjected to Road Transportation under the Hot, Humid Tropical Conditions *Int. J. Agric. Biol.*, Vol. 12, No. 6, 2010.

تأثير النقل البري على وزن الجسم ومعدلي النبض والتنفس ودرجة حرارة المستقيم في حالات التغذية حتى الشبع والصيام وتجريب المحاليل لعجول الجاموس

رعدة عادل تغيان ، فاروق محمد علام ، محمد نصرت محمود عبدالعاطي ، سيف اليزل فتحي عباس

قسم الانتاج الحيواني والدواجن كلية الزراعة جامعة اسيوط

أجريت هذه الدراسة في مزرعة بحوث قسم الإنتاج الحيواني والدواجن ، كلية الزراعة ، جامعة أسيوط.

استهدف هذا البحث دراسة تأثيرات النقل البري على بعض الصفات الفسيولوجية في عجول الجاموس. واستهدف البحث أيضا دراسة تجريب الجاموس بالمحاليل (الكتروليت) لتقليل أو منع التأثيرات الضارة لإجهاد النقل على الحيوان. تم استخدام خمسة عشر عجل جاموس من كلا الجنسين تم تقسيمها إلى ثلاث مجموعات حسب الوزن (صغير - متوسط - كبير) وتعرضت الثلاث مجموعات إلى المعاملات التالية:

معاملة المقارنة: وتم فيها التغذية حتى الشبع ثم النقل، معاملة الصيام: تم منع التغذية والماء ستة عشر ساعات ثم النقل، المعاملة بالمحاليل: وتم فيها التغذية حتى الشبع ثم تجريب كل حيوان 2 لتر محلول إلكتروليتي .

مُنعت كل الحيوانات من الغذاء والماء عند النقل وحتى أخذ آخر قراءة. وتم نقل الجاموس في لوري لمدة ثلاث ساعات لمسافة حوالي 250 كيلومتر بمتوسط سرعة 60-80 كم/ساعة (درجة الحرارة 28-34 درجة مئوية وبدأت الرحلة في الساعة صباحا) بعد ذلك تم تنزيل الحيوانات وأخذ القراءات والعينات.

في يوم النقل تم أخذ القراءات مباشرة قبل النقل (القراءة الأولى) ثم بعد الوصول مباشرة (القراءة الثانية) ثم بعد ساعتين (القراءة الثالثة) وأخيرا بعد أربع ساعات (القراءة الرابعة). وتمت دراسة النسبة المئوية لنقص وزن الجسم الحي، معدل التنفس، معدل النبض ودرجة حرارة المستقيم.

لم توجد فروق معنوية بين الثلاث مجاميع من الجاموس في معظم الصفات المدروسة. حيث أحدث النقل زيادة في النسبة المئوية لنقص وزن الجسم الحي ودرجة حرارة المستقيم و أدى النقل كذلك إلى حدوث زيادة يتبعها نقص في معدل التنفس ومعدل النبض. بعد النقل انخفض وزن الجسم الحي في مجموعة الجاموس التي غذيت حتى الشبع و المجموعة الصائمة والمجموعة التي تجرعت المحاليل. لكن المجموعة المغذاة حتى الشبع أظهرت أكبر نسبة انخفاض في وزن الجسم الحي عن المجموعة الصائمة. بينما مجموعة الجاموس التي تجرعت المحاليل قبل النقل كانت أقل تأثرا بإجهاد النقل مما يثبت أن هذه المعاملة أدت إلى تقليل الآثار الضارة للنقل. أدى النقل إلى ارتفاع درجة حرارة المستقيم وزيادة معدلي التنفس والنبض في الحيوانات المعاملة. لكن الجاموس الذي تجرع المحاليل قبل النقل كان أقل تأثرا بإجهاد النقل. وقد يرجع ارتفاع درجة حرارته إلى زيادة استجابة قشرة الأدرينال أثناء تداول وتحميل ونقل الحيوانات. لذلك فإن تجريب المحلول الألكتروليتي للحيوانات كان فعالا في منع أو تقليل الآثار الضارة للنقل البري على الحيوانات.