

EFFECT OF SOYBEAN LECITHIN ON FREEZABILITY AND FERTILIZING POTENTIALS OF BOVINE SPERMATOZOA

M.R. BADR; MARY G. ABD EL-MALAK; K.M. MOHAMMED and EBTIHAL A. IBRAHIM

Artificial Insemination and Embryo Transfer Department, Animal Reproduction Research Institute, Haram, (P.O.B. 12556), Giza, Egypt. e-mail: magdybadr69@yahoo.com

ABSTRACT

Received at: 29/3/2012

Accepted: 22/4/2012

Soybean lecithin has been attracted increasing attention and has been used to replace egg yolk in the semen extender. In the present study, effect of soybean lecithin on bovine semen freezability and in vitro fertilizing potentials were evaluated. Semen was cryopreserved in tris-based extender supplemented with different soybean lecithin concentrations (3, 5, 7, 10 and 20%) vs 20% egg yolk (control). Semen post-thawing motility, viability and acrosomal integrity, DNA damage, enzymes leakage, total antioxidant activity (TAC), lipid peroxidation and in vitro fertilizing potentials were assessed. Current results indicated that addition of 7% soybean lecithin to semen extender significantly ($P<0.05$) improved post-thawing motility, viability and acrosomal integrity ($61.25\pm1.25\%$, 172.25 ± 5.53 and $10.25\pm2.39\%$, respectively) compared with control ($47.50\pm4.78\%$, 106.25 ± 16.88 and $22.00\pm1.47\%$, respectively). At the same time, a significantly reduced ($P<0.05$) sperm DNA damage, tail length and tail moment of the cryopreserved semen ($1.48\pm0.27\%$, $2.08\pm0.36\ \mu\text{m}$ and 3.14 ± 1.32 , respectively) compared with control ($3.31\pm0.17\%$, $3.93\pm0.24\ \mu\text{m}$ and 13.09 ± 1.38 , respectively). Moreover, extender containing 7% soybean lecithin significantly ($P<0.05$) increased TAC ($0.47\pm0.04\ \text{m}\mu\text{mol/ml}$) and decreased lipid peroxidation (Malondialdehyde) of the cryopreserved spermatozoa ($9.18\pm1.47\text{nmol/ml}$) with respect to the control ($0.19\pm0.02\ \text{m}\mu\text{mol/ml}$, and $21.57\pm1.45\ \text{nmol/ml}$, respectively). Additionally, 7% soybean lecithin significantly ($P<0.05$) improved in vitro fertilization rate, cleavage rate, morula and blastocyst development (54.14 ± 6.21 , 52.81 ± 3.32 , 27.49 ± 2.78 and $18.44\pm2.11\%$, respectively) compared with the control (35.10 ± 3.23 , 31.75 ± 5.52 , 12.2 ± 4.08 and $4.82\pm2.12\%$, respectively). It was concluded that the addition of 7% soybean lecithin to the freezing extender improved freezability and enhanced in vitro fertilizing potentials of bovine spermatozoa through protection of DNA from deterioration and reduction of oxidative stress.

تأثير ليسيسين الصويا علي قابلية حيامن الأبقار للتجميد وقدرتها الإخصابية معمليا

مجدي رمضان بدر ، ماري جاد عبد الملاك ، كامل مصطفى محمد ، ابتهاج عبد الله ابراهيم

اضافة ليسيتين الصويا الى ممدات السائل المنوي كبديل عن صفار البيض لازال يثير الاهتمام ويتزايد يوما بعد يوم. وحديثا ظهرت أنواع جديدة من ممدات السائل المنوي خالية من الإضافات الحيوانية (صفار البيض أو اللبن) وبالرغم

من ذلك مازالت الممددات التي تحتوى على صفار البيض تستخدم بصورة أساسية لتجميد السائل المنوي البقري. وتهدف الدراسة الحالية إلى دراسة قدرة حيامن الأبقار للتجميد وكذلك قدرتها الإخصابية معمليا وحقليا عند تجميدها في ممدد مضاف إليه الصويا ليسيسين بتركيزات مختلفة (٣، ٥، ٧، ١٠، ٢٠ %) مقارنة باستخدام ٢٠% صفار البيض. ولقد أوضحت نتائج الدراسة الحالية أن تجميد السائل المنوي البقري في ممدد مضاف إليه صويا ليسيسين بتركيز ٧% نتج عنه زيادة معنوية كبيرة في نسبة الحركة الأمامية بعد الإسالة، معدل الحيوية ونسبة المحافظة على غشاء القلتوسية (٦١,٢٥%، ١٧٢,٢٥ و ١٠,٢٥% على التوالي) مقارنة بتلك التي تم تمديدتها في ممدد التريس الذي يحتوى على ٢٠% صفار البيض (٤٧,٥٠%، ١٠٦,٢٥ و ٢٢,٠٠% على التوالي). كما حافظ على سلامة الحامض النووي للحيامن حيث قلل من تشظي الحامض النووي وطول ذيل المذنب وكثافته (١,٤٨%، ٢,٠٨ ميكرومول و ٣,١٤ على التوالي) بالمقارنة بالمجموعة الضابطة (٣,٣١%، ٣,٩٣ ميكرومول و ١٣,٠٩ على التوالي). كذلك نتج عنه زيادة معنوية كبيرة في مستوى مضادات الأكسدة الكلية وانخفاض معنوي كبير في معدل أكسدة الدهون (٠,٤٧ ملليمول/ملي و ٩,١٨ نانومول/ملي على التوالي) مقارنة بالمجموعة الضابطة (٠,١٩ ملليمول/ملي و ٢١,٥٧ نانومول/ملي على التوالي). كما أوضحت نتائج الإخصاب المعملية أن تجميد السائل المنوي البقري في ممدد مضاف إليه صويا ليسيسين بتركيز ٧% نتج عنه زيادة معنوية كبيرة في نسبة إخصاب البويضات وكذلك قدرتها على النمو إلى الطور التوتري و طور البلاستوسيست (٥٤,١٤، ٥٢,٨١، ٢٧,٤٩ و ١٨,٤٤% على التوالي) مقارنة بتلك التي تمديدتها في ممدد التريس الذي يحتوى على ٢٠% صفار البيض (٣٥,١٠، ٣١,٧٥، ١٢,٢٠ و ٤,٨٢% على التوالي). ولهذا يمكن أن نستنتج من نتائج هذه الدراسة أن إضافة الصويا ليسيسين بتركيز ٧% إلى ممدد السائل المنوي يمكن أن تكون الاختيار الأمثل لتجميد السائل المنوي البقري في المستقبل.

Key words: Soybean lecithin, semen cryopreservation, DNA integrity, IVF, antioxidant.

INTRODUCTION

Semen cryopreservation has profound effects on spermatozoa, many of which result in sublethal damage to the cells, and subsequent reduction of fertilizing ability. The sperm plasma membrane serves as the main physical barrier to the outside environment and is a primary site of freeze-thaw damage. Such damage includes membrane destabilization due to lateral lipid rearrangement (De Leeuw *et al.*, 1990), loss of lipids from the membrane (Buhr *et al.*, 1994 and Golal *et al.*, 1998), and peroxidation of membrane lipids as a result of formation of reactive oxygen species (ROS) (Aitken 1995; Flesch and Gadella, 2000; Badr *et al.*, 2010). These events can affect sperm motility, response to osmotic stress, and signaling pathways; therefore, the fertilizing ability is compromised (Holt, 2000). Defining causes of damage to sperm during cryopreservation is further complicated because the processing of semen for cryopreservation is not standardized and there is a wide variety of freezing diluents in use. Semen extender contain some forms of lipids, the most common being egg yolk lipids (Watson, 1995). Egg yolk, a common protectant in cryopreservation media, has

been used for providing protection against cold shock in the cryopreservation of mammalian semen for over half a century (Gousset *et al.*, 2004). However, there have been frequent arguments against the use of animal-originated ingredients, egg yolk, milk or even low density lipoprotein (LDL) extracted from egg yolk, one of which is the wide variability of composition that make it difficult to analyze the beneficial effects of a particular compound on sperm cryopreservation. Furthermore, they could introduce possible sanitary risks (viruses, bacteria and fungi), with the subsequent production of endotoxins capable of damaging the fertilizing capacity of spermatozoa (Bousseau *et al.*, 1998; Jiang *et al.*, 2007).

As one of phospholipids, lecithin (or phosphatidylcholine) is distributed widely in plants and it plays an important role in the regulation of the physiological function of animal cells bio-membrane (Thun *et al.*, 2002). Soy bean lecithin has similar ingredients to egg yolk used for protection of animal spermatozoa from cold shock in semen cryopreservation (Aires *et al.*, 2003). It has been suggested that soybean lecithin may play a better protective role for

spermatozoa than egg yolk during the cryopreservation process and therefore reduce the risk of introducing bacterial and mycoplasma into freezing extenders (Fukui *et al.*, 2008). The main goal for the present study was to determine the effect of soybean lecithin as cryoprotective on bull spermatozoa, through investigating the quality parameters of the cryopreserved spermatozoa, DNA integrity, antioxidant activities and in vitro fertilizing potentials of frozen-thawed bull semen.

MATERIALS and METHODS

Semen collection and processing:

Semen samples were collected from six fertile bovine bulls. Only semen samples at least 70% initial motility and 800×10^6 sperm cells/ml were used. After collection, semen samples were pooled, split into 6 portions and diluted at 30°C with Tris-based extender supplemented with different concentrations of soybean lecithin (3, 5, 7, 10 and 20%) vs 20% egg yolk (control). The extended semen was cooled to 5°C throughout 60 minute in a cold cabinet. The cooled semen was loaded into 0.25 ml French straws (IMV, L'Aigle, France), then suspended into liquid nitrogen vapor inside foam box before immersed into liquid nitrogen. Frozen semen straws were thawed in a water bath at 37°C for 30 second. Post-thawing sperm motility, viability and acrosomal integrity were assessed according to Mohammed *et al.* (1998).

Assessment of sperm DNA integrity:

DNA integrity and the incidence of DNA strand breaks or fragmentation was detected using alkaline comet assay according to Boe-Hansen (2005). Briefly, DNA status of individual cells was determined by the neutral single cell gel electrophoresis (comet) assay. For this assay, frozen-thawed spermatozoa were diluted in phosphate buffer saline (PPS), embedded in agarose, followed by cell lysis, DNA decondensation, electrophoresis and DNA staining with 50 µl of 20 µg/ml ethidium bromide (Sigma). The cells were then visualized by fluorescent microscopy. Intact nuclei in the comet assay

appeared to have compact and brightly fluorescent heads; in contrast, strand breaks in damaged cells allow DNA migration during electrophoresis, and a tail of DNA could be seen behind the head, giving the appearance of a comet (Hughes *et al.*, 1996). After subjecting spermatozoa to the comet assay, sperm nuclei were analyzed by computer software program.

Biochemical analysis:

Extra-cellular aspartate-aminotransferase (AST); alanine-aminotransferase (ALT) and alkaline phosphatase (ALP) enzymes leakage during cryopreservation was assessed spectrophotometrically according Tietz (1976) to evaluate the membrane stability of spermatozoa. Additionally, total antioxidant capacity and membrane lipid peroxidation was estimated by the end point generation of malondialdehyde (MDA) determined by the thiobarbituric acid (TBA) test, of the cryopreserved spermatozoa were measured as described by Cortassa *et al.* (2004).

Evaluation of in vitro fertilizing potential of the treated semen:

The fertilizing potentials of the treated semen were assessed using *in vitro* fertilization technology, as demonstrated by Totey *et al.* (1992). Three straws from each treatment were thawed in a water bath at 37°C for 30 sec. The most motile spermatozoa were separated by swim up technique in the fertilization medium, modified Tyrode's Albumin-Lactate-Pyruvate (TALP) containing 6 mg/ml bovine serum albumin (BSA), for 1 hour as recorded by Parrish *et al.* (1988). The uppermost layer of the medium containing the most motile spermatozoa was collected and washed twice by centrifugation at 2000 rpm for 10 minutes. The sperm pellet was resuspended in the fertilization TALP medium containing 10 µg/ml heparin. After appropriate dilution, 2 µl (final concentration 2×10^6 sperm cell/ml) of sperm suspension was added to the fertilization drops, containing *in vitro* matured oocytes. Gametes were co-incubated in the fertilization drops under sterile mineral oil for 18 hour at 39°C in an

atmosphere of 5% CO₂ in air with maximum humidity. The inseminated oocytes were freed from extra cumulus cells and attached spermatozoa by gentle pipetting and then cultured in TCM-199 medium for seven days in the same previous conditions. The proportional of cleaved oocytes was recorded 48 hour after insemination and those developed to the morula and blastocyst stages were recorded at 5-7 day post-insemination.

Statistical analysis:

All data were analyzed by using Costat Computer Program (1986) Cottort Software, and were compared by the least significant difference least (LSD) at 5% levels of probability. The results were expressed as means $\pm$ SE.

RESULTS

The results presented in Table 1 revealed that, addition of soybean lecithin to the freezing extender improved the freezability of bull spermatozoa compared with the control semen in a dose-dependent trend. Addition of 7% soybean lecithin to semen extender, appeared to be the best concentration that increased ($P < 0.05$) significantly the post-thawing sperm motility; viability index and maintained acrosomal integrity ($61.25 \pm 1.25\%$, 172.25 ± 5.53 and $10.25 \pm 2.39\%$, respectively) compared to the control semen ($47.50 \pm 4.78\%$, 106.25 ± 16.88 and $22.00 \pm 1.47\%$, respectively).

Data regarding the effect of soybean lecithin addition to the freezing extender on the total antioxidant capacity (TAC) and lipid peroxidation of the cryopreserved semen which indicated by malondialdehyde ((MDA) are presented in table 2. In vitro provision of semen extender with 7% soybean lecithin significantly ($P < 0.05$) increased the total antioxidant and diminished lipid peroxidation of the frozen-thawed semen (0.47 ± 0.04 m μ /ml, and

9.18 ± 1.47 nmol/ml, respectively) compared with the control extender (0.19 ± 0.02 m μ /ml, and 21.57 ± 1.45 nmol/ml, respectively). Moreover, data presented in table 2 clarified that, addition of 7% soybean lecithin to the semen extender maintained sperm cell membrane integrity and this appeared through reduction of extracellular enzymes (AST, ALT and ALP) leakage (70.00 ± 4.56 , 11.5 ± 1.04 and 14.65 ± 2.79 U/L, respectively) compared with the control extender (106.00 ± 5.95 , 23.00 ± 2.48 and 23.27 ± 3.75 U/L, respectively)

With respect to the effect of soybean lecithin addition to the freezing extender on the DNA integrity of the frozen-thawed bovine spermatozoa are demonstrated in Table 3. The present data indicated that, in vitro provision of semen extender with 7% soybean lecithin significantly ($P < 0.05$) decreased the DNA fragmentation, tail length and tail moment of the frozen-thawed semen ($1.48 \pm 0.27\%$, 2.08 ± 0.36 μ m and 3.14 ± 1.32 , respectively) as compared with the control extender ($3.31 \pm 0.17\%$, 3.93 ± 0.24 μ m and 13.09 ± 1.38 , respectively).

Data concerning the effect of replenishing of semen extender with soybean lecithin on the in vitro fertilizing potentials and embryo development are presented in tables 4 and 5. The current results revealed that, addition of 7% soybean lecithin to the freezing extender had a positive effect ($P < 0.05$) on the in vitro fertilization rate and embryo developmental rate compared with the control semen. When 7% soybean lecithin was added to the freezing extender, a higher proportion of in vitro fertilized oocytes, cleavage rate, morula and blastocyst development (54.14 ± 6.21 , 52.81 ± 3.32 , 27.49 ± 2.78 and $18.44 \pm 2.11\%$, respectively) compared with the control semen (35.10 ± 3.23 , 31.75 ± 5.52 , 12.2 ± 4.08 and $4.82 \pm 2.12\%$, respectively).

Table 1: Effect of different soybean lecithin concentrations on bovine spermatozoa freezability.

Treatment	Pre-freeze motility (%)	Post-thaw motility (%)	Viability index	Acrosomal integrity (%)
Control	78.75±3.12 ^a	47.50±4.78 ^b	106.25±16.88 ^b	22.00±1.47 ^a
3% soy lecithin	78.75±1.25 ^a	53.75±2.39 ^{ab}	131.88±10.38 ^b	15.5±1.33 ^{ab}
5% soy lecithin	77.5±3.23 ^a	56.25±2.40 ^{ab}	140.5±12.04 ^{ab}	15.5±2.55 ^{ab}
7% soy lecithin	82.25±1.44 ^a	61.25±1.25 ^a	172.25±5.53 ^a	10.25±2.39 ^b
10% soy lecithin	81.25±2.39 ^a	51.25±5.54 ^{ab}	116.87±9.92 ^b	15.25±1.92 ^{ab}
20% soy lecithin	80.00±2.04 ^a	48.75±3.75 ^b	113.00±7.72 ^b	20.25±2.93 ^a

Means in the same column with different superscripts are significantly differ at P<0.0

Table 2: Effect of different soybean lecithin concentrations on biochemical activity of bovine spermatozoa.

Treatment	AST (U/L)	ALT (U/L)	ALP (U/L)	TAC (mμ/ml)	MDA (nmol/ml)
Control	106.00±5.95 ^a	23.00±2.48 ^a	23.27±3.75 ^a	0.19±0.02 ^c	21.57±1.45 ^a
3%soy lecithin	90.75±7.48 ^{ab}	17.75±1.25 ^{ab}	15.48±2.08 ^b	0.31±0.08 ^b	16.1±1.64 ^{ab}
5%soy lecithin	74.75±8.78 ^{bc}	15.25±2.75 ^{bc}	16.43±1.10 ^b	0.42±0.04 ^a	14.39± 2.6 ^{bc}
7%soy lecithin	70.00±4.56 ^c	11.5±1.04 ^c	14.65±2.79 ^b	0.47±0.04 ^a	9.18±1.47 ^c
10%soy lecithin	100.25±0.55 ^a	22.25±1.49 ^a	18.55±1.34 ^{ab}	0.29±0.02 ^b	16.79±2.59 ^{ab}
20%soy lecithin	91.25±2.17 ^{ab}	21.00±1.47 ^a	19.9±1.38 ^{ab}	0.27±0.03 ^b	17.94±1.69 ^{ab}

Means in the same column with different superscripts are significantly differ at P<0.05

AST: Aspartate-aminotransferase

ALT: Alanine-aminotransferase

ALP: Alkaline phosphatase

TAC: Total antioxidant capacity

MDA: Malondialdehyde

Table 3: Effect of different soybean lecithin concentrations on DNA integrity of bovine spermatozoa.

Treatment	DNA integrity (%)	Tail length (μm)	Tail moment
Control	3.31±0.17 ^a	3.93±0.24 ^a	13.09±1.38 ^a
3% soy lecithin	2.06±0.11 ^{bc}	2.28±0.45 ^{ab}	4.80±1.22 ^c
5% soy lecithin	2.17±0.25 ^{bc}	2.38±0.20 ^{ab}	5.31± 1.47 ^{bc}
7% soy lecithin	1.48±0.27 ^c	2.08±0.36 ^b	3.14±1.32 ^c
10% soy lecithin	2.34±0.16 ^b	2.63±0.15 ^{ab}	7.11±2.41 ^{bc}
20% soy lecithin	3.06±0.32 ^a	3.22±0.20 ^{ab}	10.05±1.72 ^{ab}

Means in the same column with different superscripts are significantly differ at P<0.05

Table 4: Effect of different soybean lecithin concentrations on the in vitro fertilizing potentials of bovine spermatozoa.


Treatment	No. of oocytes	No. of Penetrated oocytes	Penetration rate (%)	No. of fertilized oocytes	Fertilization rate (%)
Control	87	54	63.07±8.34 ^a	31	35.10±3.23 ^b
3% soy lecithin	75	49	65.03±1.72 ^a	30	41.47±5.42 ^{ab}
5% soy lecithin	63	40	63.62±5.14 ^a	28	43.84±5.63 ^{ab}
7% soy lecithin	84	62	73.55±3.66 ^a	56	54.14±6.21 ^a
10% soy lecithin	71	46	63.41±4.13 ^a	33	46.03±3.44 ^{ab}
20% soy lecithin	63	43	68.19±6.27 ^a	22	34.71±2.28 ^b

Means in the same column with different superscripts are significantly differ at $P < 0.05$

Table 5: Effect of different soybean lecithin concentrations on the bovine embryo development in vitro.

Treatment	No. of oocytes	Cleavage rate No. (%)	Morula stage No. (%)	Blastocyst stage No. (%)
Control	83	26 (31.75±5.52) ^b	10 (12.2±4.08) ^b	4 (4.82±2.12) ^c
3% soy lecithin	94	39 (41.67±1.28) ^{ab}	16 (16.59±3.16) ^{ab}	10 (9.42±1.09) ^{bc}
5% soy lecithin	70	32 (45.32±5.39) ^{ab}	13 (18.85±2.99) ^{ab}	9 (12.83±1.79) ^{ab}
7% soy lecithin	89	47 (52.81±3.32) ^a	24 (27.49±2.78) ^a	16 (18.44±2.11) ^a
10% soy lecithin	88	36 (40.97±1.52) ^b	15 (16.35±3.92) ^{ab}	7 (7.28±3.03) ^{bc}
20% soy lecithin	83	29 (34.87±8.93) ^b	7 (8.3±2.55) ^b	4 (3.53±1.73) ^c

Means in the same column with different superscripts are significantly differ at $P < 0.05$

**Fig 1:** Bovine spermatozoa cryopreserved in tris-7% soybean lecithin. The single cell gel electrophoresis (comet) assay showed reduction in the DNA fragmentation as represented by a limited amount of DNA present in the comet tail.




Fig 2: Bovine spermatozoa cryopreserved in tris-egg yolk extender. The single cell gel electrophoresis (comet) assay showed increased in the DNA fragmentation as represented by an increasing amount of DNA present in the comet tail.

DISCUSSION

In the present study, freezing extender supplemented with 7% soybean lecithin could provide better cryoprotective action for bovine spermatozoa during cryopreservation compared with control group (20% egg yolk). These results are in accordance with the previous reports in the cryopreservation of bovine (Moussa *et al.*, 2002 and Amirat *et al.*, 2005), buffalo (Badr 2008) and sheep (Fukui *et al.*, 2008) semen. Contrary, other researchers reported that the cryoprotective effect of soybean lecithin extenders on bovine sperm freezability was similar or slightly inferior to that of 20% egg yolk extender (Aires *et al.*, 2003). The precise mechanism by which soybean lecithin protects spermatozoa during cryopreservation remains unclear. It was generally accepted that cold shock and cryodamage might impair the physiological function of spermatozoa membrane due to the change of the lipid composition of its bilayer and the fluidity of the plasma membrane during freeze-thawing process as a result of formation ROS (Johnson *et al.*, 2000). Excessive generation of ROS negatively affect the fluidity of sperm plasma membrane and integrity of DNA in the sperm nucleus (Cocuzza *et al.*, 2007).

The beneficial effect of soybean lecithin extender on the sperm function may be attributed to the ability of the lecithin to maintain the integrity of cell membranes, facilitating the movement of fluids inside and outside the cell; and without lecithin cell membranes would harden and would no longer stay semi-permeable (Zeisel, 2000). It is believed that phospholipids from egg yolk or soybean lecithin might integrate with sperm membrane to form a protective film against the formation of lethal intracellular ice crystal and protect the sperm membrane from mechanical damage during freeze-thawing process (Quinn *et al.*, 1985). These beneficial effect of soybean lecithin dose dependent on plasma membrane stability leads to reduction of extracellular enzymes leakages (AST, ALT and ALP) which appeared clearly in our findings mainly at concentration of 7%. Additionally, soybean lecithin could play a protective role for sperm during cryopreservation due to its low viscosity and less debris which enhance the sperm motion characteristics and fertilizing ability, compared with egg yolk which provided higher viscosity and the presence of particulate debris in extenders (Van Wagtendonk-de Leeuw *et al.*, 2000).

In a way, soybean lecithin is also thought as a better emulsifier that might promote cryoprotectants to distribute uniformly and reduce its local concentration, which led to relieve the toxicity of cryoprotectants during the freeze-thawing process (Trotta *et al.*, 2002). Moreover, soybean lecithin might reduce the cholesterol/phospholipids ratio of sperm cell membranes by permeating into the sperm membrane, so capacitation like changes during the freezing process were restrained to improve the fertilizing ability of frozen-thawed spermatozoa (Galantino-Homer *et al.*, 2006). The present results evidently revealed that soybean lecithin decreased significantly membrane lipid peroxidation, throughout reduction of malondialdehyde (MDA) production and increased total antioxidant capacity (TAC) in the frozen-thawed spermatozoa that maintain a suitable level of ROS which play a significant role in many physiological processes of the sperm such as capacitation, hyper-activation and sperm-oocyte fusion (Sies, 1993; Lewis *et al.*, 1995). However, ROS must be continuously inactivated to keep only a small amount necessary to maintain normal cell function, as a result the excessive generation of ROS in semen can cause damage to sperm that increased by extruding of cytoplasm during the maturation process which is the major source of antioxidants. Therefore, the improved freezability and in vitro fertilizing potentials of the semen that extended in soybean lecithin extender may be attributed to the ability of the lecithin to protect the spermatozoa from the destructive effects of oxidative stress during cryopreservation (Chatterjee and Gagnon, 2001). Additionally, soybean lecithin may act as a stabilizer and protectant of proteins and cell membranes, whose fluidity decreases during temperature downshift. This may emphasize the current results which indicated that soybean lecithin provision to the freezing extender diminished the enzymes leakage.

Furthermore, the beneficial effect of soybean lecithin extender on semen cryopreservation may be attributed to the high concentration of linoleic acid in its

constituent, which is the precursor of prostaglandin E. Addition of prostaglandin E to the semen extender increased the life span of the spermatozoa and the cleavage rate of the inseminated oocytes (Kolev and Dimov 1998). Accordingly, in our results, post-thaw motility and viability index appeared significantly higher ($P < 0.05$) in soybean additive diluent particularly at 7% than 20% egg yolk addition.

In conclusion, the present study provides novel evidence that addition of 7% soybean lecithin to the freezing extender improved freezability and enhanced in vitro fertilizing potentials of bovine spermatozoa through protection of DNA from deterioration, reduction of the oxidative stress. We suggest that consistent with quality standards that should be required for cryoprotectant extender, soy lecithin-based extender in recommended dose (7%) is a viable alternative to conventional egg-yolk-based freezing diluents for cryopreserving of bovine bull spermatozoa.

Acknowledgement: The authors greatly acknowledge to colleague prof. Dr. Hani Hassan for his valuable and privileged role in the comet assay.

REFERENCES

- Aires, V.A.; Hinsch, K.D.; Mueller-Schloesser, F.; Bogner, K.; Mueller-Schloesser, S. and Hinsch, E. (2003): In vitro and in vivo comparison of egg yolk-based and soybean lecithin-based extenders for cryopreservation of bovine semen. *Theriogenology*, 60: 269-279.
- Aitken, R.J. (1995): Free radicals, lipid peroxidation and sperm function. *Reprod. Fertil. Dev.*, 7: 659-668.
- Amirat, L.; Anton, M.; Tainturier, D.; Chatagnon, G.; Battut, I. and Courtens, J.L. (2005): Modification of bull spermatozoa induced by three extenders: Biociphos, Low density lipoprotein and Triladyl, before, during, and after freezing and thawing. *Reproduction*, 129: 535-543.

- Badr, M.R. (2008):* Cryopreservation of buffalo spermatozoa in soy lecithin-based extenders. *Assiut Vet. Med. J.*, 54 (116): 272- 284 .
- Badr, M.R.; Mary G. Abd El- Malak and Hassan, H.M. (2010):* Effect of trehalose on cryopreservation, oxidative stress and DNA integrity of buffalo spermatozoa. *J. Reproduction and infertility*, 1 (2): 50-57.
- Boe-Hansen, G.B. (2005):* Hydrogen peroxide alters the physical state and function of the plasma membrane of pulmonary artery endothelial cells. *Journal of Cellular Physiology*, 20: 362-374.
- Bousseau, S.; Brillard, J.P.; Guienne, B.M.L.; Guérin, B.; Camus, A. and Lechat, M. (1998):* Comparison of bacteriological qualities of various egg yolk sources and the in vitro and in vivo fertilizing potential of bovine semen frozen in egg yolk or lecithin based diluents. *Theriogenology*, 50: 699-706.
- Buhr, M.M.; Curtis, E.F. and Kakuda, N.S. (1994):* Composition and behavior of head membrane lipids of fresh and cryopreserved bull sperm. *Cryobiology*, 31: 224-238
- Chatterjee, S. and Gagnon, C. (2001):* Production of reactive oxygen species by spermatozoa undergoing cooling, freezing and thawing. *Mol. Reprod. Dev.*; 59: 451-458.
- Cocuzza, M.; Sikka, S.C.; Athayde, K.S. and Agarwal, A. (2007):* Clinical relevance of oxidative stress and sperm chromatin damage in male infertility: An evidence based analysis. *Int. Braz J. Urol.* Vol. 33 (5): 603-621.
- Cortossa, S.; Aon, M.A.; Waston, R.L. and O'Rourke, B. (2004):* A mitochondrial oscillator dependent on reactive oxygen species. *Biophysic. J.*, 87: 2060-2073.
- Costat Computer Program Copyright (1986):* Version 3.03 copyright Cottort Software.
- De Leeuw, F.E.; Chen, H.C.; Colenbrander, B. and Verkleij, A.J. (1990):* Cold-induced ultrastructural changes in bull and bull sperm plasma membranes. *Cryobiology*, 27: 171-183
- Flesch, F.M. and Gadella, B.M. (2000):* Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochim Biophys Acta* , 1469: 197-235.
- Fukui, Y.; Kohno, H.; Togari, T.; Hiwasa, M. and Okabe, K. (2008):* Fertility after artificial insemination using a soybean-based semen extender in sheep. *J. Reprod. Dev.* 54: 286-289.
- Galantino-Homer, H.L.; Zeng, W.X.; Megee, S.D.; Dallmever, M. and Voelk, D. (2006):* Effect of 2-hydroxypropyl-beta-cyclodextrin and cholesterol on porcine sperm viability and capacitation status following cold shock or incubation. *Mol. Reprod. Dev.*; 73 (5): 638-650.
- Golal, R.L.; Georgie, R.K.; Tuli, R.K.; Dixit, V.P. and Chand, D. (1998):* Lipid peroxidation during freeze processing of washed buffalo (*Bubalus bubalis*) spermatozoa. *Annals of Biology*, 14 (2): 207-210.
- Gousset, K.; Tsvetkova, N.M.; Crowe, J.H. and Tablin, F. (2004):* Important role of raft aggregation in the signaling events of cold-induced platelet activation. *Biochim Biophys Acta* 1660: 7-15.
- Holt, W.V. (2000):* Basic aspects of frozen storage of semen. *Anim Reprod Sci.*, 62: 3-22.
- Hughes, C.M.; Lewis, S.E.; McKelvey-Martin, V.J. and Thompson, W. (1996):* A comparison of baseline and induced DNA damage in human spermatozoa from fertile and infertile men, using a modified comet assay. *Mol. Hum. Reprod.*, 2: 613- 619.
- Jiang, Z.L.; Li, Q.W.; Hu, J.H.; Li, W.Y.; Zhao, H.W. and Zhang, S.S. (2007):* Improvement of the quality of boar cryopreservation semen by supplementing with low density lipoprotein in diluents. *Cryobiology*, 54: 301-304.
- Johnson, L.A.; Weitze, K.F.; Fiser, P. and Maxwell, W.M. (2000):* Storage of boar

- semen. Anim. Reprod. Sci. 62:143-172.
- Kolev, S.I. and Dimov, K. (1998):* Influence of synthetic PGF₂ alpha and oxytocin on life time of ram's spermatozoa with and without sperm plasma at 46°C. Bulgarian J. Agr. Sci., 4: 6, 831-835.
- Lewis, S.E.; Boyle, P.M.; McKinney, K.A.; Young, I.S. and Thompson, W. (1995):* Total antioxidant capacity of seminal plasma is different in fertile and infertile men. Fertil Steril. Oct; 64(4): 868-70.
- Mohammed, K.M.; Ziada, M.S. and Darwish, G.M. (1998):* Practical trials for freezing semen of buffalo and Friesian bulls: Effect of various regimens of freezing, different milk extenders and types of straws packages on post-thawing semen characters. Assiut Vet. Med. J.; 39 (77): 70-93.
- Moussa, M.; Martinet, V.; Trimeche, A.; Tainturier, D. and Anton, M. (2002):* Low density lipoproteins extracted from hen egg yolk by an easy method: cryoprotective effect on frozen-thawed bull semen. Theriogenology, 57: 1695-1706.
- Parrish, J.J.; Susko-Parrish, J.; Winer, M.A. and First, N.L. (1988):* Capacitation of bovine sperm by heparin. Biol. Reprod.; 38: 1171-1180.
- Quinn, P.J. (1985):* A lipid-phase separation model of low-temperature damage to biological membranes. Cryobiology, 22: 128-146.
- Sies, H. (1993):* Strategies of antioxidant defense. Eur J Biochem; 215: 213-.
- Thun, R.; Hurtado, M. and Janet, F. (2002):* Comparison of Biociphos-Plus® and TRIS-egg yolk extender for cryopreservation of bull semen. Theriogenology, 57: 1087-1094.
- Tietz, N.W. (1976):* Fundamentals of clinical chemistry. W.B. Saunders Company, Philadelphia.
- Totey, S.M.; Singh, G.; Taneja, M.; Pawshe, C.H. and Talwar, G.P. (1992):* In vitro maturation, fertilization and development of follicular oocytes from buffalo (*Bubalus bubalis*). J. Reprod. Fertil.; 95: 597-607.
- Trotta, M.; Pattarino, F. and Ignoni, T. (2002):* Stability of drug-carrier emulsions containing phosphatidylcholine mixtures. Eur. J. Pharm. Biopharm., 53: 203-208.
- Van Wagendonk-de Leeuw, A.M.; Haring, R.M.; Kaal-Lansbergen, L.M.T.E. and Den Daas, J.H.G. (2000):* Fertility results using bovine semen cryopreserved with extenders based on egg yolk and soy bean extract. Theriogenology, 54: 57-67.
- Watson, P.F. (1995):* Recent developments and concepts in the cryopreservation of spermatozoa and the assessment of their post-thawing function. Reprod Fertil Dev., 7: 871-891.
- Zeisel, S.H. (2000):* Cholin: An essential for humans. Nutrition, 16: 669.