

## EVALUATION OF THE HYGIENIC QUALITY OF RAW MILK BASED ON THE PRESENCE OF *BIFIDOBACTERIA* SPP. AS AN INDICATOR OF FAECAL CONTAMINATION IN ASSIUT CITY

MARWA M.N. EL-GENDI and LAMIAA M.T. ALI  
Animal Health Research Institute, Assiut Lab.

---

### ABSTRACT

---

Received at: 25/6/2012

Accepted: 26/8/2012

This study aimed to evaluate the hygienic quality of raw buffalo's and cow's milk sold in Assiut city, Egypt; based on the presence of *Bifidobacterium* spp. as indicator of fecal contamination. *Bifidobacteria* species forms one of the most important groups of flora in the intestine of both human and animals. Species isolated in humans are different from those isolated in animals. It should therefore be possible to determine contamination origin (human or animal). Seventy samples of raw buffalo's and cow's milk (35 samples from each) were collected randomly from; some dairy farms, individual cases of dairy buffaloes and cows and dairy shops in Assiut city, all samples were examined for the presence of *Bifidobacteria* spp. Each sample was firstly inoculated in MRS & BHIMup, then two different culture media (CMup & *Bifidobacterium* media (BFM)) were used for isolation and identification of *Bifidobacterium*. *Bifidobacterium* spp. was identified in 65.71 % (23 samples) and 51.42% (18 samples) of raw buffalo's and cow's milk, respectively. Isolates were identified and differentiated and the findings revealed that raw cow's milk harboured *B. dentium* in 61.11% , *B. suis* in 27.77% and *B. bifidum* in only 2 samples (11, 11 %) of examined samples. *B. dentium*, *B. suis* and *B. bifidum* were found contaminating 47.82, 39.13 and 13.04 % of examined raw buffalo's milk samples. Presence of *B. bifidum* revealed contamination of raw cow's and buffalo's milk with human stool (adult and infants); however isolates of *B. suis* indicated the contamination with feces of piglets. Also the detection of *B. dentium* in examined samples showed that there is contamination with human discharges coming from mouth suffering dental caries; human abscesses, vaginal discharges and feces. These organisms considered potentially pathogenic and have hazards effect on human health.

---

**Key words:** *Bifidobacteria*, raw Buffalo's milk, raw Cow's milk.

---

### INTRODUCTION

The local dairy commodity chain, from farm to retail sites is informal and often escapes monitoring of quality. In addition, most stakeholders lack knowledge on hygiene and sanitary aspects of their production, which can result in poor production standards and contamination of local raw milk. *Bifidobacteria* are well known for their beneficial effects on health and are used as probiotics in certain food and pharmaceutical products. Since they constitute one of the most important bacteria group in human and animal digestive tract, they were recently considered as fecal contamination indicators in raw milk dairy products. Moreover, the *Bifidobacteria* spp. isolated in humans are different from that isolated from animals (Delcenserie, *et al.*, 2005). Fecal contamination of raw milk in farms has been shown by the detection of the same and most frequent *Bifidobacterium* spp. in milk as in cow's dung. Raw milk can be assumed to be the first critical point in an HACCP analysis of the raw milk cheese industry, but

a follow-up of contamination during the cheese-making process is also of interest. The standard in Europe for fecal contamination control of raw milk cheese is *Escherichia coli* (Beerens *et al.*, 2000).

The general objective is to define a new standard, using the *Bifidobacteria* (genus *Bifidobacterium*) as indicator organisms in order to point out unsatisfactory hygienic conditions of raw material and food products. These new indicators will be applied on raw milk and cheese made from it, as the quality of which depends on the hygiene of milking, the farm environment and the hygiene along the cheese production chain. Also meat and meat products in which *Bifidobacteria* will indicate hygienic shortcomings during slaughter, cutting and debuting as well as retail trade. Good hygienic practices must lead to the elimination of all bacterial hazards such as *Listeria*, *E. coli* (Enterohemorrhagic *E. coli* (EHEC)) or *Salmonella* spp. The knowledge of the *Bifidobacterium* spp. that contaminate raw milk, cheese made with raw milk, meat and meat products will point out the sites from which the contamination

derives the development of a fast, sensitive and generally applicable technique will enable industries to detect *Bifidobacteria* during processing and to take corrective actions immediately. Due to their intestinal ecology, milk is not the natural environment of *Bifidobacteria*. Their presence in milk is a sign of fecal contamination (Delcenserie *et al.*, 2005) and can therefore serve as a marker of the level of hygiene in the production chain (Beerens *et al.*, 2000). In addition, the species characterization and assessment of antibacterial potential could lead to its determination in milk, as a selection method of food safety strains. Moreover, an advantage in using *Bifidobacteria* instead of other fecal contamination indicators is the host specificity of *Bifidobacterium* spp in human or animal (Gavini and Beerens, 1999). Raw milk collected with proper hygiene should not contain *Bifidobacteria*. In fact, the presence of *Bifidobacteria* in raw milk indicates fecal contamination and poor farm hygiene (Beerens *et al.*, 2000; Beerens and Neut, 2005). This bacterium has been recently proposed as a fecal indicator in water (Lynch *et al.*, 2002; Gilpin *et al.*, 2003 and Nebra *et al.*, 2003) and in meat and raw milk samples (Beerens, 1998; Gavini and Beerens, 1999; Beerens *et al.*, 2000).

*Bifidobacteria* have been suggested as among the more promising alternative indicators of faecal pollution. They are present in faeces at levels 10:100 times greater than coliforms and, due to their anaerobic nature, fastidious growth requirements and inability to grow below 20°C, they are unlikely to multiply outside the intestinal tract (Mara and Oragui, 1983).

Since most bifidobacterial strains used in dairy products are common members of the human intestinal microbiota, they may behave as opportunistic pathogens, like other commensal bacteria (Saarela *et al.*, 2002). Some commensal *Bifidobacteria* have been connected with certain dental infections, pulmonary infections, bacteremia, abscesses and bloodstream infections (Green, 1978; Gasser, 1994; Saarela *et al.*, 2002). In infants aged 2 to 16 months, Brook (1996) described a prevalence of 57 *Bifidobacterium* isolates among 2033 specimens originating from chronic otitis media, abscesses, peritonitis, aspiration pneumonia and paronychia. In most cases, the non-probiotic *B. dentium* (formerly *B. eriksonii*) was implicated, this is the only bifidobacterial species classified as a dental pathogen. A few cases also implicated *B. adolescentis*, which has been recovered pure from otitis media (Brook, 1996). Despite these few cases, the recovery of *Bifidobacteria* in clinical infections is very rare in comparison to other commensals (Borriello *et al.*, 2003). The most commonly used faecal indicator organisms, faecal coliforms, including *E. coli*, denote faecal pollution but not whether it is of human or animal origin. As a consequence, alternative indicators of faecal pollution better able to identify the faecal source have been suggested, including

*Bacteriodes* spp., phages of *Bacteroides fragilis*, F-RNA phage, *Rhodococcus coprophilus*, *Bifidobacteria*, and various chemical indicators (Sinton *et al.*, 1998).

*Bifidobacteria*, previously known as *Bacillus bifidus*, are a group of microorganisms that were classified into 29 different species in the 9th Edition of Bergey's Manual of Systematic Bacteriology (Scardovi, 1986). Nowadays, there are more than 33 species that were identified from the genus *Bifidobacterium* using recent molecular and biochemical methods of which 12 have been associated with the human gastrointestinal tract (Ventura *et al.*, 2004).

*Bifidobacteria* are Gram-positive, non-motile, and non-spore-forming bacteria. They are part of normal intestinal microbiota in humans and animals, and are generally non-pathogenic. They were considered to be anaerobic (Simpson *et al.*, 2004a).

Present study has been carried out to evaluate the hygienic quality of raw buffalo's and cow's milk sold in Assiut city based on the presence of *Bifidobacterium* spp. as indicators of faecal contamination.

## MATERIALS and METHODS

**1- Raw milk samples:** 70 samples of raw buffalo's and cow's milk (35 samples from each) were collected randomly from; some dairy farms, individual cases of dairy buffaloes and cows and dairy shops in Assiut city. Samples collected directly after milking and transported to the microbiological laboratory for examination. Each sample was kept in a sterile bag containing peptone water (0.1%) to which we added 0.25% L-cysteine.

**2- Culture media:** Numerous culture-based methods for *Bifidobacteria* detection, isolation and typing have been described. Culturing of *Bifidobacterium* and its evolution within gastrointestinal flora (human or animal) was carried out according to Martineau (1999); Rada and Petr (2000) and Petr and Rada (2001). MRS & BHIMup and two different culture media (CMup & *Bifidobacterium* media (BFM)) were used for isolation and identification of *Bifidobacterium* spp.

### 1- *Bifidobacteria* isolation:

Two culture-based methods were used to detect *Bifidobacteria* spp. which comprised enrichment and isolation steps. The first method was done by inoculating 2 g of each sample of raw milk in 18 ml MRS broth as enrichment medium. 500 mg / l of L-cysteine hydrochloride has been added to MRS broth to reduce ox-reduction (ox-re) potential. The inoculated media were incubated anaerobically at 37°C for 48 hrs. In order to obtain single colonies, the cultured mMRS were streaked on the surface of

*Bifidobacterium media* (BFM) and incubated anaerobically at 37°C for 72hrs.

The second method was carried out according to isolation procedures described by Beerens (1998) and culture-based methods modified by Delcenserie *et al.* (2005). One milliliter of each sample was transferred into 9 ml of enrichment medium BHMup [BHI, 37 g/l (Oxoid, England), 5 ml/l of propionic acid, 0.5g/l Fe-citrate, 0.5 g/l cysteine hydrochloride, 5 g/l yeast extract and 2 g/l agar]. Mupirocin, 80 mg/l was added before the use of the medium. The final pH was 5.0 and obtained with the addition of 1 mol/l NaOH solution. Tubes were incubated for 48 hrs at 37°C in Jars with anaerobic conditions. From each enrichment culture, 0.1 ml was spread on isolation medium CMup (Columbia blood agar, 0.5 g/l, Fe-citrate, 5 g/l glucose and 0.5 g/l cysteine hydrochloride). Mupirocin was added (50 mg/l) before the use of the medium. Plates were incubated anaerobically at 37°C for 72 hrs. Samples which have grown on both of the two selective media considered positive while samples which have grown on only one media, have been discarded. Colonies were rounded, blue and approximately 2mm in diameter a few minutes after removal from anaerobic condition.

**2- Identification of isolates:**

**a- Phenotypic characterization:** Each strain isolated was cultured on MRS agar (MRS agar to which we added, 0.25% L-cysteine). Inoculated plates were incubated anaerobically at 37°C for 24 to 48h. Isolates observed were stained with methylene blue and examined microscopically (1000 x). Only strains with the "Y" phenotypic form were selected for the identification (Rasic and Kurman, 1983).

All strains were initially submitted to Gram staining, catalase and spore formation test. Colonies and cells morphology characteristics on TPY containing 0.2% cysteine-HCl, MRS and M17 agar were also examined. Gas production from glucose was determined in MRS broth containing inverted Durham (Grill *et al.*, 2000). Citrate utilization, in the presence of carbohydrates, was performed on the media of Kihal *et al.* (1996).

**b- Carbohydrate fermentation**

The carbohydrate fermentation was determined on TPY broth containing bromocresol purple (0.04 g/l) as a pH indicator, and supplemented with 1% of the following carbohydrates: lactose, sucrose, xylose, arabinose, sorbitol, fructose, galactose, mannose, cellobiose, raffinose, melizitose and melibiose. To ensure anaerobic conditions, each tube was supplemented with two drops of sterile liquid paraffin after inoculation (Samelis *et al.*, 1994; Saidi *et al.*, 2002). The results obtained for morphological, physiological and biochemical tests were compared with those in standard texts for identification

(Scardovi, 1986; Miyake *et al.*, 1998, Ingrassia *et al.*, 2001) and the isolates assigned to appropriate species. The isolates belonging to the genus *Bifidobacterium* were identified to species level using sugar fermentation and the profiles were compared with the reference strains in standard texts for identification.

**RESULTS**

*Bifidobacterium* spp. was identified in 65.71 % (23 samples) and 51.42% (18 samples) of raw buffalo's and cow's milk, respectively (Table, 1). Isolates were identified and findings revealed that raw cow's milk samples harbored *B.dentium* in 61.11% while *B. suis* was identified in 27.77% of examined samples (Table, 2). *B. bifidum* were isolated from 2 samples (11.11%). *B. dentium*, *B. suis* and *B. bifidum* were found contaminating 47.82, 39.13 and 13.04% of examined raw buffalo's milk (Table, 2).

**Table 1:** Percentage of positive raw Cow's and Buffalo's milk samples that contain *Bifidobacteria* spp.

| Type of examined samples | No. of examined samples | Positive samples for all isolated <i>Bifidobacteria</i> spp. |         |
|--------------------------|-------------------------|--|---------|
|                          |                         | No.  | %       |
| Raw cow's milk           | 35                      | 18   | 51.42%  |
| Raw buffalo's milk       | 35                      | 23   | 65.71 % |

**Table 2:** *Bifidobacteria* spp. isolated from the examined samples of raw cow's and buffalo's milk.

| <i>Bifidobacteria</i> spp.     | Raw Cow's milk          |            | Raw Buffalo's milk       |            |
|--------------------------------|-------------------------|------------|--------------------------|------------|
|                                | Positive sample No.= 18 | %          | Positive samples No.= 23 | %          |
| <i>Bifidobacterium dentium</i> | 11                      | 61.11      | 11                       | 47.82      |
| <i>Bifidobacterium suis</i>    | 5                       | 27.77      | 9                        | 39.13      |
| <i>Bifidobacterium bifidum</i> | 2                       | 11.11      | 3                        | 13.02      |
| <b>Total</b>                   | <b>18</b>               | <b>100</b> | <b>23</b>                | <b>100</b> |

## DISCUSSION

Presence of *B. bifidum* revealed that raw cow's and buffalo's milk may be contaminated with feces of the human (adult and infants). However, isolation of *B. suis* could be attributed to pollution with feces of piglets. Also the detection of *B. dentium* in examined samples could be attributed also to contamination with human discharges coming from; mouth suffering dental caries, human abscesses, vaginal discharges and feces. These organisms considered potentially pathogenic and have hazards effect on human health.

Commensal *Bifidobacteria* (*B. dentium*) have been connected with certain dental and other infections (Leo *et al.*, 2008). Lynch *et al.* (2002), Nebra *et al.* (2003) and Bonjoch *et al.* (2004) recorded that *B. adolescentis* or *B. dentium* are indicators of fecal pollution. As these species are dominant in human feces, they will indicate a contamination of human origin (Scardovi, 1986).

Kouamé-Sina *et al.* (2011) found that isolates belong to five different species of *Bifidobacterium* were present in 9% of milk samples. Most of the *Bifidobacterium* isolated were *B. minimum* (53%) and *B. pseudolongum* sub spp. *globosum* (24.4%). The other strains are composed of one strain of *B. thermophilum*, *B. thermacidophilum* sub spp. *Suis* and *B. magnum*.

The typical habitats of *Bifidobacteria* are human, warm-blooded animal and honeybee intestinal tract (Scardovi, 1986). Members of genus *Bifidobacterium* are among the most common microorganisms in the human gut, comprising up to 3% of the total faecal microflora of adults (Sghir *et al.*, 2000). They are more numerous in the infant gut, where they form up to 91% of the total microflora in breast-fed babies being supported by bifidogenic factors presented in human milk and up to 75% in formula-fed infants (Harmsen *et al.*, 2000). Using classical culturing methods it has been found that *B. adolescentis* and *B. longum* are major bifidobacterial species in the intestine of adults (Biavati *et al.*, 1986; Mutai and Tanaka, 1987; Gavini *et al.*, 2001). *B. infantis* and *B. breve* are predominant species in the intestinal tract of human infants (Benno *et al.*, 1984; Biavati *et al.*, 1984; Mutai and Tanaka, 1987). In addition, *B. bifidum*, *B. catenulatum*, *B. pseudocatenulatum*, *B. angulatum*, *B. gallicum*, and *B. dentium* have also been reported to be human intestinal bifidobacteria (Scardovi, 1986). Matsuki *et al.* (1999) who used for the detection of bifidobacteria in human gut species-specific polymerase chain reaction (PCR) reported, that the most common species in the breast-fed infants are *B. breve*, *B. infantis*, *B. longum*, and *B. bifidum*. In adult intestinal tracts, the *B. catenulatum* group was the most common taxon, followed by *B. longum* and *B. adolescentis*.

In the case of raw milk samples, an enrichment step was necessary because of the possible relatively low

levels of bifidobacteria ( $10$  to  $10^6$ / ml<sup>-1</sup>) compared to those in human or animal feces ( $10^7$  to  $10^{10}$ / g<sup>-1</sup>). Beerens (1998) recommended the using at the enrichment step the BHI medium with addition of propionic acid, yeast extract, iron citrate, and at the isolation step, paromomycin as selective agent. However, the high number of lactobacilli that were not inhibited by paromomycin hides bifidobacteria at low dilutions. Rada and Petr (2000) showed that bifidobacteria were resistant to mupirocin when lactobacilli were susceptible. Mupirocin (pseudomonamic acid A) was originally isolated from *Pseudomonas fluorescens* and used as a topical antibiotic (Sutherland *et al.*, 1985). In raw milk samples, addition of mupirocin at the enrichment step can eliminate most of the lactobacilli strains present that could hide *Bifidobacteria* in raw milk. If some lactobacilli strains were still present after the enrichment step, one might suppose that they would be eliminated during the isolation step by mupirocin, when present. Grand *et al.* (2003) also used mupirocin as selective agent for detection of *Bifidobacteria* in probiotic milk products, as did Mikkelsen *et al.* (2003) in gastrointestinal samples from piglets and Simpson *et al.* (2004b) in probiotic as animal feed.

Moreover, Rhodes and Kator (1999) enumerated sorbitol-fermenting *Bifidobacteria* to define human fecal pollution in estuarine watersheds. However, in raw milk cheese, the principal contamination was shown to be of animal origin (Beerens *et al.*, 2000), most likely by cow dung on farm, since the same species, *B. pseudolongum*, was isolated from both kinds of samples. Therefore, in food industries, it seems important to define the human or animal origin of the contamination.

The milk contamination by *Bifidobacteria* is also an indicator of fecal contamination (Beerens *et al.*, 2000) and is due to the lack of hygiene in farms. Its presence on cow's udder reflects environmental contamination with feces of other animals living in farms or in the vicinity. Our results indicated insignificant variation in the prevalence of *Bifidobacterium* along the dairy production chain. Contamination by *Bifidobacterium* was present at all stages of milk production. These results indicated fecal contamination at all stages of the production chain, from production to selling point of raw milk. This suggests a lack of sanitary education of farmers and milk vendors. This sanitary education of farmers is important especially on good hygiene practices through simple actions like washing and disinfecting of hands, cow's udders and milking utensils (Bonfoh *et al.*, 2003).

Finally it could be concluded that the hygienic quality of local raw milk was assessed as poor, based on the detection of the *Bifidobacterium* in raw milk (Therefore the *Bifidobacterium* isolated in this study are rather an indication of poor hygiene). In order to improve the microbiological quality of local raw milk, fermentation as well as heating is needed to

inhibit the growth of contamination germs. The safety strategy should be developed, with good hygiene practices related to cleanliness of animals and their environment as well as sanitation of the milking process (milker's hands, milking utensils).

## REFERENCES

- Beerens H. (1998):* Bifidobacteria as indicators of faecal contamination in meat and meat products: detection, determination of origin and comparison with *Escherichia coli*. *Int. J. Food Microbiol.*, 40: 203-207.
- Beerens, H.; De la Perriere, H.B. and Gravini, F. (2000):* Evaluation of the hygienic quality of raw milk based on the presence of *Bifidobacteria*: the cow as a source of faecal contamination. *Int. J. Food Microbiol.*, 54: 163-169.
- Beerens, H. and Neut, C. (2005):* Usefulness of *bifidobacteria* for the detection of faecal contamination in milk and cheese. *Lait*. 85: 33-38.
- Benno, Y.; Sawada, K. and Mitsuoka, T. (1984):* The intestinal microflora of infants: composition of fecal flora in breast-fed and bottle-fed infants. *Microbiol. Immunol.* 28: 975-986.
- Biavati, B.; Catagnoli, P. and Trovatelli, L.D. (1986):* Species of the *Bifidobacterium* in the feces of human adults. *Microbiologica.*, 9: 39-45.
- Biavati, B.; Catagnoli, P.; Crociani, F. and Trovatelli, L.D. (1984):* Species of the *Bifidobacterium* in the feces of infants. *Microbiologica.* 7: 341-345.
- Bonfoh, B.; Wasem, A.; Traoré, A.N.; Fané, A.; Spillmann, H.; Simbé, C.F.; Alfaroukh, I.O.; Nicolet, J.; Farah, Z. and Zinsstag, J. (2003):* Microbiological quality of cows' milk taken at different intervals from the udder to the selling point in Bamako (Mali). *Food Control*, 4: 495-500.
- Bonjoch, X.; Balleste, E. and Blanch, A. (2004):* Multiplex PCR with 16S rRNA gene-targeted primers of *Bifidobacterium* spp. to identify sources of fecal pollution. *Appl. Environ. Microbiol.* 70: 3171-3175.
- Borriello, S.P.; Hammes, W.P.; Holzapfel, W.; Marteau, P.; Schrezenmeir, J.; Vaara, M. and Valtonen, V. (2003):* Safety of probiotics that contain *lactobacilliorbifido bacteria*. *Clinical Infectious Diseases.* 36: 775-780.
- Brook, I. (1996):* Isolation of non-sporing anaerobic rods from infections in children. *J. of Medical Microbiol.* 45: 21-26.
- Delcenserie, V.; Bechoux, N.; China, B.; Daube, G. and Gavini, F. (2005):* A PCR method for detection of *Bifidobacteria* in raw milk and raw milk cheese: comparison with culture-based methods. *J. Microbiol. Methods.* 61: 55-67.
- Gasser, F. (1994):* Safety of lactic acid bacteria and their occurrence in human clinical infections. *Bulletin de l'Institut Pasteur.* 92: 45-67.
- Gavini, F. and Beerens, H. (1999):* Origin and identification of *bifidobacteria* strains isolated from meat and meat products. *Int. J. Food Microbiol.* 46: 81-85.
- Gavini, F.; Cayuela, Ch.; Antoine, J.M.; Lecoq, C.; Lefebvre, B.; Membré, J.M. and Neut, Ch. (2001):* Differences in the distribution of bifidobacterial and enterobacterial species in human faecal microflora of three different (children, adults, elderly) age groups. *Microb. Ecol. Health Dis.*, 13: 40-45.
- Gilpin, B.; James, T.; Nourozi, F.; Saunders, D.; Scholes, P. and Savill, M. (2003):* The use of chemical and molecular microbial indicators for faecal source identification. *Water Sci. Technol.* 47: 39-43.
- Grand, M.; Küffer, M. and Baumgartner, A. (2003):* Quantitative analysis and molecular identification of bifidobacteria strains in probiotic milk products. *Eur. Food Res. Technol.* (217): 90-92.
- Green, S.L. (1978):* Case report: fatal anaerobic pulmonary infection due to *Bifidobacterium eriksonii*. *Postgraduate Medicine.* 63: 187-188.
- Grill, J.P.; Perrin, S. and Schneider, F. (2000):* Bile salt toxicity to some bifidobacteria strains: role of conjugated bile salt hydrolase and pH. *Edge J. Microbiol.* 46 (10): 878-884.
- Harmsen, H.J.M.; Wildehoer, A.C.M.; Raangs, G.C.; Wagendorp, A.A.; Klijn, N.; Bindels, J.G. and Welling, G.W. (2000):* Analysis of intestinal flora development in breast-fed and formula-fed infants by using molecular identification and detection methods. *J. Pediatr. Gastroenterol. Nutr.* 30: 61-67.
- Ingrassia, T.; Roques, C. and Prevots, F. (2001):* Preliminary experiments for ARDRA validation on flora associated with intestinal mucosa. *Lait* 81: 263-280.
- Kihal, M.; Prevost, H.; Lhotte, M.E.; Huang, D.Q. and Diviès, C. (1996):* Instability of plasmid-encoded citrate permease in *Leuconostoc*. *J. Appl. Microbiol.* 22: 219-223.
- Kouamé-Sina, S.M.; Kohei, M.; Delia, G.; Marcellin D.; Bernard, T.; Georges, D. and Bassirou, B. (2011):* Diversity, phylogenetic relationship and antibacterial potential of *Bifidobacterium* species isolated from raw milk production chain in Abidjan (Côte d'Ivoire). *Afr. J. Microbiol. Res.* 5 (21): 3394-3403.
- Leo, M.; Gwenaelle, L. and Anne, T. (2008):* Safety assessment of dairy microorganisms: *Propionibacterium* and *Bifidobacterium*. *Int. J. Food Microb.* 126: 316-320.
- Lynch, P.; Gilpin, B.; Sinton, L. and Savill, M. (2002):* The detection of *Bifidobacterium adolescentis* by colony hybridization as an indicator of human faecal pollution. *J. Appl. Microbiol.* 92: 526-533.

- Mara, D.D. and Oragui, J.I. (1983): Sorbitol-fermenting bifidobacteria as specific indicators of human faecal pollution. *J. Appl. Bact.*, 55: 349-357.
- Martineau, B. (1999): Comparison of four media for the selection of *Bifidobacteria* in dog fecal samples. *Anaerobe*, 5: 123-127.
- Matsuki, T.; Watanabe, K.; Tanaka, R.; Fukuda, M. and Oyaizu, H. (1999): Distribution of *Bifidobacterial* species in human intestinal microflora examined with 16 S rRNA-gene-targeted species-specific primers. *Appl. Environ. Microbiol.* 65: 4506-4512.
- Mikkelsen, L.; Bendixen, C.; Jakobsen, M. and Jensen, B. (2003): Enumeration of *Bifidobacteria* in gastrointestinal samples from piglets. *Appl. Environ. Microbiol.* 69: 654-658.
- Miyake, T.; Watanabe, K.; Watanabe, T. and Oyaizu, H. (1998): Phylogenetic analysis of the genus *Bifidobacterium* and related will genera based one 16S RNA sequences. *Microbiol. Immunol.* 42: 661-667.
- Mutai, M. and Tanaka, R. (1987): Ecology of *Bifidobacterium* in the human intestinal flora. *Bifidobacteria Microflora*. 6: 33-61.
- Nebra, Y.; Bonjoch, X. and Blanch, A. (2003): Use of *Bifidobacterium dentium* as an indicator of the origin of fecal water pollution. *Appl. Environ. Microbiol.* 69: 2651-2656.
- Petr, J. and Rada, V. (2001): *Bifidobacteria* are obligate inhabitants of the crop of adult laying hens. *J. Vet. Med., B. Infect. Dis. Vet. Public Health.* 48: 227-233.
- Rada, V. and Petr, J. (2000): A new selective medium for the isolation of glucose non-fermenting *Bifidobacteria* from hen caeca. *J. Microbiol. Methods.* 43: 127-132.
- Rasic, J.L. and Kurman, J.A. (1983): *Bifidobacteria* and their role. Microbiology, nutritional physiological, medical and technological aspects and bibliography. Binkauser Verlag, Basel, Boston Stuttgart.
- Rhodes, M.W. and Kator, H. (1999): Sorbitol-fermenting bifidobacteria as indicators of diffuse human fecal pollution in estuarine watersheds. *J. Appl. Bact.* 87: 528-535.
- Saarela, M.; Mättö, J. and Mattila-Sandholm, T. (2002): Safety aspects of *Lactobacillus* and *Bifidobacterium* species originating from human oro-gastrointestinal tract or from probiotic products. *Microbial. Ecology in Health and Disease* 14: 233-240.
- Saidi, N.; Guessas, B.; Bensalah, F.; Badis, A.; Hadadji, M.; Henni, D.E.; Prevost, H. and Kihal, M. (2002): Caractérisation des bactéries lactiques isolées du lait cru de chèvre des régions arides. *J. Alg. Reg. Arides*: 101-111.
- Samelis, J.; Maurogenakis, F. and Metaxopoulos, J. (1994): Characterization of lactic acid bacteria isolated from naturally fermented Greek dry salami. *Inter. J. Food Microbiol.* 23: 179-196.
- Scardovi, V. (1986): Genus *Bifidobacterium*. In: *Bergey's Manual of Systematic Bacteriology*, Vol. 2 (Eds.: Sneath, P.H.A./Mair, N.S. / Sharpe, M.E. / Holt, J.G.). Baltimore, Williams and Wilkins, 1418-1434.
- Sghir, A.; Gramet, G.; Suau, A.; Rochet, V.; Pochart, P. and Dore, J. (2000): Quantification of bacterial groups within the human fecal flora by oligonucleotide probe hybridization. *Appl. Environ. Microbiol.* 66: 2263-2266.
- Simpson, P.J.; Ross, R.P.; Fitzgerald, G.F. and Stanton, C. (2004a): *Bifidobacterium psychraerophilum* spp. nov. and *Aeriscardovia aeriphila* gen. nov., sp. nov., isolated from a porcine caecum. *Int. J. Syst. Evol. Microbiol.* 54: 401-406.
- Simpson, P.; Fitzgerald, G.; Stanton, C. and Ross, R. (2004b): The evaluation of a mupirocin-based selective medium for the enumeration of *Bifidobacteria* from probiotic animal feed. *J. Microbiol. Methods.* 57: 9-16.
- Sinton, L.; Finlay, R. and Hannah, D. (1998): Distinguishing human from animal faecal contamination in water: a review. *New Zealand J. Marine and Freshwater Research.* 32: 323-348.
- Sutherland, R.; Boon, R.; Griffin, K.; Masters, P.; Slocombe, B. and White, A. (1985): Antibacterial activity of mupirocin (pseudomonic acid), a new antibiotic for topical use. *Antimicrob. Agents Chemother.* 27: 495-498.
- Ventura, M.; De van Sinderen, G.F.; Fitzgerald, and Zink, R. (2004): Insights into the taxonomy, genetics and physiology of *Bifidobacteria*. *Antoine van Leeuwenhoek.* 86: 205-233.

تقييم الجودة الصحية للين الخام اعتمادا على تواجد ميكروب البيفيدوبكتريا كمؤشر للتلوث بالروث  
في مدينة اسيوط

مروة محمد نبيل الجندي ، لمياء محمد طلعت علي

استهدف هذا البحث تقييم جودة حليب الابقار والجاموس وذلك من خلال وجود ميكروب البيفيدوبكتريا كمؤشر للتلوث بالروث، هذا بالإضافة الى تحديد مصدر التلوث سواء كان من الانسان او الحيوان. البيفيدوبكتريا واحدة من الميكروبات المتعايشة الطبيعية في امعاء الانسان والحيوان وتتواجد بكثرة في البراز والروث الخاصة بهما. وجود هذا الميكروب في الالبان الخام دليل على التلوث وعدم اتباع الاشتراطات الصحية اثناء الحليب. تم جمع سبعين عينة من الالبان الابقار والجاموس الحلوب لدى المربين وكذلك من بعض مزارع انتاج الالبان ومحلات بيعها في مدينة اسيوط اخذت العينات بطريقة عشوائية مباشرة بعد الحليب واشتملت على عدد 35 عينة من من الالبان الابقار وكذلك 35 عينة من الالبان الجاموس. اجريت الاختبارات اللازمة للكشف عن وجود ميكروب البيفيدوبكتريا في كل عينة من عينات البحث. بداية تم استخدام الوسط الغذائي MRS و BHIMup كبيئة مغذية غنية لنمو الميكروب وتم التحضين على درجة C 37 لمدة يومان. هذا وقد استخدم نوعين من المستنبتات الغذائية وهما CMup و Bifidobacterium media (BFM) لعزل الميكروبيوتم التحضين على C 37 لمدة 72 ساعة. أظهرت النتائج ان عينات اللين الجاموسي الخام كانت الأكثر تلوثا بالبيفيدوبكتريا بنسبة 65.71% (23 عينة) مقارنة بعينات اللين البقري والذي بلغت نسبة التلوث به الى 51.42% (18 عينة). كما أوضحت نتائج تصنيف ال *Bifidobacterium* ان النمط *Bifidobacterium dentium* هو الاكثر عزلا وبنسبة 61.11 و 47.82%، في اللين البقري واللين الجاموسي الخام على الترتيب. في حين ان *Bifidobacterium suis* وجدت في 5 ، 9 عينات من الالبان الابقار والجاموس وبنسب 27.77 ، 39.13% على الترتيب. ال *Bifidobacterium bifidum* كانت الاقل تواجدا (بنسبة 13.04%) في اللين الجاموسي وبنسبة 11.11% في اللين البقري. عزل هذه العترات من عينات الالبان تشير الي تلوثها ببراز الانسان البالغ والاطفال وذلك نتيجة وجود ال *Bifidobacterium bifidum* بينما تواجد ال *Bifidobacterium suis* تظهر التلوث بروث صغار الخنازير. أما عزل ال *Bifidobacterium dentium* فانها تبين تلوث عينات الالبان بقراوات من الفم والاسنان المتسوسة والدمامل وبراز الانسان البالغ (human dental caries; human abscesses and stools).