

**Phylogenetic Analysis of Antibiotic Producing Bacteria Isolated from Soil Contaminated by Human Wastes using Random Amplified Polymorphic DNA (RAPD) Technique**

Doaa Hassan<sup>1</sup>, Mohamed, N.A.<sup>1</sup>, Saleh, F.<sup>1</sup>, Mohamed, E.<sup>2</sup>

<sup>1</sup>Genetics Department, Faculty of Agriculture, Assiut University

<sup>2</sup>Agricultural Botany Dep. (Genetics), Faculty of Agriculture, South Valley University

---

**Abstract:**

Antibiotic is one of the most important commercially exploited secondary metabolites produced by various microorganisms (as a bacterium or a fungus). The growing recognition of the pressing need for new antimicrobial agents for the treatment of infectious diseases demands discovering new antibiotic-producing strains of bacteria.

In the present investigation, soil samples contaminated by human feces were collected from different geographical regions belonging to Cairo, Assiut and Qena cities, Egypt; to isolate some strains of antibiotic-producing bacteria and characterize the isolates using RAPD-PCR technique. One hundred and ten strains of bacterial isolates were isolated from the collected soil samples. From which five selected isolates (D110, D3, D103, D106 and D104) showed promising results for an effective antibiotic production according to the microbial sensitivity tests with two Gram positive and two Gram negative bacterial strains by forming a clear zone of inhibition. The results showed that the average of clear zones of Gram positive bacteria are wider than the average of clear zones of Gram negative. Such isolates have more ability to inhibit the growth of Gram positive bacteria than to inhibit the growth of Gram negative bacteria. The five coded selected strains were identified as, *Bacillus licheniformis*, *Bacillus pumilus*, *Staphylococcus epidermidis*, *Serratia marcescens* and *Paenibacillus lautus*, respectively.

In addition, it has been possible to differentiate between the isolated strains at the molecular level using RAPD-PCR. The results showed that these isolates can be described according to the banding-profile pattern. The five selected isolates were grouped in four clusters; based on genetic similarity. The first and the second groups were represented by D106 and D103 respectively. The third group included the two isolates; D104 and 110 and the strain D3 represented the fourth group.

---

**Keywords:** Phylogenetic, antibiotic, human wastes, RAPD

---

**Received on:** 30/10/2013

**Accepted for publication on:** 18/11/2013

**Referees:** Prof. El-Sayed N. E. Hamed

Prof. Fahmi A. Nassef

### Introduction:

Antibiotics are one of the most important commercially exploited secondary metabolites produced by bacteria. The remarkable increase in antibiotics resistant bacterial species (Motta *et al.*, 2004; and Huda Sheikh, 2010) lead to search for new sources of antibiotics through the isolation and identification of new types of microorganisms such as bacteria, fungi and actinomycetes (Alexander, 1982). *Bacillus* species producing antibiotics have been used as biocontrol agents against pathogenic fungi (Yilmaz, *et al.*, 2005). More than 1200 isolates of lactic acid bacteria isolated from different environments were screened for antifungal activity, from which approximately 10% represented inhibitory activity and 4% showed strong activity against *Aspergillus fumigatus* (Magnusson, *et al.*, 2003).

Soil is considered one of the most suitable environments for microbial growth (Cavalcanti, *et al.*, 2006). Most of the antibiotic producers used today are soil microbes. The genus *Streptomyces*, an antibiotics producer, has been isolated from the soil of Yemen (Ahmed, 2003). One hundred bacterial isolates were isolated from six different soil samples collected from Egypt (Gebrael, *et al.*, 2008). They reported that twenty of them could antagonized some selected plants and human pathogenic fungi with varying degrees. Moreover, among twenty bacterial strains isolated from soil stressed agroecological niches of Eastern Uttar Pradesh, India, eleven isolates showed strong antimicrobial activities (Singh, *et al.*, 2009). Fecal wastes from domestic, wild life animals and human applied to the soil may con-

tain a wide variety of pathogenic viruses, bacteria, and parasites.

The traditional typing methods for discriminating different bacterial isolates of the same species are essential epidemiological tools in infection prevention and control. Traditional typing systems based on phenotypes, such as serotype, biotype, or antibiogram, have been used for many years. However, the developments of molecular methods have revolutionized our ability to differentiate among bacterial isolates (Sabat, *et al.*, 2013). Random Amplified Polymorphic DNA - PCR (RAPD-PCR), which has been described by Williams, *et al.*, (1990) is the most frequently used PCR-based techniques. In this techniques single or a pairwise combination of primers, typically 10 nucleotides in length, are used to amplify target sites for the primer happen to occur within approximately 5 Kb of each other on opposite DNA strand.

The present investigation aimed to isolate some strains of antibiotic-producing bacteria from soil samples contaminated by human feces collected from different geographical regions in Egypt and characterizing the isolates using RAPD-PCR technique.

### Materials and Methods:

#### • Collection of Soil samples:

Five soil samples were collected from different locations belonging to Cairo, Assiut and Qena cities, Egypt. Soil samples were taken in sterilized polyethylene bag using sterilized spatula then transferred to the laboratory at the Department of Genetics, Assiut University. Samples were dried at air oven and then cooled to room temperature.

• **Isolation of bacterial isolates:**

Suspensions were made by adding 1g of each soil sample to 100 ml sterile saline solution. Ten fold dilutions of these suspensions were planted on Luria – Bertani (LB) agar medium and incubated at 37°C for 72 hours. Single colonies of the grown bacteria which varied in shape and color were picked up using a sterile loop and purified by streaking on nutrient LB agar. The bacterial isolates were kept on LB slants agar and stored at 4° C and recultured monthly.

• **Testing pure isolated cultures for their abilities to produce antibiotics:**


The bacterial tester strains, which were used to detect antibiotic activity of isolated strains, were delivered from Medical Microbiology and Immunity Department, Faculty of

Medicine and, Dairy Science Department, Faculty of Agriculture, Assiut University, Assiut, Egypt. The tester strains were chosen according to (Narisawa, *et al.*, 2008; and Chandrashekhara, 2010) and included the following different genus of bacteria:


I) Two Gram positive Bacteria: *Bacillus subtilis* and *Staphylococcus aureus*.

II) Two Gram negative Bacteria: *Escherichia coli* and *Pseudomonas aeruginosa*.

All isolates were tested for antibiotic production by two different methods: Filter disk assay (Fitri and Bustam, 2010) and Agar Streak Method (Basavaraj, *et al.*, 2010; and Chandrashekhara, 2010). Fig (1) shows an example of antibiotic activity of isolated strains tested by the two methods.



(a)



(b)

Fig (1) An example of antibiotic activity of isolated strains tested by:

a- Filter disk assay

b- Agar Streak

The selected isolates were characterized morphologically. The morphological features were colony form,

surface, margin, elevation, and colour in addition to cell shape and gram stain.

• **Extraction of genomic DNA for PCR:**

The extraction of total bacterial genomic DNA was performed according to procedures described by Sambrook *et al.* (1989) with some modifications. Briefly, Cells from 50-ml cultures were harvested by centrifugation and the pellets were resuspended in 2 ml TEN buffer [150 mM NaCl, 10 mM Tris-HCl, 100 (pH=8.0), 100 mM EDTA (pH=8.0) containing 4 mg/mL lysozyme. The cells were grounded in liquid nitrogen then, 800 µl of 60°C CTAB extraction buffer (100mM Tris-HCl, 1.4M NaCl, 20mM EDTA, 2% hexadecyl-tri-methyl-ammonium bromide) was added, adjusted to pH=8.0 and incubated at 60 °C for 30 min. The DNA was extracted by adding 1 volume of chloroform-isoamyl alcohol (v/v = 24:1, pH=8.0), and the mixture was mixed gently and centrifuged at 5000 rpm for 20 minutes at 10°C. Then the aqueous phase was transferred to clean tube and the nucleic acids were precipitated by adding 0.1 volume of sodium acetate (pH=5.2) and 0.6 volume of isopropanol to the supernatant. The mixture was incubated in ice for 30 min and centrifuged (10,000 rpm for 5 min at 4 °C) to recover the precipitated nucleic acids. The pellets were washed with 70% ice-cold ethanol and centrifuged at 10,000 rpm for 2 min at 4 °C). The pellets were dissolved in 300-500 µl TE buffer (10 mM Tris/HCl, 1 mM EDTA, pH 8.0) and treated with 3µl RNAase and incubated in 37°C for 30 minutes. Samples were then treated with 3µl proteinase-K and incubated at 37°C for 30 minutes. The DNA

was precipitated, washed, and dissolved in TE as above. The DNA samples were kept at -20°C.

• **RAPD-PCR and electrophoretic analysis:**

Polymerase chain reaction (PCR) was conducted using ten arbitrary 10-mer oligonucleotied primers obtained from commercial sources (Operon Technologies, Alameda, CA, USA), Table (1) gives the codes and sequences of used Primers.. PCR reactions were conducted according to Williams *et al.* (1990). The reaction conditions were optimized and mixtures (25 µl total volume) were composed of 11.7 µl dH<sub>2</sub>O, 3.0 µl 10X reaction buffer, 3.0 µl dNTP's mix, 2.0 µl primer, 4.0 µl MgCl<sub>2</sub>, 0.3 µl Taq DNA polymerase and 1 µl Template DNA. The reaction was overlaid with mineral oil and thermal cycling achieved in TECHNE thermocycler (Model FTGEN5D, TECHNE, Cambridge Ltd, Duxford, and Cambridge, U.K.) according to the following program: initial denaturation at 85°C for 3 min, followed by 40 cycles of 85°C for 1 min, 33°C for 2 min, 72°C for 2 min, and a final extension at 72°C for 10 min and subsequently cooled to 4°C. PCR products were resolved by loading 14 µl of each reaction onto a 1.4% ultra pure agarose gel containing 0.01% ethidium bromide in 1X concentration Tris- acetate (TAE) buffer (89mM Tris-borate; 2.5mM EDTA). To visualize RAPD bands, electrophoresis was carried out under constant voltage of around 60V for approximately 3-3.5 hours. The patterns were visualized on a Transilluminator (Ultra-Violet Product, Upland, CA, USA,).

**Table (1): Codes and sequences of used Primers.**

Serial No.	Primer codes	Sequence (5' to 3')
1	OPA01	5'-CAGGCCCTTC-3'
2	OPA08	5'-GTGACGTAGG-3'
3	OPA09	5'-GGGTAACGCC-3'
4	OPI09	5'-GAGTCAGCAG-3'
5	UBC01	5'-CCTGGGCTTC-3'

**Analysis of RAPD banding patterns:**

RAPD-based molecular markers were scored visually using the software package MVSP (Multi-Variate Statistical Package) and DNA bands were scored as present (1) or absent (0). The pairwise comparisons between the tested isolates were used to calculate the coefficient of genetic similarity matrix (Gs) according to Nei and Li (1979). To convert the genetic similarity into genetic distance, logarithmic transformation (-Ln Gs) was computed to linearize the distance measure. Cluster analysis was presented as the dendrogram based on similarity estimates using the unweighted pair-group method with arithmetic average (UPGMA).

$$\text{Similarity} = \frac{2 * n11}{(2 * n11) + n01 + n10}$$

Where:

n11 - designates the number of common bands for the two compared samples,  
 n10 - cases where the bands were visible only in the first sample,  
 n01 - when bands were visible in the other sample only (Dice, 1945).

**Results and Discussion:**

**• Determination of antimicrobial activity:**

A total of 110 bacterial isolates were isolated from various soil samples obtained from different geographic regions. The isolates codes and their geographical regions are given in Table (2).

The data of the sensitivity of various tester strains (Gram-positive and Gram-negative) to the soil isolates by agar streak method are shown in (Table 3). Only ten isolates out of a total 110 bacterial isolates had broad spectrum antimicrobial activity against both Gram-positive and Gram-negative tester strains. These isolates were chosen for the further molecular differentiation.

**Table (2): Isolates codes and their geographical regions.**

Code name	Geographical region of isolates
From D1 to D30	Cairo
From D31 to D49	Qena
From D50 to D62	Assiut
From D63 to D84	New Assiut City
From D85 to D110	Qena

**Table (3): Sensitivity of various tester strains to the soil isolates by Agar Streak Method.**

+++ = Very good, ++ = Good inhibition, + = Moderate inhibition, - = No inhibition.

Soil bacterial isolates	Bacterial tester isolates			
	<i>B. Subtilis</i>	<i>S. aureus</i>	<i>E. coli</i>	<i>P. aeruginosa</i>
D 3, 55, 42	++	++	+	-
D 103	+	++	-	-
D 106, 29	++	+++	++	+++
D 110, 40, 65	+++	+++	+	-
D 104	++	-	++	-
The rest of the isolated strains	Gives moderate inhibition with just one or another of the tested strains.			

As shown in Table (3) it can be concluded that the tester strains, namely; *B. subtilis* and *S. aureus* (Gram positive) were more sensitive for bacterial soil isolates than *E. coli* and *P. aeruginosa* (Gram negative). The data also indicates that the lowest sensitivity to bacterial soil isolates linked to the tester strain *P. aeruginosa*, whereas the high sensitivity of the test strains for soil isolates were noted for *B. subtilis*. Fitri and Bustam (2010) concluded that the Gram negative bacteria usually have better protection to other antimicrobial compound rather than gram positive bacteria because both kinds of bacteria have different cell wall components, cell wall of Gram positive

bacteria contains peptidoglycan while cell wall of gram negative bacteria contains peptidoglycan and lipopolysaccharides.

Five strains that showed promising results for an effective antibiotic production through the microbial sensitivity tests were selected and identified as D110, D103, D104, D106, and D3. The characteristics of such strains are shown in Table (4). The five coded selected strains were identified as, *Bacillus licheniformis*, *Bacillus pumilus*, *Staphylococcus epidermidis*, *Serratia marcescens* and *Paenibacillus lautus* as shown in Table 6 (Doaa, et al., Personal communication).

**Table (4): The characteristics of the five selected isolates D110, D103, D104, D106, and D3.**

Characteristics Strains	Form	Surface	Margin	Elevation	Colour	Cell shape	Gram stain
D3 ( <i>Bacillus pumilus</i> )	Irregular	Smooth	Undulating	Flat	Yellowish	rod	+
D 103 ( <i>Staphylococcus epidermidis</i> )	Circular	Rough	Entire	Convex	Whitish	coccus	+
D104 ( <i>Paenibacillus lautus</i> )	Circular	Rough	Entire	Convex	Whitish	rod	-
D 106 ( <i>Serratia marcescens</i> )	Circular	Smooth	Entire	Convex slightly unbonate	Red pigment	rod	-
D 110 ( <i>Bacillus licheniformis</i> )	Irregular	Rough	lobate	unbonate	Whitish	rod	+

• **Genetic variation between the isolates by using RAPD-PCR analysis:**

RAPD-PCR analysis technique has been used to differentiate among the five selected antibiotic producing isolates. Figure (2) gives the Agarose gel electrophoresis of RAPD products obtained by the different primers. RAPD-PCR produced banding pat-

terns with products ranging from approximately 95 to 977bp. The total number of bands produced was 45, from which 19 bands were polymorphic with polymorphism percentage 42.22% (Table 5). The high polymorphism may be due to different sources/ locations of the of the isolates (Table 5).




Fig. (2): Agarose gel electrophoresis of RAPD products obtained by OPA-1, OPA-8, OPA-9, OPI-9, and UBC-1 primers.

**Table (5): Polymorphism obtained by RAPD analysis among the selected isolates of antibiotic producing bacteria using five different primers.**

primer	Total number of bands (a)	Number of polymorphic bands (b)	Polymorphism -b/a * 100%
OPA-1	8	1	12.5
OPA-8	9	7	77.78
OPA-9	10	4	40
OPI-9	9	3	33.33
UBC-1	9	4	44.44
<b>Total</b>	45	19	42.22

• **Genetic similarity matrix and cluster analysis**

Data of the presence /or absence of DNA fragments (markers) phenotypically analyzed using MVSP program of Nie and Li (1979), and pairwise comparisons between the tested isolates were used to calculate the genetic similarity. Then, based on the calculated genetic similarity values

presented in Table (6), an estimation of the relationship between the different isolates was concluded. Data showed that the lowest genetic similarity was observed between isolate D3 and D106 (80.6%), while the highest value was found between D110 and D104 (90.6.1%).




**Table (6): Genetic similarity values calculated from 45 DNA fragments generated with five primers in five bacteria strains.**

	D3	D110	D103	D106	D104
D3	1				
D110	0.879	1			
D103	0.836	0.870	1		
D106	0.806	0.870	0.829	1	
D104	0.903	0.906	0.892	0.862	1

The five promising isolates of bacteria were grouped in four clusters based on genetic similarity given in Table (6), and dendrogram as in Figure (3). The first and second groups

were represented by D106 and D103 respectively. The third group included the two isolates D104 and 110. The fourth group was represented by D3 strain.



**Fig. (3):** Dendrogram of five bacteria strains developed from RAPD data using UPGMA analysis. The scale is based on Nei and Li coefficients of similarity.

The RAPD pattern technique permitted discrimination all reference isolates. The results show that RAPD method can reliably distinguished between selected isolates. Therefore, the method is well suited for discriminatory identification. RAPD technique has been described as a useful technique for both identification and typing of wine *lactobacillus* strains (Du Plessis and Dicks, 1995; Rodas, *et al.*, 2005; Sohier, *et al.*, 1999; Bartowsky, *et al.*, 2003) and *Oenococcus oeni* (Zavaleta, *et al.*, 1997). RAPD analysis makes comparable the whole chromosomal DNA exhibiting bands of genotypically

specific value, and additionally this method turned out to be a fast, reliable, highly sensitive, and convenient method (Molnár, *et al.*, 1993). Thanos, *et al.*, (1996) proved that the numbers and sizes of amplification products were characteristic for each species.

**References:**

Ahmed, A. A. (2003). Biological study on some actinomycetes Isolates. PhD. Thesis. Faculty of Science, AL Azhar University, Cairo, Egypt.  
 Alexander, M. (1982). Soil Microbiology (2<sup>nd</sup>). John Wiley and Sons, Inc., New York. Pp.49-60

- Bartowsky, E. J., McCarthy, J. M., and Henschke, P. A. (2003). Differentiation of Australian wine isolates of *Oenococcus oeni* using random amplified polymorphic DNA (RAPD). *Aus. J. Grape Wine Res.* 9: 122-126.
- Basavaraj, N. K., Chandrashekhara, S., Ali, M., S., Prakash, S., G., and Fakirappa V., M., (2010). Isolation and Morphological Characterization of Antibiotic Producing Actinomycet. *Tropical Journal of Pharmaceutical Research* June 2010; 9 (3): 231-236 .
- Cavalcanti, M.A., Oliveira, L.G., Fernandes, M.J., and Lima, D.M. (2006). Filamentous fungi isolated from soil in districts of the Xingo' region, *Braz. Acta Bot. Bras.* 20, 831-837.m
- Chandrashekhara, S. (2010). Isolation and characterization of antibiotic production from soil isolates by fermentation. PhD. Thesis. Vinayaka Missions University Salem, Tamilnadu, India.
- Dice, L. R. (1945). Measures of the amount of ecologic association between species. *Ecology*, 26: 297-302.
- Du Plessis, E. M., and Dicks, L. M. T. (1995). Evaluation of random amplified polymorphic DNA (RAPD)-PCR as a method to differentiate *Lactobacillus acidophilus*, *Lactobacillus crispatus*, *Lactobacillus amylovorans*, *Lactobacillus gallinarum*, *Lactobacillus gasseri*, and *Lactobacillus johnsonii*. *Curr. Microbiol.*, 31: 114-118.
- Fitri, L., and Bustam, B. M. (2010). Screening of antimicrobial producing strains isolated from the soil of grassland rhizosphere in Pocu Meurah Intan Forest Park, Seulawah, Aceh Besar. *Biodiversitas* 11: 129-132.
- Gebreel, H. M., El-Mehalawy, A. A., El-Kholy, I. M., Rifaat, H. M., and Humid, A. A. (2008). Antimicrobial activities of certain bacteria isolated from Egyptian soil against pathogenic fungi. *J. Agric. Biol. Sci.* 4 , 331-339.
- Huda M. A. Sheikh, (2010). Antimicrobial activity of certain bacteria and fungi isolated from soil mixed with human saliva against pathogenic microbes causing dermatological diseases. *Saudi Journal of Biological Sciences* (2010) 17, 331-339.
- Magnusson, J., Stron, K., Rooa, R., Sjogren, J., and Schniirer, J. (2003). Broad and complex antifungal activity among environmental isolates of lactic acid bacteria. *FEMS Microbiol.* 219, 129-135.
- Molnár, K., Baska, F., Csaba, G., Glávits, R., and Székely, C. (1993). Pathological and histopathological studies of the swimbladder of eels *Anguilla anguilla* infected by *Anguillicola crassus* (Nematoda. Dracunculoidea). *Dis Aquat Org* 15: 41-50.
- Motta, A.S., Olivera, F.C., and Brandelli, A. (2004). Screening antimicrobial activity among bacteria from the Amazon Basin. *Braz. J. Microbiol.* 35, 307-310.
- Narisawa, T., Nanamiya, H., Kasai, K., Nozawa, A., and Yun CS, et al. (2008). Identification and functional analysis of novel (p)ppGpp synthetase genes in *Bacillus subtilis*. *Mol Microbiol* 67: 291-304.
- Nei, M. and Li, W. H. (1979). Mathematical model for studying ge-

- netic variation in terms of restriction endonucleases. Proc. Natl. Acad. Sci. USA., 76:5269-5273.
- Rodas, A. M., Ferrer, S., and Pardo, I. (2005). Polyphasic study of wine *Lactobacillus* strains: taxonomic implications. Int. J. Sys. Evol. Microbiol. 55: 197-207.
- Sabat, A.J., Budimir, A., Nashev, D., Sa-Leao, R., van Dijk, J. M., Laurent, F., Grundmann, H., and Friedrich, A. W. on behalf of the ESCMID Study Group of Epidemiological Markers (ES-GEM). (2013). Overview of molecular typing methods for outbreak detection and epidemiological surveillance. Euro Surveill ; 18(4): 20380.
- Sambrook, J., Fritschi, E. F., and Maniatis, T. (1989) Molecular cloning: a laboratory manual, Cold Spring Harbor Laboratory Press, New York.
- Singh, V., Praveen, V., Banga, J., and Tripathi, C.K. (2009). Antimicrobial activities of microbial strains isolated from soil of stressed ecological niches of Eastern Uttar Pradesh. Ind. J. Exp. Biol. India 47 (4), 298-303.
- Sohier, D., Coulon, J., and Lonvaud-Funel, A. (1999). Molecular identification of *Lactobacillus hilgardii* and genetic relatedness with *Lactobacillus brevis*. Int. J. Syst. Bacteriol. 49: 1075-1081.
- Thanos, M., Schonian, G., Meyer, W., Schweynoch, C., Graser, Y., Mitchell, T. G., Presber, W., and Tietz, H. J. (1996). Rapid identification of *Candida* species by DNA fingerprinting with PCR. J. Clin. Microbiol. 34, 615-621.
- Williams, J. G. K., Kube l~kA, R., Livak, K. J., Rafalski, J. A., and Tingey, S. V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers Nucl Acids Res. 18: 6531-6535.
- Yilmaz, M., Soran, H., and Beyatli, Y. (2005). Antimicrobial activities of some *Bacillus spp.* strain isolated from the soil. Microbiology 161, 127-131.
- Zavaleta, A.I., Martinez-Murcia, A.J., and Rodriguez-Valera, F. (1997). Intraspecific genetic diversity of *Oenococcus oeni* as derived from DNA fingerprinting and sequence analyses. Appl. Environ. Microbiol. 63: 1261-1267.

تحليل العلاقة الوراثية للبكتيريا المنتجة للمضاد الحيوى المعزولة من التربة الملوثة بالفضلات  
الآدمية باستخدام تقنية الـ DNA ( RAPD )  
دعاء حسن<sup>1</sup>، نبيل عبدالفتاح محمد<sup>1</sup>، فتحي محمد صالح<sup>1</sup>، عبيد أحمد محمد<sup>2</sup>  
<sup>1</sup> قسم الوراثة - كلية الزراعة - جامعة أسيوط  
<sup>2</sup> قسم النبات الزراعي ( وراثته ) - كلية الزراعة - جامعة جنوب الوادي

يعتبر المضاد الحيوى الذى تنتجه الكائنات الدقيقة المختلفة (مثل البكتيريا والفطريات)-  
أحد أهم النواتج الأيضية الثانوية التى تستخدم على نطاق تجارى ، والتعرف المتامى تحت  
ضغط الحاجه كموا د جديدة مضادة للميكروبات لمعالجة الأمراض المعدية يتطلب إكتشاف  
سلالات بكتيرية جديدة منتجة للمضادات الحيوية.

وقد أستهدف البحث الحالى عزل بعض السلالات من البكتيريا المنتجة للمضادات الحيوية  
من التربة الملوثة بالمخلفات الآدمية والتفرقة بين هذه السلالات بإستخدام تحليل RAPD-PCR.  
فى هذه الدراسة، تم جمع مائة وعشرة سلالات من العزلات البكتيرية من مناطق جغرافية  
مختلفة فى مصر من التربة الملوثة بالمخلفات الآدمية . وأظهرت الخمسة عزلات المختارة  
(D110، D3، D103، D106 و D104) نتائج واعدة لإنتاج مضاد حيوي فعال من خلال  
اختبارات حساسية هذه العزلات مع سلالتين من البكتيريا الموجبة لصبغة جرام و سلالتين من  
البكتيريا السالبة لصبغة جرام من خلال تكوين منطقة رائقة نتيجة تثبيط النمو بها . وقد أظهرت  
النتائج أن متوسط إتساع هذه المناطق فى حالة إستخدام البكتيريا الموجبة لجرام أكبر مما هو فى  
حالة إستخدام البكتيريا السالبة لجرام مما يدل على أن العزلات أكثر قدرة على تثبيط نمو  
البكتيريا الموجبة جرام بالمقارنة مع تثبيطها لنمو البكتيريا السالبة جرام.

بالإضافة إلى ماسبق ، فقد أمكن أن نفرق بين هذه السلالات الخمسة المعزولة - على  
المستوى الجزيئى باستخدام تحليل RAPD-PCR ، وقد أظهرت النتائج أن هذه العزلات يمكن  
وصفها طبقا لأنماط أشكالهم الحزمية ، وقد أعيد تجميعها فى أربع مجموعات على أساس  
تشابهاتها الوراثية المعينة . وقد مثلت المجموعات الأولى والثانية العزلات D106 و D103  
على الترتيب ، بينما ضمت المجموعة الثالثة اثنين من العزلات هما D104 و D110 ، أما  
المجموعة الرابعة فقد تمثلت فى السلالة D3.