

USING SAFETY ENVIRONMENT ESSENTIAL OIL TO IMPROVE QUALITY AND STORAGE ABILITY FOR GRAPES

Mohammed H. Hatem¹ - Abd el-Aal Z. Taieb²- Ibrahim A. Shalaby³ -
Raghda Mohammed Attia⁴

ABSTRACT

Table grapes are one of the oldest fruit known their cultivation in Egypt. Gray mold caused by *Botrytis cinerea* the most important postharvest disease of table grapes. The fungus grows vigorously on harvested grapes and can spread among berries even at low temperatures (-0.5°C). The most common commercial method to control decay is the use of SO_2 during cold storage. The SO_2 is harmful to human health and environmental. The main objective is found alternative of Sulfur dioxide pad. In this study used two varieties of grapes: - 1) Sugaone Seedless: It is a white seedless, and 2) Crimson Seedless: It is a red seedless. With threetreatments. 1) Were used LifeSpan (MAP) and essential oil pad. 2) Were used LifeSpan and essential oil was sprayed before harvest with 24 hour .3) were used LifeSpan as a kind of Modified Atmosphere Packing only (control treatment). Measurements the properties which the quality of grapes depend on its like Total Soluble Solids (T.S.S %), total acidity, weight losses, shatter percentage, brown and rots percentage and diameter .storage grapes at Maintaining a low temperature is a primary consideration in securing cooling of grapes it's 1 ± 1 . However, maintaining a high relative humidity of 95% and above during the storage is very important to minimize moisture loss and keep the stems in good condition and green. From this experiment, the LifeSpan and Bail oil was the best treatment with Sugaone (white grapes) and Crimson (red grapes). It got storage period until 60 days with very good quality.

Keyword: -Grape- Modified Atmosphere Packing-essential oil - Quality.

¹Professor, Ag. Eng. Dept. Faculty of Agriculture – Cairo University

²Associate Professor, Ag. Eng. Dept. Faculty of Agriculture – Cairo University

³Associate Professor, Medicinal and Aromatic plants, Dept.

⁴Agricultural Engineer-Horticulture Research Institute- Agricultural Research Center

INTRODUCTION

Table grape is one of the fruit with the highest input of technologies but even with an intense hand, labour activity and the price on the market is one of the higher in the fruit sector. Handling and packing (labour cost and packing materials) weight upon 25-30%; the cost of refrigerated storage condition engraves for 10% per month. Costa *et al.* (2011) reported that another relevant aspect to be taken into great account for maintaining the quality of horticultural commodities is the choice of the appropriate packaging system. Different headspace conditions can be achieved in the package depending on the interactions between respiratory activity of the packaged produce and gas transfer through the polymeric matrix.

❖ **Modified Atmosphere Packaging**

McMillin (2008) defined modified atmosphere packaging (MAP) is the removal and/or replacement of the atmosphere surrounding the product before sealing in vapor-barrier materials. Jobling (2001) reported the main aim of MAP is to change the composition of the atmosphere around the product so that the storage life of the product can be extended.

❖ **Gray mold**

Gray mold caused by *Botrytis cinerea* the most important postharvest disease of table grapes. In general, *Botrytis* is an important problem to fruit and vegetables in cold storage and subsequent shipment, because the fungus is able to grow effectively at temperatures just above freezing (Mustafa *et al.*, 2009). Artés-Hernández *et al.* (2006) reported that the most common commercial method to control decay is the use of SO₂ during cold storage, either by fumigation or generators. Meng *et al.* (2010) reported that the SO₂ application usually causes injury to fruits, and is harmful to human health. Moreover, due to increased public concern on food safety, SO₂ as a fungicide is limited to use for grape storage. For those reasons alternative techniques to SO₂ for preserving postharvest table grape quality are being currently investigated.

❖ **Essential oils**

Essential oils are volatile, natural, complex compounds characterized by a strong odour and are formed by aromatic plants as secondary metabolites (Bakkali *et al.*, 2008). The essential oils are thought to play a

role in the plant defence mechanism against phytopathogenic microorganisms (Mihaliak *et al.*, 1991). Most of the essential oils have been reported to inhibit post-harvest fungi in in vitro conditions (Hidalgo *et al.*, 2002).

The main objective is to study modified atmosphere packaging and found alternative of Sulfur dioxide pad to maintain the environment during storage at low temperature and a high relative humidity of 95% and above.

MATERIALS AND METHODS

1) Materials

a) Grapes

Two types of grapes (*Vitisvinifera L.*) were investigated:

1) Sugraone Seedless

It is a white seedless variety with exceptional large berries for an early, seedless variety. It has a good shelf life. The berries have a slight muscatflavour when fully matured.

2) CrimsonSeedless

It is a red, mid- to late season seedless variety. It normally ripens one week later Sunred Seedless. Crimson tends to ripen well on the vine, extending the season for seedless grapes. The bunches are attractive, filled with slightly elongated berries of maroon-red color. Flame has a melting crispy flavour and sweet taste.

b) Packs

1. LifeSpan (Modified Atmosphere Packing)

This bag is designed for 9kg of Seedless Grape-Special for best performance. Store at 0 °C. Finger 1 shows the work methods of LifeSpan.

Warning: - bag must be opened on removal from cold chain.

Important: liner must be used as stated in packing information sheet supplied. Information supplied as guidance only. Customer must determine suitability for use.

Bag Code: - L612.

Batch No.: D161109.




Fig.1. LifeSpan work method

2. Source of essential oils

The essential oil was isolated by hydro-distillation through Clevengers apparatus. Fresh plant parts (leaves or Flower or herb) were cut into small pieces and then thoroughly washed with sterilized water. The plant material was then placed in the round-bottom flask of the Clevengers apparatus. The ratio between the plant material and water in the flask was maintained as 1:3. Water was heated to produce steam that carried the most volatile fractions of the aromatic material with it. The steam was then chilled (in a condenser) and the resulting distillate was collected. The essential oil was found to float on the top of the hydrosol (the distilled water component) and was separated off. The extracted oils were dehydrated by the addition of anhydrous sodium sulphate, followed by thorough shaking and standing for 6–8 h and filtration (Tripathi *et al.*, 2008)

2) Packing and alternative of Sulfur dioxide pad experiment

This experiment was executed in PICO farm -Badr Center -Tahrir Directorate at season 2012. This experiment was conducted as follow:

A. First Treatment

Storage two varieties of grapes seedless one is Sugraone Seedless which whiten colour, and another is Crimson Seedless which red colour. It is storage on $1\pm 1^{\circ}\text{C}$ and 95% humidity, using LifeSpan as a kind of Modified Atmosphere Packing and essential oil Pad.

B. Second Treatment

Storage two varieties of grapes seedless one is Sugraone Seedless which whiten colour, and another is Crimson Seedless which red colour. It is storage on $1\pm 1^{\circ}\text{C}$ and 95% Humidity, using LifeSpan as a kind of Modified Atmosphere Packing and essential oil was sprayed before harvest with 24 hour .

C. Third Treatment

Storage two varieties of grapes seedless one is Sugraone Seedless which whiten colour, and another is Crimson Seedless which red colour. It is storage on $1\pm 1^{\circ}\text{C}$ and 95% Humidity, using LifeSpan as a kind of Modified Atmosphere Packing only.

3) Methods of Analyses

a. Chemical Properties

1) Total Soluble Solids (T.S.S %)

A hand refractometer was used to determine the total soluble solids percentage in fruit juice.

2) Total Acidity

Total acidity was determined in terms of anhydrous malic acid percentage after titration against 0.1 N Sodium hydroxide using Phenolphthaline as an indicator (A.O.A.C., 1995)

$$\text{Acidity} = \frac{\text{Average Titration}}{\text{The volume of grapes juice}} \times \text{Titration against NaOH} \times \text{Tartaric acid Strength}$$

Where:

The volume of grapes juice is 5 ml

Titration against NaOH is 0.1

Tartaric acid strength is 75.

3) Sugar: Acid ratio

The sugar: acid ratio was determined according to the following equation:

$$\text{Sugar : Acid ratio} = \frac{\text{Total Soluble Solids}}{\text{Total Acidity}}$$

b. Physical Properties

1) Percentage of Weight Losses (%)

The percentage weight loss was determined according to the following equation:

$$\text{Weight losses \%} = \frac{W_0 - W_t}{W_0} \times 100\%$$

Where:

W_0 : Weight bunch of grapes at zero time (g).

W_t : Weight bunch of grapes at (T) time (g).

The sample weight was determined by means of a digital precision balance (± 0.1 g) (Gibertini Europe, Italy). At each sampling time, three replicates were made.

2) Percentage of Shatter (%)

The percentage of shatter was determined according to the following expression

$$\text{Shatter \%} = \frac{S_t}{W_0} \times 100\%$$

Where

W_0 : Weight bunch of grapes at zero time (g).

S_t : Weight shatter grapes berries at (T) time (g).

3) Percentage of Brown (%)

The percentage of brown was determined according to the following expression

$$\text{Brown \%} = \frac{B_t}{W_0} \times 100\%$$

Where:

W_0 : Weight bunch of grapes at zero time (g).

B_t : Weight brown grapes berries at (T) time (g).

4) Percentage of Rot (%)

The percentage of rot was determined according to the following expression

$$\text{Rot \%} = \frac{R_t}{W_0} \times 100\%$$

Where:

W_0 : Weight bunch of grapes at zero time (g).

R_t : Weight rot grapes berries at (T) time (g).

RESULTS AND DISCUSSION

1. Specifications exportbunch

- a. T.S.S % : not less than 16:17 %
- b. Berry Size: not less than 17 mm.
- c. Bunch Weight is 750:350gm.

- d. Bunch disjointed.
- e. The bunch length is 15:13 cm.
- f. Similar berries within the bunch.
- g. Free from insect disease (trips and spider) and fungal diseases (molds).

2. Analysis of essential oils

The main components of Basil oil are shown in table 1 and Fig. 2 shows the Gas Chromatography of Basil essential oil.

Table 1. The main components of Basil oil

Peak Number	components	Area %	Height %
3	β -Pinene	8.61	16.92
5	1.8 Cineol	33.63	24.60
7	Linalool	27.81	21.10
17	β - Caryophyllene	6.47	7.83




Fig.2. Gas Chromatography of Basil essential oil

3. *Effect of LifeSpan, LifeSpan and spray basil oil before 24 h of harvest and LifeSpan and basil oil pad on chemical and physical properties during storage Sugaone and crimson grapes.*

a. Total Soluble Solids (T.S.S %)

Generally T.S.S % in Sugaone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as shown in Fig.3 and 4.




Fig.3. The relationship between T.S.S% and time during storage period of Sugraone grapes using LifeSpan and basil oil treatment.




Fig.4. The relationship between T.S.S% and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

T.S.S % in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping T.S.S% of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested

b. Acidity

Generally acidity in Sugaone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as show in Fig.5 and 6.




Fig.5. The relationship between acidity and time during storage period of Sugaone grapes using LifeSpan and basil oil treatment.




Fig.6. The relationship between acidity and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The acidity in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase

with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping acidity of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

c. Sugar: Acid ratio

Generally sugar: acid ratio in Sugraone and Crimson at $1\pm 1\text{ }^{\circ}\text{C}$ and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was decreased as show in Fig.7 and 8.




Fig.7. The relationship between sugars: acid ratio and time during storage period of Sugraone grapes using LifeSpan and basil oil treatment.




Fig.8. The relationship between sugars: acid ratio and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The sugar: acid ratio in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping sugar: acid ratio of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

d. Weight losses percentage %

Generally weight losses% in Sugaone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as show in Fig.9 and 10.




Fig.9. The relationship between weight losses% and time during storage period of Sugaone grapes using LifeSpan and basil oil treatment.




Fig.10. The relationship between weight losses% and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The weight losses% in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the

with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping acidity of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

c. Sugar: Acid ratio

Generally sugar: acid ratio in Sugraone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was decreased as show in Fig.7 and 8.




Fig.7. The relationship between sugars: acid ratio and time during storage period of Sugraone grapes using LifeSpan and basil oil treatment.




Fig.8. The relationship between sugars: acid ratio and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The sugar: acid ratio in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping sugar: acid ratio of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

d. Weight losses percentage %

Generally weight losses% in Sugaone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as show in Fig.9 and 10.




Fig.9. The relationship between weight losses% and time during storage period of Sugaone grapes using LifeSpan and basil oil treatment.




Fig.10. The relationship between weight losses% and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The weight losses% in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the

increase with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping weight losses% of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

e. Shatter percentage %

Generally shatter% in Sugaone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as show in Fig. 11 and 12.




Fig. 11. The relationship between shatter % and time during storage period of Sugaone grapes using LifeSpan and basil oil treatment.




Fig.12. The relationship between shatter % and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The shatter in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase

with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping shatter % of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

f. Rot percentage %

Generally rot% in Sugaone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as show in Fig.13 and 14.




Fig.13. The relationship between rot% and time during storage period of Sugaone grapes using LifeSpan and basil oil treatment.




Fig.14. The relationship between rot% and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The rotting in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase

with control treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping rot% of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

g. Brown percentage %

Generally brown% in Sugraone and Crimson at 1 ± 1 °C and 95 % Humidity during storage in three treatments of used basil oil and LifeSpan was increased as show in Fig.15 and 16.




Fig.15. The relationship between brown% and time during storage period of Sugarone grapes using LifeSpan and basil oil treatment.




Fig.16. The relationship between brown% and time during storage period of Crimson grapes using LifeSpan and basil oil treatment.

The brown in all the tested treatments was increased during storage period. The increases with basil oil treatments were less than the increase with control

treatment. Basil oil pad treatment was the best treatment with two varieties of grapes.

The superior treatments which keeping brown % of grapes fruits was basil oil pad treatment (4000 ppm) and LifeSpan compared with the other tested.

CONCLUSION

From this experiment, the LifeSpan and Basil oil were the best treatment with Sugraone (white grapes) and Crimson (red grapes). It got storage period until 60 days with very good quality.

REFERENCES

- A.O.A.C. (1995). Official and tentative methods of analysis of the Association of Official Agricultural Chemists. pp.158, 8th ed. Washington D.C. .
- Artés-Hernández, F.; Tomás-Barberán, F.A. and Artés, F. (2006). Modified atmosphere packaging preserves quality of SO₂-free 'Superior seedless' table grapes. *Postharvest Biology and Technology*, 39(2), 146-154.
- Bakkali, F.; Averbeck, S.; Averbeck, D. and Idaomar, M. (2008). Biological effects of essential oils – A review. *Food and Chemical Toxicology*, 46(2), 446-475.
- Costa, C.; Lucera, A.; Conte, A.; Mastromatteo, M.; Speranza, B.; Antonacci, A. and Del-Nobile, M.A. (2011). Effects of passive and active modified atmosphere packaging conditions on ready-to-eat table grape. *Journal of Food Engineering*, 102(2), 115-121.
- Hidalgo, P.J.; Uebera, J.L.; Santos, J.A.; LaFont, F.; Castelanos, C.; Pálomino, A. and Roman, M. (2002). Essential oils in *Culamintha sylvatica*. *Bromf. ssp. ascedens* (Jordan) P.W. Ball wild and cultivated productions and antifungal activity. *J Essential Oil Res* 14:68–71.
- Jobling, J. (2001). Modified atmosphere packaging: Not as simple as it seems. Sydney Postharvest Laboratory Information Sheet. *Good Fruit and Vegetables Magazine* 11(5).
- McMillin, K.W. (2008). Review where is MAP Going? A review and future potential of modified atmosphere packaging for meat. *Meat Science* 80 (2008) 43–65(www.elsevier.com/locate/meatsci).
- Meng, Xiang-Hong; Qin, Guo-Zheng and Tian, Shi-Ping. (2010). Influences of preharvest spraying *Cryptococcus laurentii* combined with postharvest chitosan coating on postharvest diseases and quality of table grapes in storage. *LWT - Food Science and Technology*, 43(4), 596-601.
- Mihaliak, C.A.; Gershenzo, J. and Croteau, R. (1991). Lack of rapid monoterpene turnover in rooted plants, implications for theories of plant chemical defense. *Oecologia* 87:373–376.

- Mustafa, C.; Kalpulov, T.; Zutahy, Y.; Ish-shalom, S.; Lurie, S. and Lichter, A. (2009). Quantitative and qualitative analysis of Botrytis inoculated on table grapes by qPCR and antibodies. *Postharvest Biology and Technology* 52 (2009) 235–239.
- Tripathi, P.; Dubey, N. K. and Shukla, A. K. (2008). Use of some essential oils as post-harvest botanical fungicides in the management of grey mould of grapes caused by *Botrytis cinerea*. *World J Microbiol Biotechnol*(DOI 10.1007/s11274-007-9435-2), (2008) 2024:2039–2046.

المخلص العربي

استخدام الزيوت الطبية لتحسين الجوده والقدرة التخزينية للعنب

محمد هاشم حاتم^١ – عبد العال زكى تايب^١ - ابراهيم أحمد شلبي^٢ - رغه محمد عطية^٣

يعد العنب من السلع القابلة للتلف جدا، لذلك نجد ان عنب المائدة يصاب بمشاكل حادة. يتعرض العنب للعدوى لعدد من امراض ما بعد الحصاد. تشكل امراض ما بعد الحصاد خطورة على عنقايد العنب في جميع أنحاء العالم، حيث يسبب خسارة عظيمة ويؤثر أيضا جودة العنب. وخصوصا خلال الحصاد والتعبئة والتصدير والتخزين والتسويق. يعتبر الفطر البوتريتس المسبب لمرض العفن الرمادى واحدة من أهم الفطريات التي تهاجم العنب خلال المراحل المختلفة التي يمر بها العنب. وخصوصا عندما يتم شحنها في درجة حرارة تتراوح من صفر إلى ١ م. ويعد استخدام بطانات ثنائي أكسيد الكبريت (SO₂ Pads) أثناء التخزين الباردة هو الأسلوب الأكثر شيوعا تجاريا للسيطرة على العفن الرمادى. وفي الواقع، تشكل بقايا ثنائي أكسيد الكبريت خطرا على المستهلكين وكذلك على البيئة.

ولذلك، فإن الهدف الرئيسي هو دراسة أساليب التعبئة المستخدمة وإيجاد مواد طبيعية بديلة للبطانات ثنائي أكسيد الكبريت للمحافظة على البيئة أثناء التخزين في درجة حرارة منخفضة ورطوبة عالية نسبيا تصل إلى ٩٥%. وفي هذه الدراسة تم تخزين صنفى من العنب صنف Sugraone وهذا صنف ابيض اللون و صنف Crimson وهو صنف احمر اللون على درجة حرارة ± 1 م و رطوبة تصل إلى ٩٥% مع:

- ١- التعبئة فى اللايف اسبان LifeSpan (معاملة الكنترول)
- ٢- تعبئة العنب فى اللايف اسبان بعد رشه بزيت الريحان قبل حصاده ب ٢٤ ساعة.
- ٣- تعبئة العنب فى اللايف اسبان مع عمل بطانات من زيت الريحان (Basil oil Pad) بتركيز ٤٠٠٠ جزء فى المليون.

وتم قياس مجموعه من العوامل التي تحدد عليها جوده العنب وهى نسبة المواد الذائبة، نسبة الحموضة، نسبة الفقد فى الوزن، نسبة الفطر وكذلك نسبة الاغقان وتلون الثمار باللون البنى.

ومن هذه التجربة تم التوصل إلى ان تعبئة العنب فى اللايف اسبان مع بطانات زيت الريحان عند تركيز ٤٠٠٠ جزء فى المليون مع صنفى العنب (الأبيض و الأحمر) اعطيت افضل نتائج وتم الحصول على جوده عاليه لصنفى العنب لمدته تصل إلى ٦٠ يوما.

^١ استاذ الهندسة الزراعية – كلية الزراعة – جامعة القاهرة.

^٢ استاذ مساعد الهندسة الزراعية – كلية الزراعة – جامعة القاهرة.

^٣ استاذ مساعد النباتات الطبية والعطرية – معهد بحوث البساتين – مركز البحوث الزراعية.

^٤ مهندسة زراعية – معهد بحوث البساتين - مركز البحوث الزراعية.