

SEROLOGICAL STUDY ON BRUCELLOSIS IN CATTLE SUFFERING FROM REPRODUCTIVE DISORDERS IN COMPARISON WITH APPARENTLY HEALTH CATTLE IN ASSIUT GOVERNORATE

A.M. KORIEM; S.M. EL BERBAWY and H.A. ABD EL-KADER

Animal Health Research Institute - Assiut Provincial Lab.

Email: ayman_korciem@yahoo.com

ABSTRACT

Received at: 4/5/2014

Accepted: 17/7/2014

A total 1534 blood samples (348 from cattle with history of reproductive disorders include 48 abortion, 56 retained placenta & 244 repeat breeder and 1186 from cattle apparently health) were collected from different localities in Assiut Governorate, was tested for the detection of antibodies against *Brucella spp.* The results obtained by Buffer acidified plate antigen test (BAPAT) and Rose bengal test (RBT) as screening tests revealed a positive reactors percentage 4.02 & 0.51 % respectively. The *Brucella* positive reactors were subjected to confirmation by Tube agglutination test (TAT) and Rivanol test (Riv.T). The incidence of brucellosis was higher in cattle suffering from reproductive disorders (3.45%) in comparison with apparently health cattle (0.51%). The incidence of brucellosis according to type of reproductive disorders, (abortion, retained placenta & repeat breeder) were 8.33, 8.9 & 1.2 % respectively, this incidence reflect in addition to brucellosis, some other agents were involved in reproductive disorders.

Key words: Serological Study, Brucellosis, Cattle, Reproductive Disorders.

INTRODUCTION

Brucellosis is a chronic zoonotic disease caused by *Brucella species* which are Gram-negative and facultative anaerobic non-motile intracellular bacteria, with *Brucella abortus* being the principal cause of brucellosis in cattle (Cutler & Cutler, 2006).

Brucellosis in cattle is usually caused by biovars of *Brucella abortus*. In some countries, particularly in southern Europe and western Asia, where cattle are kept in close association with sheep or goats, infection can also be caused by *B. melitensis* (Verger, 1985 & Jimenez de Bagues *et al.*, 1991). Occasionally, *B. suis* may cause a chronic infection in the mammary gland of cattle, but it has not been reported to cause abortion or spread to other animals (Ewalt *et al.*, 1997).

Bovine brucellosis is usually caused by *Brucella abortus*, less frequently by *Brucella melitensis* and rarely by *Brucella suis*. It is characterized by abortion, with excretion of the organisms in uterine discharge and milk. Major economic losses result from abortion. Loss of calves, reduced milk yield in females and infertility in males (WHO, 1971).

Bovine brucellosis, caused by *Brucella abortus*, is a serious zoonotic disease manifested by reproductive disorders such as abortions, infertility, retention of placenta, stillbirth and calf loss in animals, and results in huge economic losses to dairy farmers (Radostits *et al.*, 2000 & Singh *et al.*, 2002).

Brucellosis is reported as a cause of abortion, retention of placenta, repeat breeding, infertility and prolonged intercalving period due to early embryonic deaths (Roberts, 1999).

Brucellosis is a highly contagious, zoonotic, and economically important bacterial disease of animals worldwide (OIE, 2000). It causes significant economic losses including abortion, loss in milk production, low fertility rates, and cost of replacement of animals (McDermott & Arimi, 2002).

Brucellosis is a widespread and highly infectious zoonosis disease, with a great economical impact on cattle farming. The infection is caused by *Brucella abortus*, and it is characterized by abortion, metritis, orchitis and epididymitis. The disease in cattle mainly affects the reproductive organs of infected animals (England *et al.*, 2004).

Cattle Brucellosis is usually caused by *Brucella abortus* biovars and occasionally by *Brucella melitensis* (OIE, 2008). In Egypt, starting from 1998 *Brucella melitensis* was reported to be the common strain isolated from cattle as reported by Shalaby *et al.* (2003); Sayour (2004) and Shehata (2004). The disease is mainly characterized by abortion, stillbirths or weak calves and lactating cows may show decrease in milk yield (Matope *et al.*, 2010).

BAPAT and RBPT serological tests revealed the highest rate of sensitivity that guide us to use these tests as screening tests on animals brucellosis. Riv test showing the highest rate of specificity that bearing in mind the BAPAT and RBT positive samples should be confirmed by this test (Montasser *et al.*, 2011).

The prevalence of brucellosis was higher in animals suffering from reproductive troubles (34.52%) than among animals apparently health (6.07%) (Ammar *et al.*, 2005).

So the objectives of the present investigation was intended to throw light on the incidence of brucellosis in cattle suffering from reproductive

disorders in comparison with apparently health cattle.

MATERIALS and METHODS

► Samples collection :

A total 1534 blood samples (348 from cattle with history of reproductive disorders include, 48 abortion, 56 retained placenta & 244 repeat breeder, and 1186 from cattle apparently health) were collected under strict hygienic condition from different localities. Blood samples were sent to the laboratory as soon as possible, were allowed to clot and the sera were obtained by centrifugation and stored at -20 C° until performing serological tests.

► Serological Examination:

All used four antigens were obtained from Veterinary Serum and Vaccine Research Institute - Abbasia, Cairo - Egypt.

All sera were screened for antibodies against *Brucella* by BAPAT as described by Angus & Barton (1984), and RBT as described by Alton *et al.* (1988). All positive serum samples were further retested by TAT and Riv T as quantitative confirmatory tests described by Alton *et al.* (1988).

RESULTS

All data obtained through the investigation were illustrated in Tables 1-3.

Table 1: Incidence of brucellosis in cattle suffering from reproductive disorders in comparison with apparently healthy cattle.

Reproductive status of examined cattle	Total number	RBT		BAPAT		TAT		Riv.T	
		+ve	%	+ve	%	+ve	%	+ve	%
Cattle apparently healthy	1186	6	0.51	6	0.51	6	0.51	6	0.51
Cattle with history of reproductive disorders	348	14	4.02	14	4.02	12	3.45	12	3.45

Table 2: Incidence of brucellosis in cattle according to type of reproductive disorders.

Type of reproductive disorders	No of examined samples	Positive seroreactive samples	
		No	%
Abortion	48	4	8.33
Retained placenta	56	5	8.9
Repeat breeder	244	3	1.2
Total	348	12	3.45

Table 3: Different serological reactions for seroreactive samples.

Reproductive status of examined cattle	Total Reactors	Serological tests														
		TAT									Rivanol					
		RBPT			BAPAT			Total			1/25	1/50	1/100	1/200	1/400	Total
		1/10	1/20	1/40	1/80	1/160	1/320	Total	1/25	1/50	1/100	1/200	1/400	Total		
Cattle Apparently Healthy	6	6	6	-	-	-	2	2	2	6	1	2	-	1	2	6
Cattle with history of reproductive disorders	14	14	14	-	2	2	2	3	5	14	1	3	1	2	5	12

DISCUSSION

Brucellosis is still a major problem in several countries including Egypt. The disease is one of the most serious disease reducing animals productivity and reproductivity and also constitute a public health hazard (Benkirane, 1997).

Surveillance of brucellosis in animals is of great importance and considered the angle stone for better understanding of the disease and improving the control and eradication policy (Fosgate *et al.*, 2003).

In the present study, the obtained results revealed that BAPAT and RBT showed high rate of sensitivity as screening tests where no missed positive cases, these findings agreed with El-Gibaly (1993) and Montasser *et al.* (2011), while Angus & Barton (1984) and Gall & Nielsen (2004) showed that BAPAT was more sensitive and accurate than the other conventional tests for detection of brucella in bovine serum. This was attributed in part to the instability of some of the antigen preparations used in the other tests and also may attributed to the pH(4) that could overcome the problem of non specific reactions, moreover Shalaby *et al.* (2003) said that BAPAT is a superior test and can detect low titres missed by TAT. The confirmation with Rivanol test is also recommended due to its high specificity and reliability in detecting the infected cases without serious number of false positive (Huber & Nicolletti, 1986).

The number of the positive cases to Rivanol test (Table 3) seemed to be less than that of those on both RBPT, BAPAT and TAT. These results are in agreement with the results reported by Shalaby *et al.* (2003), it may be due to the precipitating activity of the Rivanol solution on the IgM as recorded by Pietz and Cowart (1980).

Our findings revealed also that the dilution at which the Rivanol test gave positive results varied from

1/25 up to 1/400 and this indicates true positive cases because the test determines the agglutinating activity of the IgG after destroying the IgM (Alton *et al.*, 1988). These findings are in agreement with (Shalaby *et al.*, 2003).

The obtained result as shown in Table (1) revealed the prevalence of brucellosis in apparently health cattle was (0.51 %). This result agrees with that obtained by Abd El-Hafeez *et al.* (2001) (0.53 %) and lower than the result recorded by Seddek (1999) (2.81%). The variation in the prevalence of brucellosis may be due to animal population, their susceptibility, vaccination status and the hygienic measures applied.

Correlation between the rate of brucella infection with the reproductive status of the examined animals as shown in Table (1) revealed that the rate of brucella infection was higher among cattle with a history of reproductive disorders (3.45 %) than those among cattle apparently health without history of reproductive disorders (0.51 %). These findings are in agreement with the results recorded by Hafez (1980); EL-Gamel (2004) and Ammar *et al.* (2005) who reported that abortion, retained placenta and infertility were the principle guide accompanying infection with brucellosis. On the other hand, animals which were seropositive and had no history of reproductive disorders constitute the major sources of infection and control failure. Similar result was recorded by Njiro *et al.* (2011) they recorded 3.8 % prevalence of brucellosis in cattle with history of reproductive disorders.

According to the type of reproductive disorders, the obtained results as showed in Table (2) revealed that the incidence of brucellosis relative to abortion was 8.33 %. This result agreed with Teferi Degefa *et al.* (2011) they found that the prevalence of abortion was 8.7 % and Amin *et al.* (2004) recorded 12 %. While lower incidence 3.2 % was reported by Ibrahim (2003). This difference in prevalence rate

may be due to the variation in cattle husbandry management system.

The gained result Table (2) declared that the incidence of retained placenta was 8.9 % .Similar results were also reported by Ahmed *et al.* (1992) and Amin *et al.* (2004) they found 7.14 & 7.41 % respectively. While higher incidence 16.8 &18.3 % were recorded by Gebre-Mariam (1996) and Teferi Degefa *et al.* (2011) respectively.

The finding of the present study (Table 2) revealed that low incidence of repeat breeder 1.2 %. Nearly similar finding were noticed by Ahmed *et al.* (1992) and Amin *et al.* (2004) they recorded 3.16 & 1.64 % prevalence of brucellosis respectively.

From these obtained results, we can conclude that *Brucella* infection still higher among cattle with history of reproductive disorders, which need more efforts and an effective plan for control and eradication of the disease. The high prevalence of abortion and retained placenta together with this low incidence of brucellosis may suggest the presence of other causes of reproductive disease in addition to *Brucella*.

REFERENCES

- Abdel-Hafeez, M.M.; AbdEl-Kader, H.A.; Ali, M.M.; Bastawrows, A.F. and Seddek, S.R. (2001): Zoonotic importance of brucellosis among farm animals and veterinary field employees at Assiut Governorate. Assiut Vet. Med. J., 44(88): 119-135.
- Ahmed, J.U.; Alam, M.G.S.; Rahman, M.M. and Hossain, M. (1992): Seroprevalence of brucellosis in idigenous zebu cattle of Bangladesh. Bangladesh J. Micrbiol., 9: 21-27.
- Alton, G.G.; Jones, L.M.; Angus, R.D. and Verger, J.M. (1988): Techniques for the brucellosis laboratory. Institut National de la Recherche Agronomique, Paris, 190 pp.
- Amin, K.M.R.; Rahman, M.B.; Kabir, S.M.L.; Sarkar, S.K. and Akand, M.S.I. (2004): Serological Epidemiology of Brucellosis in Cattle of Mymensingh Districts of Bangladesh. J. Anim. Vet. Adv.,3(11): 773-775.
- Ammar, K.M.; AL-Gaabary, M.H.; Osman, S.A. and Hegazy, Y.M.A. (2005): Some studies on brucellosis among ruminants in Gharbia Governorate. 8th Sci. Cong., Egyptian Society For Cattle Diseases, 11-13 Dec., Assiut, Egypt.
- Angus, R.D. and Barton, C.E. (1984): The production and evaluation of a buffered plate antigen for use in a presumptive test for brucellosis. In Proc. 3rd International Symposium on brucellosis, Algiers, Algeria. Dev. Biol. Standard., 56, 349-356.
- Benkirane, A. (1997): Analysis of the international animals health code for brucellosis (with special focus in north africa and the middle east). J. Egypt Vet. Med. Ass., 75 (1): 5-6.
- Cutler, S.J. and Cutler, R.R. (2006): Brucellosis, the most common bacterial zoonosis. The Biomedical Scientist, 50 (4): 336-341.
- EL-Gamel, A.M. (2004): Control trials of some occupational zoonotic diseases of Egyptian farmer. M.V.Sc. Thesis (Zoon), Fac. Vet. Med., Tanta University, Kafr El-Sheikh Branch.
- El-Gibaly, S.M. (1993): Correlation between serotests and isolation of *Brucella melitensis* in an infected sheep farm. Proc. 2 Sci. Cong. Society for cattle Disease, Assiut, 194-203.
- England, T.; Kelly, L.; Jones, R.D.; MacMillan, A. and Wooldridge, M. (2004): A simulation model of brucellosis spread in British cattle under several testing regimes. Prev. Vet. Med., 63: 63-73.
- Ewalt, D.R.; Payeur, J.B.; Rhyan, J.C. and Geer, P.L. (1997): *Brucella suis* biovar 1 in naturally infected cattle: a bacteriological, serological, and histological study. J. Vet. Diagn. Invest., 10: 417-420.
- Fosgate, G.T.; Adesiyun, A.A.; Hird, D.W.; Johnson, W.O.; Hietala, S.K.; Schurig, G.G. and Diptee, M.D. (2003): Evaluation of brucellosis RB51 vaccine for domestic water buffalo in Trinidad. Prev. Vet. Med., 58: 211-225.
- Gall, D. and Nielsen, K. (2004): Serological diagnosis of bovine brucellosis: a review of test performance and cost comparison. Rev. Sci. Tech. Off. Int. Epiz., 23 (3): 989-1002.
- Gebre-Mariam, T. (1996): Survey on prepartum and postpartum reproductive problems of dairy cattle in Mekele and its environments. DVM Thesis, FVM, AAU Debre Zeit, Ethiopia.
- Hafez, E.S.E. (1980): Reproduction in Farm Animals 3rd ed. Lea and Febger, Philadelphia, chapter IV: Reproductive failure, pp. 449-503.
- Huber, J.D. and Nicolletti, P. (1986): Comparison of the results of card, rivanol, complement fixation and milk ring test with the isolation rate of *Br. abortus* from cattle. Am. J. Vet. Res., 47:1529-1531.
- Ibrahim, O. (2003): Study in major reproductive health problems of smallholder dairy farms in and around Kombolcha. DVM Thesis, FVM, AAU Debre Zeit, Ethiopia.
- Jimenez de Bagues, M.P.; Marin, C. and Blasco, J.M. (1991): Effect of antibiotic therapy and strain 19 vaccination on the spread of *Brucella melitensis* within an infected dairy herd. Prev. Vet. Med., 11: 17-24.

- Matope, G.; Behabhe, E.; Muma, J.B.; Lund, A. and Skjerve, E. (2010): Risk factors for Brucella species infection in small holder household herds. *Epidemiology and Infection*, 139: 157-164.
- McDermott, J.J. and Arimi, S.M. (2002): Brucellosis in sub-Saharan Africa: Epidemiology, Control and Impact. *Veterinary Microbiology*, 90(1-4): 111-134.
- Montasser, A.M.; Affi, M.M.; EL-Bayoumy, E.M.; Abdul-Raouf, U.M. and Mohamad, H.A. (2011): Efficiency of serological tests for detection of brucellosis in ruminant at south provinces of Egypt. *Global Veterinaria* 6(2): 156-161.
- Njiro, S.M.; Kidanemariam, A.G.; Tstotesti, A.M.; Katsande, T.C.; Mnisi, M.; Lubisi, B.A.; Potts, A.D.; Baloyi, F.; Moyo, G.; Mpfu, J.; Kalake, A. and Williams, R. (2011): A study of some infectious causes of reproductive disorders in cattle owned by resource-poor farmers in Gauteng Province, South Africa. *J.S. Afr. Vet. Ass.*, 82(4): 213-218.
- OIE (2000): "Bovine brucellosis," in *Manual of Standards for Diagnostic Tests and Vaccines*, pp. 328-345, OIE, Paris, France, 4th edition.
- OIE (2008): *Manual of the diagnostic tests and vaccines for terrestrial animals*, Vol.1, 5th Edition (Office International Des Epizooties, Paris, France).
- Pietz, D.E. and Cowart, W.O. (1980): Use of Epidemiological data and serological tests in bovine brucellosis. *J. Am. Vet. Med. Ass.*, 177:1221-1226.
- Radostits, O.M.; Gay, C.C.; Blood, D.C. and Hinchcliff, K.W. (2000): *Veterinary Medicine: a textbook of the diseases of cattle, sheep, pigs, goats and horses*. 9th ed. W. B. Saunders, London, pp. 1877.
- Roberts, S.J. (1999): *Veterinary Obstetrics and Genital Diseases (Theriogenology)*. 2nd ed. CBS publishers and distributors, New Delhi, India, chapter XIII: Infertility in the cow, pp. 499.
- Sayour, A.E. (2004): The use of recent bacteriological techniques in the differentiation of Brucella group of microorganisms. Ph.D. Thesis, Microbiology, Fac. Vet. Med., Cairo University.
- Seddek, S.R. (1999): Serological studies on Brucella infection in cattle, sheep and goat. *Assiut Vet. Med. J.*, 42 (83): 216-227.
- Shalaby, M.N.; Ghobashy, H.M.; El-Bayomy, E.M. and Saleh, W.M. (2003): Prevalence of brucellosis among farm animal species in some governorates in Egypt. *Proceedings of the Seventh Scientific Congress, Egyptian Society for Cattle Diseases, Assiut, Egypt*, pp. 271-282.
- Shehata, A.E. (2004): Immunological studies on Brucella microorganisms in farm animals in Bani Suef governorate. M.V.Sc. Thesis (Microbiology), Fac. Vet. Med., Cairo University, Bani Suef Branch.
- Singh, G.; Sharma, D.R.; Sandhu, K.S. and Dhand, N.K. (2002): Economic losses occurring due to bovine abortions in Punjab. In: 10th International Congress of Asian-Australasian Association of Animal Production Societies, 23-27 September. Ashoka Hotel, New Delhi, India.
- Teferi Degefa; Asmamaw Duressa and Reta Duguma (2011): Brucellosis and Some Reproductive problems of Indigenous Arsi Cattle in Selected Arsi zones of Oromia Regional State, Ethiopia. *Global Veterinaria*, 7(1): 45-53.
- Verger, J.M. (1985): *B. melitensis* infection in cattle. In: *Brucella melitensis*, Plommet & Verger, eds. Martinus Nijhoff Publ., Dordrecht-Boston-Lancaster. 197-203.
- WHO (1971): Technical Report Series, Joint FAO/WHO Expert Committee in Brucellosis, 5th Report: No. 464.

دراسة سيروولوجية عن البروسيلة في أبقار تعاني من مشاكل تناسلية بالمقارنة مع أبقار سليمة ظاهريا في محافظة أسيوط

أيمن منير إبراهيم كتريم ، سعد محروس البرباوي ، حسين علي عبد القادر

Email: ayman_koriem@yahoo.com

نظرا لاهمية مرض البروسيلة من الناحية الصحية والتناسلية للحيوان وما يسببه من خسائر اقتصادية نتيجة الاجهاض واحتباس المشيمة وبعض المشاكل الاخرى فكان من الضروري اجراء هذه الدراسة لتحديد نسبة الاصابة بالبروسيلة في الابقار التي تعاني من مشاكل تناسلية ومقارنتها بنسب الاصابة في الابقار السليمة ظاهريا باستخدام الاختبارات السيروولوجية المختلفة. اجريت هذه الدراسة علي عدد 1034 من الابقار (248 بقرة تعاني من مشاكل تناسلية شملت 48 حالة اجهاض و 56 احتباس مشيمة و 244 شياع متكرر و 1186 بقرة سليمة ظاهريا) حيث تم جمع عينات الدم منها. تم فحص العينات سيروولوجيا للبروسيلة باختباري الـروزينجال والمحصن المخدم الشريحي وقد خضعت العينات الايجابية لهذين الاختبارين لاختباري التلازن الانبوبي البطي والريفانول. اظهرت النتائج ان نسبة الاصابة بالمرض كانت مرتفعة في الابقار التي تعاني من مشاكل تناسلية (3.45%) عنها في الابقار السليمة ظاهريا (0.51%) بينما كانت نسبة الاصابة في الحالات التي تعاني من مشاكل تناسلية (اجهاض واحتباس مشيمة وشياع متكرر) 8.33 و 1.2 و 8.9% على التوالي.