

EFFECT OF SOME WEED CONTROL PACKAGES ON SEED COTTON YIELD AND FIBER PROPERTIES OF SOME COTTON GENOTYPES (*GOSSYPIUM BARBADENSE*, L.) AND ITS ASSOCIATED WEEDS

SOLIMAN, I. E.¹, AZZA E. KHAFFAGY¹, A. M. GHALWASH¹ and AMAL S. ABD EL-AAL²

1. Weed Research Central Laboratory, ARC, Giza, Egypt.
2. Cotton Res. Institute, ARC, Giza, Egypt.

(Manuscript received 26 November 2013)

Abstract

Two field experiments were conducted during 2011 and 2012 seasons at Sakha Agricultural Research Station, to investigate the performance of three cotton genotypes (Giza 86, Giza 88 and the new promising hybrid Giza 89 x 86) under some weed control packages (pendimethalin, butralin, fluazifop-p-butyl and sethoxydim) plus one hand hoeing for each herbicide in addition to the hand hoeing twice for controlling total weeds and their effects on some vegetative characters, yield components and fiber properties of cotton. Results indicated that the promising hybrid Giza 89 x 86 decreased dry weight of broad-leaved, grassy and perennial weeds by 27.6, 31.9 and 26.8% respectively, at second survey, also recorded the highest values of yield and its components, as compared to the genotype Giza 86, followed by Giza 88 in both seasons. Moreover, all weed control packages significantly decreased weeds parameters and increased yield components in both seasons. Also, gave highly significant increase in seed cotton yield (Kantar /Fadden). The highest values were obtained with (pendimethalin, butralin) plus one hand hoeing treatments, followed by hand hoeing twice and (fluazifop-p-butyl, sethoxydim) plus one hand hoeing by 66.7, 62.3, 63.7, 56.5 and 46.4 % and increased micronaire reading by 34.0, 34.0, 25.2, 30.5 and 28.0% respectively, as compared to the control treatment. The interaction between cotton genotypes and weed control packages exerted a significant integrated impact on weeds characters, and this reflected on increases in seed cotton yield components and fiber quality. From results of correlation analysis, the dry weight of total weeds biomass were negative correlated with cotton yield and micronaire reading. These results indicated that under heavily infested soil with weeds, it is better to grow the promising hybrid Giza 89 x 86 and Giza 88 with the application of weed control packages such as (pendimethalin or butralin) plus one hand hoeing or hand hoeing twice. These practices gave the highest reduction in weeds density and increased cotton yield and its components and improved yield cotton and fiber quality.

INTRODUCTION

Cotton (*Gossypium barbadense*, L.) is an important crop in Egypt for local consumption of fibers and oils and exportation due its high as long staple fiber. In 2013 the cultivated area with cotton arrived 520000 fadden (Anonymous 2102). Growth and yield of cotton is substantially reduced by weed competition. Bukun

(2004), in Turkey found that weeds should be eliminated from 1-2 weeks up to 11-12 weeks of plant emergence and weed control strategies should be done in these periods by enhancing the herbicides use and other methods of weed management including cultivations. Cotton plants is a weak competitor for weeds due to its prolonged season especially in early growing periods. Ferrel *et al.* (2001) in India found that the infestation of weed flora in cotton crop reduced the yield by 1.28 and 1.6 ton /ha compared to 2.41 and 2.33 ton/ha from weed free cotton field, and the severity of weed competition depending on weed densities and their compositions. Several scientists have studied the influence of different weed species exist in cotton fields. In all cases, yield has been the most sensitive indicator of weed competition. (*Echinochola crus-galli*) can be a problem for irrigated cotton grown in the Western United States (Miller *et al.* 1961). Competition from 1 to 3 plants of (*Cyperus esculentus*) per three cotton plants reduced cotton dry weight, but yield was reduced most when cotton was stressed by low fertility or low soil moisture. Seed cotton yield in hand weeded control plots averaged 14% higher than in plots where the weed remained throughout the season (Keeley *et al.* 1973 and Keeley and Thullen 1975). Competition with (*Cyperus esculentus*) for 6 to 8 weeks reduced yield 20% and full season competition cut it 34%. They also mentioned that, micronaire readings were lower for cotton samples collected from plots that contained nutsedge for longer than 4 week than from samples collected from weed free plots indicating delayed crop maturity, also growth and yield and delayed fiber maturity. (*Xanthium pensylvanicum*) also is a very serious problem weed in cotton where at densities of 1 to 10 plants per 10 cotton plants reduced yield by more than 20% up to more than 80% (Buchanan and Burns, 1971).

Concerning the use of herbicides for weed control in cottons Several workers have shown that dinitroaniline herbicides e.g. pendimethalin were more effective in controlling summer weeds and need light hoeing as complements (Fayed *et al.* 1983, and Khan *et al.* 2001). They obtained highest seed cotton yield with application of pendimethalin. Moreover, Dilbaugh *et al.* (2009) indicated that application of pendimethalin 33% on dry bed furrow before applying irrigation produced 82.5 % broad leaf and 84.1 % narrow leaf control which ultimately led towards obtaining seed cotton yield of 2689 kg ha⁻¹ which was 115.1% higher than the weedy check. El-Maghraby (1971) indicated that hand hoeing increased the number of bolls/ plant, mean while, fiber strength, elongation percent and fiber fineness were not affected by using pre-emergence herbicides at different doses. Also Ghaly (1981) found that weed control treatments had a significant effect on seed cotton yield and fiber fineness. Lint percentage, fiber length and strength were

insignificantly affected by weed control treatments. Nabil *et al.* (1983) stated that the application of Stomp before planting gave the highest lint percentage, micronaire value and oil percentage. El- Shaer *et al.* (1985) found that seed cotton yield per plant and per faddan as well as number of opened bolls per plant were increased. However, fiber length, strength and micronaire reading were not affected by weed control treatments. Hussain *et al.* (1989) compared the effects of 4 weed control treatments on cotton. They found that all treatments increased number of bolls per plant, seed cotton weight per plant and lint cotton yield. Abd El-Rehim *et al.* (1995) found that fiber length at 50% and 2.5% S.L., micronaire reading, fiber strength uniformity ratio and fiber elongation were not significantly affected by either hand hoeing or the various herbicidal treatments. However, fiber length uniformity ratio, fiber strength at both zero and 1/8 inch gauge length, fiber stiffness, fiber toughness and yarn strength were significantly increased by hand hoeing or different herbicidal treatments.

Ghourab (1990) stated that the herbicide combination Goal + Amex showed higher seed cotton yield per plant or faddan than single application of these herbicides. Micronaire reading was significantly affected by herbicidal treatments in one season only. He added that weed control treatment had insignificant effect on fiber strength, elongation and fiber length. Meanwhile, Abd El-Bary *et al.* (2010) mentioned that the cotton promising cross Giza 89x 86) exceeded the commercial genotype Giza 86 by about 10% of cotton yield. For these reasons, the objective of this investigation was to determine the degree of integration between chemical and mechanical weed control and cotton genotypes on associated weeds, cotton yield and its fiber properties.

MATERIALS AND METHODS

Two field experiments were conducted at Sakha Agricultural Research Station Center, Kafer El-Sheikh Governorate during 2011 and 2012 seasons. The aim was to study the performance of three cotton genotypes (*Gossypium barbadense*, L.) , Giza 88, Giza 86 and the promising hybrid Giza 89 x Giza 86, under six weed control packages on their associated weeds and its prabroly and fiber properties. All genotypes were planted in April, 5th in both seasons. Soil texture of the experimental site in both seasons was clay loam (Table 1).

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

Table 1. Mechanical and chemical analysis for experimental soil.

soil analysis	2011	2012
Sand (%)	16.7	18.9
Silt (%)	33.14	32.73
Clay (%)	50.2	48.4
Soil textural class	Clay	Clay
(Ph)	8.00	8.09
Soil salts E.C.(m.mhos/cm)	2.49	3.01
Organic matter (%)	1.54	1.30
Available nitrogen (ppm)	19.35	22.00
Available phosphorus (ppm)	15.00	20.00
Available potassium (ppm)	278.1	283.00

Eighteen treatments of each experiment were arranged in a split plot design with four replicates. The sub plot area was 10.5 m² contained five ridges 3.5 m length and 60 cm apart. The distance between hills was 25 cm apart. Seedlings were thinned to secure the required number of plants before the 2nd irrigation. Cotton plants were picked on twice, the firstly picked on Oct. 5th and secondly on Nov. 1st in the two seasons. The treatments were as follow:

I. Main plots (cotton genotypes):-

- 1- Giza 86. 2-Giza 88. 3-The promising hybrid Giza 89 x 86).

II. Sub plots (integrated weed control packages):-

1. Amex (butralin, 48% EC) 4-(1,1-dimethylethyl)-N-(1-methylpropyl)-2,6-dinitrobenzenamine) at rate of 2.5 l/fad., applied post sowing, followed by one hand hoeing after 45 days from sowing.
2. Stomp Extra (pendimethalin, 45.5% CS) N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine at rate of 1.7 l/fad., applied post sowing, followed by one hand hoeing after 45 days from sowing.
3. Fusilade Super (fluazifop-P- butyl 12.5% EC) butyl @-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy] henoxyl]propanoate at rate of 1.5 l/fad., applied at 21 days after sowing followed by on hand hoeing after 45 days from sowing.
4. Nabu.S (sethoxydim, 12.5% EC) (t)-(EZ)-2-(1-ethoxyminobutyl)-5-[2-(ethylthio) propyl]-3-hydroxycyclohex-2-enone) at the of 1.5 l/fad., applied at 21 days after sowing followed by one hand hoeing after 45 days from sowing.
5. Hand hoeing (twice), at 30 and 45 days from sowing.
6. Control (untreated).

Herbicides in both experiments were sprayed by knapsack sprayer CP3 with water volume of 200 liters water/fad. All recommended agronomic practices of cotton were adopted throughout both growing seasons. The collected data were as follows:

A. Dry weight of weeds (g/m²):-

Weeds were hand pulled at random from one square meter of each plot after 75 and 105 days from sowing and classified into four categories, broad leaf, grassy,

perennial and total weeds. The dry weight was determined as (g/m²) after drying in a forced draft oven at 70 C° for 48 hours.

B. Cotton vegetative characters and yield components:

At harvest time, samples of ten plants were collected at random from the central ridges of each plot to assess the following criteria: plant length (cm), number of fruiting branches / plant, number of bolls / plant, boll weight (g), seed index, lint and seed cotton yield was estimated as kantar per faddan. (Boll weight: the average weight of ten open bolls in grams, seed index: the weight of 100 seeds in grams, lint percentage: ratio of lint to seed cotton expressed as a percentage).

C. Cotton fiber properties:

The cotton fiber properties studied were: Micronaire reading, fiber length: the length at 2.5% span length were measured, uniformity ratio was calculated (Sundaram, 1980), pressely index as measured by Fibrograph (ASTM, D 1447-83, 1984), and fiber length (mm) the cotton fiber properties were estimated in Cotton Agronomy Department Cotton Res. Institute (ARC).

Correlation study:

Simple correlation matrix was carried out for the two seasons to investigate the relationship between dry weight of different weed categories and cotton genotypes yield and its components according to Steel and Torrie (1980).

Statistical analysis:

Results were analyzed as split plot design with four replicates and a combined statistical analysis for the two years following the procedure outlined by Snedecor and Cochran (1980) Data were exposed to Bartlett test and were homogenous, for this reason the combined data of the two years were presented in the following results. Means were compared according to Duncan's multiple range test (1955).

RESULTS AND DISCUSSION

A: Effect on weeds

The most dominant weeds in experimental fields were wild jute (*Corchorus olitorius* L.), cocklebur (*Xanthium brasiliicum* L.), white goosefoot (*Amaranthus album* L.), common purslane (*Portulaca olerceae* L.), and black nightshade (*Solanum nigrum* L.), as broad leaf weeds and deccan grass (*Echionchloa colona* L.), Panz (*Dinebra retroflexa* L.), and bermuda grass (*Cynodon dactylon* L.), as grassy weeds and purple nutsedge (*Cyperus rotundus* L.) as perennial weeds in both 2011 and 2012 seasons. Table (2) show mean squares of variance for the effects of cotton genotypes, weed control packages and their interactions on dry weight of weeds were statistically significant at 1% level of probability.

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

Table 2. Means squares from analysis of variance for the effect of cotton genotypes weeds control and their interaction on dry weight of weeds (g/m^2) (combined data of 2011 and 2012).

Source	DF	Days after sowing									
		75					105				
		Grasses weeds	Broadleaf weeds	perennial weeds	Total Weeds	Dry weight of weeds in (g/m^2)	Grasses weeds	Broadleaf weeds	Perennial weeds	Total weeds	Total weeds
Year (Y)	1	6611.34**	1026.8	66.9	1688.23	389.12	2957981.5**	118094.3**	4339942.96**		
R x Y	4	867.63**	2838.5*	476.02	4376.04*	5889.43	19150.1	94569	17722.73		
Genotypes (G)	2	15978.86**	49063.4**	44247**	307757.07**	127125.48**	1328060.46**	60095.47**	3057096.04**		
Y x G	2	1599.62**	10511.58**	945.82**	24604.29**	24286.04*	44346.13*	24840.51**	65682.7*		
Error Treatment	8	321.28	1640.95	502.64	2288.68	10428.7	28737.02	1239.27	38040.63		
(T)	5	36093.57**	193917.37**	64080.7**	734629.81**	470080.08**	1777848.8**	91757.78**	5229997.8**		
Y x T	5	1479.59**	22531.75**	1676.1**	24722.48**	18707.72**	94960.44**	30177.57**	365795.64**		
G x T	10	6264.25**	11528.65**	7822.04**	72449.69**	48064.48**	175254.36**	9051.53**	511907.42**		
Y x G x T	10	121.17	1355.42	51.42	2275.57	860417*	5539.4	4806.03**	13947.47		
Error	60	619.49	1887.6	488.74	3327.7	8056	21073.53	646.04	24034.8		

Values followed by * are significantly at ($p=0.05$).

1- Effect of cotton genotypes:

Results in Table (3) show that dry weight of weed groups was affected by cotton genotypes at 75 and 105 days from sowing. Giza 86 genotype plots had the heaviest dry weight of weeds while, Giza 88 had the moderate weights for grassy, broad-leaved, perennial and total weeds at 75 and 105 days from Sowing. The promising hybrid Giza 89 x 86 cotton genotype gave the broadest spectrum of weed control and decreased dry weight of weed groups by 27.6, 31.9, 18.7 and 26.8% for weeds, respectively at 75 days after sowing as compared with Giza 86 where the respective values at 105 days were 17.46, 27.25, 18.6 and 24%, respectively. These results sustained that Giza 89 x 86 hybrid genotype is more competitive genotype than both Giza 86 and Giza 88 ones. Variation in weed competition strength between the three cotton genotypes may be attributed to their variation in allelopathic potential or variation in canopy architecture. Abd El-Bary *et al* (2010) mentioned that hybridization between Giza 89 as female and Giza. 86 as male

in 1989 collected features as compact growth and the plants were highly resistant to fusarium wilt.

Table 3. Effect of cotton genotypes on dry weight of weeds (g/m²) after 75 and 105 days from sowing (combined data of 2011 and 2012).

Cotton genotypes	Days after sowing							
	75				105			
	Dry weight of weeds (g/m ²)							
	Grasses weeds	Broadleaf weeds	perennial weeds	Total weeds	Grasses Weeds	Broadleaf weeds	perennial weeds	Total weeds
Giza 86	51.9a	136.7a	90.8a	279.4a	150.1a	485.1a	110.1a	745.3a
Giza 88	44.1b	105.7b	82.3b	232.1b	131.5b	388.1b	97.5b	617.1b
promising hybrid Giza 89 x 86	37.6c	93.1c	73.8c	204.5c	123.9c	352.9c	89.7c	566.5c

Means followed by the same alphabetical letters were not statistically significant according to Duncan's multiple range test.

2-Effect of weed control treatments:

Data presented in Table (4) indicate that different weed control packages depressed the dry matter weight of different weed categories namely grasses, broad-leaved, perennial and total weeds than control treatment at 75 and 105 days from sowing. The highest control percentages were obtained from pendimethalin/hoeing, butralin/hoeing packages and then followed by hard hoeing twice treatment which were 89.96, 88.98 and 82.6% of total weeds at 75 days after sowing, respectively. These results had the same trend at 105 days from sowing.

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

These results are logic because there is no solely weed control treatment is sufficient by it self, and herbicidal hoeing integration is needed for sufficient weed control in cotton due to the longevity of plant season and its poor competition to weeds. Several investigators reported that dinitroaniline as pendimethalin were effective in controlling cotton summer weeds (Fayed *et al.* 1983 and Khan *et al.* 1994). In this respect Fayed *et al.* (1983) reported that applying one supplementary hoeing in cotton herbicidal treatments was necessary to eliminate the weed plants which survived or escaped from the herbicides and to achieve promising weed control along the growing season of cotton plants.

Table 4. Effect of weed control treatments after 75 and 105 days from sowing (combined data of 2011 and 2012).

Weed control packages	Days after sowing							
	75				105			
	Dry weight of weeds (g/m ²)							
	Grasses weeds	Broadleaf weeds	perennial weeds	Total weeds	Grasses Weeds	Broadleaf weeds	perennial weeds	Total weeds
Pendimethalin (1.7 l/f) / hand hoeing.	13.1b	18.2d	34.1b	65.3d	48.8b	130.3c	38.4bc	217.6d
Butralin (2.5 l/f) / hand hoeing	12.7b	21.8cd	30.6b	65.2d	35.9b	194.0c	39.1bc	269.0cd
Fluazifop -p-butyl (1.5 l/f) / hand hoeing	5.5b	108.7b	22.9b	137.2bc	18.6b	323.3b	23.0c	365.0bc
Sethoxydim (1.5 l/f) / hand hoeing	7.0b	116.8b	26.2b	150.0b	17.1b	354.1b	42.5b	413.7b
Hand hoeing twice	15.1b	49.8c	37.8b	102.8cd	23.1b	340.1b	40.6bc	403.9b
Control (untreated)	120.0a	295.5a	175.9a	591.3a	423.4a	1006.2a	210.8a	1640.5a

Means followed by the same alphabetical letters were not statistically significant according to Duncan,s multiple range test.

3-The effect of interaction between genotypes and weed control treatments:-

Table (5) show that the effect of interaction of cotton genotypes with weed control packages on dry weight of broad-leaved, grasses, perennial weeds and total weeds (g/m²) at 75 and 105 days from sowing was statistically significant at 5% level.

Fluazifop -p-butyl + hand hoeing package under promising Giza 89 x 86 exerted the highest percentage of controlling of grassy weeds. This result is owing to the sensitivity of grasses to Fluazifop -p-butyl herbicide in addition to the high competition strength of the promising Giza 89x 86 genotype plants against grassy weeds (Table 3). Similar results were obtained by Dilbaugh *et al.* (2009) whom indicated that pendimethalin gave 82% control of broad-leaf weeds and 84.1% of narrow leaf.

The highest controlling percentages of broad leaf weeds were obtained from pendimethalin application followed by one hand hoeing which was under studied cotton hybrid genotype as compared with the other genotypes. This might be attributed to the susceptibility of broad-leaf weeds to pendimethalin herbicide integrated with the drastic competitive effect of cotton hybrid on weeds.

The studied interaction was statistically significant on dry weight of perennial weeds at 75 and 105 days from sowing. Flusifop-p-butyl + hoeing packages, under promising G89xG86 exerted the highest controlling percentage of perennial weeds. This result is owing to the role this herbicide integrated with the role of this cotton genotype.

Concerning the effect of interaction between cotton genotypes under weed control packages on dry weight of total weeds at both 75 and 105 days from sowing, the highest controlling % was achieved with pendimethalin/ hand hoeing packages under cotton hybrid. This is may be attributed it's the integrated effects of cotton genotype, herbicide and hoeing in maximizing weed control for the aforementioned weed categories in Table (5).

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

Table 5. The effect of interaction between cotton genotypes and weed control treatments after 75 and 105 days from sowing in 2011 and 2012 summer seasons.

Cotton genotypes	Weed control packages	Days after sowing											
		75					105						
		Grasses weeds	Broadleaf weeds	perennial weeds	Total weeds	Grasses Weeds	Broadleaf weeds	perennial weeds	Total weeds				
Giza 86	Pendimethalin (1.7 l/f) / hand hoeing.	22.7b	36.3ghi	63.7c	122.7efgh	88.8cd	188.3fghi	69.8de	346.9def	Dry weight of weeds in (g/m ²)	Total weeds	perennial weeds	Total weeds
	Butralin (2.5 l/f) / hand hoeing	19.7b	40.7ghi	46.7cd	107.0efgh	71.7cd	272.3fgh	71.2de	415.2de				
	Fluzifop -p-butyl (1.5 l/f) / hand hoeing	9.3b	137.5cd	35.3de	182.2de	35.0d	461.3cde	45.0de	541.3cd				
	Sethoxydim (1.5 l/f) / hand hoeing	13.3b	140.7cd	44.2cd	198.2d	34.5d	518.3cd	73.2de	626.0c				
	Hand hoeing twice	31.8b	51.7fghi	68.2c	151.7def	42.2d	545.0c	70.5b	657.7c				
Control (untreated)	215.0a	413.2a	286.5a	914.7a	628.3a	1525.0a	331.3a	2484.7a					
Giza 88	Pendimethalin (1.7 l/f) / hand hoeing.	12.0b	16.7hi	28.3de	57.0ghi	46.7cd	142.0ghi	30.8fg	219.5efg	Dry weight of weeds in (g/m ²)	Total weeds	perennial weeds	Total weeds
	Butralin (2.5 l/f) / hand hoeing	12.3b	20.3hi	33.5de	66.2ghi	28.7d	192.8fghi	29.2fg	250.7efg				
	Fluzifop -p-butyl (1.5 l/f) / hand hoeing	5.2b	103.2def	24.5de	132.8defg	17.0d	302.2efghi	13.9fg	333.1ef				
	Sethoxydim (1.5 l/f) / hand hoeing	5.5b	120.50cde	24.7de	150.7defg	13.8d	338.5def	37.8fg	390.2de				
	Hand hoeing twice	10.3b	65.3efgh	33.0de	108.7efgh	19.7d	336.2 defg	26.0fg	381.8de				
Control (untreated)	99.3b	308.3b	169.5b	577.2b	483.3b	1017.2b	207.3c	1707.8b					
Promising hybrid Giza 89 x 86	Pendimethalin (1.7 l/f) / hand hoeing.	4.7b	1.7i	10.2e	16.5i	11.0d	60.6i	14.8fg	86.4g	Dry weight of weeds in (g/m ²)	Total weeds	perennial weeds	Total weeds
	Butralin (2.5 l/f) / hand hoeing	6.2b	4.7i	11.7e	22.5i	7.3d	116.8hi	17.2fg	141.3g				
	Fluzifop -p-butyl (1.5 l/f) / hand hoeing	2.0b	85.7defg	9.0e	96.7fgh	4.0d	206.7fghi	10.2g	220.8efg				
	Sethoxydim (1.5 l/f) / hand hoeing	2.2b	89.3defg	9.7e	101.7fgh	2.8d7	205.5fghi	16.8fg	225.1efg				
	Hand hoeing twice	3.3b	32.5ghi	12.3e	48.7hi	7.3d	139.5hi	25.6fg	172.4fg				
Control (untreated)	45.7a	165.0c	71.7c	282.3c	158.7c	476.7cde	93.8d	729.2c					

Means followed by the same alphabetical letters were not statistically significant according to Duncan's multiple range test.

B- Effect on seed cotton yield and its components:-

Data of ANOVA in Table (6) indicate that the effect of each of cotton genotypes or weed control packages on all studied characters were statistically significant except lint%, and the effects of the interaction between genotypes and weed control packages on plant length, boll weight and seed cotton yield were statistically significant where on the rest traits the effect of the interaction were not statistically significant. Also, data in Table (6) show that the effect of interaction between cotton genotypes and weed control packages on plant height (cm), number of fruiting branches per plant, number of bolls/plant and seed index was not statistically significant, meaning that the two studied factors act independent. Meanwhile, the effect of interaction between cotton genotypes and weed control packages on plant height (cm), boll weight (g), seed cotton yield quintar/fad. and lint% were statistically significant.

Table 6. Means squares of analysis of variance for the effect of cotton genotypes, weed control and their interactions on the growth, yield and yield component (combined data of 2011 and 2012).

Source	DF	Plant length (cm)	No. of fruiting branches	No. of bolls / plant	Boll weight (g)	Seed index	Seed cotton yield (quintar /fad.)	Lint %
Year (y)	1	2.676	0.13	0.021	0.388**	47.07**	81.085**	535.113**
Replications x Year	4	87057*	12.213*	1.252	0.080*	2.469*	0.106	6.313
Genotypes (G)	2	3220.75**	27.82**	54.606**	1.113**	11.713**	13.639**	9.628
YG	2	113.676	11.682*	3.165	0.005	5.754**	1.421**	12.171*
Error	8	536.41	5.39	4.411	0.041	1.444	0.147	11.88
Weed control treatments (T)	5	4029.15**	58.94**	65.95**	1.884**	7.737**	54.813**	14.27
YT	5	77.298	1.72	3.661*	0.092*	3.071**	5.708**	10.796
GT	10	248.05*	1.56	3.014	0.302**	0.945	1.264**	19.446
YGT	10	73.99	2.593	3.351*	0.015	5.173**	0.920**	10.32
Error	60	217.25	4.09	3.052	0.068	1.215	0.181	10.978

Values followed by * are significantly (P=0.05).

1. Effect of cotton genotypes:

Data recorded in Table (7) show that the promising hybrid Giza 89 x Giza 86 significantly exceeded Giza 88 and Giza 86 genotypes in all growth characters and seed cotton yield (quintar /fad) and its attributes except, number of bolls/plant. The

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

promising hybrid Giza 89 x Giza 86 recorded the highest values of plants height (cm), number of fruiting branches, boll weight / plant (g), seed index, seed cotton yield/quintar followed by Giza 88 as compared with Giza 86. These results are confirmed results obtained by Abd El- Bary *et al* (2010), they mentioned that the promising cross Giza 89x 86 characterized by high yielding and out yielded Giza 86 by about 10% in cotton yield.

Table 7. The yield and yield components as affected by cotton genotypes (combined data of 2011 and 2012).

Cotton genotypes	Plant length (cm)	No. of fruiting branches	No. of bolls / plant	Boll weight (g)	Seed index (g)	Seed cotton yield (quintar /fad)	Lint %
Giza 86	147.2b	19.12ab	16.4b	2.53b	8.66b	10.1c	36.26
Giza 88	156.8ab	13.82c	17.6a	2.45b	8.90b	10.5b	36.37
promising hybrid Giza 89x86	166.1a	19.20a	16.1ab	2.78a	9.74a	10.8a	35.43

Means followed by the same alphabetical letters were not statistically significant according to Duncan,s multiple range test.

2-Effect of weed control packages:

Data in Table (8) indicated that all used weed control packages as well as hand hoeing increased significantly seed cotton yield and its components. Pendimethalin and butralin treatments followed plus by one hand hoeing for each gave the highest values of plant length, number of fruiting branches, number of bolls per plant and seed cotton yield per (quintar /fad) in both seasons followed by fluazifop-p-butyl and sethoxydim plus one hand hoeing and hand hoeing twice. These treatments increased seed cotton yield by 66.7, 62.4, 56.5, 46.4 and 63.8 % respectively, as compared to control treatment. The influence of such treatments on seed cotton yields had the same trend of the abovementioned yield attributes traits. It is worthwhile to mention that, these treatments which gave the highest values of seed cotton yield were also show lowest dry weight of weeds. In this respect Fayed *et al.* (1983) and Khan *et al* (1994) recorded that the highest seed cotton yield with application of pendimethalin and attributed such increase to the increases in seed index, boll weight, number of boll/plant and plant heights which are attributed to minimization of weed crop competition for light, space and mineral uptake. Ikram *et al.* (2012) found that the

maximum nutrient losses of NPK were found in control and improving growth and NPK uptake of cotton under weed control by herbicides.

Table 8. The yield and yield components as affected by weed control packages (combined data of 2011 and 2012).

Weed control packages	Plant length (cm)	No. of fruiting branches	No. of bolls / plant	Boll weight (g)	Seed Index (g)	Seed cotton yield (quintar /fad)	Lint %
Pendimethalin (1.7 l/f) / hand hoeing	173.7a	25.6a	20.6a	2.9a	9.9a	11.5a	35.2b
Butralin (2.5 l/f) / hand hoeing .	168.2ab	25.5a	19.2b	2.8ab	9.7ab	11.2a	36.9a
Fluazifop -p-butyl (1.5 l/f) /hand hoeing	160.3bc	23.6ab	17.6c	2.8ab	9.2ab	10.8b	36.6a
Sethoxydim (1.5 l/f) / hand hoeing .	157.2cd	20.7b	16.9c	2.6bc	9.0ab	10.1c	36.2ab
Hand hoeing twice	149.1d	20.5b	16.5c	2.5c	8.3c	11.3a	36.6a
Control (untreated)	131.7e	15.7c	15.2d	1.9d	8.4c	6.9d	34.7c

Means followed by the same alphabetical letters were not statistically significant according to Duncan,s multiple range test.

3-The effect of interaction between genotypes and weed control treatments on yield and its components:-

Data in Table (9) indicate that all interactions between cotton genotypes and weed control treatments were statistically significant on their effects on plant height (cm), boll weight (g) seed index cotton yield quintar/fad, and lint% increased seed cotton yield and its components except, for plant length and number of fruiting branches. The tallest plants were obtained from the the promising hybrid Giza 89 x Giza 86 with pendimethalin (35.8%) as compared with the shortest plants whom resulted from untreated check.

The interaction between the promising hybrid Giza 89 x 86 with pendimethalin plus one hand hoeing gave the highest values for cotton yield (12.6 quintar/fad) while, the lowest values (7.0 quintar/fad) was obtained by Giza 86 with control treatment. The increases in such trait is attributed to the highest weed elimination % under these treatments than unweeded check, and consequently, decreasing weed/crop competition.

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

Table 9. The yield and yield components as affected by interaction between cotton genotypes and weed control treatments at harvest (combined data at 2011 and 2012).

Cotton genotypes	Weed control packages	Plant length (cm)	Boll weight (g)	Seed index (g)	Seed cotton yield (quintar/fad)	Lint %
Giza 86	Pendimethalin (1.7 l/f) / hand hoeing .	153.0def	2.8 a	9.6	11.9 a	32.2bc
	Butralin (2.5 l/f) / hand hoeing .	155.5def	2.9 a	9.3	11.0 cd	37.3a
	Fluazifop -p-butyl (1.5 l/f) / hand hoeing	154.3def	2.9 a	8.5	11.6ab	37.1a
	Sethoxydim (1.5 l/f) / hand hoeing	151.8def	2.6abcd	8.5	11.0 cd	37.0a
	Hand hoeing twice	146.7ef	2.36cd	7.9	11.2bc	36.9a
	Control (untreated)	121.8g	1.7e	8.2	7.0	36.9a
Giza 88	Pendimethalin (1.7 l/f) / hand hoeing .	178.2abc	2.9abc	9.7	11.9 a	36.8a
	Butralin (2.5 l/f) / hand hoeing .	167.2bcd	2.7abcd	9.5	11.4abc	36.3ab
	Fluazifop -p-butyl (1.5 l/f) / hand hoeing	155.2def	2.7abcd	8.5	10.9cd	37.0a
	Sethoxydim (1.5 l/f) / hand hoeing	157.2de	2.4d	8.9	10.4e	36.6a
	Hand hoeing twice	146.7ef	2.4bed	8.2	11.9a	35.9ab
	Control (untreated)	136.3fg	1.7e	8.7	7.1h	35.7ab
promising hybrid Giza 89 x 86	Pendimethalin (1.7 l/f) / hand hoeing .	189.8a	2.9a	10.5	12.6a	36.6a
	Butralin (2.5 l/f) / hand hoeing .	182.0b	2.8a	10.5	11.1bcd	37.1a
	Fluazifop -p-butyl (1.5 l/f) / hand hoeing	171.5 bcd	2.8a	10.6	9.6f	35.7ab
	Sethoxydim (1.5 l/f) / hand hoeing	162.7cde	2.9a	9.6	8.9g	34.8abc
	Hand hoeing twice	153.8def	2.7ab	8.9	10.9cde	36.9a
	Control (untreated)	136.8fg	2.6abcd	8.4	7.3i	31.5c

Means followed by the same alphabetical letters were not statistically significant according to Duncan, s multiple range test.

C- Effect on cotton fiber properties:-

Mean squares of analysis of variance for the effect of cotton genotypes, weed control treatments and their interactions on cotton fiber properties are shown in Table (10).

Table 10. Means squares of analysis of variance for the effect of cotton genotypes, weed control treatments on cotton fiber properties and their interactions on fiber properties are show (combined data of 2011 and 2012).

Source of variation	DF	Micronaire reading	Uniformity ratio	Pressely index	Fiber length
Year (Y)	1	0.311	0.078	14.16**	62.563**
RY	4	0.263	6.977**	3.451**	9.931*
Genotypes (G)	2	0.991	9.480**	0.245	15.593
YG	2	0.04	0.008	0.845*	0.023
Error	8	0.347	1.387	0.749	7.172
Treatment (T)	5	2.963**	2.095	0.535	2.844
YT	5	0.011	0.009	0.747	0.014
GT	10	0.189	1.273	2.087	3.691
YGT	10	0.009	0.01	0.714	0.01
Error	60	0.098	1.053	1.268	3.126

Values followed by * are significantly ($p= 0.05$).

1- Effect of cotton genotypes:-

It is obvious in Table (11) that off the studied four fiber quality measurements, uniformity ratio was significantly affected by genotypes. The promising hybrid gave the lowest value of uniformity ratio whereas the others two genotypes were statistically equal.

Table 11. Fiber quality measurements as affected by cotton genotypes at harvest (combined data of 2011 and 2012 summer seasons).

Cotton genotypes	Micronaire reading	Uniformity ratio	Pressely index	Fiber length (mm)
Giza 86	3.96	87.12	10.36	34.13
Giza 88	3.80	87.28	10.26	35.24
Promising hybrid Giza 89 x 86	4.19	86.32	10.19	34.08

Means followed by the same alphabetical letters were not statistically significant according to Duncan,s multiple range test.

2- Effect of weed control packages:-

Data in Table (12) indicated that micronaire reading increased significantly under various weed control packages namely pendimethalin/hoeing, butralin /hoeing, fluazifop/hoeing, sethoxydim/hoeing and hand hoeing twice by 33.9, 34.5, 30.5 and

25.16 % than untreated check respectively. The increases in micronaire reading of cotton obtained from weed control packages may be attributed to successful control weeds which reduced competition and consequently favored growth, cotton yield and lint maturity. These results are in harmony with those obtained by Keeley and Thullen (1975) whom mentioned that the micronaire reading were lower for cotton sample collected from plots contained nutsedge than weed free plots indicating delayed crop and fiber maturity. Nabil *et al.* (1983) and El-Shaer *et al.* (1983) stated that the herbicide stomp gave the highest micronaire r reading, fiber length and fiber strength.

Table 12. The combined data for cotton fiber properties as affected by weed control treatments in 2011 and 2012 seasons.

Weed control packages	Micronaire reading	Uniformity ratio	Pressely index	Fiber length (mm)
Pendimethalin (1.7 l/f) / hand hoeing .				
Butralin (2.5 l/f) / hand hoeing .	4.26a	86.93	10.38	34.51
Fluazifop -p-butyl (1.5 l/f) / hand	4.26a	86.92	10.09	35.13
hoeing Sethoxydim (1.5 l/f) / hand	4.15ab	87.01	10.27	34.07
hoeing	4.07ab	87.23	10.08	34.66
Hand hoeing twice	3.98b	86.25	10.53	34.48
Control (untreated)	3.18c	87.1	10.28	34.07

Means followed by the same alphabetical letters were not statistically significant according to Duncan,s multiple range test.

3-The effect of interaction between genotypes and weed control treatments:

The effect of interaction between cotton genotypes and weed control treatments on cotton fiber micronaire reading was statistically significant (Table 13). The highest micronaire reading was obtained with promising hybrid with pendimthalin application (4.50) while the lowest value were obtained from Giza 86 under untreated check (3.05). These results mean that cotton fiber maturity can be enhanced with weed elimination from cotton fields. Similar results were obtained by Keeley and Thullen (1975) whom mentioned that the micronaire reading were lower for cotton sample collected from plots contained nutsedge than weed free plots indicating delayed crop and fiber maturity.

Table 13. The micronaire reading as affected by interaction between cotton genotypes and weed control treatments as quality measurement of cotton genotypes.

Weed control packages	Cotton genotype		
	Giza 86	Giza 88	promising hybrid Giza 89 x 86
Pendimethalin (1.7 l/f) / hand hoeing .	4.17 b	4.10a	4.50a
Butralin (2.5 l/f) / hand hoeing .	4.45ab	4.00ab	4.32ba
Fluazifop -p-butyl (1.5 l/f) / hand hoeing	3.93b	4.07ab	4.45a
Sethoxydim (1.5 l/f) / hand hoeing	4.05ab	3.85ab	4.30ab
Hand hoeing twice	4.10ab	3.67b	4.17b
Control (untreated)	3.05c	3.13c	3.37c

Means followed by the same alphabetical letters were not statistically significant according to Duncan,s multiple range test.

Correlation between studied characters and cotton yield:

Data presented in Table (14) indicated clearly that simple correlation coefficients between dry weight of grassy weeds and broad-leaved weeds species and cotton yield was statistically significant and negative at 5% level. Such correlation was strong with broad-leaved weeds (-0.946, -0.774 and -0.903) than with grassy weeds (-0.865, -0.756 and -0.551) for the three genotype of cotton (Giza 86, Giza 88 and promising hybrid Giza 89 x 86), respectively. This mean that broad-leaved were more aggressive in their competition to cotton genotypes than grassy weeds. Correlation between dry weight of total annual weeds and seed cotton yield recorded the highest value, where negatively affected cotton yield by (-0.960,-0.783 and -0.876) for the three genotypes of cotton (Giza 86, Giza 88 and promising hybrid Giza 89 x 86), respectively.

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

Table 14. Correlation coefficient between studied characters and some cotton genotypes yield, (combined data of 2011 and 2012).

Cotton genotypes	Studied characters	Grassy weeds (g/m ²)	Total weeds(g/m ²)	Plant height (cm)	Boll weight/(g)	Pressely index	Fiber length	Seed cotton yield (quintar /fad)
Giza 86	Broad-leaved weeds (g/m ²)	0.853**	0.987**	-0.656**	-0.853**	-0.223	-0.283	-0.946**
	Grassy weeds (g/m ²)	-	0.915**	-0.376	-0.782**	-0.044	-0.383	-0.865**
	Total weeds (g/m ²)	-	-	-0.593*	-0.862**	0.204	-0.295	-0.960**
	Plant height (cm)	-	-	-	0.504*	0.324	-0.158	0.576*
	Boll weight /(g)	-	-	-	-	-0.464	-0.062	0.703**
	Pressely index	-	-	-	-	-	-0.227	-0.172
	Fiber length	-	-	-	-	-	-	0.238
Giza 88	Broad-leaved weeds (g/m ²)	0.914**	0.989**	-0.397	-0.764**	-0.138	-0.351	-0.774**
	Grassy weeds (g/m ²)	-	0.962**	-0.430	-0.616**	-0.048	-0.306	-0.756**
	Total weeds (g/m ²)	-	-	-0.413	-0.731**	0.073	-0.328	-0.783**
	Plant height (cm)	-	-	-	-0.456	0.155	-0.140	0.363
	Boll weight /(g)	-	-	-	-	-0.064	-0.047	0.612**
	Pressely index	-	-	-	-	-	-0.096	-0.112
	Fiber length	-	-	-	-	-	-	0.235
Promising hybrid Giza 89 x 86	Broad-leaved weeds (g/m ²)	0.659**	0.957**	-0.540*	-0.221	-0.091	-0.040	-0.903**
	Grassy weeds (g/m ²)	-	0.842**	-0.211	-0.499*	-0.088	-0.003	-0.551*
	Total weeds (g/m ²)	-	-	-0.474*	-0.616**	0.006	-0.026	-0.876**
	Plant height (cm)	-	-	-	0.531*	0.064	-0.109	0.614**
	Boll weight /(g)	-	-	-	-	-0.076	0.120	0.732**
	Pressely index	-	-	-	-	-	-0.333	-0.084
	Fiber length	-	-	-	-	-	-	0.076

Also, correlation analysis revealed that the yield increases due to type of weed competition were positively contributed to the increases in growth characters and yield components. The correlation between total weeds and cotton yield in three cotton genotypes was highly statistically significant. Hence, applying weed control packages play a major role in increasing cotton productivity and improve fiber maturity per unit urea, when applied at the suitable time, rate and stage of weed growth.

REFERENCES

2. Abd El-Bary, A. M. R., H. M. E. Hadmoud, W. M. B. Yehia, M.S.M. Sour, Samia E.S. Ali and Eman M. R. M. Saleh. 2010. The long staple Egyptian cotton promising cross Giza 89xGiza 86. J. Agric. Chemistry and Biotechnology. Mansoura Univ., Vol. (6): 325-339.
3. Abd El-Rehim, Salwa A., Samia A. Saad EL-Din and A. H. A. Afifi. 1995. Comparative study on the effect of hand hoeing and different herbicidal treatments on physical properties of fiber and yarn growth and yield in Egyptian cotton. J. Agric. Sci. Mansoura Univ., 21 (7): 2473-2483.
4. Anonymous. 2012. Brief of Agricultural Statistics, Arab Republic of Egypt, Ministry of Agricultural and Land Reclamation , bulletin, P 30.
5. ASTM. 1984. American Society for Testing and Materials. Standards on textile materials. D: 1447-83, D: 1448-74 and D:1445-75.
6. Buchanan, G. A. and E.R. Burns. 1971. Weed competitions in cotton. Weed Sci, 19: 580-582. 11.
7. Bukun, B., 2004. Critical periods for weed control in cotton in Turkey. Weed Res. 44, 404- 412.
8. Dilbaugh, M., N. A. Muhammad, I. Raza and M. A. Mian. 2009. Growth and development of cotton (*Gossypium hirsutum* L.) as affected by different methods of pendimethalin application, Pak. J. Weed Sci. Res. 15(1): 11-17,
9. Duncan, B.O. 1955. Multiple range and multiple range F test. Biometrics, 11: 1-42.
10. El- Shaer, M. H., M. S. Abd El- Raouf, A. M. Samar and H. A. Ali. 1985. Effect of plant population density and weed control treatments on weed dry weight and cotton yield and quality. Annals of Agric. Sci. Ain. Shams Univ., 28 (3): 1291-1307.
11. Fayed, M.T, M.T. Mostafa and E.E. Hassanein. 1983. Increasing the efficiency of herbicides in controlling cotton weeds by one light hoeing .Proc. First Conf. of Agron. Egyptian Soc. of Crop Sci., 2: 679-688.
12. Ferrel, J.A., G.E. Mac Donald and B. J. Brecke. 2001. Weed management in cotton. University of Florida IFSA Extension, pp, 1-32.(C.F. Asian j. of plant sci. 6(2): 344-348,
13. Ghourab, M.H.H. 1990. Effect of some herbicides on Egyptian cotton. Ph.D. Thesis, Fac. Agri., Al-Azhar Univ., Egypt.
14. Ikram R. M., M. A. Nadeem, A. Tanveer, M. Yasin, A. U. Mohsin, R. N. Abbas, H. Rehman, M. Sibtain and M. Irfan. 2012. Comparative efficacy of different pre-

EFFECT OF SOME WEED CONTROL PACKAGES
ON SEED COTTON YIELD AND FIBER PROPERTIES
AND ASSOCIATED WEEDS

- emergence herbicides in controlling weeds in cotton (*Gossypium hirsutum L.*) Pak. J. Weed Sci. Res. 18(2): 209-222.
15. Keeley, P.E., C.H. Carter and J.H. Miller. 1973. Yellow nutsedge and cotton response to several herbicides. Weed Sci. 21:327-329.
 16. Keeley, P.E., and R.J. Thullen. 1975. Influence of yellow nutsedge competition on furrow-irrigated cotton. Weed Sci. 23:171-175.
 17. Khan, N.U., S.U. Khan, G. Hassan, I. H. Shah and Q. Nawaz. 2001. Studies on weed control in cotton (*Gossypium hirsutum L.*) online, J. Bid. Sci. 1 (3): 143-145.
 18. Nabil, Z.S., A. M. Samara and S. H. Yousef. 1983. Effect of different pesticide treatments on fiber and seed characteristics. (Proc. of the first Conf. of Agron., 96: 699-707).
 19. Snedecor, G. W. and W. G. Cochran. 1980. " Statistical Methods "6th ed., Iowa State Univ. Press, Ames. Iowa, U.S.A, 325-330.
 20. Steel, R. G. D. and Jour. H. Torrie. 1980. Principles and Procedures of Statistics. MC. Graw Hill Book Company Inc. New York, 481 pp.
 21. Sundaram, V. 1980. Hand book of Methods of Tests for Cotton Fibers, Yarn and Fabrics. Indian Council of Agric. Res., P. 71-76.

تأثير بعض حزم مكافحة المتكاملة للحشائش على نمو ومحصول القطن وصفات التيلة لبعض التراكيب الوراثية للقطن والحشائش المصاحبة

ابراهيم السيد سليمان^١، عزة السيد خفاجي^١،
عادل مصطفى غلوش^١ و أمل سامى عبد العال^٢

١. المعمل المركزى لبحوث الحشائش - مركز البحوث الزراعية - الجيزة- مصر

٢. معهد بحوث القطن- مركز البحوث الزراعية - الجيزة- مصر

• أجريت تجربتان حقليتان فى مزرعة محطة البحوث الزراعية بسخا - كفر الشيخ خلال موسمى الزراعة ٢٠١١ و ٢٠١٢م - لدراسة تأثير ثلاثة أصناف من القطن المصرى (هجين ٨٩X٨٦ ، جيزة ٨٨ ، جيزة ٨٦) وبعض حزم مكافحة المتكاملة للحشائش (مبيد بنديمتالين وبيوترالين وفلوزيفوب-ب-بيونائل وسيثوكسيديم) متبوعة بعزقة يدوية واحفظكل منهم بالإضافة الى معاملتى العزيق مرتين وبدون معاملة وتأثير تلك الحزم على الحشائش و المحصول وصفات التيلة للقطن.

• أظهرت النتائج ان هجين ٨٦ X٨٩ قلل الوزن الجاف لكل من الحشائش الحولية (عريضة وضيقة الأوراق) والحشائش المعمرة فى المتر المربع وذلك بنسبة ٢٧.٦ و ٣١.٩ و ٢٦.٨% على الترتيب وذلك مقارنة بالصنف جيزة ٨٦ وزيادة معنوية لمكونات محصول القطن مثل طول النبات ومتوسط وزن اللوزة ومحصول بذرة القطن/ نبات. ومحصول القطن بالقنطار / فدان . كذلك حقق هذا الصنف زيادة معنوية فى صفات التيلة لمحصول القطن ممثلة فى قراءة الميكرونيير وتجانس ومثانة و طول التيلة. أيضا أعطت جميع حزم مكافحة الحشائش نقص معنوى فى جميع أنواع الحشائش الحولية (عريضة و ضيقة الأوراق) والحشائش المعمرة وزيادة معنوية فى جميع صفات مكونات المحصول وصفات التيلة للقطن خلال موسمى الزراعة. أيضا أعطت جميع حزم مكافحة الحشائش زيادة فى محصول القطن للفدان وسجلت حزم مكافحة الحشائش المكونة من (مبيد بنديمتالين أوبيوترالين متبوعة بعزقة يدوية واحدة) أعلى نسبة زيادة يليهما معاملة العزيق مرتين ثم معاملة مبيد فلوزيفوب-ب-بيونائل - وسيثوكسيديم متبوعين بعزقة يدوية واحدة وذلك مقارنة بمعاملة الكنترول حيث كانت نسبة الزيادة ٦٦.٧ و ٦٢.٣ و ٦٣.٧ و ٥٦.٥ و ٤٦.٤ % وزيادة قراءة الميكرونيير بمقدار ٣٤.٠ و ٣٤.٠ و ٢٥.٢ و ٣٠.٥ و ٢٨.٠% على التوالي مقارنة بمعاملة الكنترول.

• أدى التفاعل بين الأصناف وحزم مكافحة الحشائش الى نقص معنوى فى كل من الحشائش الحولية (عريضة وضيقة الأوراق) والحشائش المعمرة وتم الحصول على أعلى محصول قطن (قنطار للفدان) بزراعة صنف القطن هجين ٨٩ X٨٦ مع استخدام مبيد بنديمتالين أو بيوترالين متبوعة بعزقة يدويه واحدة أو معاملة العزيق مرتين يليه الصنف جيزة ٨٨ مع المعاملات السابقة وذلك مقارنة بالصنف جيزة ٨٦ مع معاملة الكنترول. كذلك أظهر تحليل الارتباط ان الوزن الجاف للحشائش يرتبط ارتباطا سلبيا مع محصول القطن كما ارتبطت صفات المحصول ارتباطا ايجابيا مع محصول القطن للفدان. لذا فان مكافحة الحشائش تلعب دورا كبيرا فى رفع انتاجية الفدان من محصول القطن.

• توصى هذه الدراسة باستخدام أصناف القطن المصرى الأكثر تأثيرا على الحشائش مثل (هجين ٨٩X٨٦ ، جيزة ٨٨) مع استخدام أي من حزم مكافحة الحشائش المكونة من (مبيدات للحشائش / عزقة يدوية واحدة) لتلافى منافسة الحشائش لمحصول القطن و بالتالى تحسين انتاجية وصفات التيلة لمحصول القطن المصرى.