



EFFECT OF USING BIO – AND ORGANIC FERTILIZERS ON PRODUCTIVITY AND QUALITY OF CUCUMBER CROP UNDER PLASTIC HOUSES

[2]

Al-Hmoudi¹, A.S.; M.H. Mohamed²; S.M. Al-Menaway²
and S.A. Hussain¹

1- Ministry of Environment and Water United Arab Emirates

2- Hort, Dept. Fac. of Agric, Ain Shams Univ., Cairo, Egypt

Keywords: Organic, Fertilizers, Bio-organic Cucumber, Plastic House. Productivity

ganic fertilizers can be used a source of nutrition instead of chemical fertilizers for cucumber plants.

ABSTRACT

An experiment was carried out to study the effect of Bio and organic fertilizer on cucumber growth and yield during two successive seasons of 2010 and 2011 at Northern region in united Arab Emirates. The Queen cucumber hybrid was fertilized with different organic fertilizers i.e. Horse manure, compost, chickens manure and cows manure. Also using the bio fertilizer i.e. phosphobactein, Azotobacter and mycorrhizal. The results showed that cucumber yields differed in their response to the bio and organic fertilizer.

The results showed that the compost was higher nitrogen percent and lower in C/N ratio, and pH, than the others organic fertilizer especially horse manure. The study demonstrated that the average cumulative cucumber yield was higher with compost + 22 (gm) mycorrhizol/ plots treatments compared to other treatments throughout the experiment during the two successive winter seasons of 2010 and 2011. The plant height and plant fresh weight (g), with N100% mineral (control) decreased by 28% and 40% compared to N 100% organic (Compost) respectively during the winter season of 2010. The nitrogen, phosphorus and potassium content of cucumber fruits (%) significantly increased, as did the soil with the increase of organic fertilizes applied. The experimental results confirmed the combination of bio-and organic fertilizers could increase plant growth, yield and quality. It also confirmed that composted or-

INTRODUCTION

It is very important depended on natural crops. we should be de creased the chemical fertilizers, but need to be use Bio- & organic fertilizers. Protected cultivation is now representing a significant contributes and expanding markedly in the newly reclaimed lands especially in united Arab Emirates. Cmjucumber is one of the most important vegetablegrown under plastic houses and produce high total yields.

Also it's a good source of vitamins, minerals, fibers. Cucumber flesh is rich in potassium and vitamin C.

Cucumber contains calcium (20gm/100g), iron (0.7mg/100g), thiamin (0.3mg/100g), niacin (0.2 mg / 100g) and riboflavin (0.01mg/ 100g).

In this studies includes two different resources for the fertilizers, bio-and organic fertilizers. **Hekal, A.I.A. (2001)**. Reported that the fungicides and pesticides increased the product costs and same time decreased the profit, cucumber was found in subtropical and tropical region favorable environmental condition for humid topics.

Regab (1984) find that the average fruit weight was controlled by 2-3 genes and this number of genes was highly affected by environmental conditions.

In soil **Masloand Gamaynnov (1989)** reported that the organic manure 40 ton/hectar for cucumber increased total yield 57 : 136%.

(Received 14 December, 2014)

(Accepted 21 December, 2014)

Eissa (1996) observed from his studies on the application organic manure increased significantly the fresh weight, also increased phosphorus in cucumber fruits while nitrogen content in cucumber shoot was increased. Vogtman et al (1993) reported that, compost treatments resulted in lower vegetable yields in the first two years but there were no different after the third year. Kostov et al (1995) reported that it was more economical to use composting vegetable residues for greenhouse cucumber production than cattle manure. They also reported that greenhouse cucumbers (*curcumas sativus* L.) grown on a medium containing composted vegetables or vegetable waste with the addition of synthetic nutrients produced fruit 10 to 20 days earlier and had a yield 48 to 79% higher than those grown in soil mixed with cattle manure at a 2.1 ratio (dry weight basis).

Roe et al (1997) studied cucumbers in a sandy soil fertilized with compost or mineral fertilizers. They found that, yields were usually higher when compost was combined with mineral fertilizers.

Esawy et al (2009) reported that the average nitrate content of cucumber fruits was only detected in the plots treated with 100% organic N from the composted tested.

The objectives of this study were to assess the effect of four types from organic fertilizers (Horse manure, compost, chickens manure and cows manure) and three bio-organic fertilizers (*phosphobacterin*), *Azotobacter* and *mycorrhizal*), on cucumber (*cucumis sativa*) plants during the two seasons 2010/2011 and sandy soil properties.

MATERIALS AND METHODS

An experiment was carried out in an un heated plastic house at the ministry of water and Environmental, experimental farm, Northern region in UAE during winter seasons of 2010 and 2011. The hybrids Queen F1 were used. This hybrid are already grown as minicucumbers on very large scales by most of the growers in the UAE. Cucumbers were seeded in cell flats (cell size 3x3x10 cm) filled with peat: perlite mixture, placed on benches under the plastic house conditions, for a period of 6 weeks. Seedlings were then transplanted (November 8th 2010 and November 15th 2011) to soil and grown under controlled conditions in an experimental plastichause.

The physical and chemical characteristic of the experimental soil sites are presented in Table (1). A split plot design with three replicates was used and the area of each replicate plot was 4.8m². The organic fertilizers were arranged in main ppls,

while the bio-organic fertilizer, plotted at random in sub plots.

Table 1. The physical and chemical analysis of the experimental soil in 2010 and 2011

Soil properties	2010 seasons	2011 seasons
A-Physical analysis		
Coarse sand %	8.6	8.7
Fin Sand %	63.8	64.1
Silt %	12.8	12.9
Clay %		
Soil type	Sandy loam	Sandy loam
B-Chemical analysis		
pH (1 : 5)	8.2	8.5
EC (mmhos/cm at 250)	8.11	8.61
Total N (ppm)	135.2	159.1
Available P (ppm)	6.5	6.9
Available K (ppm)	69.2	80.3
Iron	3.8	4.0
Zinc	0.7	0.91
Manganese	2.9	3.2
Copper	0.22	0.31

Plants were spaced at 50cm a part in 5 rows plastic house in rows of 40 m long. 50cm deep and 50cm wide. The soil was sandy is texture and drip irrigation system was used.

Four different organic fertilizers were applied + three bio-organic fertilizer were applied.

- 1- Control full dose chemical 100% NPK.
- 2- 100% dose NPK organic (Compost) + Phosphobacterin.
- 3- 100% dose NPK organic (Compost) + Azotobacter.
- 4- 100% dose NPK organic (Compos+) + Mycorrhizal.
- 5- 100% dose NPK Horse manure+ Phosphobacterin.
- 6- 100% dose NPK Horse manure + Azotobacter.
- 7- 100% dose NPK Horse manure + Mycorrhizal.
- 8- 100% dose NPK chickens manure + Phosphobacterin.
- 9- 100% dose NPK chickens manure + Azotobacter.
- 10-100% dose NPK chickens manure + Mycorrhizal.
- 11-100% dose NPK cowesmanure + Phosphobacterin.
- 12-100% dose NPK Cowesmanure + Azotobacter.
- 13-100% dose NPK Cowes manure + Mycorrhizal.

**Effect of Using Bio – and Organic Fertilizers on Productivity and Quality of Cucumber Crop 15
Under Plastic Houses**

Table 2. The chemical analysis of the experimental organic fertilizers

Sampel	N	P2O5	pH	EC	M.C.%	O.M.%	NaCl%	C/N ratio
Horse Manure	1.4	1.14	6.35	4.80	24.50	80.58	1.18	33.46
Compost	3.39	1.95	5.84	12.77	5.78	72.37	1.09	12.41
Chickens manure	2.77	3.00	7.02	19.11	35.03	79.57	2.56	16.70
Cows manure	1.51	1.88	8.05	14.26	9.27	46.34	2.18	17.84

Other cultural practices such as, irrigation, pesticide were applied according to the recommendation of ministry of water & Environmental. Plant height and Fresh weight of plants were estimated at the end of the harvesting season. Number of fruits/plant, average fruit weight (gm) and, total yield per plant (Kg) were calculated at the acceptable marketing stage.

Nitrogen, phosphorous and potassium content in cucumber fruits were analyzed according to the methods described by (Miller and Keeny, 1982), (Arnold et al 1992) and (Naseem, 2003) respectively.

RESULTS AND DISCUSSION

Vegetative growth

Plant height

Fertilization had a significant effect on the plant height of cucumber plants during the two successive winter seasons of (2010 and 2011). (Table 3). Comparing treatments with 100% organic fertilizers and bio-organic fertilizers, the highest plant height of cucumber were in the plants treated with the compost compared with the rest of organic fertilizer and control (Table 2). No significant differences were found between chicken and cows manure treatments since bio-organics applied at the harvest stages. In addition control did not reflect any negative effect on plant height at two successive seasons.

Mycorrhizal treatments and compost adversely affected plant height with highly significant results compared with phosphobacterin and Azotobacter treatments.

However, using bio-organics treatments had a positive effects on plant height compared with control.

Plants grown with compost +mycorrhizal treatments were superior in plant height compared with any other treatments included in the experiment and control (Table 3).

This treatment gave 69.2 and 70.5 cm compared with control 53.7 % 50.3 cm increment in plant height over seasons 2010 and 2011 respectively. This means that compost + mycorrhizal fertilizers are quite favorable for the plant height on cucumber plants growing in winter seasons.

These results agreed with those obtained by Abdel-Maniem et al (2002) and Abou-Hadid et al (2003).

Plant fresh weight (Kg)

Bio-and organic fertilization had a significant effect on the average plant fresh weight of cucumber plants for the two successive winter seasons of 2010 and 2011 (Table 4).

Plant fresh weight treated with 100% compost + mycorrhizal gave the highest results than in the other treatments of organic fertilization especially in the plote treated with chemical fertilizers only (control). The vules were 2.06 & 2.10kg/plant in two successive seasons respectively. This might be due to the availability of water and nitrogen provided by compost to the root system of the plants. These results were in accordance with those Abou-Hadid et al (1993) and Frank, J. (1976).

Number of fruits/plant

Interaction between the compost and microbial is the key objective to solve the conflict between high yields and environmental protection. The fruits number / plant in the 100% compost + 22 grame-mycorrhizal in cucumber plot treatment were higher than in the other treatments throughout the experiment (Table 5).

Role of organics in increasing number of fruits/plant of cucumber was attributed to supply of all essential nutrients due to continus mineralization of organic manures. Our results also showed that the cucumber responded more to compost than the other organic manure + bio-organics has been treated in our experiments.

Table 3. Effect of using bio- and organic fertilizers on plant height (cm) of Queen cucumber cv (*Cucumis saliva*.) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	55.8 e-f	56.1 ef	58.2 de	56.70 C
Compost	65.4 ab	66.0 ab	69.2 a	66.99 A
Chickens manure	64.4 be	63.2 be	62.5 be	63.39 B
Cows manure	62.7 be	60.5 cd	63.3 be	62.20 B
Control	51.9 gh	50.4 h	53.7 f-g	52.00 D
Mean	60.1 AB	59.26 B	61.41A	
Second season				
Horse manure	60.30 a-c	58.40 b-d	63.20 ab	60.63 AB
Compost	66.50 ab	63.00 ab	70.50a	66.67 A
Chickens manure	57.40 b-d	55.70 b-d	62.40 ab	58.50 B
Cows manure	61.20 a-c	60.20 a-c	65.00 ab	62.13 AB
Control	47.30 d	48.50 d	50.30 cd	48.70 C
Mean	58.54 AB	57.16 B	62.28 A	

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal.

Table 4. Effect of using bio- and organic fertilization on plant fresh weight (Kg) of Queen cucumber cv (*Cucumis sativa*.) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	1.67 ef	1.67 ef	1.77 d-f	1.70 C
Compost	2.08 ab	1.92 c-d	2.18 a	2.06 A
Chickens manure	1.85 c-e	1.63 f	1.92 b-d	1.80 C
Cows manure	1.93 b-d	1.86 c-e	2.02 a-c	1.93 B
Control	1.21 g	1.17 g	1.30 g	1.23 D
Mean	1.75 B	1.65 C	1.84 A	
Second season				
Horse manure	1.53 d-f	1.74 cd	1.62 de	1.63 B
Compost	2.00 a-c	2.00 a-c	2.30 a	2.10 A
Chickens manure	1.77 cd	1.53 d-f	1.85 b-d	1.72 B
Cows manure	2.00 a-c	1.76 cd	2.17 ab	1.98 A
Control	1.11 kg	1.31 e-g	1.27 fg	1.23 C
Mean	1.68 B	1.67 B	1.84 A	

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal

**Effect of Using Bio – and Organic Fertilizers on Productivity and Quality of Cucumber Crop 17
Under Plastic Houses**

Table 5. Effect of using bio- and organic fertilizers on number of fruit / plant of Queen cucumber cv (*Cucumis sativa.*) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	24.74 ef	21.76 fg	25.67 de	24.04 C
Compost	29.71 ab	28.25 b-d	31.67 a	29.90 A
Chickens manure	26.80 b-e	25.46 de	26.42 b-e	26.25 B
Cows manure	26.08 c-e	26.50 b-e	29.21 a-c	27.22 B
Control	21.09 g	20.88 g	20.42 g	20.80 D
Mean	25.68 AB	24.57 B	26.67 A	
	Season 2011			
Horse manure	26.33 c-e	23.50 e-g	24.67 e-f	24.83 C
Compost	28.67 b-d	27.50 b-e	33.00 a	29.72 A
Chickens manure	31.00 ab	23.50 e-g	27.33 b-e	27.28 B
Cows manure	25.00 d-f	27.33 b-e	30.00 a-c	27.44 AB
Control	21.00 fg	19.67 g	19.33 g	20.00 D
Mean	26.40 A	24.30 B	26.87 A	

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal.

Average Fruit weight (gm)

Growing cucumber plant in the compost + 22 gram mycorrhizal / plote gave significantly greater fruit weight compared to the other organic treatments used in both successive seasons (Table 6).

Similarly, other levels gave significantly higher fruit weight in the control which included the chemical mineral only. This might be due to the availability of water and nitrogen provided to the roots system of the plants by using the bio-and organic fertilization. The best results was on compost + mycorrhizal treatments in the second season than the first season, values were 120.3 and 115(gm) respectively.

These results could be explained through the effect of the availability of organic fertilization on average fruit weight (Bradley and Flemig, 1958). Also these results agree with the findings of Shahein and El-Habbasha (1988), which indicated that the average fruit weight significantly differed within different evaluated cucumber hybrids.

Total yield / plant (kg)

The lowest total yield was obtained when cucumber plants were grown in control level (Table 7).

It seems that bio - and organic fertilizers are essential factors for cucumber plants production. The compost + 22 (gm) mycorrhizal/plote had positive effect on total yield of cucumber, it gave higher yield than other treatments and control. Using the chemical mineral only (control), decreased the total yield during the two successive seasons. The highest total yield was obtained from compost + micorrhizal on second season 2010 which gave 3.96 kg/plant.

Role of organics increasing yield of cucumber was attributed to supply of all essential nutrients due to continuous mineralization of organic manures and compost.

Our results also showed that the cucumber responded more to plant compost than the animal manures. This because of the fact that the low C/N ratio, phosphorous and EC of the plant compost compared to the other used manures.

Table 6. Effect of using bio- and organic fertilization on average fruit weight (g) of Queen cucumber cv (*Cucumis sativa.*) during seasons 2010 and 2011

Treatments	Season 2010			Mean
	Phos.	Aoz.	Mr.	
Horse manure	91.6 c-e	87.4 c-e	94.5 b-e	91.17 C
Compost	105.1 ab	95.0 b-e	115.0 a	105.0 A
Chickens manure	90.8 c-e	85.0 de	96.3 b-d	90.70 C
Cows manure	97.8 be	91.6 c-e	104.5 ab	97.97 B
Control	86.7 c-e	83.3 e	88.9 c-e	86.30 C
Mean	94.39 B	88.47 C	99.85 A	
Season 2011				
Horse manure	96.30 b-e	85.40 ef	93.30 c-e	91.67 B
Compost	100.8 be	93.40 c-e	120.3 a	104.8 A
Chickens manure	87.53 d-f	91.10 c-f	99.40 b-d	92.68 B
Cows manure	90.50 c-f	90.70 c-f	107.2 b	96.13 B
Control	80.50 f	79.40 f	80.10 f	80.00 C
Mean	91.13 B	88.00 B	100.1 A	

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal

Table 7. Effect of using bio- and organic fertilization on total yield/plant (Kg) of Queen cucumber cv (*Cucumis sativa.*) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	2.27 c-e	1.90 ef	2.43cd	2.20C
Compost	3.11 b	2.70 be	3.64 a	3.15A
Chickens manure	2.45 cd	2.16 d-f	2.54cd	2.39C
Cows manure	2.56 cd	2.43 cd	3.05 b	2.69 B
Control	1.83 ef	1.73 f	1.82 ef	1.80 D
Mean	2.45 B	2.20 C	2.70 A	2.45 B
Second season				
Horse manure	2.41 cd	2.05ed	2.35 c-e	2.27ce
Compost	2.89 b	2.61 be	3.96 a	3.15 A
Chickens manure	2.71 c	2.13dR	2.67 cd	2.50 c
Cows manure	2.26 d	2.48 cd	3.21 b	2.65 B
Control	1.69 c	1.62 f	1.61ef	1.64 d
Mean	2.39 B	2.18 c	2.76 A	2.44

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal

These results were in accordance with those of **Takeshita et al (1978), Shaheennd El-Habbasha (1988)**. They found that the growing cucumber under condition of high humidity, moisture and fertilizer had positive effect on total yield of cucumber plants.

Chemical Analysis

Nitrogen

**Effect of Using Bio – and Organic Fertilizers on Productivity and Quality of Cucumber Crop 19
Under Plastic Houses**

Table 8. Effect of using bio- and organic fertilization on Nitrogen content in fruits (%) of Queen cucumber cv (*Cucumis/ sativa.*) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	2.70 b-e	2.30 d-f	3.07 a-d	2.69 B
Compost	3.40 ab	2.90 a-e	3.30 a-c	3.20 A
Chickens manure	3.10 a-d	2.60 b-e	3.20 a-c	2.97 AB
Cows manure	3.60 a	3.00 a-d	3.30 a-c	3.30 A
Control	2.10 of	1.70 f	2.50c-e	2.10 C
Mean	2.90 A	2.50 B	3.07 A	
Second season				
Horse manure	2.5cd	2.5cd	3.0b	2.67 B
Compost	3.6a	3.1b	3.5a	3.40 A
Chickens manure	3.0b	2.7bc	2.8bc	2.83 B
Cows manure	3.8a	3.1b	3.0b	3.30 A
Control	2.4cd	2.2d	2.7cd	2.33 C
Mean	3.06A	2.72A	2.94A	

Means having the same letter (S) are not significant at 5% level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal

Table (7) shows nitrogen content of cucumber fruits as a result of bio-and organic fertilizer treatments.

Adding bio-and organic fertilizers to cucumber plant significantly increased the fruits nitrogen contents as compared to the control in the two successive seasons. The results showed that, with using the cows manure + phosphobacterin (22 gm/plote). Gaves the highest fruits nitrogen contents during the two seasons, the values was 3.6% and 3.8% from every 100 fruit dry weight respectively.

It can be due either to the effect of cows manure and phosphobacterin on improving soil physical properties, or to a higher mineralization of cows manure which it due to mineral N in puts. It increased plant availability of macro-and micro nutrients which led to high vegetative growth and more absorption of fruits nitrogen content. The lowest fruits nitrogen content was obtained when plants were grown in control without bio-and organic fertilization.

These results hold good during the two successive seasons. **Hunter, (1977)** reported that nitrogen nutrition appears to be an important factor affecting fruit quality, but perhaps not in total yield.

Phosphorus

The average of phosphorus content in cucumber fruits of bio - and organic fertilizers treatments increased by 80% in cows manure mixed with mycorrhizal 50% mixed with Azotobacter and 45% mixed with phosphobacterin compared with control treatment for the two successive winters seasons of 2010 and 2011 (**Table 9**).

The highest P content in fruits was obtained with cows manure + 22 (gm) / plotemycorrhizal.

The P content in fruits decreased by using control only without any organic fertilizers used. The increase of P content on cucumber fruits appeared to be more obvious when the bio-and organic fertilization was mixed and added to the sandy soil which growing cucumber plants.

These results confirmed those obtained by **Huang and Lin (2001); Zhang and Fang, (2007)** who found that application of organic fertilizers improved the yield of crops, also can correct any deficit of P in the cucumber plants.

Potassium content

Potassium content on cucumber fruits in the two successive seasons are presented in **Table (10)**.

Table 9. Effect of using bio- and organic fertilization on Phosphorus content in fruits (%) of Queen cucumber cv (*Cucumis sativa*.) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	0.79 bc	0.67 bc	0.86 a-c	0.77 B
Compost	0.73 bc	0.83 a-c	0.98 ab	0.85AB
Chickens manure	0.80 be	0.80bc	0.80 be	0.80 AB
Cows manure	0.87 a-c	0.91 a-c	1.12 a 0.97 A	0.97A
Control	0.60 c	0.64 c	0.62	0.72B
Mean	0.76B	0.77B	0.87	
Second season				
Horse manure	0.80bc	0.72b-d	0.91ab	0.81BC
Compost	0.80 be	0.90ab	1.1a	0.92AB
Chickens manure	0.79 b-d	0.75 b-d	0.75 b-d	0.76C
Cows manure	0.79 b-d	0.96 ab	1.1 a	0.95 A
Control	0.54 d	0.59 cd	0.61 cd	0.58 d
Mean	0.74 B	0.78 B	0.89 A	

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal

Table 10. Effect of using bio- and organic fertilization on Potassium content in fruits (%) of Queen cucumber cv (*Cucumussativa*.) during seasons 2010 and 2011

Treatments	First season			Mean
	Phos.	Aoz.	Mr.	
Horse manure	4.9 c-e	4.8c-e	5.7 be	5.13 B
Compost	5.5 b-d	5.1 c-e	6.7 ab	5.77 AB
Chickens manure	5.0 c-e	5.1c-e	6.0 be	5.37 B
Cows manure	5.6 bc	5.4 cd	7.2 a	6.07A
Control	4.1 e	4.0 e	4.3 de	4.13 C
Mean	5.02 B	4.88 B	5.98 A	
Second season				
Horse manure	5.3 ab	5.0 ab	6.1 ab	5.5AB
Compost	5.7 ab	5.4 ab	5.9 ab	5.7AB
Chickens manure	5.3 ab	5.0 ab	6.3 ab	5.5 AB
Cows manure	6.0 ab	5.7 ab	6.9 a	6.2 A
Control	4.7 ab	4.4 ab	4.1b	4.4 B
Mean	5.4 B	5.1 C	5.9 A	

Means having the same letter (S) are not significant at 5 % level (Phos) phosphobacterin & (Aoz) Azotobacter & (Mr.) mycorrhizal

Effect of Using Bio – and Organic Fertilizers on Productivity and Quality of Cucumber Crop 21 Under Plastic Houses

The average percent of K content on cucumber fruits of cows manure + 22 (gm) plotemycorrhizal increased by 68% and 69% in two successive winter seasons (2010 and 2011) respectively.

The highest k content on fruits was obtained with cows manure + 22 (gm/ plote) mycorrhizal compared with the other treatments on the first season. The k content on fruits was decreased with control treatment which using chemical mineral only. Using bio-and organic fertilizers increased plant availability of macro and micro nutrients which led to high vegetative growth and more absorption of potassium.

Conclusion

The use of chemical fertilizers for cucumber plants can lead to nitrate an accumulation and ground water pollution. This study confirmed that the application of bio-and organic fertilizers increased accumulation of organic nitrogen, phosphorous and potassium more than application of minerals fertilizers. Compost and cows manures +mycorrhizal fertilizers were the best management system for increasing soil fertility, cucumber yield and quality and decrease the cost of minerals fertilizers.

REFERENCES

- Abdel-Kawy, T.F.A. 2003. The effect of some organic and in organic fertilizer on plant growth and fruit quality of cucumber plant under plastic house, M.Sc. Thesis faculty of Agric. Ain Shams Univ. Egypt. p. 53.
- Abedel-Moniem, E.M.; Abedel-Kawy, T.F.; El-Behairy, U.A. and Abou-Hadid, A.F. 2002. Influence of some organic fertilizers on the yield of cucumber plants under plastic house, *Egyptian Jour. Of hort.* 2002. Publ 2004 29 (3/4) pp. 499-511.
- Abou-Hadid, A.F.; Gul, A.; Tuzel, I.H. and Ongun, A.R. 2003. Organic cucumber production under green house conditions. *ActaHorti*: (608) pp.149-157.
- Abou-Hadid, A.F.; Medany, M.; Khalifa, H.; El-Beltagy, A. 1993. Response of growth the development of cucumber and sweet pepper to heating effectiveness *Egyptian Journal of Horticulture*, 18(2): 185-197.
- Arnold, E.; Greenberg, A.P.H.A.; Lenore, G.; Weff, A. and Eaton, W.A. 1992. Standard methods for the exanination of water and waste water p. 115.
- Bradly, G.A. and Fleming. J.W. 1958. The effect of nitrogen, phosphorus and potassium application on the quality of water, *J. Am. Soc. Hort. Sci. No. 73 p. 341.*
- Esawy, M.; Abd El-Kader, N.; Robin, P.; Akkal, N. and Abd El-Rahman, L. 2009. world *Journal of Agric Sci.*, 5(4): 408-414.
- Frank, J. 1976. European Greenhouse cucumbers, growing in spring proceedings of the annual, western green house, vegetable growers conference, July 1978. **Fresno California USA. No. 9 pp. 43-51.**
- Hekal, A.I.A. 2001. Evaluation of some cucumber hybrids for resistance to powdery mildew disease under Egyptian environmental conditions. Msc, thesis environ, Sci, Agric., Institute of Environ, studies and Res, Ain Shams Univ., Egypt p. 1.
- Huang, S.N. and Lin, J.C. 2001. Current status of organic materials recycling in southern Taiwan. *Soil and fertilizer experiment bulletin*, 3, pp. 43-48.
- Hunter, J.C. 1977. Studies on fruit curcature in European green house cucumbers proceeding of the annual, western green house vegetable growers conference. July 1978, fresno. California USA. No. 5 pp. 23-29.
- Kostov, O.; TZ vetkov, Y.; Kaloianova, N. and van cleemput, O. 1995. cucumber cultivation on some wastes during their aerobic composting. *Biore Source Technology*, 53(3): 237-242.
- Maslo, A.V. and Gamayunov A.T. 1989. The effect of prolonged application of fertilizers on the fertility of grey prodcolized soil and the productivity of a vegetable crops. *Agrokhimiya No. (1) pp. 73-76.*
- Miller, R.H. and Keeney, D.R. 1982. Method of soil analysis part (2).
- Naseem, M.G. 2003. Methods of soil analysis Book shop mainsheet El-Maarf, Alex., Egypt. p. 200.
- Regab, M.A. 1984. Studies on the inheritance of some economic characters in cucumber "*cucumissativus L.*" ph.D. Thesis. Faculty of Agriculture Mansoura University, Mansoura, Egypt.
- Roe, N.; Stoffella, P.J. and Graetz, D. 1997. Composts from various municipal solid waste feeds tocks affected vegetable crops 2: Growth, yields and fruit quality. *J. Amer. Soc. Hort. Sci.*, 122: 433-437.
- Shaheen, A.M. and El-Habasha, K.M. 1988. Biological and marketable yield often cucumber "*cucumissativus L.* Ph.D. Thesis. Faculty of

Agriculture Mansoura University, Mansoura, Egypt.

Takeshita, S.; Kato, M. and Kato, K. 1978. The chemical properties and microflora of the soil associated with the yield of vegetable in a plastic green house. Bulletin of the Kanagawa Horticultural Experiment Station No. 25: pp. 69-75.

Vogtmann, H.; Matthies, K.; Kehres, B. and Aeiroploeger, A. 1993. Enhanced food quality; Effects of compost on the quality of plant foods. **Compost Sci. Util. 1: pp. 82-100.**

Zhang, M. and Fang, L. 2007. Effect of tillage, fertilizer and green manure cropping on soil quality at an abandoned brick making site. **Soil and Tillage Research 93: 87-93.**



تأثير استخدام الأسمدة العضوية والحيوية على إنتاجية وجودة محصول الخيار داخل الصوب البلاستيكية

[٢]

على حسن الحمودي^١ - محمد هاشم محمد^٢ - صلاح الدين محمود المنياوي^٢ -
صلاح عبد المنعم حسين^١

١- وزارة البيئة والمياه - دولة الإمارات العربية المتحدة

٢- قسم البساتين - كلية الزراعة - جامعة عين شمس - القاهرة - مصر

كان أعلى قيم للوزن الطازج باستخدام الكمبوست مخلوط مع الميكروهيذا يليه الكمبوست المخلوط بالأزوتوباكتري.

وبالنسبة لعدد الثمار على النبات أعطت المعاملة بالكمبوست المخلوط بالميكروهيذا أعلى القيم (٣١,٧ ، ٣٣ ثمرة / نبات) خلال موسمي الزراعي الأول والثاني على التوالي. بينما أعطت معاملة الكنترول أقل القيم خلال موسمي الزراعة.

بالنسبة لصفة وزن الثمرة أعطت المعاملة بالكمبوست المخلوط مع الميكروهيذا أعلى القيم (١٢٠,٣ ، ١١٥ جرام/ثمرة) خلال موسمي الزراعة الثاني والأول على التوالي.

وأوضحت النتائج تفوق استخدام الكمبوست المخلوط بالميكروهيذا في الحصول على أعلى محصول من الخيار للنبات يلي ذلك استخدام الكمبوست المخلوط بالفسفوبكتريين خلال موسمي الزراعة.

بالنسبة لمحتوى الثمار من النيتروجين كانت أعلى القيم المتحصل عليها من خلال استخدام سماد الأبقار المخلوط بالفسفوبكتريين كنسبة مئوية (٣,٦% ، ٣,٨%) على التوالي خلال موسمي الزراعة ٢٠١٠، ٢٠١١م. وأوضحت النتائج أن استخدام سماد البقر المخلوط بالميكروهيذا اعطى أعلى قيمة بالنسبة لمحتوى الثمار من الفسفور والبوتاسيوم كنسبة مئوية.

الكلمات الدالة: أسمدة عضوية - أسمدة حيوية - صوب بلاستيكية - خيار

الموجز

يعتبر الخيار من أهم محاصيل الخضار التي تزرع في العالم وفي دولة الإمارات العربية المتحدة من أجل الاستهلاك المحلي أو التصدير للخارج حيث انه يعتبر مصدر جيد من الفيتامينات والمعادن والألياف. ولقد أجريت الدراسة بمحطة تجارب المنطقة الشمالية التابعة لوزارة البيئة والمياه لمدة موسمين زراعيين متتاليين خلال عامي ٢٠١٠، ٢٠١١م بهدف دراسة تأثير استخدام الأسمدة العضوية والبيولوجية على إنتاجية وجودة الخيار كبديل للزراعة الكيماوية داخل الصوب البلاستيكية.

وتم استخدام أربعة أنواع من الأسمدة العضوية وهي سماد الخيول، سماد الأبقار، سماد الدواجن، بالإضافة إلى الكمبوست. بالإضافة إلى ثلاثة أنواع أسمدة بيولوجية وهي الفوسفوبكتريين، أزوتوباكتري والميكروهيذا. كما استخدم السماد الكيماوي للمقارنة. وخلصت النتائج المتحصل عليها في زيادة طول النبات باستخدام السماد العضوي كمبوست مخلوط مع الميكروهيذا يليها الكمبوست مع الفوسفوبكتريين بينما

تحكيم: أ.د. أحمد أبو اليزيد

أ.د. فتحى أبو النصر سديرة