

**EFFECT OF TIMES AND METHODS OF POTASSIUM FERTILIZER ON
YIELD AND YIELD COMPONENTS OF NEW HYBRID COTTON (GIZA 86 X
10229) UNDER EARLY AND LATE SOWING**

Abdel-Aal, S.A. Amal; M.A.A Emara and S.A.F. Hamoda

Department of Agronomy, Cotton Research Institute, Agriculture Research Center,
Giza, Egypt

ABSTRACT

Two field experiments were carried out at Sakha Agricultural Research Station at Kafr El-Sheikh Governorate, Egypt, on new hybrid cotton (Giza 86 X 10229) during 2013 and 2014 seasons to study the effect of sowing dates (15 April and 15 May) as main plots and five methods and time of potassium sulphate application as sub plots i.e:

- 1- Soil application of 24 kg K₂O/fed. as potassium sulphate (48% K₂O) after thinning.
- 2- Soil application of 24 kg K₂O/fed. as potassium sulphate (48% K₂O) at two weeks after thinning.
- 3- Foliar application of 5 kg potassium sulphate (48% K₂O)/fed. sprayed two times at the square and start flowering stages.
- 4- Foliar application of 5 kg potassium sulphate (48% K₂O)/fed. sprayed two times at the start and peak of flowering stages.
- 5- Foliar application of 5 kg potassium sulphate (48% K₂O)/fed. sprayed three times at square, start and peak of flowering stages. The most important results obtained could be summarized as follows:
 - 1) Results obtained revealed that early sowing (15 April) significantly increased no. of internodes and sympodia on the main stem, no. of open bolls/plant, boll weight, seed cotton yield/plant and /feddan, dry weight of vegetative and fruiting organs of plant, days to first flower, days to first open boll and earliness percentage, while plant height, no. of more monopodial/plant and first sympodium node were decreased.
 - 2) Results indicated the positive effect of potassium when it was applied at the proper time as soil application at two weeks after thinning which resulted in less no. of monopodia and higher no. of sympodia, higher dry weight of vegetative and fruiting organs, heaviest bolls, and higher no. of open bolls as well as seed cotton yield/plant and /feddan. Potassium fertilization showed no effect on no. of main stem internodes/plant, first sympodium node, days to first flower, days to first open boll, seed index, earliness percentage and lint percentage.
 - 3) The interaction of planting date by K treatments was not significant for all traits studied in this investigation.
 - 4) Generally, results obtained revealed that early sowing (15 April) and the soil application of potassium sulphate (24 kg/feddan) fertilizer at two weeks after thinning for obtaining high productivity of new hybrid cotton (Giza 86 x 10229).

KEY WORDS: Cotton, Sowing dates, Potassium sulphate, Fertilizer, Growth, Earliness and Yield.

INTRODUCTION

Cotton is not only the king of fibers and crucial crop used for fiber production all around the world (**Killi and Aloglu, 2000**) but also it is a vital source of foreign exchange earnings. The suitable sowing date and nutrients play a vital role in cotton production, where the early sowing date is one of the most important management factors involved in producing high yielding and quality (**Dong et al., 2006** and **Bozbek et al., 2006**). The second one provides the cotton plants with one of the major feeding elements.

Applying different sowing dates expose the cotton plants to different temperature responses, which affecting the optimum requirements of the whole plant, (**Woodward and Sheely, 1983**). **Young et al (1980)** cleared that early sowing fits the cotton plant to full season in order to obtain complete thermal units regime. **McMahon and Low (1972)** mentioned that planting cotton in a suitable time leads to forming the first fruiting branch at a lower node on the stem and only an optimum height, increasing no. of bolls/plant and seed cotton yield, escaping from leaf and boll-worms and aphids at the end of the season and early picking. **Boquet et al. (2003)** showed that the excessive plant height at late planting date was partly responsible for lower yield as the crop used a larger portion of its energy budget for vegetative growth and the excess plant height caused lodging. Seed cotton yield/fed. was significantly decreased with delayed planting.

Early sowing appears higher yield potential and alternately, late planting of cotton crop shows very vegetative and difficult to manage resulting in lower seed cotton yield as well (**Ali et al., 2009**). However, several reports have indicated that early sown cotton produces taller plants with higher no. of branches, no. of bolls, seed cotton yield (**Pettigrew, 2006, Arshad et al., 2007** and **Bange et al., 2008**). These findings are also supported by other researchers **Emara et al., (2006)**, **El-Shahawy and Hamoda (2011)**, **Abdul Wahab et al., (2014)** and **Hamoda et al., (2014)** where they found that early planting date significantly increased seed cotton yield/fed. due to the increase of no. of open bolls/plant and boll weight. The planting date treatments did not exhibit significant effect on lint %.

Potassium (K) is a unique plant nutrient for cotton because of its continuous need through all growth stages and its relatively high uptake rate. Yet the cotton plant is a relatively inefficient as a K absorber. Potassium is an important nutrient that has favorable effects on the metabolism of nucleic acids, proteins, vitamins and growth substances. Furthermore, K plays an important role in the translocation of photosynthates from sources to sinks (**Bednarz and Oosterhuis, 1999** and **Morteza et al., 2005**). However, recently positive response of cotton plants to potassium fertilizer has been documented comparing with unfertilized plants with potassium fertilization (**El-Sayed and El-**

EFFECT OF TIMES AND METHODS OF POTASSIUM FERTILIZER 15
 Menshawy 2001, Kassem and Ahmed 2005, El-Sayed *et al.* 2006, Sary *et al.* 2008 and Abou-Zaid *et al.*, 2009). Many studies have shown increased yield and cotton productivity in response to potassium fertilization as reported by Emara (2012), Sema *et al.*, (2012), Abdel-Aal *et al.*, (2014), Sawan (2014), Gomaa *et al.*, (2014) and Emara (2014).

The aim of this investigation is to evaluate the response of the new hybrid cotton (Giza 86 X 10229) to different application methods and time of potassium sulphate under early and late sowing dates.

MATERIALS AND METHODS

Two field experiments were carried out at Sakha Agricultural Research Station at Kafr El-Sheikh Governorate, Egypt, during 2013 and 2014 seasons using the new hybrid cotton (Giza 86 X 10229) belonging to *Gossypium barbadense*, L. Characterized the new hybrid cotton are showed in Table (1).

Table (1): Characterized the new hybrid cotton (Giza 86 x 10229)

Hybrid name	New promising hybrid cotton (Giza 86 x 10229).
Species	Barbadense.
Category	Long staple and extra fine.
Pedigree	Crossing between G86 x 10229.
Characteristics	Long staple characterized by high yielding, early maturity, resistance to Fuzarium and high lint (%).
Botanical distinguishing characters	The stem has a medium length with polygon shape also has green color mixed by dim red with medium length internodes. The leaves have palmate shape with large size with no deep lobes and leather fell. The node of the first fruiting branch ranged from 8 - 9. A flower petal has tubular shape. The boll size is large and pyramid shape with drawn summit. Seed is big-sized and the fuzz covers about fuzz less to ¼ from the whole size and fuzz color is gray-greenish.
Hybrid bred by	Breeding Res. Section, Cotton Res. Inst., Agric. Res. Center, Giza, Egypt.

Each experiment was laid out in a split plot design with four replications. The main plots were assigned to the two sowing dates i.e.; April 15 (early) and May 15 (late). While, the sub-plots were allocated to four applications of potassium i.e.;

- 1) Soil application of 24 kg K₂O/fed. as potassium sulphate (48% K₂O) after thinning.
- 2) Soil application of 24 kg K₂O/fed. as potassium sulphate at two weeks after thinning.
- 3) Foliar application of 5 kg K₂O/fed. as potassium sulphate sprayed two times at the square and start flowering stages.

- 4) Foliar application of 5 kg K₂O/fed. as potassium sulphate sprayed two times at the start and peak of flowering stages.
- 5) Foliar application of 5 kg K₂O/fed. as potassium sulphate sprayed three times at square, start and peak of flowering stages.

The sub-plot size was 18 m² including 6 rows (5 m long and 60 cm width). Nitrogen fertilizer in the form of ammonium nitrate (33.5%N) was added in bands and divided into two equal doses, the first, one was applied after thinning just before the first irrigation and the second part before the second irrigation. All other cultural practices were done as recommended in cotton production that is involved as basic dose of 150 kg calcium superphosphate (15.5% P₂O₅) at land preparation. Potassium was added to sub-main plots according to the experimental treatments (type, rate and time of application). Soil samples were taken in the two seasons before planting cotton to estimate the soil characters using the standard methods as described by Chapman and Parker, (1981). The results are shown in Table (2). In both seasons, the soil texture was clay loam, low content of organic matter, very low calcium carbonate and non-saline. The available amounts of macro- elements were low for nitrogen, phosphorus and potassium.

Table (2): Mechanical and chemical analysis of the experiment soil in 2013 and 2014 seasons.

Seasons	Properties Texture	pH	Organic Matter (%)	TSS (%)	Bicarbona te (%)	Available element (ppm)		
						N	P	K
2013	Clay loam	7.70	1.69	0.64	1.82	12.10	9.21	131.2
2014	Clay loam	8.38	1.74	0.69	1.81	11.95	9.50	126.2

Five representative hills were chosen by random from the four inner rows in order to study the following characters:

- A. **Growth characters:** Final plant height (cm), no. of main stem internodes/plant, no. of monopodia/plant and no. of sympodia/plant.
- B. **Dry weight:** A random sample of five plants were taken at 130 days from sowing. All plants were carefully uprooted, washed hard then floated in water bath for final separation from the muddy medium. All plant parts were dried in an air forced oven at 90°C to a constant weight. The following data were recorded; dry weight of leaves/plant (g), dry weight of stem/plant (g) and dry weight of reproductive organs of plant; squares, flowers and bolls (g).
- C. **Earliness measurements:** First sympodial node, days to first flower, days to first open boll and earliness percentage.
- D. **Yield components:** No. of open bolls/plant, boll weight (g), seed cotton yield/plant, no. of plants at harvest/fed., lint percentage, seed index (g) and

EFFECT OF TIMES AND METHODS OF POTASSIUM FERTILIZER 17
 seed cotton yield (kantar/feddan) were estimated from picking all plants of four inner rows of each plot.

The climatic records included minimum and maximum air temperatures (°C) and relative humidity (%) for Sakha Weather Station through the two growing seasons are given in Table (3).

Table (3): Monthly air temperature (°C) and relative humidity (%) during 2013 and 2014 seasons

Month	2013 season				2014 season			
	Air temp. (°C)		RH (%)		Air temp. (°C)		RH (%)	
	Max.	Min	Max.	Min	Max.	Min	Max.	Min
March	26.8	11.8	86.3	49.5	27.0	12.0	85.9	49.0
April	31.5	15.0	85.0	47.7	30.9	14.6	86.0	48.0
May	34.0	18.0	76.7	38.0	33.8	17.9	70.0	39.0
June	37.0	22.2	82.5	46.5	38.1	23.1	82.9	47.1
July	38.0	24.4	79.5	49.2	39.2	24.5	78.7	48.9
August	37.5	24.2	84.0	50.0	38.3	24.7	83.7	51.1
September	38.2	22.7	88.0	49.6	39.2	23.1	87.3	50.0
October	33.0	19.0	82.0	48.0	32.8	20.1	81.3	48.5

The data obtained were subjected to statistical analysis according to procedure outlined by **Snedecor and Cochran (1981)** by using LSD at 5% level.

RESULTS AND DISCUSSION

A. Plant growth characters:

The data presented in Table (4) showed that, final plant height and no. of monopodia/plant were significantly increased in favour of late sowing treatment, while no. of internodes on the main stem and no. of sympodia/plant were significantly decreased. Regarding to potassium fertilizer treatments, the data illustrated in Table (4) showed that, final plant height, no. of monopodia/plant and no. of sympodia/plant were significantly affected by potassium treatments in both seasons. On the other hand, it is apparent that potassium application treatments had insignificant effect on no. of main stem internodes/plant. Adding potassium as soil application at two weeks after thinning produced the tallest plants with high no. of sympodia. The results in the same table indicate that the early application of potassium 24 kg K₂O/feddan at two weeks after thinning relatively prolonged the vegetative growth period. **Halvey (1976)** demonstrated that maximum amount of the K is taken up by cotton plants between the period of 57 to 84 days after emergence (from early squaring to early blooming stage). Similar results were obtained by **Makram et al., (1994)** for plant height and **Ali and El-Sayed (2001)**, **Pettigrew (2006)**, **Arshad et al., (2007)** and **Bange et al., (2008)** for no. of main stem internodes/plant, no. of monopodia/plant, and no. of sympodial/plant.

Table (4): Means of some growth characters as affected by sowing dates, time and application methods of potassium and their interaction of 2013 and 2014 seasons.

Treatments Characters	Seasons	Sowing dates (S)			Potassium treatments (K)					Inter S x K	
		15 April	15 May	L.S.D	Soil after thinning	Soil 2w after thinning	2 Sprays Sq + F	2 Sprays F + 2F	3 Sprays Sq + F + 2w		L.S.D
Final plant height (cm)	2013	145.8	153.5	3.20	148.30	151.80	150.20	149.30	148.30	1.82	NS
	2014	146.2	153.9	4.10	150.00	151.70	148.00	150.50	150.00	0.87	NS
No. of main stem internodes/plant	2013	21.50	18.60	1.20	20.70	20.80	19.70	19.80	20.00	NS	NS
	2014	22.70	17.52	2.70	18.90	19.10	20.10	19.90	19.80	NS	NS
No. of monopodia/plant	2013	0.97	1.15	0.08	1.08	1.03	1.05	1.08	1.05	0.01	NS
	2014	1.16	1.25	0.01	1.22	1.10	1.20	1.20	1.22	0.02	NS
No. of sympodia/plant	2013	17.90	15.20	1.21	17.00	17.10	16.20	16.20	16.30	0.18	NS
	2014	18.16	14.82	2.13	16.55	16.87	16.58	16.13	16.32	0.19	NS

B. Dry matter weight:

Data presented in Table (5) showed that planting date and potassium treatments had a significant effect on the dry matter plant organs in the two seasons, early fertilized plants with potassium application as soil potassium 24 kg K₂O/feddan at two weeks after thinning increased the dry matter of plant organs. Dry weight production in early and late sowing as well as K absorption was affected in all plant parts. The dry matter accumulation is a good result to the favourable effect of this element on the photosynthesis activity of leaves, promotion of CO₂ assimilation and the translocation of carbohydrates from the leaves to the reproductive organs (Hart, 1969). It is worth to be mentioned here that Halvey (1976) reported that the maximum K⁺ uptake rate in cotton plants occurred at 57 to 84 days after emergence (early squaring to early blooming stage). Thus, the period of peak K⁺ uptake preceded the peak rate of dry matter production. Basset *et al.*, (1970) came to same conclusion. This may explain why K application before the second irrigation was found to be the proper time to provide the plant with K for building up the maximum dry matter.

Leaves dry weight/plant was significantly increased due to early sowing in both seasons. Late sowing resulted in rapid vegetative growth as compared to early sown plants which exposed relatively to low air temperatures (McMahon and Low, 1972). The increase in leaves dry weight is mainly due to the increase of leaves number of plant. From other point of view, cotton plants of earlier sowing harvested the highest amount of heat units through the growing season (Young *et al.*, 1980). This situation resulted in much formation of leaf assimilates necessary for increasing the fruiting capacity (Makram *et al.*, 2001).

Table (5): Means of dry weight of vegetative and fruiting organs as affected by sowing dates, time and application methods of potassium and their interaction of 2013 and 2014 seasons.

Treatments Characters	seasons	Sowing dates (S)			Potassium treatments (K)						Inter S x K		
		15 April		15 May	Soil after thinning		2 Sprays Sq +F		2 Sprays F +2F			3 Sprays Sq +F +2w	
		25.80	26.10	22.10	24.32	27.80	22.95	23.10	20.82	21.33		23.75	24.20
Leaves dry weight/plant (g)	2013	25.80	26.10	22.10	24.32	27.80	22.95	23.10	20.82	21.33	23.75	24.20	1.80
	2014	26.10	26.10	22.80	26.10	28.00	23.10	23.10	21.33	24.20	24.20	24.20	1.70
Stem dry weight/plant (g)	2013	27.00	28.40	25.31	26.30	27.20	25.80	27.60	24.00	25.60	24.50	26.80	2.30
	2014	28.40	28.40	26.92	28.00	28.85	27.60	27.60	25.60	26.80	26.80	26.80	1.10
Dry weight of fruiting organs/plant (g)	2013	12.19	12.25	10.25	12.32	13.80	10.60	11.00	9.65	10.55	10.11	9.75	1.72
	2014	12.25	12.25	10.60	12.00	13.95	11.00	11.00	10.55	10.55	9.75	9.75	2.10

NS = Not significant.

2w = Two weeks after thinning.

Sq = Square.

F = Flowering.

2F = Two weeks after flowering.

C. Earliness

Data given in Table (6) showed the effect of sowing dates, time and application methods of potassium and their interaction on some earliness measurements.

Table (6) cleared that, node of the first sympodium was significantly lowered in favour of early sowing while number of days from planting to first flower and open boll was significantly decreased by delaying the date of sowing but it still late in the calendar date of appearance for both characters than earlier sowing. This might be due to relatively low temperature of air at the beginning of the season for early sowing which induced the balance between vegetative growth and fruiting capacity, while in late sowing the increase of air temperatures directed the cotton plant development to vegetative growth. At harvest (first pick) most bolls in case of early sowing reached maturity age, in the same time few bolls of late planting reached maturity which was reflected on increasing earliness percentage in early sowing. These results are in line with those obtained by *Emara et al.*, (2006), *Ali et al.*, (2009), *El-Shahawy and Hamoda* (2011), and *Hamoda et al.*, (2014).

Table (6) cleared that, application of potassium gave insignificant effect on position of first sympodium, days to first flower and days to first open boll and earliness percentage. These results were in good agreement with those reported by *El-Sayed*, (2005).

D. Yield and yield components:

Results presented in Table (7) cleared the effect of sowing dates, time and application methods of potassium and their interaction on yield and yield components.

Results presented in Table (7) showed that both no. of open bolls/plant, boll weight and seed cotton yield/plant were significantly increased in favour of earlier sowing. Also, the yield of seed cotton/feddan was significantly increased as a result of increasing no. of open bolls/plant and boll weight, where the plant stand means at harvest remained in narrow range of differences, while lint percentage and seed index were not significantly affected in the two seasons. However, previous results obviously cleared that early sowing fits the cotton plants to full season in suitable climatic window in order to obtain complete thermal units requirements and gave enough time to develop a heavy boll load with large seeds (*Young et al.*, 1980), besides the exposure of cotton plant at different stages to suitable air temperature (*Makram et al.*, 2001). Similar results were obtained by *Emara et al.*, (2006), *Arshad et al.*, (2007), *Bange et al.*, (2008), *El-Shahawy and Hamoda* (2011), *Abdul Wahab et al.*, (2014) and *Hamoda et al.*, (2014).

Data in Table (7) show that no. of open boll/plant, boll weight and seed cotton yield/plant and /feddan were affected significantly by time and application methods of potassium, while lint percentage and seed index were not significantly affected in the two seasons. Data of seed cotton yield and yield components showed that the fertilization with soil potassium 24 kg K₂O/feddan at two weeks after thinning were highly significant. The increment in the two characters, i.e., seed cotton yield/plant and boll weight may be due to the role of potassium fertilizer in encouraging root hairs to grow early and increasing its elongations well as early appearance of bolls of cotton plants.

Table (6): Means of some earliness measurements as affected by sowing dates, time and application methods of potassium and their interaction of 2013 and 2014 seasons.

Treatments Characters	Seasons		Sowing dates)S(Potassium treatments)K(Inter S x K
	2013	2014	15 April	15 May	L.S.D	Soil after thinning	Soil 2w after thinning	2 Sprays + F	2 Sprays Sq + F	2 Sprays F + 2F	3 Sprays +F +2w	L.S.D		
			5.31	6.81	0.95								6.02	
First symodial node	5.24	6.79	1.01	6.02	6.02	6.02	6.02	5.93	5.98	6.13	6.03	NS		
Days to first flower	73.57	69.07	2.20	70.92	71.03	71.28	71.03	71.38	71.08	71.14	71.05	NS		
	73.19	69.05	1.85	71.03	71.03	71.03	71.03	71.08	71.08	71.14	71.05	NS		
Days to first open boll	120.45	115.87	1.92	117.93	118.27	118.27	118.27	118.08	118.08	118.10	118.42	NS		
	119.79	116.11	2.10	118.07	117.95	117.95	117.95	118.12	118.12	118.55	117.75	NS		
Earliness percentage (%)	79.67	70.64	3.81	75.73	75.97	75.97	75.97	74.23	74.23	74.52	75.32	NS		
	79.72	70.71	3.72	76.05	75.88	75.88	75.88	74.35	74.35	74.43	75.37	NS		

NS =Not significant.
 2w =Two weeks after thinning.
 Sq =Square.
 F =Flowering.
 2F =Two weeks after flowering.

Table (7): Means of seed yield (kentar/fed) (.and yield components as affected by sowing dates, time and application methods of potassium and their interaction of 2013 and 2014 seasons.

Treatments Characters	Seasons	Sowing dates)S(Potassium treatments)K(Inter S x K
		15 April	15 May	L.S.D	Soil after thinning	Soil 2w after thinning	2 Sprays +F	2 Sprays F +2F	3 Sprays +F +2w	L.S.D			
		17.00	14.03	1.31	15.50	15.60	15.35	15.50	15.40	0.08			
No .of open bolls/plant	2013	17.00	14.03	1.31	15.50	15.60	15.35	15.50	15.40	0.08	NS		
	2014	17.24	14.22	1.72	15.97	16.22	15.70	15.33	15.63	0.09			
Boll weight (g)	2013	3.02	2.68	0.07	2.89	2.95	2.78	2.78	2.67	0.12	NS		
	2014	3.10	2.81	0.02	3.00	3.06	2.90	2.86	2.95	0.09			
Seed cotton yield/plant (g)	2013	51.34	37.60	3.80	45.08	45.73	42.67	43.59	41.12	1.83	NS		
	2014	53.44	39.96	4.30	47.91	49.02	45.53	43.84	46.11	2.30			
Lint percentage (%)	2013	39.64	38.69	NS	39.48	39.31	38.84	39.09	39.11	NS	NS		
	2014	39.68	38.58	NS	39.18	39.17	39.04	39.05	39.22	NS			
Seed index (g)	2013	12.60	12.39	NS	12.41	12.48	12.55	12.52	12.53	NS	NS		
	2014	12.63	12.41	NS	12.54	12.53	12.43	12.48	12.58	NS			
No .of plants at harvest/fed.	2013	38.010	38.100	NS	38.030	38.799	37.999	38.056	38.040	NS	NS		
	2014	38.025	38.180	NS	38.100	38.200	37.913	38.225	38.080	NS			
Seed cotton yield/fed. (kentar)	2013	12.30	9.07	1.72	11.01	11.12	10.39	10.28	10.61	0.08	NS		
	2014	12.97	9.65	1.11	11.66	11.96	11.03	10.70	11.17	0.92			

NS =Not significant.
 2w =Two weeks after thinning.
 Sq =Square.
 F =Flowering.
 2F =Two weeks after flowering.

Regarding the proper time of potassium application, it had been found that soil application with potassium 24 kg K₂O/feddan at two weeks after thinning seemed to be the best timing. This might be due to the fact that most amount of potassium which needed by cotton plant is laying after flowering, while at early stages of growth, the seeding requirements is very low (Etourneand, 1995). Eatone and Ergle (1957) proved that total amount of K needed by cotton plants early in its growth is small. Similar results were obtained by El-Sayed and El-Menshawi (2001), Kassem and Ahmed (2005), Abou-Zaid *et al.*, (2009), Sema *et al.*, (2012), Sawan (2014), and Emara (2014).

The interaction between sowing dates and potassium treatments had no significant effect on all traits under study in both seasons.

CONCLUSION

Generally, the results obtained in this study could lead us to a package of recommendations, which seemed to be useful for increasing the cotton yield production in quantity and quality. It could be concluded the early sowing (15 April) and the soil application with potassium sulphate (24 kg/feddan) at two weeks after thinning for obtaining high productivity of new hybrid cotton (Giza 86 x 10229) under Sakha location.

REFERENCES

- Abdel-Aal, Amal, S.A.; S.A.F. Hamoda and M.A.A. Ibrahim (2014). Effect of different sources of potassium on growth and productivity of cotton. Egypt. J. of Appl. Sci., 29 (11): 562 – 572.
- Abdul Wahab, S.; F. Hussain, Ch. Abdul Razzaque, Z. Muhammad, S. Muhammad, K. Fakhar and B. Karim (2014). Effects of sowing time on yield, got and fiber traits of upland cotton. International J. of Scientific & Engineering Res., December, 5(12): 194 – 198.
- Abou-Zaid, M.K.; M.A. Emara and S.A. Hamoda. (2009). Future of Egyptian cotton production in the newly reclaimed desert land of Egypt: 10-Cotton response to soil, foliar potassium application and potassium dissolving bacteria (KDB). J. Adv. Agric. Res., (Fac. Agric. Saba Basha), 42(1): 73 - 80.
- Ali, H.; M. Afzal, S. Ahmad and D. Muhammad (2009). Effect of cultivars and sowing dates on yield and quality of *Gossypium hirsutum* L. Crop. J. Food Agri. Environ., 7: 244 - 247.
- Arshad, M.; A. Wajid, M. Maqsood, K. Hussain, M. Aslam and M. Ibrahim (2007). Response of growth, and quality of different cotton cultivars to sowing dates. Pak. J. Agric., 44(2): 208 - 212.
- Bange, M.P.; S.J. Caton and S.P. Milroy (2008). Managing yields of high fruit retention in transgenic cotton (*Gossypium hirsutum* L.) using sowing date. Australian J. Agric. Res., 59(8): 733 - 741.
- Basset, D.M.; W.D. Anderson and C.H. Werkhoven (1970). Dry matter production and nutrient uptake in irrigated cotton (*G. hirsutum*). Agron. J., 62: 299 - 303.

quality of new hybrid cotton [Giza 83 (Giza 75 x 5844)] x Giza 80. Egypt. J. Agric. Res., 92(1): 125 – 137. Special Issue. The 1st International Cotton Conference "Challenges to Sustainable Cotton Production & Quality". 25 – 26 Feb. 2014.

- Hart, C.E. (1969).** Effect of potassium deficiency upon translocation of G14 in attached blades and entire plants of sugarcane. *Plant Physiol.*, 44: 1461 - 1469.
- Kassem, M.A. and F.A. Ahmed (2005).** Effect of potassium source and level on cotton productivity and soil chemical properties. *Assiut J. Agric. Sci.*, 36(3): 77 – 90.
- Killi, F. and K. Aloglu (2000).** Determination of yield, yield components and technological properties of some cotton genotypes. *Proceedings of FAO Interregional Cooperative Research Network on Cotton, Adana Turkey*, 20 – 24 September, pp: 88 - 90.
- Makram, E.A.; H.A. Abd El-Aal; A.A. Darwish and W.M. El-Shazly (2001).** Air thermal units in relation to growth and development of cotton plants through different sowing dates. *Minufiya J. Agric. Res.*, 26(3): 659 - 671.
- Makram, E.A.; M.I. El-Shahawy; S.F. El-Gahel and R.R. Abd El-Malik (1994).** Effect of hill spacing, soil fertilization and its interaction on growth, yield and earliness in Egyptian cotton cultivar. *Giza 70. J. Agric. Sci., Mansoura Univ.*, 19(1): 1 - 13.
- McMahon, J. and A. Low (1972).** Growing degree-days as a measure of temperature effects on cotton. *Cotton Grow. Rev.*, 49: 39 - 49.
- Morteza, M.; A. Slaton, E. Evans, J. McConnell, M. Fred and C. Kennedy (2005).** Effect of potassium fertilization on cotton yield and petiole potassium. *Summaries of Arkansas Cotton Res.*, pp: 74 - 78.
- Pettigrew, W.T. and J.J. Adamezyk (2006).** Nitrogen fertility and planting date effects on lint yield and Bt Endotoxin production. *Agro. J.*, 98: 69 - 697.
- Sary, G.A.; A. Roshdy, O.M. Wassel and S.Sh Abd El-Gayed (2008).** Effect of potassium fertilizer under two planting dates on yield and yield components of Giza 80 cotton cultivar. *Annals of Agric. Sci. Moshtohor*, 46 (4): 327 – 332.
- Sawan, Z.M. (2014).** Cottonseed yield and its quality as affected by mineral fertilizers and plant growth retardants. *Agric. Sci.*, (5)3: 186 – 209.
- Sema, B.; T. Bahar, E. Remzi, Y. Mustafa and B. Yüksel (2012).** The determination effects of potassium and zinc application to rate of photosynthesis, fiber yield and quality on cotton. 11th Meeting of inter-regional cooperative research network on cotton for the Mediterranean and Middle East regions. November 5 - 7, Antalya, pp: 25.
- Snedecor, G.W. and W.G. Cochran (1981).** *Statistical Analysis Methods*. Seventh Edition, Iowa State Univ., Ames, Iowa, USA, pp: 225 - 269.
- Woodward, F.I. and J.E. Sheehy (1983).** *Principles and Measurements in Environmental Biology*. pp: 263. Butterworth & Co (Publishers) Ltd., UK.

Young, E.F.; R.M. Teyler and H.D. Petarson (1980). Day-degree units and time in relation to vegetative development and fruiting for three cultivars of cotton. *Crop Sci.*, 20: 370 - 375.

تأثير طرق ومواعيد التسميد البوتاسي على المحصول ومكوناته لهجين القطن الجديد (جيزة ٨٦ × ١٠٢٢٩) تحت مواعيد الزراعة المبكرة والمتأخرة

أمل سامي على عبد العال ، مصطفى عطية أحمد عمارة ، سعيد عبد التواب فرج حمودة

قسم بحوث المعاملات الزراعية - معهد بحوث القطن - مركز البحوث الزراعية - الجيزة - مصر

أجريت تجربتان حقليتان بمحطة البحوث الزراعية بسخا، محافظة كفر الشيخ خلال موسمي ٢٠١٤، ٢٠١٤ وذلك بهدف دراسة تأثير طرق ومواعيد إضافة سماد البوتاسيوم على نمو وإنتاجية القطن تحت ظروف الزراعة المبكرة والمتأخرة (١٥ أبريل، ١٥ مايو) لهجين القطن الجديد جيزة ٨٦ × ١٠٢٢٩ وهذا الهجين انتج من قسم بحوث تربية القطن وفي مرحلة اعداد التوصيات الفنيه له، وكانت معاملات إضافة سماد البوتاسيوم كالآتي:

- ١- إضافة أرضية بسلفات البوتاسيوم بمعدل ٢٤ وحدة بوحدة بوحدة للفدان عند الخف.
- ٢- إضافة أرضية بسلفات البوتاسيوم بمعدل ٢٤ وحدة بوحدة للفدان بعد الخف بأسبوعين.
- ٣- إضافة سلفات البوتاسيوم بمعدل ٥ كجم للفدان رشا مرتين عند ظهور الوسواس وعند بداية التزهير.
- ٤- إضافة سلفات البوتاسيوم بمعدل ٥ كجم للفدان رشا مرتين عند بداية التزهير وبعد التزهير بأسبوعين.
- ٥- إضافة سلفات البوتاسيوم بمعدل ٥ كجم للفدان رشا ثلاث مرات عند بداية الوسواس وعند بداية التزهير ثم بعد التزهير بأسبوعين.

وقد صممت التجربة بتصميم القطع المنشقة في أربع مكررات حيث وضعت مواعيد الزراعة (١٥ أبريل، ١٥ مايو) في القطع الرئيسية ووضعت طرق ومواعيد إضافة السماد البوتاسي في القطع الشقية. وتتخصص أهم النتائج المتحصل عليها فيما يلي:

- ١- أدت الزراعة في الميعاد المبكر (١٥ أبريل) إلى زيادة معنوية في عدد السلاميات علي الساق الرئيسي، عدد الأفرع الثمرية علي النبات، عدد اللوز المتفتح علي النبات متوسط وزن اللوزة، ومحصول القطن الزهر بالقطار للفدان، والوزن الجاف للأعضاء الخضرية والثمرية للنبات وزيادة عدد الأيام لتفتح أول زهرة وأول لوزة وكذلك زيادة نسبة التكبير، بينما أدت إلى نقص الطول النهائي للنبات وعدد الأفرع الخضرية علي النبات وإنخفاض موقع أول فرع ثمرى علي النبات.
- ٢- لم تؤثر مواعيد الزراعة علي تصافي الحليج، معامل البذرة وعدد النباتات للفدان عند الجني.
- ٣- أدى إضافة السماد للبوتاسي إلى نتائج إيجابية حينما أضيف في الميعاد المناسب أرضاً بمعدل ٢٤ وحدة بوحدة للفدان بعد الخف بأسبوعين، حيث أدت تلك المعاملة إلى نقص عدد الأفرع للخضرية علي النبات وإلى زيادة عدد الأفرع الثمرية علي النبات وزيادة الوزن الجاف لأجزاء النبات المختلفة وزيادة وزن اللوزة وعدد اللوز علي النبات ووزن المحصول للنبات والفدان.
- ٤- لم يؤثر إضافة البوتاسيوم علي عدد السلاميات للساق الرئيسي وعلى موقع أول فرع ثمرى وعدد الأيام لتفتح أول زهرة وأول لوزة ووزن ١٠٠ بذرة وتصافي الحليج وعدد النباتات للفدان عند الجني.
- ٥- لم يكن للتفاعل بين مواعيد الزراعة ومواعيد وطرق إضافة السماد البوتاسي أي تأثير معنوي علي الصفات تحت الدراسة.

من النتائج المتحصل عليها في هذه الدراسة فإنه يمكننا التوصية بزراعة هجين القطن الجديد (جيزة ٨٦ × ١٠٢٢٩) مبكراً في (١٥ أبريل) مع إضافة لرضية لسلفات البوتاسيوم بمعدل ٢٤ كجم K_2O /فدان مرة واحدة بعد الخف بأسبوعين وذلك لزيادة إنتاجية محصول القطن تحت ظروف منطقة سخا.