

## YOGHURT ENRICHED WITH CONJUGATED LINOLEIC ACID USING PROBIOTIC BACTERIA BIO PRODUCTION AND ADDITION OF AVOCADO FRUIT

El-Abd M.M<sup>1</sup>; Wafaa. B. Elsabee<sup>2</sup>; E.M. Hamad <sup>3</sup>and Hagar.F.Saad<sup>4</sup>

1, 3 Dairy Dept faculty of agric, Cairo university

2, 4 Dairy Chemistry Dept, Animal Prod. Res. Inst. ARC

### ABSTRACT

This study was performed to characterize the ability of active Bifidobacteria strains and Avocado to produce conjugated linoleic acid (CLA). Using Bifidobacteria such as *B. breve*, *B. bifidum* and *B. lactis* showed an increase in CLA that convert linoleic acid (LA) to conjugated linoleic acid (CLA).

In first part set yogurt was manufactured using buffalo milk 6% fat, with adding *B. breve*, *B. bifidum* and *B. lactis*. Each of the prepared yoghurt assessed for chemical, microbiological and organoleptic properties when fresh and during refrigerated storage at  $6 \pm 1^\circ\text{C}$  for 14 days.

Results indicated that using probiotic bacteria (*B. breve*, *B. bifidum* and *B. lactis*) was effective in producing conjugated linoleic acid (CLA) in yoghurt treatments during fermentation as compared with the control yoghurt. Contents of probiotic bacteria (*B. breve*, *B. bifidum* and *B. lactis*) were decreased in yoghurt treatments during storage, Firmness, Syneresis and pH values were also decreased while T.S, Fat and acidity increased.

In second part stirred yoghurt was made using the same Bifidobacteria species and 10% or 15% avocado fruit. Each of the prepared yoghurt was assessed for chemical, microbiological and organoleptic properties when fresh and during refrigerated storage at  $6 \pm 1^\circ\text{C}$  for 14 days.

Obtained results indicated that adding avocado fruit to probiotic yoghurt highly increased CLA content in all yoghurt treatments till the end of storage period. Also Bifidobacteria counts gradually increased during storage period.

**Keywords:** yoghurt, CLA, bifidobacteria bacteria, Avocado.

### INTRODUCTION

Conjugated linoleic acid (CLA), refers to a group of positional and geometric isomers of octadecadienoic acid (C18: 2). conjugated double bonds (i.e., the two double bonds are separated by one single bond) in either cis (c) or trans (t) configuration are present predominantly in positions 8 and 10, 9 and 11, 10 and 12, or 11 and 13. The cis9 - trans 11 isomer is the main isomer constituting 90% of the total CLA found in dairy and beef lipids so called rumenic acid. Linoleic acid is converted to CLA by linoleic acid isomerase enzyme of rumen bacteria. For these reasons, CLA occurs naturally as a mixture of isomers in dairy foods such as milk and meats derived from ruminant animals. The most abundant isomer of CLA in nature is the cis9 - trans 11 (cis9 - trans 11) isomer. Commercially available CLA is usually a mixture of cis9 - trans11 and trans-10, cis-12 (t10c12) isomers with other isomers as minor components. Conjugated linoleic acid isomer mixture and c9t11 and t10c12 isomers alone have been attributed to provide several health benefits that are largely based on animal and in vitro studies.

CLA has been shown to exert various physiological functions such as anti carcinogenic, anti diabetic, anti obesity and prevention of atherosclerosis (Arunabh Bhattacharya, 2006) and (Yanagitab, 2013). The daily intake of CLA is almost 200 mg /day are reviewed by Marialice P.C. Silvestre, (2013) and Microwaving caused a significant decrease in CLA content in milk .The loss was up to 53 % (Walter Bisig., etal 2007).

Bifidobacterium species as *Bifidobacterium bifidum* LMG 10645, *Bifidobacterium animalis subsp lactis* BB12 and *Bifidobacterium breve* LMC520 were used for increasing CLA by microbial conversion of linoleic acid to conjugated linoleic acid and in part 2 using Bifidobacterium species as *Bifidobacterium bifidum* LMG 10645, *Bifidobacterium animalis subsp lactis* BB12 and *Bifidobacterium breve* LMC520 with different ratios from Avocado (*Persea Americana* ).

Many researchers investigated the role of common lactic acid bacteria in CLA production. Several strains of food grade micro-organisms have been identified as potential producers of CLA, including strains of *Bifidobacterium*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Propionibacterium* and *Streptococcus*. These cultures may be very important for the food industry, in particular to the dairy industry, as they can be used for the production of traditional and or novel CLA enriched foods through in situ fermentation or used to obtain CLA as a food-ingredient through a biotechnological process. Probiotic or symbiotic dairy foods with potential contents of CLA have been reported by (Rodrigues D, 2011), Silva L I., (2011) and Rodrigues D., (2012).

Avocado, or *Persea American* (P. americana Mill.) is a kind of evergreen fruit plants of the genus *Persea* family .Avocado contains a variety of essential nutrients as monounsaturated fatty acids (MUFA)-rich fruit oil with 71% MUFA, 13% polyunsaturated fatty acids (PUFA) and 16% saturated fatty acids (SFA). As the avocado fruit ripens, the saturated fat decreases and the monounsaturated fatty acids. Avocados can fit into a heart healthy dietary pattern such as the DASH diet plan USDA and HHS, 2010, Jakobsen etal, 2009, de Souza.,etal, 2008 and Appel .,etal (2005).

Slater.,etal., (1975), Moreno .,etal (1980) and Lu. et al (2009) showed that the use of avocado dips and spreads as an alternative to more traditional hard, SFA rich spreads or dips can assist in lowering dietary SFA intake Cited from <http://www.avocadocentral.com/nutrition/avocado-spread-comparison>. This study aims to increase CLA content in yogurt using some spp of Bifidobacteria and avocado fruit (*Persea Americana*).

## MATERIALS AND METHODS

### Materials:

Fresh buffalo's milk: was obtained from Dairy technology Unit, Animal, Production Research Institute, Agric. Res. Center, Ministry of Agriculture.

Yoghurt culture: Mixed culture consisted of *Lactobacillus delbrueckii sub sp. bulgaricus* and *Streptococcus salivarius sub sp thermophilus* was obtained from Chr. Hansen's Lab A/S Copenhagen, Denmark.

Bifidobacterium strains: *Bifidobacterium bifidum* LM 04, *Bifidobacterium breve* LMC 017 and *Bifidobacterium animalis subsp. Lactis* LMG 10645 were obtained from Chr. Hansen Lab., Copenhagen Denmark.

Microbiological media: Microbial media used in this study were ready made using Oxid Division. Oxoid Ltd London.

Persea Americana (Avocado) fruit: was obtained from local market

#### Methods of analysis

Moisture content: The moisture content was determined gravimetrically according to AOAC (1990) using a Vacuum Oven, Haraeus T5042EK, Germany.

pH: pH values of yoghurt samples were estimated using a digital pH meter (Jenway 3505 pH meter).

Titrate acidity: The acidity was determined by titration following the method described by A.O.A.C. (1990).

Viscosity : Viscosity of yoghurt samples was measured using a Brookfield DV- E viscometer in 100 ml yoghurt at 20°C.

Fat content: Fat content was determined according to A.O.A.C. (2007).

Curd tension (Firmness): Curd tension (Firmness) was measured according to chanderosek hara.,etal (1975).

Syneresis: Syneresis was measured as described by lowerence (1959).

Quantification of CLA production by UV spectroscopy: Total CLA determination was carried out at a wavelength of 233 nm in a Perkin-Elmer spectrophotometer (Lambda650 model, Beaconsfield, UK) with scan program (190–350 nm). Measurements were obtained in triplicate from 2 ml of the lipid extract in hexane placed into quartz cuvettes. In order to verify the suitability of this method, a standard curve was constructed for the absorbance at 233 nm versus CLA (C18:2 c9, t11) concentration (0–30 lg/ml).

Lipid extraction: Lipid isolation from culture media was carried out as a variation of the method described by Alonso, Cuesta, and Gilliland (2003). Briefly: 10 ml of culture media were centrifuged at 7500 rpm, 5 min, and 4 °C. Three ml from the resulting supernatant were added with 6 ml of isopropanol and vortexed for 1 min. Addition of hexane (5 ml) followed which were vortex for 1 min and finally centrifuged at 2000 rpm, 5 min, and 4 centigrade.

Microbiological analysis: Microbiological analysis was done for *B. bifidum*, *B. breve* and *B.lactis* enumerated (Dave and Shah, 1996) using the modified MRS agar supplemented with 0.05% L. cysteine-HCL. The antibiotic mixture NPML (Neomycin sulphate, paramonycin sulphate, Nalidixic acid and lithium chloride) as a selective agent was sterilized by filtration through 0.22 µm Millipore filter (Gellman Sci., England) then added to the medium at a rate of 50 ml/L medium just before pouring into the plates. The L. cystein-HCL solution which sterilized by filtration (0.05% final concentration) was

also added using MRS-Salccin agar. The plates were incubated at 37°C for 48 hrs. Total bacterial count was measured as suggested by The American Public Health association APHA (1992).

Preparation of probiotic set yoghurt: Fresh buffalo's milk of 6% fat was heated to 80°C for 20 min, and cooled to 42°C. Such milk was divided into 4 equal portions. The 1st was inoculated with 3% yoghurt culture (control), the 2nd inoculated with 3% yoghurt culture and *Bifidobacterium bifidum* (1:1), the 3rd inoculated with 3% yoghurt culture and *Bifidobacterium breve* (1:1), and the 4th inoculated with 3% yoghurt culture and *Bifidobacterium lactis* (1:1). The prepared probiotic set yoghurt and their control were assessed for chemical, microbiological and organoleptic properties when fresh and during refrigerated storage at 6± 1 °C for 14-day.

Preparation of probiotic stirred yoghurt fortified with Avocado: Fresh buffalo's milk of 6% fat was heated to 80°C for 20 min and cooled to 42°C. Such milk was divided into 9 equal portions. The 1st was inoculated with 3% yoghurt culture, 2nd inoculated with 3% yoghurt culture with 100 gm Avocado /L milk, 3rd inoculated with 3% yoghurt culture + *B.bifidum* (1:1) and 100 gm Avocado /L milk, the 4th inoculated with 3% yoghurt culture and *Bifidobacterium breve* (1:1) + 100 gm Avocado /L milk, the 5th inoculated with 3% yoghurt culture and *B.lactis* (1:1)+ 100 gm Avocado /L milk, the 6th inoculated with 3% yoghurt culture +150 gm avocado, the 7th inoculated with 3% yoghurt culture and *B. B.Bifidum* (1:1)+ 150 gm Avocado/L milk; the 8th inoculated with 3% yoghurt culture + *B.breve* (1:1) and + 150 gm Avocado /L milk and the 9th inoculated with 3% yoghurt culture and *B. lactis* (1:1) + 150 gm Avocado /L milk and stirred. The prepared probiotic yoghurt were assessed for chemical, microbiological and organoleptic properties at fresh and during refrigerated storage at 6± 1 °C for 15-day.

Sensory evaluation: All samples were evaluated according to Clark et al. (2009)

## RESULTS AND DISCUSSION

1-Use of some *Bifidobacteria* species for Conjugated linoleic acid bio production in set yoghurt

**Table 1. Effect of using different *Bifidobacteria* species on some properties of set yoghurt during cold storage (6±1)**

reatments	% Fat		% T.S		Acidity						pH			
	0	14	0	14	0	3	7	10	14	0	3	7	10	14
	(Days)													
1	6.00	6.03	15.2	16.0	0.63	0.73	0.82	0.91	1.00	4.63	4.54	4.46	4.33	4.21
2	6.00	6.03	15.1	16.0	0.66	0.76	0.88	0.99	1.15	4.54	4.44	4.33	4.25	4.10
3	6.00	6.03	15.1	16.3	0.68	0.80	0.92	1.08	1.21	4.57	4.49	4.34	4.21	3.94
4	6.00	6.03	15.3	16.5	0.64	0.75	0.84	0.94	1.10	4.57	4.47	4.32	4.19	4.06

1- Control yoghurt culture (3 %)

2- Yoghurt culture + *Bifidobacterium bifidum* (3 %) 1:1.

3- Yoghurt culture + *Bifidobacterium breve* (3 %) 1:1.

4- Yoghurt culture + *Bifidobacterium lactis* (3 %) 1:1.

Data presented in table (1) show an increase in fat and total solids (T.S) contents of yoghurt samples after 14 days of cold storage ( $6\pm1^\circ\text{C}$ ). These increases could be attributed to water evaporation during storage period. Gradual increasing in yoghurt titratable acidity (as lactic acid %) was also observed during cold storage period while pH values of the same samples showed the opposite trend.

**Table 2. Conjugated linoleic acid content of set yoghurt treatments fortified with some Bifidobacteria spp during cold storage ( $6\pm1$ )**

Treatments	CLA content (mg/ 100 gm fat)							
	0	%	3	%	10	%	14	%
		Increasing		Increasing		Increasing		Increasing
	(Days)							
1	0.18	0000	0.23	0000	0.30	0000	0.41	0000
2	0.33	88.00	0.52	127.5	0.64	112.5	0.87	110.0
3	0.37	109.5	0.48	112.7	0.75	148.5	1.02	147.5
4	0.28	60.00	0.44	95.50	0.60	99.6	0.74	77.90

Data tabulated in table 2 clearly show a noticeable increase in fresh yoghurt CLA contents when Bifidobacteria species were incorporated with yoghurt starters comparing to control treatment (1). CLA concentrations of probiotic yoghurt treatments 2, 3 and 4 (containing Bifidobacterium species) were nearly the twice. On the other hand gradual increases in CLA content of all yoghurt treatments during cold storage period were observed till 14 days. The highest CLA content was recorded in treatments (3) samples (*Bifidobacterium breve*) after 14 days of storage (the end of storage period). Samples of treatments (2 and 4) (*Bifidobacterium bifidum* and *Bifidobacterium lactis*) showed less CLA content than samples of treatments (3). Obtained results are in accordance with those found by Kianoush *et al* (2014).

**Table 3. Changes in some rheological properties of set yoghurt treatments during cold storage ( $6\pm1$ )**

Treatments	Firmness \ gm				Syneresis \ ml			
	0	3	10	14	0	3	10	14
	(Days)							
1	50.6	49.2	45.0	43.0	21	20	18	16
2	48.6	46.6	42.0	39.0	23	23	21	19
3	45.7	44.0	39.9	37.8	23	25	22	20
4	44.6	42.8	40.0	36.9	24	25	23	21

Changes in yoghurt sample firmness and syneresis are presented in table (3). The obtained results show gradual decrease in both firmness and syneresis values till the end of storage period. In general probiotic yoghurt samples (treatments 2, 3 and 4) showed lower firmness values and higher syneresis volumes as compared with control samples (treatment 1) during storage period. These findings are in agreement with those obtained by Ali *et al* (2013) and Moayednia (2014).

Table 4.Changes of Bifidobacteria counts in set yoghurt treatments during cold storage (6±1)

Treatments	Changes in Bifidobacteria spp counts (10 <sup>7</sup> )						
	0	%	3	%	10	%	14
	Decreasing		Decreasing		Decreasing		
	(Days)						
1	nd	000	nd	000	nd	000	nd
2	196	3.00	189	13.5	169	12.5	171
3	168	8.00	154	11.0	148	14.8	143
4	176	5.80	166	5.50	168	7.30	163

Table 4 showed that *bifidobacterium breve*, *bifidobacterium bifidum* and *bifidobacterium lactis* count decreased gradually during storage period for 14 days because of increase of acidity, Low oxygen content and low redox potentials have been cited as important factors for the viability of bifidobacteria during storage of fermented milk products (Brunner .etal 1993a and 1993b).

Table 5. Sensory evaluation of fresh set yoghurt treatments

Treatment	Flavor(20)	Color(10)	Texture(10)	Sensory evaluation(40)
1	19	10	9	38
2	18	10	8	36
3	17	10	8	35
4	16	10	8	34

Sensory evaluation (total score 40) of fresh yoghurt samples are shown in table (5). Tabulated data show that control samples gained the highest score reaching 38, while probiotic yoghurt samples gained less score specially treatment (4) (with *B.lactis*).

Π-Enrichment of probiotic stirred yoghurt with avocado fruit

Table (6) Effect of adding avocado fruit on some properties of probiotic stirred yoghurt during cold storage (6±1)

reatments	% Fat		% T.S		Acidity								pH			
	0 14		0 14		0 3		7 10		14 0		3 7		10 14			
	(Days)															
1	6.00	6.10	15.2	16.0	0.64	0.73	0.82	0.92	1.00	4.62	4.54	4.46	4.33	4.21		
2	6.90	7.20	20.6	21.7	0.69	0.79	0.88	0.96	1.10	4.42	4.31	4.22	4.11	4.01		
3	6.90	7.20	20.6	21.6	0.71	0.83	0.90	1.07	1.21	4.35	4.21	4.11	4.00	3.87		
4	6.90	7.20	20.6	21.7	0.73	0.88	0.99	1.13	1.28	4.32	4.17	4.03	3.89	3.78		
5	6.90	7.10	20.8	21.9	0.69	0.82	0.90	1.00	1.17	4.37	4.28	4.10	3.97	3.86		
6	7.30	7.60	20.8	21.7	0.71	0.84	0.92	1.11	1.24	4.27	4.18	4.07	3.91	3.88		
7	7.30	7.60	20.9	21.8	0.73	0.88	1.00	1.16	1.32	4.36	4.29	4.16	4.02	3.92		
8	7.30	7.60	20.692	21.668	0.74	0.90	1.02	1.21	1.37	4.30	4.18	4.05	3.88	3.71		
9	7.30	7.70	20.686	21.635	0.72	0.88	0.96	1.14	1.26	4.34	4.20	4.10	3.91	3.81		

1-Control yoghurt culture (3 %).

2-Yoghurt culture (3 %) +100 gm avocado/1 liter milk.

3-Yoghurt culture + *bifidobacterium bifidum* (1:1) +100 gm avocado / 1 liter milk.

4-Yoghurt culture + *bifidobacterium breve* (1:1) +100 gm avocado / 1 liter milk.

5-Yoghurt culture + *bifidobacterium lactis* (1:1) +100 gm avocado / 1 liter milk.

6- Yoghurt culture (3%) +150 gm avocado / 1 liter milk.

7- Yoghurt culture + *bifidobacterium bifidum* 1:1+ 150 gm avocado / 1 liter milk.

8-Yoghurt culture + *bifidobacterium breve* (1:1) +150 gm avocado / 1 liter milk.

9- Yoghurt culture + *bifidobacterium lactis* (1:1) +150 gm avocado / 1 liter milk.

Effect of adding avocado fruit (10 or 15%) on fat, T.S, acidity and pH of probiotic yoghurt samples during cold storage is shown in table (6). Tabulated data showed obvious increase in fat and T.S content due to avocado addition in fresh yoghurt samples. During cold storage fat and T.S content were increased as mentioned before in table (1) due to water evaporation. Titratable acidity of yoghurt sample gradually increased through out storage period while pH values decreased till the end of storage. Addition of avocado fruit clearly affected titratable acidity and pH of fresh yoghurt samples.

**Table7.Content of CLA in stirred yoghurt fortified with some Bifidobacteria spps and different ratios of avocado during cold storage (6±1)**

Treatments	CLA content (mg/ 100 gm fat)							
	0	%	3	%	10	%	14	%
	Increasing (Days)							
1	0.18	0000	0.28	000	0.30	000	0.41	000
2	1.10	513.4	1.22	437	1.82	500	2.56	518
3	1.18	562.9	1.46	542	2.40	691	2.80	575
4	1.67	838.2	1.70	649	2.72	797	3.09	647
5	1.04	485.3	1.20	430	1.85	510	2.31	458
6	1.37	671.3	1.75	672.6	2.56	744	2.86	591
7	1.57	783.7	1.99	777.9	2.83	834	3.33	705
8	1.93	987.0	2.35	937.4	3.25	972	3.64	780
9	1.40	687	1.80	691	2.62	764	3.10	650

Results in table (7) indicated that CLA contents gradually increased during storage period. Also these results obviously showed significant increase in CLA content when avocado fruit was added to yoghurt samples. In other hand CLA content increasing was positively correlated with avocado concentration used. On the other hand CLA content increasing was positively correlated with avocado concentration used.

**Table 8.changes in viscosity of stirred Yoghurt treatments fortified with avocado during cold storage (6±1)**

Treatment	Viscosity(cp)			
	0	3	10	14
	(Days)			
1	6000	6600	8800	9800
2	9000	9800	12400	13200
3	8000	9000	11700	12400
4	8500	9300	11800	12300
5	8700	9700	12000	12800
6	10100	11000	11700	13700
7	9700	10300	11400	12900
8	9700	10800	11800	12900
9	10000	11000	12300	13400

In general CLA increase could be attributed to avocado addition and conversion of LA to CLA during biohydrogenation by bifidobacteria species. (Milutal, 1992). The highest CLA content was recorded in treatments (9) sample (*Bifidobacterium breve* + 150 gm avocado fruit) after 14 days of storage (the end of storage period). Samples of other treatments showed less CLA content than samples of treatments (9).

Data presented in table (8) show gradual increases in viscosity values of Yoghurt treatments during storage period. Viscosity increase could be due to acidity development and avocado fortification. (Davenport,2013).

**Table 9. Changes of Bifidobacteria counts in stirred yoghurt treatments fortified with avocado during cold storage (6±1)**

Treatments	Changes in Bifidobacteria spp counts(10 <sup>7</sup> )						
	0	%	3	%	10	%	14
	Increasing (Days)						
1	nd		nd		nd		nd
2	nd		nd		nd		nd
3	364	19.7	436	35.7	494	44.0	525
4	240	29.0	310	67.0	401	113	512
5	286	15.7	331	51.0	432	92.3	550
6	nd		nd		nd		nd
7	389	28.0	498	58.0	616	82.0	708
8	261	38.0	361	73.0	452	120	575
9	314	20.0	377	51.0	475	92.0	605

Table 9 showed that *Bifidobacterium breve*, *Bifidobacterium bifidum* and *Bifidobacterium lactis* counts gradually increased during storage period for 14 days due to avocado addition. The highest increase of Bifidobacterium counts was recorded with *Bifidobacterium bifidum* treatments as it was previously mentioned (table4). Increase of probiotic bacterial counts could be ascribed by the simultaneous presence of inhibitory compounds such as antibacterial poly phenolics (Ripa.,etal ,2009).

Results in table (10) showed sensory evaluation of fresh probiotic yoghurt treatments fortified with avocado fruit. Almost no differences were recorded between all treatments except control sample (1).

**Table 10. Sensory evaluation of fresh stirred yoghurt treatments fortified with avocado**

Treatments	Flavor(20)	Color(10)	Texture(10)	Sensory evaluation(40)
1	19	10	9	38
2	15	8	9	32
3	15	8	9	32
4	15	8	9	32
5	15	8	9	32
6	16	7	9	32
7	16	7	9	32
8	15	7	9	31
9	16	7	9	32

## REFERENCES

- A.O.A.C. (1990). Official Methods of Analysis Chemists, 15th ed. association of Official Analytical Chemists AOAC, USA, Virginia, Arlington.
- A.O.A.C. (1997). Official Method of Analysis, 15th ed. AOAC, Washington, DC, USA.
- A.O.A.C. (2007). Official methods of analysis of the Association of the Analytical Chemists. 17 ED. Published by the Association of official analytical chemists. PO Box 540. Benjamin Franklin station Washington DC20044.
- A.P.H.A. (American Public Health Association)(1992). Standard Methods for examination of Dairy products. American Public Health Association. Inc 16th ED Publ. Avico Inc. Westport Connecticut USA.
- Ali, F. S.; Saad O. A.O and Salwa,A .G. Hussein (2013). Probiotic stability of yoghurts during refrigerated storage. Egypt. Egyptian Academic Journal of Biological Sciences 5(2): 9-19.
- Alonso, L., Cuesta, E. P., & Gilliland, S. E. (2003). Production of free conjugated linoleic acid by lactobacillus acidophilus and lactobacillus casei of human intestinal origin. Journal of Dairy Science.86: 1941–1946.
- Appel L. J., Sacks F. M., Carey V. J., Obarzanek E., Swain J. F., Miller E. R., Conlin P. R., Erlinger B. A., Rosner B. A., Rosner B. A., Laranjo N. M., Chatleston J., McCarron P., Bishop L. M.(2005). Effect of protein, monounsaturated fat and carbohydrate intake on blood pressure and serum lipids: Results of the Omni Heart randomized trial. Journal of the American Medical Association. 294:2455–2464.
- Arunabh Bhattacharya, J. B., Mizanur Rahman, Jennifer Causey, Gabriel Fernandesemail (2006). Biological effects of conjugated linoleic acids in health and disease. Journal of Nutritional Biochemistry. 17(12): 19.
- Brunner JC, Spillman H, Puhon Z. (1993a). Metabolism and survival of bifidobacteria in fermented milk during cold storage. Milchwirtschaftliche 22:19–25.
- Brunner JC, Spillman H, Puhon Z. (1993b). Changes in pH, free sulphydryl groups, oxygen and redox potential during fermentation of milk with Bifidobacterium longum. Milchwirtschaftliche 22: 19–25.
- Candrosek hara.M.R., Bhajawan. RK. Swaminan.M. and Subrahmanyam ,V.(1975). The use of mammalian milk food processed with foods in feeding of infants. Indian journal of child health.701.
- Clark,S.,Drake,M.A.,Cotello,M. and Body felt ,F.(2009).The sensory evaluation of dairy products .221-222.Springer science + Business media, LLC, New York.
- Davenport,et al. (2013). Hass Avocado Composition and Potential Health Effects. Critical Reviews in Food Science and Nutrition 53: 738–750.

- De Souza R. J., Swain J. F., Appel L. J., Sacks F. M. (2008). Alternatives for macronutrient intake and chronic disease: A comparison of the OmniHeart diets with popular diets and with dietary recommendations. *American Journal Clinical Nutrition*. 88:1–11. <http://www.avocadocentral.com/nutrition/avocado-spread-comparison>.
- Jakobsen, M. U., O'Reilly, S., Heitmann, B. L., Pereira, M. A., Balter, K., Fraser, G. E., Goldbourt, U., Hallmans, G., Knekt, P., Liu, S., Pietinen, P., Spiegelman, D., Stevens, J., Virtamo, J. and Willett, W. C. (2009). Major types of dietary fat and risk of coronary heart disease: A pooled analysis of 11 cohort studies. *Am. J. Clin. Nutr.* 89:1425–1432.
- Kianoush Khosravi-Darani, Fatemeh Seyed Reihani and Reza Feili (2014). Bioproduction of Conjugated Linoleic Acid in Yogurt by Probiotic Bacteria. *International Journal of Biotechnology for Wellness Industries* 3: 62-68.
- Lawrences, A.J. (1959). Syneresis of rennet curd. *Journal of dairy technology* 14:166.
- Lu Q.-Y., Zhang Y., Wang Y., Lee R.-P., Gao K., Byrns R., Heber D. (2009). California Hass Avocado: Profiling of carotenoids, tocopherols, fatty acids, and fat content during maturation and from different growing areas. *Journal of Agricultural and Food Chemistry*. 57:10408–10413.
- Marialice P.C. Silvestre, D. C. F. L., Mauro R. Silva<sup>3</sup> and Viviane D.M. Silva (2013). Availability of Food Sources of Conjugated Linoleic Acid in Brazil. *Pakistan Journal of Nutrition*. 12(1): 08-11.
- Milutal, Postal Bag Galil Hamaaravi. (1992). An Assessment of Processing Potential of Avocado Fruit. *California Avocado Society Yearbook* . 76: 137-144.
- Moayednia, N. (2014). Quality Evaluation of New Developed Symbiotic Yogurt over the Storage at Refrigerator. *Journal of Food Biosciences and Technology* 4(1): 57-64.
- Moreno A., Dorantes L., Galindez J., Guzman R. I. (1980). Effect of different extraction methods on fatty acids, volatile compounds, and physical and chemical properties of avocado (*Persea americana* Mill.) oil. *Journal of Agricultural and Food Chemistry*. 51:2216–2221.
- Ripa, F.A., Haque, M., Nahar, L., Islam, M., (2009). Antibacterial, cytotoxic and antioxidant activity of *Passiflora edulis* Sims. *European Journal of Scientific Research* 31, 592–598.
- Rodrigues D, R.-S. T. A. P., Gomes A M, Goodfellow B J and Freitas A C (2012). Lipolysis in probiotic and synbiotic cheese: the influence of probiotic bacteria, prebiotic compounds and ripening time on free fatty acid profiles. *Food Chemistry* 131(4)1414-1421.
- Rodrigues D, R.-S. T. A. P., Pereira C I, Gomes A M, Malcata F X and Freitas A C (2011). The potential effect of FOS and inulin upon probiotic bacterium performance in curdled milk matrices *Food Science and Technology*. 44:100-108.

- Silva L I, R. D., Freitas A C, Gomes A M, Rocha-Santos T A P, Pereira M E and Duarte A C (2011). Optical fiber based methodology for screening the effect of probiotic bacteria on conjugated linoleic acid (CLA) in curdled milk. Food Chemistry 127(1): 222–227.
- Slater G., Shankman S., Shepherd J. S., Alfin-Slater R. B. (1975). Seasonal variation in the composition of California avocados. Journal of Agricultural and Food Chemistry. 23:468–474.
- USDA and HHS. (2010). Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans (Part B. Section 2: The Total Diet. B2:11). U.S. Department of Agriculture and U.S. Department of Health and Human Services. Washington, DC.
- Walter Bisig, Pius Eberhard, Marius Collomba, Brita Rehberger (2007). Influence of processing on the fatty acid composition and the content of conjugated linoleic acid in organic and conventional dairy products - a review. INRA, EDP Sciences 87: 1-19.
- Yanagita, K. K. a. T. (2013). Review Health benefits of conjugated linoleic acid (CLA). Obesity Research & Clinical Practice 8(6):525-532.

اليوجورت المدعم بحمض اللينوليك المرتبط باستخدام الانتاج الحيوى لبكتريا البروبيوتك بالاضافه الى فاكهة الأفوكادو  
منير محمود العبد ١، وفاء بديع السبع ٢، عصام محمد حمد ٣ وهاجر فتحى سعد ٤  
١، ٣ قسم الالبان كلية الزراعة جامعة القاهرة  
٢، ٤ - قسم كيمياء الالبان معهد بحوث الانتاج الحيوانى - مركز البحوث الزراعيه

أجريت هذه الدراسة لمعرفة قدرة سلالات البفيدوبكتيريا النشطة والأفوكادو لإنتاج حمض اللينوليك المرتبط (CLA) وقد أظهرت البفيدوبكتيريا B. Breve, B. bifidum and B. lactis زيادة في CLA حيث يتم تحويل حمض اللينوليك (LA) إلى حمض اللينوليك المرتبط (CLA).  
في الجزء الأول تم تصنيع اليوجورت من لبن جاموسى ٦% دهن يحتوى على B.breve, B.bifidum and B.lactis حيث وتم تقييم اليوجورت الناتج كيميائيا و ميكروبيولوجيا وحسباً وذلك لليوجورت الطازج وخلال فترة التخزين على ٦ ± ١ درجة مئوية لمدة ١٤ يوم.  
وقد أوضحت النتائج ان استخدام بكتريا البروبيوتك B.breve, B. bifidum and B. lactis كان لها تأثير على زيادة انتاج حمض اللينوليك المرتبط اثناء التخمير مقارنة بالكنترول .  
وقد انخفضت اعداد بكتريا البروبيوتك B.breve, B. bifidum and B. lactis كما انخفض معدل طرد الشرش و معدل صلابة الخثرة و قيم الرقم الهيدروجينى pH بينما ارتفعت قيم كل من الجوامد الكلية و الدهن و الحامضه.  
وفي الجزء الثانى تم تصنيع اليوجورت باستخدام نفس السلالات البكتيرييه مع اضافته نسب ١٠ او ١٥ % من فاكهة الأفوكادو و تم تقييم الناتج كيميائيا و ميكروبيولوجيا وحسباً و ذلك لليوجورت الطازج وخلال فترة التخزين على ٦ ± ١ درجة مئوية لمدة ١٤ يوم.  
و اذت اضافة الأفوكادو لليوجورت المقلب المدعم ببكتريا البروبيوتك الى زيادة نسبة حمض اللينوليك المرتبط فى جميع المعاملات حتى نهاية مدة التخزين كما ادت الى زياده اعداد بكتريا البروبيوتك خلال مدة التخزين.