

INFLUENCE OF DIFFERENT COTTON VARIETIES AND GENOTYPES ON THE INFESTATION BY THE SPINY BOLLWORM *Earias insulana* (BOISD.) AT KAFR EL-SHEIKH GOVERNORATE

Somaa, H.M.H. and Asmaa M.A. El-Ghobary

Plant Protection Research Institute- Agric. Res. Center, Egypt

ABSTRACT

The present work was carried out during two successive seasons (2012 and 2013) at Farm of Sakha Agricultural Research Station, Kafr El-Sheikh Governorate, Egypt to evaluate the influence of different cotton varieties and genotypes on the infestation by the spiny bollworm *Earias insulana* (Boisd.) at Kafr El-Sheikh Governorate. Results during the two experiments showed the infestation percentage of SBW to six cotton varieties and genotypes of both seasons were (8.14, 7.64, 7.66, 6.08, 7.15 and 8.36%) for Giza 86, Pima S6 x Giza 77, Giza 92, Giza 94, Giza 86 x 10/229 and Giza 88, respectively. The highest means of infestation % were recorded in the G.88 and G.86 in both seasons, while the lowest means of infestation% were recorded in the G.94 and G.86 x 10/229 in both seasons. The differences among means of infestation % may be due to the differences of gossypol % in wall thickness of green boll. The means of gossypol% in green boll wall were (1.78, 1.32, 1.44 0.91, 1.75 and 1.16%) and the means of wall thickness were (2.77, 2.38, 2.07, 2.82 , 3.04 and 2.94 mm) for above cotton varieties and genotypes during both seasons, respectively. There were highly significant differences among the cotton varieties and genotypes in some characters.

INTRODUCTION

The Egyptian cotton lint is highly valued worldwide and is one of the major sources of foreign currency to the Egyptian national income. Cotton plants are subjected to attack by a wide range of insect pests throughout growing stages until near maturity. Spiny bollworm, is one of the most important pests of cotton and other crops in Egypt. This pest attacks the fruit bodies of cotton and cause heavy losses in the yield. This insect caused about 80% damage to cotton in south Khorrasan region of Iran (Fasell, 1977). So, far, the main method of controlling spiny bollworm on cotton has been the use of insecticides. Chemical control of this insect is expensive and serious peripheral problems have emerged, these include the development of insecticide-resistant insect strains, reduction of pest insect natural enemies, resurgence of pest populations in the absence of natural enemies, occurrence of secondary pests and environmental contamination. Therefore, it is strictly necessary to select resistant varieties as one of the simplest and useful tactics in integrated pest management programme. Differences in the susceptibility of cotton varieties to spiny bollworm infestation have been previously studied by Hassanein *et al.* (1969); Abd El-Rahim *et al.* (1976); Abdel-Bary *et al.* (1980); Abdel-Halim *et al.* (2000); El-Mezayyen (2004) and Al-Ameer *et al.* (2010). Therefore, the present study was initiated to determine the relative susceptibility of six cotton varieties and genotypes to infestation by *E. insulana* under field agro-conditions.

MATERIALS AND METHODS

Six Egyptian cotton varieties and genotypes were selected for this study namely: Giza 86, Pima S6 x Giza 77, Giza 92, Giza 94, Giza 86 x 10/229 and Giza 88. This experiment was conducted at Sakha Agricultural Research Station, Kafr El-Sheikh Governorate during two successive seasons 2012 and 2013 seasons.

Plot size measured 42 m² (1/100 feddan) with three replicates for each variety and genotype arranged in a complete randomized block. Therefore, the experimental field contained 18 plots. Cotton was sown on March 29th 2012 and March 20th, 2013, adopting the normal agricultural practices of irrigation and fertilization including three recommended insecticidal treatments as follows: 1st treatment: applied in July with Sylian 72% EC (0-4-bromo-2-chlorophenyl-O-ethyl-S-propyl phosphorothioate) at rate of 750 cm/feddan, and Dimeuron 10% EC, 1-[3,5-dichloro-4-(1,1,2,2-tetra-fluoroethoxy) phenyl]-3-(2,6-difluorobenzoyl) urea, at rate of 200 cm/feddan, 2nd treatment: in July and August (two weeks after the 1st treatment) with killeforon 5% EC (RS)-1-[2,5-dichloro-4-(1,1,2,3,3,3-hexafluoroproxy) phenyl]-3-(2,6-difluorobenzoyl), urea (at rate of 160 cm/feddan), and Cyperco 20% EC (RS)-a-cyano-3-phenoxy (1RS-3RS, 1RS-3SR)-3-(2, 2-dichlorovinyl)2, 2-dimethyl cyclopropane carboxylate, (at rate of 150 cm/feddan), and 3rd treatment: in August and September with Sylian 72% EC and Dimeuron 10% EC. Every insecticide was sprayed at 2 weeks intervals starting from July using the recommended rate of the proper insecticide diluted with 400 liters water. All sprays were conducted using knapsack sprayer (CP3 model). Weekly random samples of 50 green bolls were collected from each plot (150 green bolls each variety) and kept in tightly closed polyethylene bags, then dissected and examined in the laboratory of Plant Protection Research Institute at Sakha, Kafr El-Sheikh from the second week of July until the second week of October. Percentages of infestation by spiny bollworm were determined in each cotton variety and genotype. So, cotton yield g/m² of each variety and genotype was evaluated. 40 bolls each variety and genotype samples for estimating gossypol ratio in green boll wall were analyzed in the chemical laboratory of Chemical Research Department, Cotton Research Institute in Giza. The actual measurement of green boll wall thickness in millimeters was done as the distance across the fertile lemma and palea at the widest point (IRRI Stand, 1996). The statistical analysis was conducted using the software programme MSTATC.

RESULTS AND DISCUSSION

Inspections of infestation by spiny bollworm, *Earias insulana* in during the two seasons 2012 and 2013:

Results in Table (1) show the mean numbers of infested green boll by SBW larvae per 50 bolls for each of six varieties and genotypes of cotton, mean number of infested bolls were (5.28, 3.57, 4.22, 3.71, 3.29 and 5.71 bolls) for G.86, Pima S6 x Giza 77, G. 92, G. 94, G. 86 x 10/229 and G.88, respectively. The highest mean numbers of infestation bolls during this

season was estimated in G.88 and G.86 (5.71 and 5.28 bolls), respectively. While, the lowest was in G.86 x 10/229 and Pima S6 x G.77 (3.29 and 3.57 bolls), respectively.

Data presented in Table (2) indicate that the mean number of infested bolls by SBW larvae in season 2013, were (2.86, 4.07, 3.43, 2.36, 3.86 and 2.64 bolls) for the same varieties and genotypes, respectively. The highest mean number of infested bolls during this season was estimated in Pima S6 x G.77 and G.86 x 10/229 (4.07 and 3.86 bolls), respectively. While, the lowest was in G.94 and G.88 (2.36 and 2.64 bolls), respectively.

Table (1): Weekly mean number of infested green bolls by spiny bollworm, *E. insulana* (Boisd.) per 50 green bolls during season 2012.

Variety Date	G.86	Pima S6 x G.77	G.92	G.94	G.86 x 10/229	G.88
26/7	0	0	0	0	0	0
2/8	0	1	0	1	0	0
9/8	1	1	0	0	0	1
16/8	0	0	0	1	0	1
23/8	1	1	0	0	0	0
30/8	3	0	0	0	2	2
6/9	2	2	3	1	3	2
13/9	3	3	3	3	1	3
20/9	3	7	6	4	3	6
27/9	11	9	8	4	4	17
4/10	11	8	12	11	10	14
11/10	10	5	6	6	5	11
18/10	18	7	12	17	11	18
25/10	11	6	9	4	7	5
Total	74	50	59	52	46	80
G.M.	5.28	3.57	4.22	3.71	3.29	5.71

General means of infested green bolls from six varieties and genotypes of cotton during both seasons:

In both seasons (2012 and 2013), data in table (3) show that the general means of infestation ratios were (8.14, 7.64, 7.66, 6.08, 7.16 and 8.36%). The highest and lowest average means of data over both seasons (8.36 and 8.14%) and (6.08 and 7.16%) for (G.88 and G.86) and (G.94 and G.86 x 10/229), respectively. Data in Table (4) indicate the insignificant differences among varieties and genotypes (0.99) and insignificant differences between years (10.86), but between varieties and years was significant differences (3.34*). Many authors found differences between varieties and genotypes (early bloomer varieties) were most resistant for infestation by SBW larvae, such as (G.76, G.89, G.80 and G.31), while (late bloomer varieties and genotypes) were the most susceptible, such as (G.45, Bahtim 101, G.85, G.88, G.75 and G.70) (Hassanein *et al.*, 1969; Abdel-Rahim *et al.*, 1976; Abdel-Bary *et al.*, 1980;; Abou Toor *et al.*, 1989; Abdel-Halim *et al.*, 2000; and El-Mezayyen, 2004). So, Al-Ameer *et al.* (2010) found Karshensky 2 and G.70 (late bloomer) were most susceptible varieties, while Pima S6 x G.89 and Seuvin (early bloomer variety) were the most resistant during 2007 and 2008 seasons at Kafr El-Sheikh region.

Table (2): Weekly mean number of infested green bolls by spiny bollworm, *E. insulana* (Boisd.) per 50 green bolls during season 2013.

Variety Date	G.86	Pima S6 x G.77	G.92	G.94	G.86 x 10/229	G.88
25/7	0	0	0	0	0	0
1/8	0	0	0	0	0	0
8/8	2	0	0	0	0	0
15/8	5	1	1	2	3	1
22/8	4	5	3	2	8	4
29/8	2	3	1	2	5	0
5/9	5	4	4	1	3	3
12/9	1	4	3	2	2	4
19/9	1	1	4	2	3	2
26/9	0	1	3	2	1	3
3/10	2	4	3	2	3	2
10/10	4	8	6	4	6	4
17/10	8	16	12	8	12	8
24/10	6	10	8	6	8	6
Total	40	57	48	33	54	37
G.M.	2.86	4.07	3.43	2.36	3.86	2.64

Gossypol concentration in green boll wall during the two seasons 2012 and 2013:

Results in Table (3) show that the means of gossypol in green boll wall of varieties and genotypes of cotton were (1.78, 1.33, 1.44, 0.88, 1.75 and 1.16%) for G.86, Pima S6 x G.77, G.92, G.94, G.86 x 10/229 and G.88, respectively. The highest average of gossypol % during 2012 season was estimated in G.86 and G.86 x 10/229 (1.78 and 1.75%), respectively. While, the lowest was in G.94 and G.88 (0.88 and 1.16%), respectively.

Data presented in Table (3) indicate that the means of gossypol% in green boll wall in season 2013 were (1.77, 1.31, 1.43, 0.93, 1.74 and 1.15%) for the same varieties and genotypes of cotton, respectively. The highest average of gossypol% during this season was estimated in G.86 and G.86 x 10/229 (1.77 and 1.74%), respectively. While, the lowest was in G.94 and G.88 (0.93 and 1.15%), respectively.

Gossypol percentage from mean data during two seasons:

In both seasons (2012 and 2013) the averages of gossypol percentage as shown in Table (3) were (1.78, 1.32, 1.44, 0.91, 1.75 and 1.16%). The highest and lowest averages of gossypol% during the two both seasons, (1.78 and 1.75%) and (0.91 and 1.16%) for (G.86 and G.86 x 10/229) and (G.94 and G.88), respectively. Previous results mean that infestation of green bolls, correlated with gossypol %, so it showed that the highest varieties and genotypes of gossypol % was lowest in infestation by SBW larvae. While, the lowest varieties and genotypes of gossypol % was the highest infestation by SBW larvae. Results in Table (4) showed significant differences between varieties (0.685**), but it was insignificant differences between years (0.001^{NS}) and between (varieties and years) was (0.001^{NS}).

Table (3): Mean performance of varieties and genotypes of cotton under two years and their means for six traits:

Varieties	Infestation			Gossypol %			Boll wall thickness (mm)			Weight of bolls (g)			Yield (g)		
	2012	2013	Mean	2012	2013	Mean	2012	2013	Mean	2012	2013	Mean	2012	2013	Mean
G.86	5.28 a	2.86 b	4.07 a	1.78 a	1.77 a	1.78 a	2.75 ab	2.78 a	2.77 b	13.81 cd	13.22 ab	13.52 bc	825.00 a	303.33 bc	564.17 bc
Pima S6 x G.77	3.57 c	4.07 a	3.82 a	1.33 b	1.31 bc	1.32 c	2.37bc	2.38 b	2.38 c	13.66 cd	13.68 ab	13.67 b	460.00 c	445.00 ab	452.50 d
G.92	4.22 b	3.43ab	3.83 a	1.44 b	1.43 b	1.44 b	2.08 c	2.05 c	2.07 d	14.07 c	12.25 c	13.16 bc	910.00 a	363.33abc	636.67 ab
G.94	3.71 bc	2.36 b	3.04 c	0.88 d	0.93 d	0.91 e	2.83 a	2.81 a	2.82 ab	15.59 b	12.29 c	13.94 b	800.00 a	216.67 c	508.34 cd
G. 86x 10/229	3.29 c	3.86 a	3.58 b	1.75 a	1.74 a	1.75 a	3.02 a	3.06 a	3.04 a	17.50 a	14.83 a	16.17 a	920.00 a	513.33 a	716.67 a
G.88	5.71a	2.64 b	4.18 a	1.16 c	1.15 c	1.16 d	2.82 a	3.06 a	2.94 ab	12.87 d	11.74 c	12.31 d	675.00 b	215.00 c	445.00 d
Grand mean	4.30	3.20	3.75	1.39	1.39	1.39	2.65	2.69	2.67	14.58	13.00	13.80	765.00	342.78	553.89
LSD 0.05	0.60	0.73	0.40	0.11	0.16	0.09	0.42	0.29	0.24	0.97	0.83	0.60	122.00	188.00	104.85

In the same column, means followed by the same letter are not significantly different at 0.05 level of significance according to GMRT.

(4): Mean squares of years, varieties and genotypes of cotton and interaction for six traits.

Source of variance	df		Infestation			Gossypol %			Boll weight thickness (mm)			Weight of bolls (g)			Yield (g)		
	Single	Comb	2012	2013	Comb.	2012	2013	Comb.	2012	2013	Comb.	2012	2013	Comb.	2012	2013	Comb.
Years (Y)	-	1	-	-	10.86	-	-	0.001 ^{NS}	-	-	0.001 ^{NS}	-	-	22.56 ^{**}	-	-	1604449.4 ^{**}
Error (a)	-	4	3.48	0.914	2.20	0.012	0.004	0.008	0.07	0.06	0.03	0.52	0.07	0.29	7537.50	9684.72	8611.11
Varieties (V)	5	5	2.91	1.427	0.99	0.360*	0.325 ^{**}	0.685 ^{**}	1.055 ^{**}	1.093 ^{**}	2.146 ^{**}	8.50 ^{**}	3.89 ^{**}	10.02 ^{**}	90600.00 ^{**}	44248.88*	69199.44 ^{**}
V x Y	-	5	-	-	3.34*	-	-	0.001 ^{NS}	-	-	0.002 ^{NS}	-	-	2.37 ^{NS}	-	-	65649.40 ^{**}
Error (b)	10	20	0.114	0.109	0.112	0.004	0.008	0.006	0.054	0.026	0.040	0.29	1.79	0.251	4567.5	10679.72	7623.61
CV%			7.87	10.31	8.91	4.76	6.27	5.57	7.80	5.38	6.70	3.74	10.30	7.41	8.83	30.15	16.00

*, ** significant at 0.05 and 0.01 levels of probability, respectively.

Data in Table (5) show that the correlation coefficient was negative and significant between infestation green bolls and concentration of gossypol (-0.451*).

Table (5): The correlation coefficient among all studied characters

Traits	Infestation %	Gossypol %	Bollwall thickness(mm)	Weightof bolls (g)	Yield (g)
Infestation%	-	-0.451*	-0.487**	-0.085 ^{NS}	-0.540**
Gossypol %		-	-0.466**	0.307 ^{NS}	0.222 ^{NS}
Boll wall thickness (mm)			-	0.194 ^{NS}	-0.129 ^{NS}
Weight of bolls (g)				-	0.613**
Yield (g)					-

These previous results were in agreement with Al-Ameer *et al.* (2010) who found that the correlation between concentration of gossypol and insect infestation was significant and negative (-0.551*). Also, Abou-Toor (1986) estimated that the correlation was negative and significant between resistance to bollworms infestation and number of glands/cm² of boll and total gossypol contents. According to Bottger (1964) gossypol is also toxic to cotton bollworm, furthermore Shaver and Lukefahr (1969) showed the effect of gossypol (concentration) on bollworms and budworms. Also, Vilkova (1989) reported that even though high gossypol lines had weight when compared to those on low gossypol lines, the larval from the high gossypol lines that survived had a higher pupal weight because of their apparent resistance to gossypol, but fecundity of these survivors was significantly reduced. Abd El-Hamid and Helw (1973) and Meisner *et al.* (1977) suggested that gossypol content may be one of the factors associated with resistance to cotton leafworm, so these genotypes can be used as a stock in breeding programs or using in the direct and general agriculture.

Green boll wall thickness (mm) during the two seasons 2012 and 2013:

Results in Table (3) show that the means of green boll wall thickness of 6 varieties and genotypes of cotton, were (2.75, 2.37, 2.08, 2.83, 3.02 and 2.82 mm) for G.86, Pima S6 x G.77, G.92, G.94, G.86 x 10/229 and G.88, respectively. The highest mean of green boll wall thickness during this season was estimated in G.86 x 10/229 and G.94 (3.02 and 2.83 mm), respectively. While, the lowest was in G.92 and Pima S6 x G.77 (2.08 and 2.37 mm), respectively.

Data presented in Table (3) indicate that the means of green boll wall thickness were (2.78, 2.38, 2.05, 2.81, 3.06 and 3.06 mm) for the same varieties and genotypes of cotton, respectively. The highest mean of green boll wall thickness was estimated in G.86 x 10/229 and G.88 (3.06 and 3.06 mm), respectively. While, the lowest was in G.92 and Pima S6 x G.77 (2.05 x 2.38 mm), respectively.

Green boll wall thickness from mean data during both seasons:

In both seasons (2012 and 2013), the green boll wall thickness from mean data (Table 3) were (2.77, 2.38, 2.07, 2.82, 3.04 and 2.94 mm). The highest and lowest average means of green boll wall thickness from mean data over both seasons (3.04 and 2.94 mm) and (2.07 and 2.38 mm) for (G.86 x 10/229 and G.88) and (G.92 and Pima S6 x G.77), respectively.

Results in Table (4) reveal that the significant difference among varieties was (2.146**), but it insignificant differences between years (0.001^{NS}) and between (varieties and years) (0.002^{NS}). Also, data in Table (5) show that the value of correlation was negative significant (-0.485**) between infestation of green boll by larvae SBW and green boll wall thickness.

Boll weight (g) during the two seasons 2012 and 2013:

Results in Table (3) show the means of boll weight of 6 varieties and genotypes of cotton were (13.81, 13.66, 14.07, 15.59, 17.50 and 12.87 g) for G.86, Pima S6 x G.77, G.92, G.94, G.86 x 10/229 and G.88, respectively. The highest means of boll weight were estimated in G.86 x 10/229 and G.94 (17.50 and 15.59 g), respectively. While, the lowest ones were in G.88 and Pima S6 x G.77 (12.87 and 13.66 g), respectively.

The data presented in Table (3) indicated that the means of boll weight, were (13.22, 13.68, 12.25, 12.29, 14.83 and 11.74 g) for the same varieties and genotypes of cotton, respectively. The highest means of boll weight were estimated in G.86 x 10/229 and Pima S6 x G.77 (14.83 and 13.68 g), respectively. While, the lowest ones were in G.88 and G.92 (11.74 and 12.25 g), respectively.

Boll weight from mean data during both seasons:

In both seasons (2012 and 2013), the boll weight from mean data (Table 3), were (13.52, 13.67, 1.16, 13.94, 16.17 and 12.31 g). The highest and lowest average of boll weight were (16.17 and 13.94 g) and (12.31 and 13.16 g) for (G.86 x 10/229 and G.94) and (G.88 and G.92), respectively. The results presented in Table (4) showed significant differences among varieties (10.02**), so, it was between years (22.56**), but it was insignificant difference between years and varieties (2.37^{NS}). Previous results in Table (5) showed the values of correlation was negative insignificant (-0.045^{NS}) between infestation green bolls and boll weight.

Yield/m² of varieties and genotypes of cotton during the two seasons 2012 and 2013:

Results in Table (3) show that the means of yield of 6 varieties and genotypes of cotton were (825, 460, 910, 800, 920 and 765 g) for G.86, Pima S6 x G.77, G.92, G.94, G.86 x 10/229 and G.88, respectively. The highest means of yield were estimated in G.86 x 10/229 and G.92 (920 and 910 g), respectively. While, the lowest ones were Pima S6 x G.77 and G.88 (460 and 675 g), respectively.

Data presented in Table (3) indicate that the means of yield, were (303.33, 445.00, 363.33, 216.67, 513.33 and 215.00 g) for the same varieties and genotypes of cotton, respectively. The highest means of yield during this season was estimated in G.86 x 10/229 and Pima S6 x G.77 (513.33 and 445.00 g), respectively. While, the lowest ones were in G.88 and G.94 (215.00 and 216.67 g), respectively.

Yield/m² of varieties and genotypes of cotton from mean data during both seasons:

In both seasons (2012 and 2013), the yield/m² from mean data (Table 3) were (564.17, 452.50, 636.67, 508.34, 716.67 and 445.00 g/m²). The highest and lowest average of yield/m² from mean data during both seasons were (716.67 and 636.67 g/m²) and (445.00 and 452.50 g/m²) for (G.86 x

10/229 and G.92) and (G.88 and Pima S6 x G.77), respectively. Data in Table (5), show the value of correlation was negative insignificant (-0.263^{NS}) between infestation green bolls and yield/m². But, Al-Ameer *et al.* (2010) found the value of correlation was negative significant between infestation and lint yield/m² (-0.684^*).

REFERENCES

- Abdel-Bary, A.A.; M.A. Omer and M.M. Shahbander (1980). Evaluation of bollworm resistance in some Egyptian cotton cultivars and strains. *Alex. J. Agric. Res.*, 28(1): 91-96.
- Abdel-Halim, A.; K.A.M. El-Khawass and H.A.F. Salem (2000). Boll infestations with pink bollworm *Pectinophora gossypiella* (Saund.) and spiny bollworm (*Earias insulana* (Boisd.) on certain cotton varieties in Damietta Governorate. *Al-Azhar J. Agric. Res.*, 31: 149-155.
- Abdel-Hamid, A.M. and M.R. Helw (1973). Cotton leaf worm on different genotypes of cotton. *Cott. Gr. Rev.*, 50(4): 338.
- Abd El-Rahim, W.A.; S.M.I. Metwally and F. El-Dakrouy (1976). Susceptibility of some Egyptian cotton varieties to the infestation by *Pectinophora gossypiella* (Saund.) and *Earias insulana* (Boisd.). *J. Agric. Res. Tanta Univ.*, 2(2): 332-338.
- Abou-Toor, H.B. (1986). Genetical studies on resistance to insects in Egyptian cotton. Ph.D. Thesis, Fac. Agric., Tanta Univ., 210 pp.
- Abou-Toor, H.B.; M.M. Abou-Kahla; A.A.S. El-Zanan and I.A.I. Helal (1989). The susceptibility of nine Egyptian cotton cultivars to infestation with cotton insect pests. The 7th Arab Pesticide Conference, Tanta Univ., Sept., 11-12.
- Al-Ameer, M.A.; M.A. Abd El-Salam; W.M.B. Yehia and I.A.I. Saad (2010). Evaluation of some cotton genotypes for ability to infestation tolerance to bollworm for improving of some important economical characters. *J. Agric. Res., Kafr El-Sheikh Univ.*, 36(2): 147-172.
- Bottger, G.T. (1964). Relation of gossypol content of cotton plants to insect resistance. *J. Econ. Entomol.*, 57: 283-285.
- El-Mezayyen, G.A. (2004). Evaluation of some Egyptian cotton varieties of infestation of piercing sucking insects, cotton leafworm and bollworms at Kafr El-Sheikh region. *J. Agric. Sci., Mansoura Univ.*, 29(1): 433-441.
- Fasell, M.D. (1977). Investigation on the biology, ecology and control of *Earias insulana* (Boisd) (Noctuidae). *Entomol. Phytopath. Appl.*, 43: 39-54.
- Hassanein, M.H.; M. Hafez and G.A. Rizk (1969). The susceptibility of certain cotton varieties to bollworms infestation. *Bull. Soc. Ent. Egypte*, 53: 261-269.
- IRRI (1996). Standard evaluation system for rice INGER Prog. International Rice Research Institute July Edition, p. 12, 13, 43, 47.

- Meisner, J.; M. Zur; E. Kabonci and K.R.S. Ascher (1977). Influence of gossypol content of leaves of different cotton strains on the development of *Spodoptera littoralis* larvae. J. Econ. Entomol., 70(6): 714-716.
- Shaver, T.N. and M.J. Lukefahr (1969). Effects of flavonoids pigments and gossypol on growth and development of the bollworm, tobacco budworm, and pink bollworm. J. Econ. Entomol., 62: 643-646.
- Vilkova, N.A. (1989). Effect of cotton cultivars with high content of gossypol on development of the cotton bollworm *Heliothis armigra* (Hbn) (Lepidoptera: Noctuidae). J. Econ. Entomol. Rev., 68: 129-137.

تأثير اصناف وطرز القطن المختلفة للإصابة بدودة اللوز الشوكية فى محافظة كفر الشيخ

حسن محمد حسن صومع و أسماء محمد على الغبارى
معهد بحوث وقاية النباتات - مركز البحوث الزراعية - الدقى - مصر

أجريت هذه التجارب خلال موسمين ٢٠١٢، ٢٠١٣ فى مزرعة محطة البحوث الزراعية - بسخا - محافظة كفر الشيخ بهدف دراسة تأثير اصناف وطرز القطن المختلفة للإصابة بدودة اللوز الشوكية فى محافظة كفر الشيخ وهى جيزة ٨٦، بيما س ٧٧، جيزة ٧٧، جيزة ٩٢، جيزة ٩٤، جيزة ٨٦ × ٢٢٩/١٠، جيزة ٨٨. تم فحص ثلاث مكررات لكل صنف أسبوعيا كل مكررة عبارة عن ٥٠ لوزة بداية من أواخر شهر يوليو حتى أواخر شهر أكتوبر لتقدير نسبة الإصابة بدودة اللوز الشوكية وأشارت النتائج إلى نسب الإصابة التالية (٨.١٤%، ٧.٦٤%، ٧.٦٦%، ٦.٠٨%، ٧.١٦%، ٨.٣٦%) للأصناف السابقة بالترتيب وكانت أعلى نسبة إصابة مسجلة فى صنفى جيزة ٨٨ وجيزة ٨٦ فى كلا الموسمين، بينما أقل نسبة إصابة سجلت فى صنفى جيزة ٩٤ وجيزة ٨٦ × ٢٢٩/١٠ فى نفس الموسمين. وهذه الاختلافات بين الأصناف نتيجة الاختلافات فى نسبة مادة الجوسيبول فى جدار لوزة القطن فى العامين وهى (١.٣٢%، ١.٤٤%، ٠.٩١%، ١.٧٥%، ١.١٦%) للأصناف السابقة فى كلا الموسمين بالترتيب، وكان متوسط جدار اللوزة (٢.٧٧، ٢.٣٨، ٢.٠٧، ٢.٨٢، ٣.٠٤، ٢.٩٤ مم) للأصناف السابقة فى الموسمين بالترتيب مع وجود اختلافات بين الأصناف فى صفات أخرى مدروسة ولكنها ليست معنوية بالنسبة للإصابة.