

EFFICACY OF SOME BIOAGENTS AND NEMASTOP COMPOUND IN CONTROLLING ROOT KNOT DISEASE ON PEANUT.

Howida A. Metwaly¹ and Hanaa S. Zawam²

1- Central Laboratory of Organic Agriculture, Agricultural Research Center, Giza, Egypt.

2- Nematode Research Department, Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt.

ABSTRACT

Biological control of plant diseases especially root knot nematode has been handled in many scientific papers. *Bacillus subtilis*, *Trichoderma harzianum* (T1), *T. viride* (T2), mixture of both (T1 and T2) in addition to the commercial product Nemastop were used under field conditions to control root knot nematode, *Meloidogyne javanica* Chitwood on peanut plants which was the most frequent extracted from soil and roots.

The efficacy of the treatments at different concentrations was also assayed under laboratory conditions as percentage of juvenile mortality and inhibition of egg hatching. All treatments revealed good effect in controlling root-knot at the highest concentrations, Nemastop was the most effective at (1:10), whereas, *B. subtilis* was the lowest one. Results obtained from field experiments were in harmony with those obtained from laboratory compared with control. Efficacy of the treatment also positively correlated with number of application time. The plot treated with any one of the bioagents tested three times achieved highest effect in controlling root knot nematode and increased peanut yield. Under field condition mixture of *Trichoderma* isolates was the most effective followed by Nemastop.

Keywords: Biological agents, Root-knot nematode, *Meloidogyne javanica*, Peanut, *Bacillus subtilis*, *Trichoderma harzianum*, *T. viride*.

INTRODUCTION

Peanut (*Arachis hypogaea* L.), is very important annual herb legume crop of tropical and subtropical areas of the world. The importance of peanut as a food and oil crop and as a cash source in the semi-arid tropics is well known. Peanuts are rich in energy and contain health benefit nutrients, minerals, antioxidants and vitamins that are essential for optimum health (Umesh 2009). The production of peanut is negatively affected by soil borne fungi and nematodes. Root-knot nematodes (*Meloidogyne* spp.) are considered the most important plant parasitic nematodes attacking more than 2000 plant species (Jung and Wyss 1999). Reduction in yield of peanut was recorded due to more than 55 pathogens including fungi, bacteria, viruses, mycoplasma, nematodes and parasitic flowering plants (Podile and Kishore 2002). Plant parasitic nematodes cause significant damage to agriculture throughout the world. The damage caused by the root-knot nematodes are much higher in tropical and sub-tropical countries compare with cold ones (Sasser and Freckman 1987). The presence of root-knot nematodes in the roots induces galls that restrict nutrient, water uptake and peanut growth, while also facilitating fungal infections. Root-knot nematodes are the most

significant nematode pest of Egyptian peanut and can cause up to 60% yield loss (Sharma and McDonald 1990).

In recent years, the awareness of the fungicides and nematicides hazards to human and environment has directed the attention towards soil-borne antagonists and natural products to control nematodes in replacing of chemicals. Biological control is gaining increase role throughout the world for nematode suppression. The genus, *Trichoderma* is common saprophytic filamentous imperfect fungus and considered as the most common fungi in the rhizosphere and found in almost all soil type. Sharon *et al.*, 2001 reported that *T. harzianum* was antagonistic organism to root knot pathogen *M. javanica* in soil. Other *Trichoderma* species and isolates have also exhibited significant biocontrol activity against *M. javanica* in growth chamber experiments.

Biological control is gaining increasing role throughout the world for soil pathogens and nematodes suppression. Mixing antagonists with each others may be lead to antagonistic effect consequently decrease efficacy of treatment or lead to synergistic effect and increase the efficacy (Robinson *et al.*, 2009).

Bacillus subtilis is reported as a biocontrol agent against root-knot nematodes. (Khan *et al.*, 2002, Huang *et al.*, 2005 and Huang *et al.*, 2009). Several reports clarified that the basic mechanisms of *B. subtilis* included production of extracellular antibiotics metabolites or enzymes (e.g. proteases, chitinases and glucanases), stimulation of host defenses, incensement of plant growth, induced systemic resistance in plants, suppression of the plant diseases and secreting volatile nematicidal substances (Abd- El- Moneim, 2005, Ji *et al.*, 2006, Kloepper and Ryu 2006 and Lahlali *et al.*, 2013).

The present study is conducted to control the root knot nematode on peanut plants, different biocontrol agents *T. harzianum*, *T. viride*, mixture of both and *B. subtilis* as well as Nemastop are used to improve protection effect against this disease and to increase the peanut yield. Also, to determine the effect of these biological treatments compared with control under laboratory as well as field conditions.

MATERIALS AND METHODS

Bioagents and biocide:

Different biocontrol agents, *B. subtilis*, *T. harzianum* (T1), *T. viride* (T2), mixture of both T1 and T2 and a commercial biocide, Nemastop were kindly obtained from central lab of Organic Agricultural Research Center, Giza, Egypt. *B. subtilis* was grown on nutrient glucose broth (NGB) suggested by Dowson (1957) *T. harzianum* (T1), *T. viride* (T2) were grown in liquid gliotoxin fermentation medium (GFM) (Brian and Hemming 1945). The bioagents were allowed to grow under complete darkness for nine days just to stimulate toxin production at 28°C (Abd-El-Moity and Shatla 1981). Different bioagents were formulated as suspension using method developed by Abd-El-Moity (1985). Prepared suspension was adjusted to contain 30×10^6 cfu /ml and mixture of them was added as (1:1). Nemastop was used as

commercial biocide to compare its effect with other bioagents against root knot nematode on peanut. Peanut shoot weight, shoot length, root weight, root length and yield were determined.

Extraction and classification of Nematodes:-

Nematode infestation was verified by analysis of soil samples by a laboratory specialist. Different kinds of nematode infested the soil samples associated with rhizosphere of roots and root galls which showed identical galls root knot symptoms from peanut plants collected through the experimental duration in two seasons in El-Bustan Research Station, Nubaria–Behira Governorate, Egypt. An aliquot of 250cm³ from each soil sample was processed for nematode extraction by sieving and modified Bearman technique (Goodey, 1957). After 48 hours the extracted nematodes were counted using 1 ml Hawksly counting slide under a stereo microscope and classified according to (Mai and Lyon 1975 and Siddiqi 1986) females of root knot nematode were excised from large galls on the roots of peanut plants and individually macerated and accomplished with automated apparatus(Phast System Pharmacia, Uppsala,Sweden) to have esterase phenotype (Tomaszewski,et al.,1994)

Effect of biocontrol agents on *M. javanica* under laboratory and field conditions

I- Laboratory experiment:-

To study the efficacy of biocontrol agents on *M. javanica* under laboratory conditions, one ml of each biocontrol agents (*B.subtilis*, *T. harzianum* (T1), *T. viride* (T2) and mixture of (T1and T2) were containing (30 x 10⁶) cfu /ml and Nemastop were added to nematode suspension at different concentrations (1:10, 1:25 and1:50) in glass vials for each of them. Each of biocontrol agent suspension was added to five handpicked egg-masses and to 500 active nematode juveniles to detect their effect on hatching and juveniles mortality of *M. javanica*. The same number of egg-masses and juveniles received distilled water and served as control. Each treatment was applied in three replicates. The percentage of inhibition in egg hatching was recorded after 3 days and the percentage of Juveniles mortality was recorded after 48 hours under a stereoscopic microscope.

II- Field experiments:-

Field experiment was carried out in naturally heavy infested field with *M. javanica* at El-Bustan Research Station, Nubaria–Behira Governorate, Egypt in April 2012 / 2013 growing seasons. A randomized complete block design with 5 replications was used in each season. A field experiment consisted of plots (7 × 6 m); each comprised of 10 rows 20 cm distance and 30 holes/row at which each treatment was conducted with five plots as replicates as well as check treatment. Gliotoxin fermentation medium (GFM) used to grow the bioagents. *B. subtilis*, *T. harzianum* (T1), *T. viride* (T2), mixture (T1andT2) and Nemastop were used as seed coating before sowing and sprinkled as a liquid at rate of 10 L/fed. after 15 days of sowing(two times applications) and after month of sowing (three times applications).

Plants only treated with water act as control treatment. Peanut seeds Giza 6 obtained from Field Crop Research Institute, Agricultural Research

Centre, Giza. All treatments received the same agricultural treatment such as amount of water, number of seeds /plot and amount of fertilizers.

Effect of applications time using biocontrol agents against (*M. javanica*) in peanut.

Different bioagents were used as one, two and three applications with interval 15 days between applications. Biocontrol agents were used in liquid form and the first treatment was applied as seed coating in the suspension for 20 minutes, where the second and third treatments were applied as soil drench, at five liters of each bioagent / plot. Treated plants were examined at the end of seasons. The roots were then washed to get rid of the adhering and particles to determine the number of root knot galls, juveniles in soil, eggs /5 gm roots, developmental stages in the roots. The number of nematode larvae (in 250 cm³ soil) and RF (Reproduction Factor) was determined according to Norton (1978).

Reproduction Factor (RF) =

$$\frac{\text{No. Eggs + Developmental stages + Free Nematode in soil+ adult females}}{\text{Initial Population}}$$

Developmental stages (DS) =

number of developed juveniles (third and fourth stages) embedded in the roots.

Efficiency of each treatment recorded according to the equation:-

$$\text{Efficiency} = \frac{\text{RF in control} - \text{RF in treatment}}{\text{RF in control}} \times 100$$

At harvest some peanut parameters such as shoot length (cm), shoot weight (g), root length (cm), root weight (g) and yield of pods with kilogram per plot were determined and the efficacy of each bioagents were recorded as average of two seasons.

Statistical Analysis:-

Data obtained were subjected to statistical analysis of variance for completely randomized design (CRD) and randomized complete block design (RCBD) as outlined by Steel and Torrie (1980).

RESULTS AND DISCUSSION

Nematode associated with peanut :-

Data in Table (1) show the percentage frequency of the nematodes recovered species from El-Bustan Research Station, Nubaria-Behira governorate, *Criconemella* sp., *M. javanica*, *Paratrichodorus minor*, *Pratylenchus* sp. and *Tylenchorhynchus* sp. Lamberti (1981) stated that, plant parasitic nematodes account worldwide, for an average estimated by 10-20% yearly loss of agricultural peanut plants which multiplication and spread over thousands of feddans under best conditions. The most dominant root knot nematode was *M. javanica*. Root knot nematodes cause varying degrees of stunting, chlorosis depending on the initial population. (Taylor and Sasser, 1978) showed that more than 80% of the major *Meloidogyne* population belonged to *M. javanica*. Root-knot nematodes (*M. javanica*) are among the most damaging nematodes in agriculture, causing an estimated US\$100 billion loss/year on peanut plants worldwide (Oka *et al.*, 2000). *M. javanica* Chitwood

and other root-knot nematodes cause galls in roots of many crops impeding normal uptake of water and nutrients.

Table (1): The percentage frequency of nematodes attacked peanut plants.

Nematode genera	Frequency of occurrence
<i>Criconemella</i> sp.	1.37
<i>Meloidogyne javanica</i>	70.09
<i>Paratrichodorus minor</i>	3.50
<i>Pratylenchus</i> sp.	3.43
<i>Tylenchorhynchus</i> sp.	21.61


Data in Table (1) indicate that, *M. javanica* was the most frequent species of overall the seasons of peanut with frequent percentages of (70.09%) while, *Criconemella* sp. was the least frequent species with frequent percentages of (1.37%).

Effect of different bioagents on the viability of *M. javanica*:-

The morphological change of *M. javanica* juveniles was examined using an inverted microscope during 7 days of incubation at 27 °C. Deforming of juveniles was showed in Figures (1: 5), for treatments and the figure (6) for untreated control. Some juveniles appeared thin with destruction in some parts, but no deforming was observed in the control. Observations through the inverted microscope demonstrated that the microorganisms were widely attached the juveniles of *M. javanica*. The morphological changes of *M. javanica* juveniles agreed with the study of Westcott and Kluepfel (1993) who reported that chitinases produced by bacteria was potent in attacking the cuticle of *M. javanica*. Regarding to *Trichoderma* species (Ghisalberti 2002) reported that chitinolytic enzymes produced by these fungi are thought to be responsible for the degradation of cell walls; they also had ability to produce a wide range of secondary metabolites with diverse biological actions including mycoparasitism also through production of antibiotic substances. *T. viride* and *T. harzianum* destructed and lysis the juveniles body after seven days. The effect of Nemastop on root knot nematode might be due to alkyl cysteine sulphoxides which released a mixture of volatile alkyl thiols and sulphides, these volatile compounds override the inhibitory effect on parasitic nematode (Coley-Smith, 1976)). Results of an antagonism test showed in (Table 2) had significant inhibition and mortality as affected by treatments of *B. subtilis*, *T. harzianum*, *T. viride*, mixture of them (T1,T2) and Nemastop compound .




1- *Tricoderma harzianum* (T1)



2- *Trichoderma viride* (T2)




3- (T1 and T2)



4- Nemastop



5- *Bacillus subtilis*



6- untreated control

Fig. (2): Interaction between bioagents and juvenile of pathogenic nematode

These bioagents had various degrees of effectiveness toward the nematode juveniles. Moreover, the percentage of mortality increased as increase of the concentration. Results also revealed there is positive correlation between different concentrations of treatments and efficacy of application. The obtained results might be due to the increase in the amount of the active component of bioagent. Nemastop was the most effective treatment of the three concentrations. The highest concentration (1:10) for all the tested treatments achieved the highest percentage of juvenile mortality and inhibition of eggs hatching. The inhibition in egg-hatching caused by all the used treatments ranged from 100% for both Nemastop and *T. harzianum* to 56% for *T. viride*. The lowest concentration (1:50) recorded the lowest percentage of egg hatching inhibition. Rahman Khan *et al.*, (2005) mentioned that *B. subtilis* produce nematotoxic metabolites that involved in the nematode suppression. The morphological change of *M.javanica* juveniles agreed with the study of Westcott and Kluepfel (1993) who reported that chitinases activity produced by PGPR were more potent in attacking eggs than the cuticle of *M.incognita*, this might have resulted from the direct damage caused by the bacterial chitinase.

Table (2): The inhibition and mortality of *M. javanica* as affected by biocontrol agents.

Treatments	Concentrations	Inhibition of egg hatching%	Mortality of juvenile %
<i>B. subtilis</i>	1/10	59	75
	1/25	42	47
	1/50	19	30
<i>T. harzianum</i> (T1)	1/10	100	96
	1/25	96	79
	1/50	79	68
<i>T. viride</i> (T2)	1/10	56	87
	1/25	33	61
	1/50	20	36
Mixture of (T1 +T2)	1/10	94	95
	1/25	87	89
	1/50	73	78
Nemastop	1/10	100	98
	1/25	95	95
	1/50	88	83
Control		0	0
L.S.D. at 5%(TREATMENT)		2.625	1.977
L.S.D. at 5%(CONCN.)		1.856	1.398

Effect of applications time using biocontrol agents against (*M. javanica*) in peanut.

To determine the suitable time for adding different antagonist to get the highest effect, different bioagents were added at three different times. These adding times were at planting time, fifteen days after planting and one month after planting. The early treatment showed the highest effect in plant protection and vice versa. Time of application was reflected on the yield and percentage of dry matter in treated plants. This is due to that early treatment,

at planting time, allows propagules of used antagonist to spread out and surround newly developed roots causing protection against root knot nematode that may be found in court of infection. On the contrary, adding antagonist fifteen days and one month after planting this means that, the developed roots before this period, already attacked by pathogens which negatively affect plant growth and developing (Mannanov and Sattar 2009). That is why plants treated very early, at planting time, gave the highest yield and highest percentage in dry matter.

Data in Table (3) showed that, significant differences in the number of developmental stages and eggs /5g roots, Juveniles/ 250 cm³ soil between peanut plants treated one time and those treated two or three times. In plants treated at three times, the percentage of nematodes was found to be less than those recorded after one or two times. This might be due to appearance of new roots on the treated plants free from the nematode infection. Obtained data also showed that, all antagonists significantly reduced the number of root galls, developmental stages and juveniles in soil. Mixture of *Trichoderma* spp. isolates showed good effect in controlling nematode. Mixing different isolates may increase the scope on mode of action consequently increase efficacy of the treatment this agree with Zhang and Zhang (2009) who showed that combination of isolated *T. viride* strain and *T. harzianum* reduced root galling and the number of new infection by nematode. *T. harzianum* parasitized eggs and larvae of nematode. The hyphae penetrate the eggs and larval cuticle by dissolving the chitin layer through enzymatic activity. They proliferate within the organism and produce toxic metabolites (Dos Santos *et al.*, 1992). Thus, the enzymes produced by *Trichoderma* spp. such as chitinases, glucanases and proteases seem to play an important role in parasitism (Haran *et al.*, 1996). *Trichoderma* has not only been proved to parasitize nematodes and inactivate pathogen enzymes but also help in tolerance by enhanced root development (Harman, 2000). Sharma and Pandey (2009) found that *T. harzianum* alone can reduce gall formation and improved plant growth. As known by Li *et al.* (2007) and (Kim *et al.*, 1997) that *B. subtilis* produce antibiotics and antifungals that performed as bio-pesticide against nematode, as bacillomycin, iturin, surfactin, and agrocin. The reduction of nematode plant parasitic nematodes associated with *B. subtilis* may be attributed to diverse mechanisms which involve phytohormones production, mineral solubilisation, reduction of the activity of egg hatching factors, alteration of root exudates and inhibition of nematode penetration into the roots thereby interfering with host finding process and reducing galls

Table (3): Effect of applications on number of using biocontrol agents against (*Meloidogyne javanica*) in peanut plants.

Treatments	Application time	Number of developmental stages/5 g roots		Eggs /5 g roots		Juveniles /250 cm ³ soil		Galls/ root		Rf		Effecacy%	
		2012	2013	2012	2013	2012	2013	2012	2013	2012	2013	2012	2013
<i>B.subtilis</i>	One time	312	199	184	159	2100	1800	275	459	3.65	3.23	85.5	87.6
	Two time	122	163	92	89	1120	730	113	153	1.34	.83	94.7	96.8
	Three time	55	63	43	31	640	250	64	99	.53	.30	97.9	98.8
<i>T.harzianum</i> (T1)	One time	98	213	95	113	2730	840	256	147	7.69	2.59	69.5	90.0
	Two time	67	109	39	39	1250	560	164	109	3.15	1.23	87.5	95.3
	Three time	31	89	27	24	840	280	98	104	1.72	0.36	93.2	98.6
<i>T.viride</i> (T2)	One time	156	218	189	152	1450	1750	184	152	3.14	3.75	87.6	85.6
	Two time	107	115	114	102	860	950	87	124	1.44	1.70	94.3	93.5
	Three time	52	55	76	41	580	620	63	31	0.61	0.56	97.6	97.8
Mixture of (T1 and T2)	One time	81	179	95	103	1920	380	215	203	5.68	1.50	77.5	94.2
	Two time	43	97	60	71	620	190	109	83	2.57	0.89	89.8	96.6
	Three time	22	18	51	21	210	83	38	21	0.68	0.38	97.3	98.5
Nemastop	One time	64	317	236	234	1500	350	378	234	1.89	1.05	92.5	95.9
	Two time	27	121	80	62	380	220	121	127	1.55	1.03	93.9	96.0
	Three time	11	23	40	29	130	90	34	29	1.36	0.52	98.6	98
Control		942	498	265	218	2400	3900	681	664	25	25.98		
LSD at 5%		5.6	2.751	5.5	1.275	15.28	58.9	4.293	2.126				

Data in Table (4) showed that positive correlation between numbers of application and plant growth improvement. All treatments either bioagent alone or in combination with each other as well as commercial biocide led to high significant effects on yield, shoots and roots (weight and length). Increase number of application to be two or three times led to significant increase in efficacy due to increase establishment of the antagonist which led to increase in yield of treated plants. This may be due to repeated treatment with bioagents which increase growth regulators and increase growing roots and also may increase plant nutrients uptake. Data also recorded that the mixture of *Trichoderma* isolates occupied the highest rank (20.71kg/plot) compared with control treatment at the third application. This may be explain as compatible relation between mixture of *Trichoderma* isolates, led to synergistic effect between them and increasing the uptake of macro and micro nutrients and improving all vegetative characters. This increase in yield may be due to that rhizosphere organisms produced greater amount of organic acids, such as tartaric, citric acid and lactic acid which may improve plant productivity and increase the root system growth.

Table(4):Effect of applications number using biocontrol agentsin improving agronomic characteristic on peanut plants.

Treatments	Applicatio n time	Shoots		Shoot		Roots		Root		Yield	
		Weight(g.)		Length(cm.)		Weight(g.)		Length (cm.)		kg / plot	
		2012	2013	2012	2013	2012	2013	2012	2013	2012	2013
<i>Bacillus subtilis</i>	One time	32.98	73.7	27	29	25	39.03	25	22	9.28	11.2
	Two time	26.89	147.4	29	30	27	42.32	18	25	12.41	55.4
	Three time	53.84	203.1	43.3	44	30	57.52	30	25	14.17	72
<i>Trichoderma harzianum</i> (T1)	One time	41.9	119.2	25	30	20	44.39	25	29	8.12	42.8
	Two time	42.87	140.3	26.8	38	20	66.05	27	30	9.47	53
	Three time	37.5	142.1	28.3	39	30	71.47	20	30	11.64	55.34
<i>Trichoderma viride</i> (T2)	One time	39.94	100.6	27	31	20	50.4	30	28	3.59	44.41
	Two time	36.2	121.2	30	35	30	50.49	20	30	7.81	49.14
	Three time	48.76	182.3	37.7	39	36.7	71.03	30	33	11.69	91.2
Mixture of (T1+T2)	One time	31.5	138.4	27.7	30	25	36.24	20	30	13.12	66.5
	Two time	62.04	148.6	32	34	30	41.88	36.7	35	13.57	67.8
	Three time	47.28	162.3	45	44	35	59.07	25	40	20.71	79.13
Nemastop	One time	55.31	140	22.7	31	30	43.82	35	30	12.61	70.15
	Two time	38.01	153.1	35	39	30	50.64	30	32	13.34	81.49
	Three time	79.33	192.5	42	47	35	84.03	35	45	14.25	107.69
Control		18	33.5	33.7	27.5	29	33.5	33.5	21	4.33	13.5
LSD at 5 %		2.92	1.774	0.747	1.131	0.90	0.689	0.88	1.077	0.45	1.171

REFERENCES

- Abd-El-Moity, T. H. and M. N. Shatla (1981). Biological control of white rot disease of onion (*Sclerotium cepivorum*) by *Trichoderma harzianum*. *Phytopathology, Z.*, 100: 29- 35.
- Abd-El-Moity, T.H. (1985). Effect of single and mixture of *Trichoderma harzianum* isolates on controlling three different soil borne pathogens. *Egypt. J. Microbial., Special Issue*, 111-120.

- Abd-El-Moneim, Maisa ; L . (2005). Effect of some non-chemicals, seed and foliage applications on cotton damping off disease incidence and yield characteristics. Zagazig, J. Agric. Res., 32 (6):1829 – 1853.
- Brian, P.W. and H.G.Hemming (1945). Gliotoxin as fungistatic metabolic product of *Trichoderma viride*. Ann.Appl. Biol., 32:214-220
- Coley-Smith, J. R. (1976). Some interaction in soil between plants, sclerotium-forming fungi and other microorganisms. (In Biochemical aspects of Plant Parasite Relationships. Friend, J. and D. R. Threlfall (Eds.) . Academic Press London, New York, San Francisco. pp.11-23.
- Dos Santos, M.A.; S .Ferraz and J.J. Muchovej (1992). Evaluation of 20 species of fungi from Brazil for biocontrol of *Meloidogyne incognita* race-3. Nematologica 22: pp.183-192.
- Dowson, W. J. (1957). Plant diseases due to bacteria second Ed., Cambridge the university Press, London, pp23.
- Ghisalberti, E .L. (2002). Anti-infective agents produced by the hyphomycetes Genera *Trichoderma* and *Gliocladium*. Current Medicinal Chemistry- Anti infective Agents ,1(4) 343-374 .
- Goodey, J.B. (1957). Laboratory methods for work with plant and soil nematodes. Tech. Bull. No. 2. Min. Agric. Fish Ed. London, 47 pp.
- Haran, S.; H. Schickler and I.Chet (1996). Molecular mechanisms of lytic enzymes involved in the biocontrol activity of *Trichoderma harzianum*. Microbiol. 142: 2321-2331.
- Harman GE (2000). The myths and dogmas of biocontrol: changes in perceptions derived from research on *Trichoderma harzianum* strains T-22. Plt Disease 84(4):377-393.
- Huang, X. W.; N. H. Zhao and K. Q. Zhang. (2005). Extracellular enzymes serving as virulence factors in nematophagous fungi involved in infection of the host. Res. Microbiol. 115: 811–816.
- Huang, Y., C. Xu, L. Ma, K. Zhang, C. Duan and M. Mo (2009). Characterisation of volatiles produced from *Bacillus megatherium* YFM3.25 and their nematocidal activity against *Meloidogyne incognita*. Euro. J. Plant Pathol. 26: 417-422.
- Ji, P., H. Campbell; J. Kloepper; J. Jones; T. Suslow and M. Wilson (2006). Integrated biological control of bacterial speck and spot of tomato under field conditions using foliar biological control agents and plant growth promoting rhizobacteria. Biol. Cont. 36: 358–367.
- Jung, C. and U. Wyss (1999). New approaches to control plant parasitic nematodes. Appl. Microbiol. Biotechnology. 51: 439-446.
- Khan, M.R.; K. Kounsar and A. Hamid (2002). Effect of certain rhizobacteria and antagonistic fungi on root-knot nodulation and root-knot nematode disease of green gram. Nematologia Mediterranea, 30(1): 85-89.
- Kim, H.S.; D.Y. Byung; H.L. Chang; H.S. Hyun; M.O. Hee; K. Tohoru and T. Yoshiki(1997). Production and properties of a lipopeptide biosurfactant from *B. subtilis* C9. J Ferment Bioeng 84: 41–46.

- Kloepper, J. W. and C. M. Ryu (2006). Bacterial endophytes as elicitors of induced systemic resistance. In: Microbial root endophytes, Heidelberg, (Schulz, B., C. Boyle and T. Sieber eds.), Springer, Verlag, 33–51 pp.
- Lahlali, R., G. Peng; B. D. Gossen; L. McGregor; F. Q. Yu; R. K. Hynes; S. F. Hwang; M. R. McDonald and S. M. Boyetchko (2013). Evidence that the Biofungicide Serenade (*Bacillus subtilis*) Suppresses Clubroot on Canola via Antibiosis and Induced Host Resistance. Biol. Cont. 103: 245-254.
- Lamberti, F. (1981). Plant nematode problems in Mediterranean region. *helminthological Abstracts* 50:145-166.
- Li, L., Mo, M.H., Qu, Q., Luo, H., and Zhang, K.Q. (2007). Compounds inhibitory to nematophagous fungi produced by *Bacillus* sp. strain H6 isolated from fungistatic
- Mai, W.F. and H.H. Lyon (1975). Pictorial Key to genera of plant parasitic nematodes. Fourth edition, revised. Ithaca, Ison: Comstock Publishing Associates, Cornell Univ. Press, 219 PP.
- Mannanov, R. and R. Sattarova (2009). Inhibition of cotton pathogens by natural antagonists. *IOBC/WPRS Bulletin*, 43: 255-258.
- Norton, D.C. (1978). Ecology of plant parasitic nematode. John Wiley and sons, New York, USA. 238 pp.
- Oka, Y., H. Koltai, M. Bar-Eyal, M. Mor, E.C. Sharon and I. Y. Spiegel (2000). New strategies for the control of plant parasitic nematodes. *Pest Management Science.*, 56:983-988.
- Podile, A.R. and G.K. Kishore (2002). Biological control of peanut diseases. In *Biological control of crop disease* edited by Samuel S. Gnanamanickam. CRC.
- Rahman Khan, M.; Shahana, M. Khan and Faya, Mohide (2005). Root-knot nematode problem of some winter ornamental plants and its biomanagement. *Journal of Nematology*, 37: 198–206.
- Robinson, L., M. J. Jeger and P. Xiangming (2009). Management of strawberry gray mould using mixtures of biocontrol agents with different mechanisms of action. *Biocontrol Science and Technology*, 19: 1051-1065.
- Sasser, J.N. and D. W. Freckman (1987). A World Perspective on Nematology: The Role of the Society. In: *Vistas on Nematology*, Veech, J.A. and D.W. Dickson (Eds.). Society of Nematologists, Hyattsville, Maryland, United States, pp: 7-14.
- Sharma, P. and R. Pandey (2009). Biological control of root-knot nematode, *Meloidogyne incognita* in the medicinal plant, *Withania somnifera* and the effect of biocontrol agents on plant growth. *Afr. J. Agric. Res.*, 4: 564-567.
- Sharma, S.B. and D. McDonald (1990). Global status of nematode problems of groundnut, pigeonpea, chickpea, sorghum and pearl millet, and suggestions for future work. *Crop protection* ., 9:453-458
- Sharon, E.; M. Bar-Eyal ; I. Chet; A. Herra-Estrella; O. Kleifeld and Y. Spiegel (2001). Biological control of the root-knot nematode, *Meloidogyne javanica* by *Trichoderma harzianum*. *Phytopathology*, 91:687-693.

- Siddiqi, M.R.(1986).Tylenchida, parasites of plants and insects. Slough, UK. Common W.Inst. Parasite.645pp. soil. European Journal of Plant Pathology, Vol. 117, No. 4, pp. 329-340.
- Steel, R.and G.D. Torrie (1980). Principles and procedures of statistics.2nd ed. McGraw-Hill, New York.
- Taylor, A.L. and J.N. Sasser (1978).Biology, Identification and Control of Root-Knot Nematodes (*Meloidogyne* species). North Carolina State University, Raleigh, NC. USA. Pages: 111.
- Tomaszewski, E.K.; M.A.M. Khalil; A.A. El-Deeb; T.O. Powers and J.L. Starr (1994). *Meloidogyne javanica* Parasitic on Peanut. J. Nematol., 26: 436-441.
- Umesh Rud€rappa,copy right (2009-14) www.nutrition-and-you.com.Find us on Google+.
- Westcott, S.W. and D.A. Kluepfe (1993).Inhibition of Criconemellaxenoplax egg hatch by *Pseudomonas aureofaciens*. Phytopathology. 83: 1245-1249.
- Zhang, S. and X. Zhang (2009). Effects of two composted plant pesticide residues, incorporated with *Trichoderma viride* on root-knot nematode in balloon flower. J. Agric. Sci. China, 8: 447-454.

تأثير بعض العوامل الحيوية ومركب نيماتوب في مكافحة مرض تعقد الجذور علي الفول السوداني

هويدا عبد الوهاب متولي^١ و هناء سيدهم زوام^٢

١-المعمل المركزي للزراعة العضوية- مركز البحوث الزراعية الجيزة - مصر

٢- قسم بحوث النيماتودا معهد بحوث أمراض النباتات-مركز البحوث الزراعية الجيزة-مصر

المكافحة الحيوية لمرض نيماتودا تعقد الجذور تم تداولها في أبحاث علمية عديدة. استخدمت باسيلس ستلس وتريكودرما هارزيانم وتريكودرما فيردي وخليط منهما ونيماتوب بتركيزات مختلفة لمكافحة نيماتودا تعقد الجذور بميلودوجيني جافينكا (والتي كانت أكثر استخلاصا من التربة والجذور المصابة) في المعمل وفي الحقل علي الفول السوداني. وكان أفضلهم تأثيرا هو النيماتوب وأقلهم باسيلس ستلس في مكافحة تحت ظروف المعمل وقد أعطت كل المعاملات تأثيرا جيدا عند اعلي تركيز (١٠ /) علي نسبة موت الطور اليرقي الثاني ونسبة فقس البيض. وجدت علاقة ايجابية بين زيادة التركيز وتأثير المعاملة عند التقييم. كما توافقت النتائج المتحصل عليها في الحقل مع النتائج المتحصل عليها من المعمل حيث أعطت المعاملات الحيوية أفضل تأثيرا بالمقارنة بالكنترول. سجلت علاقة طردية بين خفض نسبة المرض و عدد مرات الاضافة حيث وجد ان المعاملة بعد ثلاث مرات من الاضافة كان لها أعلى تأثير في تقليل اعداد النيماتودا في الجذور والتربة و زيادة وزن محصول الفول السوداني. وأعطت المعاملة بخليط من عزلات التريكودرما أفضل تأثير سواء علي وزن المحصول أو خفض اعداد النيماتودا في التربة و الجذور يليها النيماتوب تحت ظروف الحقل بالمقارنة بالكنترول.