

# BIOLOGICAL CHARACTERISTICS AND HEAT REQUIREMENTS FOR *Coccinella undecimpunctata* – *Sitobion avenae* AND *Coccinella 9-punctata* – *Aphis craccivora* FEEDING SYSTEMS AT VARYING TEMPERATURE REGIMES



Bayoumy, M. H.; A. M. Abou-Elnaga; A. A. Ghanim and Gh. A. Mashhoot

Economic Entomology Department, Faculty of Agriculture, Mansoura University,

B. O. Box 35516, Mansoura, Egypt

\*Corresponding Author; mhmohamed@mans.edu.eg

## ABSTRACT

Laboratory experiments were carried out to examine some biological characteristics and heat requirements of the two coccinellid species; viz. *Coccinella undecimpunctata* L. and *Coccinella 9-punctata* L. (Coccinellidae: Coleoptera) when reared on *Sitobionavenae*(F.) and *Aphis craccivora* Koch (Aphididae: Hemiptera) at varying constant temperature regimes of 18, 23, 28, and 33 °C, respectively. The obtained results showed that the developmental time of *C. undecimpunctata* and *C. 9-punctata* immature stages declined as temperature degrees increased with the shortest developmental time was recorded at 33 °C which lasted  $15.29 \pm 0.98$  and  $14.00 \pm 1.35$  days, respectively. Daily larval consumption increased as both temperature and larval age increased with the highest consumption for the fourth instars at 33 °C which recorded by  $99.71 \pm 14.49$  and  $105.40 \pm 10.54$  for *C. undecimpunctata* and *C. 9-punctata*, respectively. Female and male longevities for both predator species increased as temperature decreased, whereas the females' fecundity increased as temperature increased. Furthermore, the total consumption of female and male of both species increased as temperature increased with the highest consumption recorded during the oviposition period. The total consumption of females was higher than that of males for both predator species, and the corresponding daily consumption values were higher for females than those of males at varying temperatures. The developmental rates for all stages of both species declined as temperature decreased. The lower developmental threshold ( $T_0$ ) recorded the lowest for pupal and larval stages of *C. undecimpunctata* and *C. 9-punctata*, respectively. The corresponding amount of heat units which required to completing the development of *C. undecimpunctata* and *C. 9-punctata* averaged  $380.73 \pm 27.71$  and  $363.83 \pm 14.44$  dd's, respectively. This study showed that *C. undecimpunctata* and *C. 9-punctata* successfully complete their development in a wide range of temperatures from 18 to 33 °C, indicating their high potential for use in biological control programs against *S. avenae* and *A. craccivora*, respectively.

**Keywords:** Developmental rate, degree-days, heat units, lower developmental threshold.

## INTRODUCTION

Many coccinellids, are well known as insect predators, playing an important role as biological control agents in regulating the population of several insect pests especially aphids, coccids, and other soft bodies' insects.

Aphidophagous ladybirds are considered to be a great economic in agro-ecosystem through their successful employment in the biocontrol of aphids, and have been received increasing attention from ecologists all over the world due to some of their characteristics, such as: ability to feed on a wide range of prey, to be very voracious, and to have a rapid numeric response (e.g., Agarwala et al., 1988; Hodek and Honěk, 1996; Bari and Sardar, 1998; Pervez and Omkar, 2005). Prior to introduce any predator in biological control program, it should be estimated its efficiency under different environmental factors, among them the predation activity of varying developmental stages, female reproductive success, body size, prey species and density, plant architecture, prey type, temperature, and relative humidity are considered the most important factors (Skirvinet et al., 1997; Kajita and Evans, 2010; Koch et al., 2003; Sarmiento et al., 2007; Bayoumy and Michaud, 2012; Bayoumy et al., 2014; Bayoumy et al., 2015).

*Coccinella undecimpunctata*L. and *Coccinella 9-punctata*L. (Coleoptera: Coccinellidae) are dominant coccinellid species in Mansoura region (Ghanim and El-Adl, 1987). Both species are euriphagous predators, which prefers to feed on aphids (Hodek and Honěk, 1996), and *C. undecimpunctata* offers interesting potential as bio-control agents in the context of integrated pest management, IPM (Cabral et al., 2011). Several investigators in different places of the world studied the predation activity of certain coccinellid predators (Sethi and Atwal, 1964; Smith, 1965; Ghanim and El-Adl, 1987; El-Serafi et al., 2004; Ghanim et al., 2009; and Bayoumy, 2011; Osman and Bayoumy, 2011; Bayoumy and Michaud, 2012; Bayoumy et al., 2014). Temperature controls the development rate of many organisms, plants, and invertebrate animals, including insects and nematodes which require certain amounts of heat to develop from one stage in their cycles to another one. This measure of accumulated heat is known as degree-days (dd's). Growth and development of insect are dependent on temperature where the temperature increase, development time decreases until the temperature become high enough to have a negative effect. This limit is defined as temperature threshold. The lower development threshold ( $T_0$ ) for a species is the minimum temperature at which development can continue and below it the development fails. Numerous entomologists in different parts of the world estimated the heat requirement of certain insect pests including the coccinellid predators either in the lab (Zalomet et al., 1983; Uygun and Atlhan 2000; Omkar and Pervez, 2004; Ghanim et al., 2014; Abd El-Halim, 2015) or in the field (Awadalla et al., 2014) to estimate the amount of heat for completing their development, to predicate with their abundance, and/or to protect them from the extensive use of chemical pesticides. Although the biology and consumption of these species have been investigated, no study aimed to theoretically estimated the thermal requirements for these predator species specifically *C. 9-punctata* (e.g., Ghanim and El-Adl, 1987; Abdel-Salam, 2004; Mari et al., 2005; Cabral et al., 2006; Solangiet al., 2007; Mohamed and Ghanim, 2008; El-Heneidy et al., 2008). Therefore, the present investigation aimed to study some biological attributes of *C. undecimpunctata*- *S. avenae* and *C. 9-punctata*-*A. craccivora* feeding

systems to use it in the estimation of the lower thermal development and then the amount of heat required to development of these coccinellid predators.

## MATERIALS AND METHODS

### **Biological characteristics of *C. undecimpunctata* and *C. 9-punctata***

A permanent culture of coccinellid predators and aphid species were maintained in the laboratory of the Economic Entomology Department, Faculty of Agriculture, Mansoura University. Experiments were carried out under four constant temperature regimes,  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$ , and  $70 \pm 5$  % R.H. The used predator species, *C. undecimpunctata* and *C. 9-punctata* were obtained from the laboratory culture. Twenty newly hatched first instars of each predator species were isolated into Petri-dishes (9.0 cm in diameter). The bottom of each dish was covered with a filter paper to facilitate the predator's larvae movement. A known number of different stages of each aphid species (*S. avenae* and *A. craccivora*) was offered daily at 10 AM and the devoured individuals were recorded. A small plant leaflet was introduced daily in each Petri-dish as a food source for the aphids for keeping them alive as long as possible. The remained aphids and their parts were removed daily from each Petri-dish before a new food. The total number of aphid species consumed by each predator larva was estimated. The developmental times for the larval and pupal stages were also estimated. Each adult predator received the same type of prey as that of its larval instars. Immediately after emergence from the pupal stage, each predator individuals were sexed and the isolated into the Petri-dishes. Known numbers of each aphid species were provided daily on a plant leaflet to each predator. Counting the consumed aphids and removing the non-devoured aphids in each Petri-dish were practiced daily before providing the new aphids. After five days of emergence, copulation took place and the two sexes were immediately separated and isolated into the Petri-dishes. The daily number of the laid eggs per each predator female during her oviposition period was counted. The total number of each aphid species consumed daily by a male or female and the total daily number of deposited eggs for each predator female species were recorded. The daily means of food consumption during longevity of each predator species were calculated.

### **B. Heat requirements (degree-days) of *C. undecimpunctata* and *C. 9-punctata***

The coccinellid predators, *C. undecimpunctata* and *C. 9-punctata* were reared under four constant temperature degrees ( $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C) to estimate the thermal requirements as a predicating tool for the annual generation of these predators. The lower developmental threshold temperature ( $t_0$ ) was estimated from the x- intercept point of the linear regression analysis (with  $t_0 = -a/b$ ) to determine the relationship between development rate (calculated as the inverse of the number of days required for development at that temperature,  $1/d$ ) and incubation temperatures for values within the linear range, resulting in the regression equation  $y = a + bx$ ; where  $x$  is the temperature and  $y$  is the development rate (Arnold, 1960;

Campbell et al., 1974). Regression lines were determined by using the SigmaPlot 11.0 (Systat, 2008).

The thermal constant (K) required for development of each stage on each temperature (the amount of thermal energy required for completion of development of 50% of individuals at constant temperatures, expressed as degree-days (dd's) accumulated above the lower developmental threshold) were estimated according to Fletcher (1981) and Obrycki and Tauber (1981) equation as follows:

$$K=D(T-T_0)$$

Where: D is the development duration in days, T is the temperature (°C) at which development occurs and  $T_0$  the lower developmental temperature threshold.

#### Statistical analysis

Data for larval, pupal, and total development durations, total larval consumption, female and male longevities, female fecundity, total female and male consumptions, and thermal units (dd's) were analyzed by One-way ANOVA and means separated using Duncan's Multiple Range Test ( $\alpha = 0.05$ )

## RESULTS AND DISCUSSION

### 1. Biological characteristics of *C. undecimpunctata* and *C. 9-punctata* *Coccinella undecimpunctata* L.

The obtained results in Table (1) showed that the incubation period decreased as temperature increased with the lowest period ( $3.1 \pm 0.52$ ) at 33 °C. Statistical analysis revealed that there was significant effect for temperature on larval and pupal durations and total development. The larval duration of the predator averaged  $20.74 \pm 1.10$ ,  $16.74 \pm 0.95$ ,  $10.61 \pm 0.87$ , and  $8.99 \pm 0.74$  days when fed on *S. avenae* under constant temperature degrees of  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C, respectively, whereas the pupal stage period lasted an average of  $7.53 \pm 0.8$ ,  $7.00 \pm 0.56$ ,  $5.30 \pm 0.40$ , and  $3.70 \pm 0.32$  days at the same temperatures, respectively. The total development time of *C. undecimpunctata* averaged  $36.17 \pm 1.85$ ,  $29.04 \pm 1.5$ ,  $15.29 \pm 0.89$ , and  $19.01 \pm 1.16$  at  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C, respectively.

Statistical analysis revealed that there was significant effect for temperature on larval stage consumption. The consumption of larval instars of *C. undecimpunctata* when fed on *S. avenae* increased as temperature and their growth increased with the highest consumption for the fourth instars at 33 °C. The daily consumption per an individual larva averaged  $25.57 \pm 5.15$ ,  $41.82 \pm 12.41$ ,  $76.12 \pm 11.08$ , and  $99.74 \pm 14.49$  at  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C, respectively (Table 2).

Statistical analysis showed that there was significant effect for temperature on male and female longevities. The female and male longevities of *C. undecimpunctata* decreased as temperature increased with the shortest longevity averaged  $50.95 \pm 2.10$  and  $40.22 \pm 0.89$  for female and male at 33 °C, respectively, whereas the daily number of deposited eggs per female

increased as temperature increased with the highest number ( $30.11 \pm 8.66$  eggs/female/day) at  $33\text{ }^{\circ}\text{C}$  (Table 3).

Statistical analysis showed that there were significant effects for temperature on total consumption for males and females during their longevity. The total consumption of female *C. undecimpunctata* was the highest during oviposition period followed by postoviposition and preoviposition periods. As presented in Table (4), *C. undecimpunctata* males consumed more aphids (*S. avenae*) than females during their longevity only at  $33\text{ }^{\circ}\text{C}$ . Furthermore, the total consumption for female and male during their longevities and the daily consumption per female and male increased with increasing temperature with the highest total female consumption of  $6943.57 \pm 39.64$  aphids, the highest total male consumption of  $7397 \pm 32.46$  aphids, the highest daily female consumption of  $136.28 \pm 18.9$  aphids, and the highest daily male consumption of  $183.83 \pm 38.46$  aphids at  $33\text{ }^{\circ}\text{C}$ .

The developmental rates for completion of embryogenesis, larval, pupal, and total development of *C. undecimpunctata* when fed on *S. avenae* increased as the temperature increased. The lower developmental threshold ( $T_0$ ) derived from the linear relationship between developmental rate (1/d) and the four tested temperatures (Fig. 1) was the lowest for pupal stage ( $5.42\text{ }^{\circ}\text{C}$ ), whereas the egg stage was the more tolerant stage for lower temperature ( $11.38\text{ }^{\circ}\text{C}$ ). The thermal units, expressed as degree-days (DD's), required for each developmental stage showed that the larval stage needs to more heat unit to develop at each temperature tested compared to other stages. The corresponding thermal unites required for each stage of *C. undecimpunctata* to complete its development on a given temperature was  $55.40 \pm 3.99$ ,  $223.29 \pm 16.95$ ,  $109.85 \pm 11.88$ , and  $380.73 \pm 27.71$  for egg, larval, and pupal stages, and total life cycle, respectively. Furthermore, statistical analysis revealed that there was significant effect for temperature on thermal units required for complete the development (Table 5).

**Table (1): Developmental time in days ( $\pm$  SE) of *C. undecimpunctata* immature stages when reared on *S.avenae* at four constant temperature regimes.**

Immatures T (°C)	Egg	Larval instars					Pupa	Egg- Adulthood
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	Total		
18	7.9 $\pm$ 0.81	4.71 $\pm$ 0.73	3.85 $\pm$ 0.61	5.84 $\pm$ 0.81	6.34 $\pm$ 0.40	20.74 $\pm$ 1.10 a	7.53 $\pm$ 0.80 a	36.17 $\pm$ 1.85 a
23	5.3 $\pm$ 0.75	3.17 $\pm$ 0.65	3.10 $\pm$ 0.57	4.72 $\pm$ 0.72	5.75 $\pm$ 0.68	16.74 $\pm$ 0.96 b	7.00 $\pm$ 0.56 a	29.04 $\pm$ 1.50 b
28	3.1 $\pm$ 0.52	1.90 $\pm$ 0.40	1.68 $\pm$ 0.31	2.93 $\pm$ 0.57	4.10 $\pm$ 0.51	10.61 $\pm$ 0.87 c	5.30 $\pm$ 0.40 b	19.01 $\pm$ 1.16 c
33	2.6 $\pm$ 0.26	1.53 $\pm$ 0.35	1.40 $\pm$ 0.26	2.56 $\pm$ 0.37	3.50 $\pm$ 0.29	8.99 $\pm$ 0.74 d	3.70 $\pm$ 0.32 c	15.29 $\pm$ 0.98 d

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (2): Consumption ( $\pm$  SE) of *C. undecimpunctata* larval stages and daily consumption per individual larva when reared on *S.avenae* at four constant temperature regimes.**

Consumption T (°C)	Larval instars					Daily consumption <sup>-1</sup>
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	Total larval stage	
18	45.70 $\pm$ 1.46	62.36 $\pm$ 1.61	186.45 $\pm$ 2.46	235.72 $\pm$ 2.86	530.23 $\pm$ 05.67 c	25.57 $\pm$ 05.15
23	56.11 $\pm$ 1.75	73.57 $\pm$ 1.90	219.72 $\pm$ 2.75	350.68 $\pm$ 2.96	700.08 $\pm$ 06.95 b	41.82 $\pm$ 12.41
28	66.76 $\pm$ 1.80	78.96 $\pm$ 2.10	265.42 $\pm$ 2.80	396.43 $\pm$ 3.50	807.59 $\pm$ 09.64ab	76.12 $\pm$ 11.08
33	70.64 $\pm$ 1.96	82.45 $\pm$ 2.50	296.54 $\pm$ 3.11	446.75 $\pm$ 4.20	896.38 $\pm$ 10.72 a	99.71 $\pm$ 14.49

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (3): Longevity in days ( $\pm$  SE) of *C. undecimpunctata* adults when reared on *S.avenae* at four constant temperature regimes.**

Adults T (°C)	Oviposition periods			Female longevity	Male longevity	Fecundity/ Female	Eggs/Female/day
	Pre-oviposition	Oviposition	Inter + Post - oviposition				
18	6.75 $\pm$ 0.92	48.95 $\pm$ 2.40	32.95 $\pm$ 1.42	88.65 $\pm$ 3.70 a	61.78 $\pm$ 1.18 a	629 $\pm$ 10.72 d	12.85 $\pm$ 4.47
23	4.97 $\pm$ 0.73	40.45 $\pm$ 1.96	28.50 $\pm$ 1.50	73.92 $\pm$ 3.10ab	55.14 $\pm$ 1.10ab	770 $\pm$ 10.96 c	19.03 $\pm$ 5.59
28	4.57 $\pm$ 0.58	35.70 $\pm$ 1.50	25.70 $\pm$ 1.37	65.97 $\pm$ 2.96 c	50.91 $\pm$ 0.90 b	950 $\pm$ 12.18 a	26.61 $\pm$ 8.12
33	4.10 $\pm$ 0.52	28.10 $\pm$ 1.36	18.75 $\pm$ 1.10	50.95 $\pm$ 2.10 d	40.22 $\pm$ 0.84 b	846 $\pm$ 11.78 b	30.11 $\pm$ 8.66

**Table (4): Daily and total consumption ( $\pm$  SE) of *C. undecimpunctata* adults when reared on *S.avenae* at four constant temperature regimes.**

Adults T (°C)	Oviposition periods			Female consumption		Male consumption	
	Pre-oviposition	Oviposition	Inter + Post-oviposition	Total	Daily	Total	Daily
18	641.25 $\pm$ 5.86	4876.19 $\pm$ 26.42	946.80 $\pm$ 6.42	6464.24 $\pm$ 30.70b	72.92 $\pm$ 8.3	3970.22 $\pm$ 26.40ab	64.26 $\pm$ 22.37
23	586.17 $\pm$ 4.76	5290.88 $\pm$ 31.17	723.11 $\pm$ 5.31	6600.16 $\pm$ 36.51b	89.29 $\pm$ 11.8	3996.44 $\pm$ 28.51ab	72.48 $\pm$ 25.92
28	610.18 $\pm$ 5.80	5563.44 $\pm$ 35.24	615.14 $\pm$ 4.75	6788.76 $\pm$ 38.17ab	102.91 $\pm$ 12.9	4370.65 $\pm$ 31.71a	85.85 $\pm$ 35.23
33	636.15 $\pm$ 6.32	5780.16 $\pm$ 37.56	527.26 $\pm$ 3.96	6943.57 $\pm$ 39.64a	136.28 $\pm$ 18.9	3397.69 $\pm$ 32.46b	84.47 $\pm$ 38.64

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (5): Developmental rates (1/d), thermal requirements (DD's), and lower developmental threshold (T<sub>0</sub>) for various developmental stages of *C.undecimpunctata* reared on *S. avenae* at four constant temperature regimes.**

Temp. (°C)	Developmental rate				Temperature (°C)	DD's			
	Egg	Larva	Pupa	Egg- Adulthood		Egg	Larva	Pupa	Egg-Adulthood
18	0.127	0.048	0.133	0.0277	18	52.29	206.57	94.60	348.32 c
23	0.189	0.059	0.143	0.0344	23	61.59	250.43	123.06	424.86 a
28	0.323	0.094	0.189	0.0526	28	51.52	211.78	119.67	373.16 b
33	0.385	0.111	0.270	0.0654	33	56.21	224.39	102.05	376.59 b
T <sub>0</sub>	11.38	8.04	5.42	8.37	Mean ( $\pm$ SE)	55.40 $\pm$ 3.99	223.29 $\pm$ 16.95	109.85 $\pm$ 11.88	380.73 $\pm$ 27.71

Values labeled by the same letters are not significantly differed at the 5% probability level

***Coccinella 9-punctata* L.**

The obtained results in Table (6) showed that the incubation period decreased as temperature increased with the lowest period ( $3.0 \pm 0.37$ ) at 33 °C. Statistical analysis showed that there was significant effect for temperature on larval and pupal durations and total development. The larval duration of the predator averaged  $16.64 \pm 1.35$ ,  $11.89 \pm 1.1$ ,  $8.56 \pm 0.92$ , and  $7.00 \pm 0.63$  days when fed on *Aphis craccivora* under constant temperature degrees of  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C, respectively, whereas the pupal stage period lasted an average of  $6.10 \pm 0.44$ ,  $4.21 \pm 0.30$ ,  $4.21 \pm 0.30$ , and  $4.00 \pm 0.26$  days at the same temperatures, respectively. The total development time of *C. undecimpunctata* averaged  $27.54 \pm 1.89$ ,  $20.14 \pm 1.63$ ,  $29.04 \pm 1.5$ , and  $36.17 \pm 1.85$  at  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C, respectively.

Statistical analysis showed that there was significant effect for temperature on larval stage consumption. The consumption of larval instars of *C. 9-punctata* when fed on *A. craccivora* increased as temperature and their growth increased with the highest consumption for the fourth instars at 33 °C. The daily consumption per an individual larva averaged  $25.05 \pm 3.84$ ,  $48.62 \pm 4.32$ ,  $76.73 \pm 5.89$ , and  $105.40 \pm 10.54$  at  $18 \pm 1$ ,  $23 \pm 1$ ,  $28 \pm 1$ , and  $33 \pm 1$  °C, respectively (Table 7).

Statistical analysis revealed that there was significant effect for temperature on male and female longevities. The female and male longevities of *C. 9-punctata* decreased as temperature increased with the shortest longevity averaged  $45.74 \pm 1.50$  and  $30.17 \pm 0.68$  for female and male at 33 °C, respectively, whereas the daily number of deposited eggs per female increased as temperature increased with the highest number ( $29.03 \pm 8.71$  eggs/female/day) at 33 °C. The female fecundity increased with increasing temperature to record the highest fecundity of  $890.56 \pm 11.32$  at 33 °C (Table 8).

Statistical analysis revealed that there were significant effects for temperature on total consumption for males and females during their longevity. The total consumption of female *C. 9-punctata* was the highest during oviposition period followed by postoviposition and preoviposition periods. As presented in Table (9), *C. 9-punctata* females consumed more aphids (*A. craccivora*) than males during their longevity. Furthermore, the total consumption for female and male during their longevities and the daily consumption per female and male increased with increasing temperature with the highest total female consumption ( $5189.36 \pm 35.20$  aphids), the highest male total consumption ( $2750.657 \pm 26.70$  aphids), the highest daily female consumption ( $113.45 \pm 23.47$  aphids), and the highest daily male consumption ( $91.17 \pm 9.52$  aphids) at 33 °C.

The developmental rates for completion of embryogenesis, larval, pupal, and total development of *C. 9-punctata* when fed on *A. craccivora* increased as the temperature increased. The lower developmental threshold ( $T_0$ ) derived from the linear relationship between developmental rate (1/d) and the four tested temperatures (Fig. 2) was the lowest for larval stage ( $7.64$  °C), whereas the tolerant stage was the egg stage ( $10.72$  °C). However, the thermal units, expressed as degree-days (DD's), required for each



developmental stage showed that the larval stage needs to more heat unit to develop at each temperature tested compared to other stages. The corresponding thermal unites required for each stage of *C. 9-punctatato* complete its development on a given temperature was  $50.03 \pm 12.58$ ,  $176.71 \pm 3.88$ ,  $66.53 \pm 15.8$ , and  $363.83 \pm 14.44$  for egg, larval, and pupal stages, and total life cycle, respectively (Table 10).

The obtained results for both predator species regarding their development, larval and adult stage consumptions, fecundity, and female and male longevities are partially confirmed by works of Ghanim and El-Adl (1987), Abdel-Salam (2004), Mari *et al.* (2005), Cabral *et al.* (2006), Solangiet *al.* (2007), Mohamed and Ghanim (2008), El-Heneidy *et al.* (2008). However, results of lower developmental threshold and degree-days are disagreed with that of Jalaliet *al.* (2014). The lower developmental thresholds for total development (egg to adult) of *Coccinellaundecimpunctataaegyptica*(Reiche) recorded  $14^{\circ}\text{C}$  and the degree-day (dd's) requirements for total development were 166.67dd's. This may be attributed to the different prey species used as food sources in both studies. Nevertheless, these results were relatively closed to those of Skouras *et al.* (2015) for *Coccinella undecimnotata* Schneider to the tobacco aphid, *M. persicaenicotianae* at five constant temperatures (17, 20, 23, 26, and  $29^{\circ}\text{C}$ ).The obtained results for total fecundity, daily oviposition per female, and lower developmental threshold for *C. undecimpunctata* are closed to those obtained by Xia *et al.* (1999) for *Coccinellaseptempunctata* L. reared on *Aphis gossypii* Clover at 15, 20, 25, 30, and  $35 \pm 0.5^{\circ}\text{C}$ .

**Table (6): Developmental time in days ( $\pm$  SE) of *C. 9-punctata* immature stages when reared on *A. craccivora* at four constant temperature regimes.**

Immature stages T(°C)	Egg	Larval instars				Total	Pupa	Egg- Adulthood
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>			
18	4.8 $\pm$ 0.65	4.37 $\pm$ 0.71	2.90 $\pm$ 0.40	3.45 $\pm$ 0.62	5.92 $\pm$ 0.82	16.64 $\pm$ 1.35a	6.10 $\pm$ 0.44a	27.54 $\pm$ 1.89a
23	3.53 $\pm$ 0.47	3.18 $\pm$ 0.52	1.96 $\pm$ 0.32	2.18 $\pm$ 0.50	4.57 $\pm$ 0.73	11.89 $\pm$ 1.10b	4.72 $\pm$ 0.37b	20.14 $\pm$ 1.63b
28	3.18 $\pm$ 0.41	2.46 $\pm$ 0.37	1.27 $\pm$ 0.30	1.64 $\pm$ 0.31	3.19 $\pm$ 0.60	8.56 $\pm$ 0.92c	4.21 $\pm$ 0.30b	15.95 $\pm$ 1.50c
33	3.00 $\pm$ 0.37	2.11 $\pm$ 0.33	1.01 $\pm$ 0.27	1.25 $\pm$ 0.24	2.63 $\pm$ 0.35	7.00 $\pm$ 0.63d	4.00 $\pm$ 0.26b	14.00 $\pm$ 1.35c

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (7): Consumption rates ( $\pm$  SE) of *C. 9-punctata* larval stages and daily consumption per individual larva when reared on *A. craccivora* at four constant temperature regimes.**

Consumption T (°C)	Larval instars				Total consumption	Daily consumption <sup>-1</sup>
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>		
18	31.92 $\pm$ 0.97	45.57 $\pm$ 1.42	153.16 $\pm$ 2.10	186.15 $\pm$ 2.40	416.80 $\pm$ 4.10 d	25.05 $\pm$ 3.04
23	40.32 $\pm$ 1.10	59.16 $\pm$ 1.53	188.46 $\pm$ 2.34	290.17 $\pm$ 2.73	578.11 $\pm$ 4.75 c	48.62 $\pm$ 4.32
28	52.71 $\pm$ 1.33	62.14 $\pm$ 1.74	240.13 $\pm$ 2.64	301.80 $\pm$ 2.96	656.78 $\pm$ 5.42 b	76.73 $\pm$ 5.89
33	60.18 $\pm$ 1.62	69.10 $\pm$ 1.94	248.12 $\pm$ 3.15	357.42 $\pm$ 3.26	737.82 $\pm$ 6.64 a	105.40 $\pm$ 10.54

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (8): Longevity in days and total and daily fecundity ( $\pm$  SE) of *Coccinella 9-punctata* adults when reared on *Aphis craccivora* at four constant temperature regimes.**

Parameters T(°C)	Oviposition periods			Female longevity	Male longevity	Fecundity/ Female	Eggs/Female/ day
	Pre- oviposition	Oviposition	Inter + Post - oviposition				
18	6.32 $\pm$ 0.48	40.58 $\pm$ 2.15	16.84 $\pm$ 0.96	63.74 $\pm$ 3.22 a	55.17 $\pm$ 1.6a	650.43 $\pm$ 9.40 c	16.03 $\pm$ 4.37
23	5.15 $\pm$ 0.45	37.46 $\pm$ 1.94	13.40 $\pm$ 0.80	56.01 $\pm$ 2.61ab	40.44 $\pm$ 1.11 b	801.50 $\pm$ 10.36 b	21.39 $\pm$ 5.34
28	4.75 $\pm$ 0.41	34.81 $\pm$ 1.52	11.39 $\pm$ 0.74	50.95 $\pm$ 1.94 b	35.18 $\pm$ 0.9c	920.40 $\pm$ 12.15 a	26.44 $\pm$ 7.99
33	4.56 $\pm$ 0.35	30.68 $\pm$ 1.30	10.50 $\pm$ 0.62	45.74 $\pm$ 1.50 c	30.17 $\pm$ 0.86 d	890.56 $\pm$ 11.32 a	29.03 $\pm$ 8.71

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (9): Daily and total consumption ( $\pm$  SE) of *Coccinella 9-punctata* adults when reared on *Aphis craccivora* at four constant temperature regimes.**

Adults T(°C)	Oviposition periods			Female consumption		Male consumption	
	Pre-oviposition	Oviposition	Inter + Post- oviposition	Total	Daily	Total	Daily
18	420.26 $\pm$ 3.70	2653.10 $\pm$ 26.17	817.10 $\pm$ 4.50	3890.46 $\pm$ 30.19 c	61.04 $\pm$ 9.38	2412.15 $\pm$ 19.16b	43.72 $\pm$ 6.42
23	486.19 $\pm$ 4.60	2994.17 $\pm$ 30.15	915.42 $\pm$ 6.42	4395.78 $\pm$ 33.16 b	78.48 $\pm$ 12.70	2105.19 $\pm$ 22.60c	52.03 $\pm$ 7.50
28	557.83 $\pm$ 4.96	3453.18 $\pm$ 31.80	930.18 $\pm$ 6.54	4941.29 $\pm$ 34.19ab	96.99 $\pm$ 17.62	2240.62 $\pm$ 25.78b	63.69 $\pm$ 8.15
33	580.26 $\pm$ 5.50	3670.90 $\pm$ 33.15	938.20 $\pm$ 6.80	5189.36 $\pm$ 35.20 a	113.45 $\pm$ 23.47	2750.65 $\pm$ 26.70a	91.17 $\pm$ 9.52

Values labeled by the same letters are not significantly differed at the 5% probability level

**Table (10): Developmental rates (1/d), thermal requirements (DD's), and lower developmental threshold (T<sub>0</sub>) for various developmental stages of *Coccinella 9-punctata* reared on *Aphis craccivora* at four constant temperature regimes.**

Temp. (°C)	Developmental rate				Temp. (°C)	DD's			
	Egg	Larva	Pupa	Egg- Adulthood		Egg	Larva	Pupa	Egg-Adulthood
18	0.21	0.06	0.16	0.03	18	34.94	172.39	46.79	349.48 b
23	0.28	0.08	0.21	0.05	23	43.35	182.63	57.77	356.28 b
28	0.31	0.12	0.24	0.06	28	54.95	174.28	72.58	361.91ab
33	0.33	0.14	0.25	0.07	33	66.86	177.55	88.96	387.66 a
T <sub>0</sub>	10.72	7.64	10.33	5.31	Mean ( $\pm$ SE)	50.03 $\pm$ 12.58	176.71 $\pm$ 3.88	66.53 $\pm$ 15.8	363.83 $\pm$ 14.44

Values labeled by the same letters are not significantly differed at the 5% probability level




Fig. 1: Relationship between developmental rate (1/d) and various temperatures for various developmental stages of *C.undecimpunctata* reared on *S.avenae* to mathematically extract the lower developmental thresholds ( $T_0 = -a/b$ ).




Fig. 2: Relationship between developmental rate (1/d) and various temperature for various developmental stages of *C. 9-punctata* reared on *A.s craccivora* to mathematically extract the lower developmental thresholds ( $T_0 = -a/b$ ).

## REFERENCES

- Abdel-Salam, A. H. (2004): Comparative biological and life table parameters of *Coccinella undecimpunctata* L. and *Cydoniavicinasis* Cr. (Coleoptera: Coccinellidae) reared on cereal aphid *Sitobion avenae* (Fabr.) (Hemiptera: Aphididae). J. Agric. Sci., Mansoura University 29(12): 7465-7480.
- Agarwala, B. K.; Das, S. and M. Senchowdhuri (1988): Biology and food relation of *Micraspis discolor* (F.) an aphidophagous coccinellid in India. J. Aphidol. 2(1-2): 7-17.
- Arnold, C. Y. (1960): Maximum-minimum temperatures as a basic for computing heat units. Proc. Amer. Soc. Hort. Sci. 76, 682-692.
- Awadalla, S. S.; Bayoumy, M. H.; Khattab, M. A. and A. H. Abdel-Wahab (2014): Thermal Requirements for Development of *Bemisia tabaci* (Hemiptera: Aleyrodidae) Biotype 'B' and Their Implication to Field Sample Population Data. Acta Phytopath. Entomol. Hung. 49 (1): 287-302.
- Bari, M. N. and M. A. Sardar (1998): Control strategy of bean aphid with predator, *Menochilus sexmaculatus* (F.) and insecticides. Bang. J. Entomol. 8 (1&2): 21-29.
- Bayoumy, M. H. (2011): Foraging behavior of the coccinellid *Nephus includens* (Coleoptera: Coccinellidae) in response to *Aphis gossypii* (Hemiptera: Aphididae) with particular emphasis on larval parasitism. Environ. Entomol. 40, 835-843.
- Bayoumy, M. H. and J. P. Michaud (2012): Parasitism interacts with mutual interference to limit foraging efficiency in larvae of *Nephus includens* (Coleoptera: Coccinellidae). Biol. Control 62, 120-126.
- Bayoumy, M. H. and J. P. Michaud (2014): Female fertility in *Hippodamia convergens* (Coleoptera: Coccinellidae) is maximized by polyandry, but reduced by continued male presence. Euro. J. Entomol. 11(4): 513-520.
- Bayoumy, M. H.; Osman, M. A. and J. P. Michaud (2014): Host plant mediates foraging behavior and mutual interference among adult *Stethorus gilvifrons* (Coleoptera: Coccinellidae) preying on *Tetranychus urticae* (Acari: Tetranychidae). Environ. Entomol. 43(5): 1309-1318.
- Campbell, J. P.; Bownas, A. D.; Peterson, N. G. and Dunnette, M. D. (1974): The Measurement of Organizational Effectiveness: A Review of Relevant Research and Opinion. Final Technical Report. NPRDC TR 75-1, Navy Personnel Research and Development Center, San Diego, CA.
- Duncan, D. B. (1955): Multiple range and multiple F. tests. Biometric (11): 1-42.
- El-Heneidy, A. H.; Hafez, A. A.; Shalaby, F. F. and I. A. B. El-Din (2008): Comparative biological aspects of the two coccinellid species; *Coccinella undecimpunctata* L. and *Hippodamia convergens* Guer. under laboratory conditions. Egypt. J. Biol. Pest Control 18(1): 51-59.

- Fletcher, G. L. (1981): Effects of temperature and photoperiod on the plasma freezing point depression, Cf concentration, and protein 'anti-freeze' in winter flounder. *Can J. Zool.* 59, 193-201.
- Ghanim, A. A. and M. A. El-Adl (1987). The feeding capacity and duration of the larval instars of three ladybird beetles fed on different aphid species under natural weather conditions at Mansoura, Egypt. *J. Agric. Sci., Mansoura Univ.* 12 (4): 981-97.
- Ghanim, A. A.; Abd Allah, F. A. and A. A. Abd El-Aziz (2014): Influence of nutrition's on longevity and fecundity, the coccinellid predator, *Coccinella undecimpunctata* L. (Coleoptera: Coccinellidae). *J. Plant Prot. Path., Mansoura Univ.* 5(7): 827-835.
- Ghanim, A. A.; El-Naggar, M. E.; Abdel-Baky, N. F. and Abd El-Halim, E. A. S. (2009): Effect of prey types on certain biological aspects of *Chrysoperla carnea* (Steph.) (Neuroptera: Chrysopidae) under constant temperature. *J. Agric. Sci., Mansoura University* 34(6): 6883-6889.
- El-Serafi H. A.; Ghanim, A. A.; El-Heneidy, A. H. and M. K. El-Sherbeni (2004): Biological characteristics of *Chilocorus bipustulatus* L. and *Chrysoperla carnea* (Steph.) reared on soft scale insects under laboratory conditions. *Egypt. J. Biol. Pest Control* 14(1): 87-95.
- Hodek, I. and A. Honěk(1996): Ecology of Coccinellidae. Kluwer Academic Publishers, The Netherlands.
- Jalali, M. A.; Mehrejjad, M. R. and D. C.Kontodimas(2014): Temperature-Dependent Development of the Five Psyllophagous Ladybird Predators of *Agonoscenapistaciae* (Hemiptera: Psyllidae). *Ann. Entomol. Soc. Am.* 107(2): 445-452.
- Kajita, Y. and E. W. Evans (2010): Relationships of body size, fecundity, and invasion success among predatory lady beetles (Coleoptera: Coccinellidae) inhabiting alfalfa fields. *Ann. Rev. Entomol. Soc. Am.* 103, 750-756.
- Koch, R. L.; Hutchison, W. D.; Venette, R. C. and G. E. Heimpel (2003): Susceptibility of immature monarch butterfly, *Danaus plexippus* (Lepidoptera: Nymphalidae: Danainae), to predation by *Harmonia axyridis* (Coleoptera: Coccinellidae). *Biol. Control* 28, 265-270.
- Mari, J. M.; Rizvi, N. H.; Nizamani, S. M.; Qureshi, K. H. and M. K. Lohar(2005): Predatory efficiency of *Menochilus sexmaculatus* Fab. and *Coccinella undecimpunctata* L. (Coccinellidae: Coleoptera) on alfalfa aphid, *Therioaphis trifolii* (Monell.). *Asian J. Plant Sci.* 4(4): 356-369.
- Mohamed, N. E. and A. A.Ghanim(2008): Biological characteristics of three predatory insects preyed on the silverleaf whitefly, *Bemisia argentifolii* Bellows and Perring under controlled conditions. *Egypt. J. Biol. Pest Cont.* 18(2): 309-313.
- Mohammed, N. E.; Ghanim, A. A. and A. A. Saleh (2011): Influence of certain prey types on certain biological characteristics of *Hippodamia tredecimpunctata* L. under constant temperature and relative humidity. *J. Plant Prot. Path., Mansoura Univ.* 2(11): 1023-1035.

- Obrycki, J. J. and M. J. Tauber (1981): Phenology of three coccinellid species: thermal requirements for development. *Ann. Entomol. Soc. Am.* 74, 31–36.
- Omkar, A. and A. Pervez (2004): Temperature-dependent development and immature survival of an aphidophagous ladybeetle, *Propyleadissecta* (Mulsant). *J. Appl. Entomol.* 128(7): 510-514.
- Osman, M. A. and M. H. Bayoumy (2011): Effect of Prey Stages of the Two-Spotted Mite *Tetranychus urticae* on Functional Response of the Coccinellid Predator *Stethorus gilvifrons*. *ActaPhytopath.Entomol. Hung.* 46 (2): 277–288.
- Pervez, A. and Omkar (2005): Functional responses of coccinellid predators: an illustration of logistic approach. *J. Ins. Sci.* 5 (5): 1–6.
- Sarmiento, R.A.; Pallini, A.; Venzon, M.; Souza, F. F.; Molina-Rugama, A. J. and C. L. Oliveira (2007): Functional response of the predator, *Eriopisconnexa* (Coleoptera: Coccinellidae) to different prey types. *Braz. Arch. Biol. Tech.* 50, 121–126.
- Sethi, S. L. and Atwal, A. S. (1964): Influence of temperature and relative humidity on the development of different stages of lady-bird beetle *Coccinellaseptempunctata* (Coleoptera: Coccinellidae). *Indian J. Agric. Sci.* 34(3): 166-171.
- Skirvin, D. J.; Perry, J. N. and R. Harrington (1997): The effect of climate change on an aphid–coccinellid interaction. *Global Change Biology* 3, 1–11.
- Smith, B. C. (1965): Effects of food on the longevity, fecundity and development of adult coccinellids (Coccinellidae: Coccinellidae). *Can. Entomologist* 97(9): 910-919.
- Solangi, B. K.; Hulno, M. H. and N. Baloch (2007): Biological parameters and prey consumption by zigzag beetle *Menochilussexmaculatus* Fab. against *Rhopalosiphum maidis* fitch, *Aphis gossypii* glov. and *Therioaphis trifolii* monell. *Sarhad J. Agric.* 23 (4): 1097-1101.
- Systat (2008): Sigmaplot 11.0. Systat Software Inc., Chicago, USA.
- Uygun, N. and R. Atlhan (2000): the effect of temperature on development and fecundity of *Scymnuslevallanti*. *BioControl* 45(4): 453-462.
- Xia, J. Y.; Van der Werf, W. and R. Rabbinge (1999): Temperature and prey density on bionomics of *Coccinellaseptempunctata* (Coleoptera : Coccinellidae) feeding on *Aphis gossypii* (Homoptera : Aphididae) on cotton. *Environ. Entomol.* 28(2): 307-314.
- Zalom, F. G.; Goodell, P. B.; Wilson, L. T. and W. J. Bentley (1983): Degree days, the calculation and use of heat units in pest management. University of California, Division of Agriculture and Natural Resources leaflet 21373. Cited in <http://www.ipm.ucdavis.edu/weather/ddconcepts.html>.



الخصائص البيولوجية والاحتياجات الحرارية لنظامي التغذية ابو العيد ١١ نقطة -  
من الحبوب و ابو العيد ٩ نقاط - من البقوليات علي درجات حرارة متعددة  
محمد حسن محمد بيومي ، احمد محمود ابو النجا ، عبد البديع عبد الحميد غاتم و  
غسان عيسى مشحوت  
قسم الحشرات الاقتصادية- كلية الزراعة- جامعة المنصورة- مصر

اجريت هذه الدراسة تحت الظروف المعملية لدراسة الخصائص البيولوجية والاحتياجات الحرارية لكل من مفترس ابو العيد ١١ نقطة وابو العيد ٩ نقاط عند التغذية علي من الحبوب ومن البقوليات علي التوالي علي درجات الحرارة ١٨، ٢٣، ٢٨، ٣٣ م°. اوضحت النتائج المتحصل عليها ان الوقت اللازم لنمو الاطوار غير الكاملة لابو العيد ١١ نقطة وابو العيد ٩ نقاط انخفضت كلما زادت درجة الحرارة حيث سجلت اسرع نمو لها علي درجة حرارة ٣٣ م° والذي كان ١٥.٢٩ ± ٠.٩٨ و ١٤.٠٠ ± ١.٣٥ يوم علي التوالي. كما ازداد معدل استهلاك اليرقة اليومي مع زيادة كل من درجة الحرارة وعمر اليرقة مع اعلي استهلاك للعمر اليرقي الرابع تم تسجيله علي درجة حرارة ٣٣ م° والتي سجلت ٩٩.٧١ ± ١٤.٤٩ و ١٠٥.٤٠ ± ١٠.٥٤ لكل من ابو العيد ١١ نقطة وابو العيد ٩ نقاط علي التوالي. عمر الانثي والذكر لكلا المفترسين ازداد كلما انخفضت درجة الحرارة، في حين ازادت خصوبة الانثي كلما زادت درجة الحرارة. علاوة علي ذلك فان معدل الاستهلاك الكلي لكل من الانثي والذكر لكلا المفترسين ازداد مع زيادة درجة الحرارة مع اعلي استهلاك تم تسجيله خلال فترة وضع البيض. سجلت اناث المفترسين معدلات اعلي من الذكور في الاستهلاك الكلي والاستهلاك اليومي للفريسة علي جميع درجات الحرارة المختبرة. انخفضت معدلات النمو لجميع اطوار المفترسين مع انخفاض درجات الحرارة. سجل طور العذراء واليرقة اقل حد حرج للنمو مقارنة بالاطوار الاخرى لكل من ابو العيد ١١ نقطة وابو العيد ٩ نقاط علي التوالي. سجلت كمية الوحدات الحرارية اللازمة لاكمال نمو كل من المفترس ابو العيد ١١ نقطة وابو العيد ٩ نقاط من طور البيضة الي خروج الحشرة الكاملة في المتوسط ٣٨٠.٧٣ ± ٢٧.٧١ و ٣٦٣.٨٣ ± ١٤.٤٤ وحدة حرارية. اوضحت الدراسة التالية ان كل من مفترس ابو العيد ١١ نقطة وابو العيد ٩ نقاط اكملوا نموهم بنجاح في مدي واسع من درجات الحرارة يتراوح من ١٨- ٣٣ م° مما يشير الي كفاءتهم العالية وامكانية استخدامهم في برامج المكافحة الحيوية لكل من من الحبوب ومن البقوليات علي التوالي.