

EFFECT OF STORAGE CONDITION AND PACKAGING MATERIAL ON INCIDENCE OF STORAGE FUNGI AND SEED QUALITY OF MAIZE GRAINS



Shabana, Y. M.¹; N. A. Ghazy²; S. A. Tolba² and E. A. Fayzalla¹.

¹Plant Pathology Department, Faculty of Agriculture, Mansoura University, El-Mansoura, Egypt. ²Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt.

ABSTRACT

In general, the highest frequency of seed-associated fungi was *A. niger* and *A. flavus*, followed by *Fusarium* spp. and *Alternaria* spp., and then *Penicillium* spp. and *Rhizopus stolonifer*. The storage temperature has no effect on the incidence of *Alternaria* spp., *Fusarium* sp., *Penicillium* spp., and *R. stolonifer*, while it has a significant effect on *A. flavus* and *A. niger*. The best storage conditions for avoiding *A. flavus* incidence on maize grains were storage in refrigerator (10°C) in packages made from cotton cloth, high- or low-density polyethylene whereas the least incidence of this hazardous fungus was recorded (0.10%). Generally, storage in refrigerator (10°C) in packages made from low-density polyethylene was favorable against grain contamination/infection with all storage fungi that may cause seed rot of maize. Storage of maize grain in cold conditions (10°C) caused a significant increase in kernel weight and seed germination while decreased seed moisture content regardless of the packaging material (with an exception with woven polyethylene as packaging material on seed germination) comparing with storage in room conditions. In general, the paper packaging material was favored for all grain quality parameters when grains are stored in cold conditions, whereas it recorded the highest kernel weight (34.86 g) and seed germination (100%) and moderate grain moisture content (8.33%). Under room conditions, grains stored in cotton+polyethylene packages had the highest oil and protein contents (19.67 and 26.20%, respectively).

In conclusion, this study recommends storing maize grains at low temperature inside packages made of low-density polyethylene in order to preserve the vitality of seeds and keep oil content at high levels in addition to reducing the incidence of storage fungi especially those producing mycotoxins/aflatoxins to maintain public health.

Keywords: Maize, storage fungi, cool storage, package type.

INTRODUCTION

Maize (*Zea mays* L.) belongs to family Poaceae (Graminae) (Waniet al., 2014 and Khan et al., 2014) is considered the third most important cereal crop all over the world (FAO, 2013). It is used mainly for human, animal and poultry feeding. Also, there are industrial uses of maize such as the industrial corn products: ethanol and hydrosorb materials, production of corn oil, dextrose and high fructose corn sweetness (Gwirtz and Garcia-Casal, 2014). Proper storage may help in alleviating problems of seed viability. Seeds of most plant species may be safely stored for several months by careful control

of temperature and relative humidity (Lacerdaet *al.*, 2003 and Chatthaet *al.*, 2012).

In some parts of the world, especially in the tropics, conditioned storage is necessary in order to maintain high viability of some seeds from harvesting to planting (Harrington, 1973; Santosoe *al.*, 2015; Tripathi and Lawande, 2014). The storage fungi, mainly comprising several species of *Aspergillus* and *Penicillium*, do not invade grains to any appreciable degree or extent before harvest (Tuite, 1961). Seed quality is a multiple criterion that encompasses several important seed attributes: which include chemical composition, physiological germination and seed vigor and presence of seed-borne pathogens. During storage, seed quality can remain at the initial level or decline to a level that may make the seed unacceptable for planting purpose, what is related to many determinants: environment conditions during seed production, pests, diseases, seed oil content, seed moisture content, mechanical damages of seed in processing, storage longevity, packaging, pesticides, air temperature and relative air humidity in storage, biochemical injury of seed tissue (Al-Yahya, 2001; Šimicet *al.*, 2004; Guberacet *al.*, 2003; Heatherly and Elmore, 2004). Field fungi, *Alternaria*, *Cladosporium*, *Curvularia*, *Fusarium* and *Helminthosporium* usually do not continue to grow in grains after harvest, but may remain alive for years in grains stored at low moisture content and low temperature. While storage fungi i.e., *Aspergillus* spp. and *Penicillium* can grow in stored grain under bad storage conditions and cause serious losses (Mehrotra, 1983).

Changes in grain germination, chemical composition, acidity and oil content of three maize genotypes, due to infection by different fungi of ear and kernel rots, were affected by storage periods (6 and 18 months), storage temperature (room temperature and 10°C) (Timóteo and Marcos-Filho, 2013) and package materials (paper, woven polyethylene and high density polyethylene) (Chathaet *al.*, 2012 and Rahraw, *et al.*, 2013). The present study aimed to determine the effect of storage conditions in combination with packaging materials on the incidence of maize storage fungi and on grain quality.

MATERIALS AND METHODS

The present experiment was carried out in the centrallaboratory and the laboratory of plant pathology at Sakha Agricultural Research Station (SARS), Kafr El-Sheikh, Egypt.

The experiment was conducted during 2011. Visual symptomless seed samples (12 kg) of tested maize cultivar "Three Way Cross 324" (TWC324) were taken at the harvest time and dried using hot air dried methods to 14% moisture content by moisture tester (model 8400, serial No.4, Michigan USA) and packaged in 6 types of packages made from different materials i.e., paper, cotton clothes, woven polyethylene, cotton+polyethylene, high density polyethylene, and low density polyethylene. Each package was filled with one kg of tested maize seeds and stored under two different conditions: cold room (10°C) and at room

conditions. Random seed samples with three replicates were taken from each package after 6 months of storage. The presence of storage fungi in each treatment was assessed according to ISTA (2004): whereas 200 seeds were surface sterilized with 5% sodium hypochlorite for 5min., washed thoroughly with sterile distilled water, and then dried in a laminar flow. The seeds were plated in potato dextrose agar (PDA) plates (5seeds/plate) and incubated for 7 days at 27°C. The developed fungi were recorded and the incidence percentage of each fungal genera was estimated. The developed fungi were purified by using astereoscopic microscope; hyphal tips from fungi were transfer it to PDA plates and later to slants for identification. Identification of fungi was carried out using their morphological and microscopic features according to Barrant and Hunter (2004). Barron, 1968; Hunter and Barnett, 1973; Hunter *et al.*, 1978 and Alexopoulos *et al.* 1996 were provide additional information on many aspects of the morphology, sporulation, growth, ecology of imperfect fungi. Identification was reconfirmed by the help of staff at the Department of Maize and Sugar Beet Disease, Sakha Agriculture Research Station, A. R. C., Giza, Egypt. Germination percentage, 100-grain weight, grain moisture content, and oil and protein contents were determined according to methods described by (AOAC, 1999). All obtained data were statistically analyzed according to the technique of analysis of variance (ANOVA) for the split-plot design to each experiment.

RESULTS AND DISCUSSION

Effect of storage conditions and packaging material on incidence of fungi during storage.

In general, the highest frequency of seed-associated fungi was *A. niger* and *A. flavus*, followed by *Fusarium* spp. and *Alternaria* spp., and then *Penicillium* spp. and *R. stolonifer* (Table 1). The effect of storage package types and conditions on frequency of fungi associated with stored maize grain is shown in Table (1). The obtained data clearly show that, the storage temperature has no significant effect on the incidence of *Alternaria* sp., *Fusarium* sp., *Penicillium* sp., and *Rhizopus stolonifer* (Table 1), whereas there were no significant differences among all packages types regarding these fungi under both storage conditions (at room temperature and at 10°C). Thus, the storage temperature has a significant effect only on *A. flavus* and *A. niger* (Table 1).

The best storage conditions for avoiding *A. flavus* incidence on maize grains were storage in refrigerator (10°C) in 2 tested types of packages made from cotton cloth, high- or low-density polyethylene whereas the least incidence (0.10%) of this storage fungus was recorded (Table 1).

Table (1): Effect of storage package material and storage condition on incidence of fungi associated with maize grain (TWC.324 hybrids cv.) after six months of storage.

Fungus	Storage conditions	Packaging material type						Mean
		Paper	Cotton clothes	Woven polyethylene	Cotton+ polyethylene	High-density polyethylene	Low-density polyethylene	
<i>Alternaria</i> sp.	At room Temperature (25±2°C)	13.33 ^a	6.70 ^a	3.40 ^a	10.07 ^a	10.07 ^a	6.70 ^a	8.37 ^a
	In refrigerator (10°C)	6.70 ^a	16.67 ^a	10.00 ^a	23.37 ^a	13.33 ^a	3.40 ^a	12.24 ^a
Mean		10.02 ^a	11.68 ^a	6.70 ^a	16.71 ^a	11.70 ^a	5.05 ^a	10.31
<i>A. flavus</i>	At room Temperature (25±2°C)	13.37 ^{de}	43.33 ^{abc}	3.40 ^{de}	60.00 ^a	36.67 ^{bc}	50.00 ^{ab}	34.46 ^a
	In refrigerator (10°C)	23.33 ^{cd}	0.100 ^e	13.33 ^{de}	3.40 ^{de}	0.10 ^e	0.10 ^e	6.73 ^b
Mean		18.35 ^{ab}	21.72 ^{ab}	8.37 ^b	31.70 ^a	18.38 ^{ab}	25.05 ^a	20.60
<i>A. niger</i>	At room Temperature (25±2°C)	23.37 ^{bc}	30.03 ^b	10.03 ^{bc}	30.00 ^b	66.67 ^a	30.03 ^b	31.69 ^a
	In refrigerator (10°C)	3.40 ^c	13.33 ^{bc}	13.37 ^{bc}	23.33 ^{bc}	6.70 ^{bc}	6.73 ^{bc}	11.14 ^a
Mean		13.38 ^b	21.68 ^{ab}	11.70 ^b	26.67 ^{ab}	36.68 ^a	18.38 ^b	21.42
<i>Fusarium</i> sp.	At room Temperature (25±2°C)	20.00 ^a	10.03 ^a	10.03 ^a	6.70 ^a	0.10 ^a	6.70 ^a	8.93 ^a
	In refrigerator (10°C)	23.33 ^a	23.33 ^a	16.67 ^a	16.67 ^a	13.37 ^a	6.73 ^a	16.68 ^a
Mean		21.67 ^a	16.68 ^{ab}	13.35 ^{ab}	11.68 ^{ab}	6.73 ^b	6.72 ^b	12.8
<i>Penicillium</i> sp.	At room Temperature (25±2°C)	0.10 ^a	3.40 ^a	6.70 ^a	0.10 ^a	0.10 ^a	3.40 ^a	2.30 ^a
	In refrigerator (10°C)	6.70 ^a	10.03 ^a	0.10 ^a	0.10 ^a	3.40 ^a	3.40 ^a	3.96 ^a
Mean		3.40 ^a	6.72 ^a	3.40 ^a	0.10 ^a	1.75 ^a	3.40 ^a	3.13
<i>Rhizopus stolonifer</i>	At room Temperature (25±2°C)	0.10 ^a	10.10 ^a	0.10 ^a	0.10 ^a	0.10 ^a	3.40 ^a	2.31 ^a
	In refrigerator (10°C)	0.10 ^a	0.10 ^a	0.10 ^a	0.10 ^a	10.10 ^a	0.10 ^a	1.76 ^a
Mean		0.10 ^a	5.08 ^a	0.10 ^a	0.10 ^a	5.08 ^a	1.75 ^a	2.04

*Values for each fungus followed by the same letter(s) are not significantly different, according to LSD test at $P = 0.05$.

These results are supported with the findings of El-Sayed and Tolba (2005) who reported that storage under 10°C in high-density polyethylene packages led to reducing infection of maize grains by pathogens including *A. flavus*. Storage in refrigerator (10°C) in packages made from low-density polyethylene was favorable against grain infection with all storage fungi that cause seed rot of maize. Also, Mehrotra (1983) reported that the storage fungi i.e., *Aspergillus* spp. and *Penicillium* can grow in stored grain under bad storage conditions and cause serious losses

Effect on maize grain quality (seed weight, germination, and moisture content):

Data in Table (2) show that storage of maize grain in cold conditions (10°C) caused a significant increase in kernel weight and seed germination while decreased seed moisture content regardless of the packaging material (with an exception with woven polyethylene as packaging material on seed germination) comparing with storage in room conditions (25±2°C). In general, the paper packaging material was favored for all grain quality parameters when grains are stored in cold conditions, whereas it recorded the highest kernel weight (34.86 g) and seed germination (100%) and moderate grain moisture content (8.33%) (Table 2). Similar results were obtained by El-Sayed and Tolba (2005) who found that seed germination and weight of 100 kernels were affected by storage temperature and packaging material. They found that germination percentages and the weight of 100 grain were significantly decreased under warehouse conditions. Our results were also supported by those of El-Sayed *et al.* (2004) who reported that germination decline was more rapidly at warehouse, but less at low temperature, while the lowest value of the weight of 100 grain was recorded at warehouse storage conditions. The results in Table (2) showed clearly that, germination percentage, weight of 100 kernels Malakeret *et al.* (2008) who tested types of containers [viz., 'dole' (bamboo made), earthen pitcher, tin container, polyethylene bag] and storage conditions [cool (10 C) and room temperature] for their effect on quality of wheat seeds during storage. They found that highest germination percentage was observed under storage in refrigerator and polyethylene bag followed by tin container and earthen pitcher. The results also similar with the obtained by Mettananda *et al.*, 2001. Who found that the maize grain stored in woven poly-ethylene under cold room have good quality during storage period comparing with maize grain stored poly-ethylene under room temperature.

Table (2): Effect of storage packaging material and storage conditions on some seed quality parameter of maize grain (TWC.324 hybrids cv.) after six months storage.

Fungus	Storage conditions	Packaging material type						Mean
		Paper	Cotton clothes	Woven polyethylene	Cotton+ polyethylene	High-density polyethylene	Low-density polyethylene	
Weight of 100 kernels (g)	At room Temperature (25±2°C)	33.13 bc	31.61 d	31.50 d	31.08 de	30.13 ef	29.48 f	31.16 b
	In refrigerator (10°C)	34.86 a	34.03 ab	34.88 a	31.36 de	32.15 cd	34.56 a	33.64 a
Mean		33.99 a	32.82 bc	33.19 ab	31.22 d	31.14 d	32.02 cd	32.40
Germination (%)	At room Temperature (25±2°C)	40.00 c	0.100 e	90.00 b	23.33 d	0.100 e	0.100 e	25.61 b
	In refrigerator (10°C)	100.100 a	96.67 ab	90.00 b	100.00 a	96.67 a	90.00 b	96.11 a
Mean		70.00 b	48.38 d	90.00 a	61.67 c	50.05 d	45.05 d	60.86
Moisture content (%)	At room Temperature (25±2°C)	14.10 c	14.17 c	14.10 c	14.23 c	15.43 a	15.73 b	14.63 a
	In refrigerator (10°C)	8.33 e	7.27 g	7.57 f	7.73 f	9.83 d	9.77 d	8.42 b
Mean		11.22 b	10.72 d	10.83 cd	10.98 c	12.63 a	12.75 a	11.53

*Values for each fungus followed by the same letter(s) are not significantly different, according to LSD test at $P = 0.05$.

Effect on oil and protein contents

Concerning the effect of storage packages and conditions on oil content of stored maize grain, data presented in Table (3) show that, under room conditions, grains stored in cotton+polyethylene packages had the highest oil and protein contents (19.67 and 26.20%: respectively). No significant difference was found among the other different packages stored in refrigerator. Similar results were obtained by El-Sayed and Tolba (2005) who reported a significant reduction of oil and protein contents in grains stored at warehouse conditions, while storage under 10°C produced the highest values of oil content. Also they found that crude protein decreased in grains stored at 10°C while increased in case of storage at laboratory conditions.

In conclusion, this study recommends storing maize grains at low temperature inside packages made of low-density polyethylene in order to preserve the vitality of seeds and keep oil content at high levels in addition to reducing the incidence of storage fungi especially those producing mycotoxins/aflatoxins to maintain public health.

Table (3): Effect of packaging material and storage conditions on oil and protein contents of maize grains (TWC.324 hybrids cv.) after six months storage.

Fungus	Storage conditions	Packaging material type						Mean
		Paper	Cotton clothes	Woven polyethylene	Cotton+ polyethylene	High-density polyethylene	Low-density polyethylene	
Seed oil content (%)	At room Temperature (25±2°C)	6.20 a	3.46 a	5.04 a	19.67 a	4.12 a	3.46 a	6.99 a
	In refrigerator (10°C)	4.65 a	4.21 a	5.12 a	5.48 a	4.96 a	5.11 a	4.92 a
Mean		5.43 a	3.84 a	5.08 a	12.57 a	4.54 a	4.29 a	5.96
Seed protein content (%)	At room Temperature (25±2°C)	17.30 d	17.46 d	20.98 c	26.20 a	23.77 b	21.28 c	21.17 a
	In refrigerator (10°C)	14.26 e	17.34 d	14.38 e	14.90 e	10.63 f	11.10 f	13.77 b
Mean		15.78 d	17.40 b	17.68 b	20.55 a	17.20 bc	16.19 CD	17.47

*Values for each fungus followed by the same letter(s) are not significantly different, according to LSD test at $P = 0.05$.

REFERENCES

A.O.A.C. 1999. Official Methods of Analysis of the Association of Official Analytical Chemists 15th Edition, published by Association of Official Analytical Chemists, Arlington, Virginia, USA.

Alexopoulos, C. J., Mims, C. W. and Blackwell, M. 1996. Introductory Mycology: John Wiley & Sons, New York.

Al-Yahya, S. A. 2001. Effect of storage conditions on germination in wheat. *J. Agro. and Crop Sci.* 186: 273- 279.

Barnett, H.L. and B.B. Hunter. 2004. Illustrated Genera of Imperfect Fungi. Fourth edition. APS PRESS, MN, USA. 218 pp.

- Barron, G. L. 1968. *The Genera of Hyphomycetes from Soil*. Williams & Wilkins, Baltimore, MD.
- Chattha, S. H., Jamali, L. A., libupoto, K. A. and Mangi, H. R. 2012. Effect of different packing materials and storage conditions on the viability of wheat seed (td-1 variety). *Sci., Tech. and Dev.* 31 (1): 10-18.
- El-Sayed, S., A., Kineber, M. A. and El-Kady, E. A. 2004. Effect of storage environment and package material on storability, yield and its quality of flax seeds. *J. Agric. Res. Tanta Univ.* 30(3): 616-638.
- El-Sayed, S., and Tolba, S.A. E. 2005. Effect of storage conditions and package on germination, chemical composition, infection by ear and kernel rot disease and oil quality in some maize grain genotypes. *Alex. J. Agric. Res.* 50 (2): 23-33.
- FAO, 2013. *FAO Statistics Division*. May 2013.
- Guberac, V., Maric, S., Lalic, A., Drezner, G. and Zdunic, Z. 2003. Hermetically sealed storage of cereal seeds and its influence on vigor and germination. *J. Agronomy and Crop Science* 189: 54-56.
- Gwartz, J. A. and Garcia-Casal M. N. 2014. Processing maize flour and corn meal food products *Annals of the New York Academy of Sciences* 1312: 66-75.
- Harrington, J. F. 1973. Packaging seeds for storage and shipment. *Seed Science and Technology* 1: 701-709
- Heatherly, L. G. and Elmore, R. W. 2004. Managing Inputs for Peak Production. Pages 451-536, in: *Soybeans: Improvement, Production and Uses*. Boerma H. R., Specht, J. E. (eds), 3rd Edition, Agronomy N-16, ASA, CSSA, SSSA, Madison, Wisconsin, USA.
- Hunter, B. B. and Barnett, H. L. 1973. Deuteromycetes (Fungi Imperfecti), In: *Handbook of Microbiology: (Vol. 1), Organismic Microbiology*. Laskin, A. I. and H. A. Lechevalier, Eds. CRC Press, Cleveland, OH.
- Hunter, B. B. and Barnett, H. L. and Buckelew, T. P. 1978. Deuteromycetes (Fungi Imperfecti), In: *Handbook of Microbiology: (Vol. 2), Fungi, Algae, Protozoa, and Viruses*. Laskin, A. I. and Lechevalier, H. A., Eds. CRC Press, West Palm Beach, FL.
- ISTA. 2004. *International Rules for Seed Testing*. 4th edition. The International Seed Testing Association, Zurich. 2004.
- Khan, F., Khan, S., Fahad, S., Faisal, S., Hussain, S. and Ali, S. 2014. Effect of different levels of nitrogen and phosphorus on the phenology and yield of maize varieties. *American J. Plant Sci.* 5: 2582-2590.
- Lacerda, A. D. S., Lazarini, E. and Filho, W. V. V. 2003. Storage of desiccated soybean seed and the evolution of physiologic, biochemical and sanitary characteristics. *Braz. J. of Seeds* 25: 97-10.
- Mettananda, K. A., weerasena, S. L. and Lianage, Y. 2001. Effect of storage environment, package material and seed moisture content on storability of maize (*zea mays* L.) seeds. *Ann. of the Sri Lanka Department of Agric.* 3: 131-142.
- Malaker, P. K., Mian, I. H., Bhuiyan, K. A., Akanda, A. M. and Reza, M. M. A. 2008. Effect of storage containers and time on seed quality of wheat. *Bangladesh J. Agric. Res.* 33 (3): 469-477.

- Mehrotra, B. S. 1983. The impact of fungal infestation of cereal grains in field and storage. Pages 185-200, in: Recent Advances in Plant Pathology. Husain, A., Singh, K., Singh, B. P. and Agnihotri, V. P. (eds.), Lucknow Print House. India.
- Rahraw, R., Rai, P. K., Kumar, A., Singh, B. A., Chaurasia, A. K. and Neeru Bala, N. 2013. Influence of polymer seed coating, biocides and packaging materials on storability of Wheat (*Triticum aestivum* L.). Rep. Opinion 5 (12): 51-55. (ISSN: 1553-9873). <http://www.sciencepub.net/report>.
- Santoso, B. B., Arya Parwata, I. G. M. and Jaya, I. K. D. 2015. Seed viability and oil content of castor bean (*Ricinus communis* L.) as affected by packaging materials during storage. International J. Appl. Sci. and Tech. 5(2): 56-61.
- Šimic, B., Popovic, S. and Tucak, M. 2004. Influence of corn (*Zea mays* L.) inbred lines seed processing on their damage. Plant, Soil and Environment 50: 157-161.
- Timóteo, T. S. and Marcos-Filho, J. 2013. Seed performance of different corn genotypes during storage. J. Seed Sci., 35 (2, p): 207-215.
- Tripathi, P. C. and Lawande, K. E. 2014. Effect of seed moisture and packing material on viability and vigour of onion seed. J. Engineering Computers and Appl. Scie. (JECAS) 3 (7): 1-5.
- Tuite, J. F. 1961. Low incidence of storage molds in freshly harvested seed of soft red winter wheat. Plant Dis. Repr. 43:470.
- Wani, A. A., Joshi, J., Titov, A. and Tomar, D. S. 2014. Effect of seed treatments and packing materials on seed quality parameters of maize (*Zea mays* L.) during storage. Indian J. Appl. Res. 4 (4): 2249-555.

تأثير ظروف التخزين ونوع عبوات التعبئة على فطريات التخزين وجودة الحبوب في الذرة الشامية

ياسر محمد نور الدين شبانه^١، نصر أحمد غازي^٢، صبحى عبد العزيز طلبه^٢ و السيد عبد المجيد فيظ الله^١

^١ قسم أمراض النبات-كلية الزراعة - جامعة المنصور- المنصورة - مصر
^٢ معهد بحوث أمراض النباتات- مركز البحوث الزراعية - الجيزة - مصر

أجريت هذه الدراسة لتقدير تأثير درجة حرارة التخزين (درجة حرارة الغرفة $25 \pm 5^{\circ}\text{C}$)، والتخزين البارد بالتلاجة على 10°C)، وكذلك نوع عبوات التخزين (الورق، القطن، البولي إيثيلين المنسوج، القطن+البولي إيثيلين، البولي إيثيلين عالي الكثافة، والبولي إيثيلين منخفض الكثافة) لحبوب الذرة الشامية لمدة ستة أشهر وذلك على الإصابة بفطريات الأعفان (الفيزاريوم، أسبرجلس فلافوس، أسبرجلس نيجر، بنيسيليم، الترناريا، الريزوبس) وصفات جودة الحبة (نسبة الإنبات، وزن حبة، الرطوبة النسبية في الحبوب، نسبة البروتين، ونسبة الزيت في الحبوب) وذلك باستخدام الصنف الهجين الثلاثي ٣٢٤ خلال الفترة من ١ سبتمبر ٢٠١١ وحتى ١ مارس ٢٠١٢، ويمكن تلخيص النتائج كالآتي:

دللت النتائج المتحصل عليها أنه بشكل عام، كان أعلى معدل تلوث بالفطريات المصاحبة لبذور الذرة الشامية كان للفطر أسبرجلس نيجر والفطر أسبرجلس فلافوس، تلاها فطريات الفيزاريوم والألترناريا، ثم البنسليوم والرايزوبس. وأوضحت الدراسة أن درجة حرارة التخزين لم يكن لها تأثير معنوي على الإصابة بفطريات الألترناريا والفيزاريوم والبنسليوم والرايزوبس، في حين كان لديها تأثير معنوي على الإصابة بفطري أسبرجلس فلافوس وأسبرجلس نيجر. كانت أفضل ظروف التخزين لتجنب الإصابة بفطر أسبرجلس فلافوس هي التخزين في التلاجة في عبوات مصنوعة من قماش القطن، أو البولي إيثيلين عالي أو منخفض الكثافة حيث سجلت أقل حالات الإصابة بهذا الفطر الخطير (١٠.١٠٪). وبصفة عامة، كان التخزين في التلاجة في عبوات مصنوعة من البولي إيثيلين منخفض الكثافة مناسباً لحماية حبوب الذرة الشامية من الإصابة بفطريات أعفان البذور. وقد أدى تخزين حبوب الذرة في التلاجة إلى زيادة كبيرة في وزن المائة حبة وكذلك نسبة إنبات البذور، كما أدى إلى انخفاض نسبة الرطوبة في البذور بغض النظر عن نوعية مادة عبوات التخزين (باستثناء البولي إيثيلين المنسوج على إنبات البذور) مقارنة مع التخزين في ظروف الغرفة. بشكل عام، كان التخزين في عبوات ورقية في التلاجة مناسباً لصفات الجودة في الحبوب، حيث سجلت أعلى وزن للمائة حبة (٣٤.٨٦ جم) وأعلى نسبة إنبات للبذور (١٠٠٪) ومحتوى معتدل للرطوبة في الحبة (٨.٣٣٪). وفي ظل ظروف الغرفة، سجلت الحبوب المخزنة في أكياس من القطن + البولي إيثيلين أعلى نسبة من محتوى الزيت والبروتين (١٩.٦٢ و ٢٦.٢٠% على التوالي).

توصى الدراسة بتخزين حبوب الذرة في درجات حرارة منخفضة داخل عبوات مصنوعة من البولي إيثيلين منخفض الكثافة وذلك حفاظاً على حيوية التقاوى ومحتواها من الزيت بالإضافة إلى خفض نسبة الإصابة بفطريات التخزين خاصة المنتج المسموم الفطرية للحفاظ على الصحة العامة.