

INFLUENCE OF ORGANIC COMPOST, PROLINE AND THE BIOFERTILIZER (SALINITY DURABLE BACTERIA) ON BARLEY GROWTH AND NUTRIENTS UPTAKE UNDER HIGH SALINITY CONDITIONS

Abbas, Y.M* and Ewees, M. S. A.**

*Soil, Water and Environment Res. Institute, Agric. Res. Center, Giza, Egypt.

** Soil and Water Department, Fac. of Agric., El-Fayoum University.

ABSTRACT

A filed experiment was carried out on salt affected soil at Kasr El-Basel village, south Etsa district, El-Fayoum Governorate, Egypt, during the winter season 2013/2014. Objective of this work was to study the effects of applied local compost at a rate of 20 m³ fed⁻¹, amino acid (proline) sprayed at rate of 3 mg/L fed⁻¹ at 20, 45, and 60 days after sowing) and biofertilizer (salinity durable bacteria) as either solely or combined treatments on barley (*Hordeum vulgare*, c.v. Giza 123) growth and yield parameters. The experimental field was irrigated with saline water (a mixture of the fresh Nile water and agricultural drainage water). The quality of the used irrigation water was classified as C2S1 (EC_{iw} = 1.66 dS/m and SAR 5.35). The influence of treatments on some soil properties (soil pH, EC_e, ESP and available macro and micronutrient contents) was studied.

Obtained results indicated that, the values of EC, ESP and pH, decreased however, the organic matter and CEC increased with the application of compost, proline and biofertilizer. The best treatment was found to be (Compost + Proline + Biofertilizer). The application of (Compost + Proline + Biofertilizer) also, decreased soil bulk density, while increased hydraulic conductivity, total porosity and soil moisture content. Plant height, number of grains/ spike, number spikes / m², 1000 grains weight, and grain and straw yields were also improved with treatments. The greatest values were associated with the triple combined treatment (Compost + Proline + Biofertilizer) as compared to the other combined or solely ones.

It could be recommended that compost, proline and the biofertilizer (salinity durable bacteria) could be used to alleviate the hazardous effects of either soil or water salinity, which negatively affected barley seed yield and quality.

Key words: Compost, Amino acids, Proline, Biofertilizers, Salinity durable bacteria, Barley, plant growth and quality parameters.

INTRODUCTION

Soil management is usually carried out through the addition of natural soil amendments and biofertilizers that have become one of the most important

practices for improving soil hydrophysical, chemical and biological properties and in turn enhancing its productivity for different vegetable crops.

Salinity is one of the major problems facing agriculture in arid and semi-arid regions. Egypt is one of the countries that suffer severe salinity problems. About 33% of the cultivated land, which comprises only 3% of total land area in Egypt are saline. Such salinity is mainly due to low precipitation (< 25 mm annual rainfall), high temperature (that ranges from 35 to 45°C), high surface evaporation (1500- 2400 mm/year), poor drainage in about 98% of the cultivated land under irrigation, high water table (less than one meter below the soil surface), and irrigation with low quality saline water (up to 4.5 dS/m). Salt stress generally leads to a reduction in biomass production owing to a diminution of the water potential, specific ion toxicities, or nutrient deficiencies (Parida and Das, 2005).

Reduction in salt affected soils productivity is due to the high osmotic potential in solution within the crop root zone, which causes disturbances in nutrients balance, reduces either soil available nutrients or water uptake by roots of growing plants and consequently reduces the quality and yield of crops (Ayers and Westcot, 1985).

The harmful effect of salinity stress is also attributed to an ionic imbalance in plant cells due to the excessive accumulation of Na^+ and Cl^- that result in a reduction in K^+ , Ca^{2+} and Mn^{2+} uptake (Tester and Davenport, 2003). Plant response to fertilizers depends on severity of salt stress in the root zone and fertilizers application to saline soils may exacerbate soil salinization (Maas and Grattan, 1999).

Barley is one of the salt-tolerant crops that tolerate adverse conditions such as salinity, heat, drought, and low soil fertility under arid and semiarid conditions.

Several investigators studied the effect of compost, proline and bio-fertilizers (salinity durable bacteria) in decreasing soil salinity effects. Khaled *et al.*, (2011) reported that the role of compost is vital in salt-affected soils because the organic source is ultimate opportunity to improve soil physical properties, which have been deteriorated to the extent that water and air passage become extremely difficult in such soils. Tea compost has been used to improve the properties of soil and reduce salinity problems, as well as to improve plant growth (Sunjeong *et al.*, 2010).

Proline amino acid plays an adaptive role in the tolerance of plant cells to salinity by increasing the concentration of cultural osmotic components in order to equalize the osmotic potential of the cytoplasm. (Wareing and Phillips, 1978, and Wated *et al.*, 1983). The increase in proline content in plant tissues with the increase in salinity retards protein synthesis, and consequently accumulates free amino acids, including proline (Wated *et al.*, 1983, Ouerghi *et al.*, 1991, Zidan and Malibari, 1993, Barakat and Abdel-Latif, 1995, Yurekli *et al.*, 1996, and El-Leboudi *et al.*, 1997). In this connection, Wageeh

(1994) reported that the best treatments that gave the most favorable response for growth by wheat plants were seed soaking for 12 hours interval in solutions of 5 ppm of each of the following amino acids: proline, glutamic acid and aspartic acid compared with soaking in distilled water.

Torello and Rief (1986) and **Tipiramaz and Cakirlar (1990)** found that the accumulation of proline was rapid in barley.

Beneficial soil microorganisms such as PGPR showed positive effects in plants, particularly on parameters such as the rate of germination, tolerance to drought and salinity and the weight of stems and roots. (**Silini et al., 2012**).

The inoculation of soils with salt-tolerant strains improves plant growth as compared with the effect of salt-sensitive strains (**Zou et al., 1995**).

Objective of the present work was to study the possibility of alleviating the harmful effects of soil salinity on barley plants growth and yield by the application of compost, proline amino acid and inoculation with salinity durable bacteria.

MATERIALS AND METHODS

A field experiment was carried out on salt affected soil at kasr El-Baseel village, south Etsa district, El-Fayoum Governorate, Egypt, during the winter season 2013/2014. Compost was applied at a rate of 20 m³ fed⁻¹, as individual or combined with proline sprayed at rate of 3 mg/L fed⁻¹ at 20, 45, and 60 days after sowing. Salinity durable bacteria was provided by the Bio-fertilizer Production Unit, Department of Microbiology, Soils, Water and Environment Research Institute, Agric. Res. Center, Giza. The seeds were soaked with *Azospirillum* and *Azotobacter* at the rate 400 gm/fed.

The experimental soil was irrigated with saline water (a mixture of the fresh Nile water and agricultural drainage water) which could be classified as (C2S1). Increased problems for soil salinity (C2) is expected. The chemical characteristics of irrigation water were carried out according to the described methods and suitability criteria for irrigation after **Page et al. (1982)** and **Ayers and Westcot (1985)**, respectively, as shown in Table (1).

Chemical analysis of compost used are presented in Table (2). The experimental plots were arranged in a combined split plots design with three replicates. The area of each plot was 10.5 m² (3.0 m width x 3.5 m length). Plots were ploughed twice in two ways after the addition of superphosphate fertilizer (15.5 % P₂O₅) at a rate of 100 kg fed⁻¹. All treatments received a similar fertilization with recommended dose of nitrogen in the form of ammonium nitrate (33.5 % N) at the rate of 134 kg N/fed for barley in to equal doses during the growing period, i. e., after 15 & 40 days from plantation. Potassium sulphate (48 % K₂O) was added at a rate of 50 kg fed⁻¹ in two equal doses, after 15 and 40 days from planting.

Treatments were as follows:

1. Control (c)
2. Compost at rate of 20 m³/fed.

3. Proline sprayed at the rate of 3 mg/L at 20, 45, and 60 days after sowing.
4. Biofertilizer (salinity durable bacteria): the seeds were soaked with Azospirillum and Azotobacter at the rate 400 gm/fed.
5. Compost + Proline.
6. Compost + Biofertilizer.
7. Proline+ Biofertilizer.
8. Compost + Proline + Biofertilizer.

Table (1): Chemical properties of used irrigation water of Baher El-Ghark

pH	EC dSm ⁻¹	Soluble ions (meq L ⁻¹)							SAR	*Irrigation water quality
		Ca ²⁺	Mg ²⁺	Na ⁺	K ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ²⁻		C2S1
8.40	1.66	3.07	4.29	8.16	0.41	3.83	6.74	5.36	4.25	

*According to Ayers and Westcot (1985) scale.

Table (2): Physical and chemical properties of the compost used.

EC dSm ⁻¹ (1:10)	pH (1:10 water suspension)	Total NPK (%)			C/N ratio	Organic matter (%)	Ammonium bicarbonate- DTPA- extractable micronutrients (mg kg ⁻¹)			
		N	P	K			Fe	Mn	Zn	Cu
2.45	7.6	1.51	0.66	1.86	16/1	35.7	79.63	36.42	24.83	9.75

Barley was planted in the winter season 2013/2014 and harvested at maturity stage to determine the yields of grains and straw. Harvest Of barley crop was done after 140 days from sowing. At harvest, grains were separated from the vegetative part (straw) and the weights of 1000 grain and straw per plots were recorded as dry weight. The obtained straw and grain from 1.0 m² central area of all experimental plots were separately analyzed for N, P, and K.

Soil samples were collected from the surface layer (0-30 cm) before starting treatments and at the end of vegetative growth (80 day after plantation), then dried, crushed and sieved through a 2 mm screen. Samples were analyzed to measure the electrical conductivity (EC_e) and pH (Jackson, 1973). Particle size distribution and calcium carbonate were determined according to (Piper, 1950). Soil organic matter was determined according to Walkley-Black method (Black et al., 1965). Cation exchange capacity was determined by using method of (Richards, 1954). Physical and chemical analyses of the studied soil before cultivation are shown in Table (3). Plant samples (grain and straw) were taken after harvest and digested to determine their contents of N, P, K according to Chapman and Prrate, (1961). Available macronutrients of N, P and K in soil were extracted by 1% potassium sulphate, 0.5 M sodium bicarbonate and 1 N ammonium acetate, respectively (Soltanpour and Schwab, 1977) and their contents in soil were determined according to Jackson (1973). Available micronutrients of Fe, Mn, Zn and Cu in soil were extracted using ammonium bicarbonate-DTPA extract according to Soltanpour and Schwab, (1977) and their contents in soil were measured by using the Atomic Absorption Spectrophotometer.

Data obtained of the tested plant characters were subjected to statistical analysis according to **Snedecor and Cochran (1980)** to define the least significant difference test (L.S.D. at $p=0.05$ level), which was used to verify the differences between the tested treatments.

Table (3): Some physical and chemical characteristics of the experimental soil

Soil characteristics	Value	Soil characteristics.	Value			
<i>Particle size distribution %</i>		ESP%	12.46			
Coarse sand	5.80	<i>Soluble ions in soil paste extract (m molc L⁻¹):</i>				
Fine sand	14.80					
Silt	30.10		Ca ⁺⁺	31.24		
Clay	49.30		Mg ⁺⁺	22.17		
Soil texture class	Clayey		Na ⁺	57.47		
CaCO ₃ %	2.48		K ⁺	1.60		
Organic matter %	0.86		CO ₃ ⁻	0.00		
E _{Ce} in dSm ⁻¹ (Soil paste):	11.33		HCO ₃ ⁻	2.78		
pH (Soil paste extract):	7.87		Cl ⁻	61.81		
			SO ₄ ⁻	47.89		
<i>Available macro and micronutrients (mg/kg soil)</i>						
N	P	K	Fe	Mn	Zn	
80.00	4.50	152	4.32	0.92	1.46	
<i>Critical levels of nutrients after Lindsay and Norvell (1978) and Page et al. (1982)</i>						
Limits	N	P	K	Fe	Mn	Zn
Low	<40.0	<5.0	<85.0	<4.0	<2.0	<1.0
Medium	40.0-80.0	5.0-10.0	85.0-170.0	4.0-6.0	2.0-5.0	1.0-2.0
High	>80.0	>10.0	>170	>6.0	>5.0	>2.0

RESULTS AND DISCUSSION

I. A general view on the experimental soil:

The results obtained of particle size distribution, Table (1), reveal that the studied soil is fine texture (clayey), and low contents of both CaCO₃ and organic matter.

II. Response of some soil chemical properties and nutrients contents availability to treatments:

a. Soil physical and chemical characteristics:

Data in Table (4) indicated that the application of compost and/or biofertilizer (salinity durable bacteria) resulted in decreases in the values of soil bulk density, E_{Ce}, pH and ESP. On the other hand, each of total porosity%, field capacity%, wilting point%, available water%, hydraulic conductivity, organic matter% and CEC were increased with the application of either compost or biofertilizer separately or in combination. The application of (Compost + proline + Biofertilizer) resulted in the greatest effect on each of the studied properties in comparison with rather the control and the or each of them alone. The results are

in agreement with those obtained by **Sunjeong et al., (2010)** who reported that tea compost has been used to improve the soil properties of the soil and reduce salinity problems.

b. Soil available macro and micronutrient contents:

The magnitudes of soil available nutrients extracted before treatments are shown in Table (2). Data showed that the studied nutrients (N, P, K, Fe, Mn and Zn) lay within the low-medium range, according to the critical levels of nutrients reported by **Lindsay and Norvell (1978)**. In general, this is true since soil is not only poor in the nutrient-bearing minerals, but also in organic matter content, which are considered as storehouse for the essential plant nutrients. On the other hand, data in Table (4) indicated that available concentrations of the studied macro- (N, P and K) and micronutrients (Fe, Mn and Zn) in the studied soil irrigated with the tested saline water were drastically severely affected by the excess salt content in soil but nutrients contents gradually increased with applied organic compost and biofertilizer. **Humax (2006)** pointed out that humic acid has a high complexation ability with ions in the environment due to the high carbon content (60 %) of both aliphatic and aromatic character and the richness in oxygen-containing functional groups such as carboxyl, phenolic, alcoholic and quinoid groups, which is beneficial for plant nutrition.

The relative increase in available nutrient concentrations may be attributed to the modified suitable air-moisture regime that control the availability of nutrients, in addition to the effect of applied organic compost in alleviating the depressive effect of salinity stress on released nutrients from either organic residues or nutrient bearing minerals. **Hegazi (1999)** found a negative correlation between salinity and available plant nutrients in soil. In addition, the suitable air-moisture regime in such sand soil positively affected biological activity and the supply of available nutrients, particularly from the organic source.

The integrated role of applied organic compost with bio-fertilizer could be also due to the released active organic acids during microbial activity that enhance the solubilization of nutrients from the native and added sources, also may be attributed to their slow release during the decomposition and mineralization processes as well as minimizing their possible lose by leaching throughout the studied relatively coarse texture soil (**Nader and Ewees, 2011**).0

Table (4): Effect of treatments on some soil properties and available nutrients concentrations.

Soil properties & nutrients status	Applied treatments									Statistical analysis (L.S.D. at 0.05)
	Control	Compost	Proline	BF	Comp + Proline	Comp+ BF	Porline + BF	Comp+BF + Proline	Mean	
Bulk density (g/cm ³)	1.33	1.26	1.32	1.29	1.25	1.22	1.28	1.21	1.27	0.01
Hydraulic conduct. (cm/hr)	0.44	1.14	0.45	0.65	1.17	1.56	0.67	1.58	0.96	0.06
Total porosity (%)	54.75	58.46	54.80	55.32	59.09	62.81	55.48	63.19	57.99	0.94
F.C. (%)	37.40	38.67	37.37	37.67	39.07	40.17	38.43	40.23	38.63	1.42
W.P. (%)	17.30	16.95	17.23	17.09	16.74	16.59	17.02	16.25	16.90	0.77
A.W. (%)	20.10	21.72	20.14	20.58	22.35	23.58	21.41	23.98	21.73	1.57
E _{Ce} (dS/m)	11.33	9.16	11.33	10.61	9.09	8.28	10.57	8.24	9.83	0.73
pH	7.87	7.63	7.86	7.79	7.62	7.51	7.77	7.49	7.69	0.10
OM%	0.86	2.05	0.87	1.26	2.12	2.36	1.30	2.39	1.65	0.06
ESP%	12.46	9.32	12.43	11.26	9.29	8.21	11.24	8.17	10.30	0.87
CEC (Meq/100g soil)	40.17	45.53	40.00	41.97	46.29	50.90	42.20	51.30	44.80	3.72
Available macro and micronutrients (mg kg⁻¹)										
N	118	165	114	133	170	193	135	196	153.62	7.16
P	4.5	11.7	4.60	6.80	11.80	13.80	6.90	13.90	9.25	0.89
K	152	187	153	165	190	214	168	217	180.50	6.49
Fe	4.32	10.94	4.33	6.53	11.11	11.58	6.65	6.78	7.78	0.66
Mn	0.92	2.05	0.93	1.30	2.14	3.09	1.34	3.18	1.87	0.1
Zn	1.46	1.84	1.48	1.58	1.87	2.13	1.61	2.16	1.77	0.07

F.C= Field capacity, W.P= Welting point, A.W= Available water, Comp=Compost and BF=Bio-fertilizer

On the Other hand, application of proline had a slightly affected on soil properties. These results are in accordance with those obtained by **Torello and Ricf (1986)** who mentioned that accumulation of proline was rapid in barley that adapted to applied salinity.

Data in Table (4) indicated that the superiority of combined effects of applied organic compost, bio-fertilizer and proline treatments for the noticeable reduction in the values of soil pH, E_{Ce} and ESP vs a pronounced increase in soil organic matter content, CEC and soil available nutrient concentrations and biological conditions that enhancing nutrients uptake by plants could be interpreted as follows:

i. Organic compost decomposition tends to accelerate in the presence of microbial media of bio-fertilizer, and in turn produces active organic and inorganic acids that may led to decrease soil pH as well chelate metals (Fe, Mn and Zn). These chelated metal cations are not sensitive to the restriction or the adverseable effects of alkaline side, consequently they are found as strategic storehouse in organo-metalic compounds that are more suitable for uptake by plant roots.

ii. The effective role of microbial activity to reduce soil salinity stress, particularly in combination with either organic or biofertilizer, could be interpreted according to many opinions outlined by Ashmayer *et al.*, (2008) reported that many strains produce several phytohormones (i.e., indole acetic acid and cytokinins) and organic acids. Such products reduce the deleterious effect of Na-salts, and simultaneously improve soil structure, i.e., increasing aggregate stability and drainable pores. Consequently, these created conductive pores enhance the leaching process of soluble salts through irrigation fractions.

III: Plant parameters as affected by treatments:

a. Plant growth characters, grain and straw yields:

Data presented in Table (5) indicate that the achieved favourable soil conditions due to the applied treatments, particularly the combination ones of compost with either bio-fertilizer (salinity durable bacteria) or foliated with proline, were positively reflected on the studied values of barley plants growth parameters (i.e. plant height, No of grains/ spike, and No spikes/m²), biological yield (grain and straw yields) and some parameters of grain quality (1000 grain weight) of barley plants grown in salt affected soil as compared to the applied solely ones.

It could be noticed from data in Table (5) that plots that received the combination of (Compost + Proline + Biofertilizer) resulted in higher growth parameters (plant height, number of grains/spike and number of spikes/m²) than the control and the previous materials with corresponding values of 102.40 cm for plant height, 46.00 grains /spike and 287 spikes /m². Increases in these characters due to the application of (Compost + Proline + Biofertilizer), the percentage of these values reached to 40.27, 53.33 and 32.87 % for plant height, number of grains/ spike and number of spikes/m² respectively, compared with that of the control. No significant differences were observed between (Compost + Proline + Biofertilizer) application and without proline supplement.

Data presented in Table (5) revealed that the, biological yield (grain and straw yields) and some parameters of grain quality (1000 grain weight) were substantially improved by the application of compost in combination with either (salinity durable bacteria) or foliar sprayed proline.

Results presented in Table (5) showed that grain, straw yields and 1000 grain weight were significantly increased by the application of different materials as solely or in combination, with no significant differences between OM + BF and (Compost + Proline + Biofertilizer). The highest yields of grain, straw and 1000 grain weight were associated with barley plants received (Compost + Proline + Biofertilizer) treatments, values were 2378.4 kg/fed, 5.63 ton/fed and 52.06 g, respectively. These values represented 156.73, 155.90 and 20.59% of that of the control, respectively. Either organic compost addition or biofertilizer with proline resulted in a significant increase on grain, straw yields and 1000 grain weight (Table 5). These results are also in line with those obtained by Nader and Ewees (2011) who stated that arbuscular mycorrhizal

(AM) fungi is capable to produce some hormones which induces the proliferation roots and root hair that increase nutrient absorbing surfaces as well as produce organic acids, which solublize inorganic and organic forms of mineral elements. **Wated *et al.*, (1983)** reported that proline amino acid plays an adaptive role in the tolerance of plant cells to salinity by increasing the concentration of cultural osmotic components in order to equalize the osmotic potential of the cytoplasm.

Table (5): Effect of treatments on growth parameters, grain and straw yields of barley grown on salt affected soil.

Growth parameters and yield	Applied treatments									Statistical analysis (LSD at 0.05)
	Control	Compost	Proline	Biofertilizer	Comp + Proline	Comp + BF	Porline + BF	Comp+BF+ Proline	Mean	
Plant height (cm)	73	101.20	97.20	100.80	101.39	102	101.30	102.40	97.41	3.82
No. of grains/spike	30	38	34	38	43	44	39	46	39.04	7.92
No. Spike/m ²	216	257	246	252	267	275	267	287	258.40	10.44
1000-grains weight (g)	43.17	48.52	46.22	48.40	48.81	50.35	49.05	52.06	48.32	3.35
Grain yield (kg/fed)	926.40	2059.20	1623.6	1707.6	2174.4	2347.2	1780.8	2378.4	1874.4	0.83
Straw yield (ton/fed)	2.20	4.93	3.87	4.06	5.16	5.58	4.25	5.63	4.46	0.50

Comp= compost and BF=Bio-fertilizer

b. Nutrient contents in barley grains:

Data of the studied macro-nutrients (N, P and K) and micronutrients (Fe, Mn and Zn) contents in barley grains are presented in Table (6). The obtained results exhibited pronounced concentrations increases for the studied macro and micronutrients due to the applied compost as a solely treatment, the greatest values were observed when it was combined with both proline and biofertilizer, followed by the combined treatments of (Compost+BioFertilizer) and (Compost+proline) as compared to the control treatment (untreated soil). Undoubtedly, the applied solely and some combined treatments were useful for releasing available nutrients, and in turn their contents in plant tissues. Such superior effect of organic compost in the combined treatments is more associated with the relatively high contents of both essential macro- and micro-nutrients (N, P, K, Fe, Mn and Zn), the released active organic acids that enhance more released micronutrients or their solubilization from both native and added sources.

In general, the improving effect of the combined treatments attained organic compost or byiofertilizer was commonly achieved may be due to lowering soil pH that improve nutrients availability, mobility and ability to uptake by plant roots. In addition, the superiority of applied treatments attained (Compost+ proline + Biofertilizer) were more attributed to their richness in organic substances that ameliorate soil-moisture regime and the biological soil condition. This beneficial effect could be explained by many aspects, *i.e.*,

increasing released macro- or micro-nutrient contents through the decomposition of applied compost, reduction of nutrient fixation and forming the stable complexes of micronutrients-humic substances supplied from such manures and keeping them in available forms for extended period (Ewees, 2012).

On the other hand, the significant response of nutrients contents in barley grain to biofertilizer and soil application of compost may be due to increased root growth that enable the grown plants to absorb more nutrients. Kloepper (2003) pointed out that phytohormones producer bacteria causes pronounced increases for plant root elongation by then uptake of more nutrients via the root system, and hence utilization of N as a result of bio-inoculation. Nader and Ewees (2011) reported that biofertilizer increase uptake of N, P, K, Fe, Zn, and Mn by plants.

It could be concluded that, the combined treatment of (Compost + proline +Biofertilizer) exhibited asuperior effect due to improving soil physico-chemical properties that positively affect nutrients availability as well as maintaining a suitable soil moisture regime. It is noteworthy to mention that the nutrient contents in plant tissues were, in general, extending parallel close to the corresponding available nutrient contents in the studied soil, as shown in Tables (4).

Table (6): Effect of treatments on nutrient contents of Barley grown on salt affected soil.

applied treatments	Grain content of macro and micro nutrients					
	Macronutrients (mg kg-1)			Micronutrients (mg kg-1)		
	N	K	P	Fe	Mn	Zn
Cotrol	1.63	1.11	0.41	142	57.80	47.50
Compost	1.91	1.31	0.50	171	71.00	63.00
Proline	1.73	1.21	0.42	159	62.40	50.90
Biofertilizer	1.82	1.28	0.48	163	66.70	54.60
Comp + Proline	1.85	1.33	0.51	189	77.70	70.00
Comp + BF	2.10	1.38	0.64	210	88.00	82.00
Proline + BF	1.76	1.30	0.49	175	70.00	60.00
Comp + Proline + BF	2.16	1.44	0.68	216	93.00	87.00
Mean	1.87	1.30	0.52			
L.S.D, at (0.05)	0.06	0.05	0.06	0.09	0.56	0.62

Comp= compost and BF=Bio-fertilizer

C. Crude protein and carbohydrates in barley grains:

Data in Table (7) showed markedly positive and significant effects due to the application of both combined treatments of (Compost + proline + Biofertilizer), (Compost + Biofertilizer) and (Compost). Such effect was achieved upon the significance of L.S.D. values at 0.05.

Relative to the control, the single treatments Compost, proline and Biofertilizer resulted in 12.51, 15.25 and 24.36% increases in crude protein (%) percentage, and gave 10.28, 0.21 and 2.50% carbohydrate content (%), respectively (Table 7). Relative to control, combination treatments Compost +

INFLUENCE OF ORGANIC COMPOST, PROLINE..... 79

proline+ Biofertilizer, Compost + Biofertilizer, Compost + proline and proline + Biofertilizer caused increases of 19.53, 33.47 and 15.03% for crude protein (%) and 5.46, 4.24 and 3.30% for carbohydrate content (%), respectively .

Table (7): Effect of applied materials on Crude Protein (%), Carbohydrate content in Barley plants grown on salt affected soil.

Applied treatments	Carbohydrate content (%)	Crude Protein (%)
Cotrol	13.90	9.11
Compost	15.33	10.25
Proline	13.93	10.50
Biofertilizer	14.25	11.33
Comp + Proline	14.49	10.89
Comp + BF	14.66	12.16
Proline+ BF	13.44	10.48
Comp + Proline + BF	15.29	13.13
Mean	14.41	10.98
L.S.D, at (0.05)	0.73	0.78

Comp= compost and BF = Bio-fertilizer

Results of the present work emphasized the possibility of alleviating the harmful effects of high soil salinity on barley plants growth, yield, grain quality and absorption of nutrients by the application of compost, proline amino acid and inoculation with salinity durable bacteria solely or in combination.

REFERENCES

- Ashmaye, S.H.; Shaban, Kh.A. and Abd El-Hader, M.G. (2008). Effect of mineral nitrogen, sulphur, organic and bio-fertilizers on maize productivity in saline soil of Sahl El-Tina. *Minufiya, J. Agric. Res.*, 33 (1): 195-209.
- Ayers, R.S. and Westcot D.W. (1985). Water quality for agriculture, irrigation and drainage. Paper No. 29, FAO, Rome, Italy.
- Barakat, M.N. and Abdel-Latif, T.H. (1995). In vitro selection for salt-tolerant lines in wheat. II. In vitro characterization of cell lines and plant regeneration. *Alex. J. of Agric. Res.* 40: 139-165.
- Black, G.R., Evans, D.D., Ensminger, L.E., White, J.L. and Chark, F.E. (1965). "Methods of soil analysis" Agron. Series , AM. Agron., Madison, Wisconsin, USA.
- Chapman, H. D. and Pratt, P. F. (1961). Methods of Analysis for Soils, plant and water. Univ. California, Division of Agric. Sci., Berkeley, CA 94720.
- El-Leboudi, A.E., Gawish, Sh.M., Abdel-Aziz, S.M. and Ahmed, M.R.M. (1997). Some metabolic aspects in wheat plants subjected to salinity. *Annals of Agric. Sci. Cairo.*42: 585-596.

- Ewees, M. S. A (2012). The Possile Response of Tomato Growth and Fruit Quality For Applxing a water soaking solution of poulty wastes through irrigation water. *J. Soil Sci.* 52, No. 1, pp: 119-139.
- Hegazi, I.M.A. (1999). Factors affecting utilization and development in lands and water resources of the northern area of El Fayoum. Ph. D. Thesis, Fac. of Agric. at El Fayoum, Cairo Univ., Egypt.
- Humax (2006). Role humic acid increasing plant growth. [htt://www. jhbiotech.com/info/humax info.htm](http://www.jhbiotech.com/info/humax%20info.htm).
- Jackson, M. L. (1973). "Soil Chemical Analysis", Constable Co., Ltd., London.pp. 521.
- Khaled, A.S.; Mona, G.A. and Seham, M.E. (2011).Evaluation of organic farm and compost combined with urea fertilizers on fertility and maize productivity in newly reclaimed. *Research J. of Agric, and Biol. Sci.*, 6(5): 388 – 397.
- Kloepper, J.W. (2003). A review of mechanisms for plant growth promotion by PGPR. 6th international PGPR workshop. 65-10 October 2003, Calcutta, India.
- Lindsay, W.L. and Norvell, W.A. (1978). Development of DTPA soil test for Zn, Fe, Mn and Cu. *Soil Sci. Soc. Am. J.*, 42, pp. 421.
- Nader, R. H and Ewess, m.s.a (2011). Improving productivity of Zucchini Squash Grown Under Moderately saline soil Msing Gypsum, organo-stimulants and A M-Fungi, *Journal of applied Sci. Res.*, 7 (12): 2112-2126.
- Ouerghi, Z., Zid, E. and Ayadi, A. (1991). Sensitivity to NaCl and exclusion of Na⁺ in sunflower (*Helianthus annuus*). *Agric. Mediterranea.* 121:110-114.
- Page, A.I.; Miller, R.H. and Keeney, D.R. (Eds.) (1982). *Methods of Soil Analysis. Part 2: Chemical and Microbiological Properties.* 2nd Edition, Amer. Soc. of Agron., Madison, Wisconsin, U.S.A.
- Parida. A. K and A. B. Das (2005). Salt tolerance and salinity effects on plants. *Ecotox. Envirion. Saf.* 60, 324-349.
- Piper, C. S. (1950). "Soil and Plant Analysis" A monograph from the wails agric. research Inst., University of Adetaide Australia.
- Richards L.A.(1954). *Diagnosis and Improvement of Saline and Alkali soils.* United. States Department of Agriculture. Handbook No. 60. Gov. print off.
- Silini, A.; Silini-Cherif, H. and Ghoul, M. (2012). Effect of *Azotobactervinelandii* and compatible solutes on germination wheat seeds and root concentrations of sodium and potassium under salt stress. *Pak. J. of Biol. Sci.* 15(3): 132-140.
- Snedecore, G. W. and W. G. Cochran (1980). *Statistical methods.* 7th ed. Iowa State Univ., Press, Amer., Iowa, U.S.A.

- Sunjeong P.; Chunxue C. and Brain B. M.G. (2010).** Inoculants and soil amendments for organic growers. *Agriculturals and Natural resources.* 10- 17.
- Soltanpour, P.N. and A.B. Schwab (1977).** A new soil test for simultaneous extraction of macronutrients in alkaline soils. *Comm. Soil Sc. and Plant Annal.*, 8: 195.
- Tester, M. and Davenport, R. (2003).** Na tolerance and Na transport in higher plants. *Annals of Botany*, 91: 503-527.
- Tipiramaz, R. and Cakirlar, H. (1990).** Effect of water and salinity stress on relative water content, proline and betaine composition in two varieties of wheat plants cultivated in Turkey, *Doga. Bigoloja. Serisi.* 14: 124 -148 .
- Torello, W.A. and Rief, L.A. (1986).** Effects of NaCl stress on proline and cation accumulation in salt sensitive and tolerant turfgrass. *Plant and Soil.* 93:241-247.
- Wageeh, M. G. M. (1994).** Effect of some treatments on salt tolerance of plants. Ph.D. Thesis, Fac. of Agric. Moshtohor. Zagazig. Univ.
- Wareing, P.E. and Phillips, I.D.J. (1978).** The control of growth and differentiation in plant. Second Edition, Chap. 3 and 4, Great Britain, by William Clowes and Sons Limited, London.
- Wated, A., Reinhard, E.L. and Lerner, H.R. (1983).** Comparison between a stable NaCl selected *Nicotiana* cell line and the wild type, Na, K and proline pools as a function of salinity. *Plant physiol.* 73: 624-629.
- Yurekli, F., Topcuoglu, S.F. and Bozcuk, S. (1996).** Effects of salt stress on proline accumulation in sunflower leaves in relation to salt concentration and duration of stress. *Turkish. J. of Biology.* 20: 163-169.
- Zou, N.; Dart, P.J. and Marcar, N.E. (1995).** Interaction of salinity and rhizobial strain on growth and N₂-fixation by *Acacia ampliceps*. *Soil Biol. Biochem.* 27 (415): 409-413.
- Zidan, M.A. and Malibari, A.A (1993).** The role of K in alleviating stress affecting growth and organic and mineral components of wheat. *Arab Gulf. of Sci. Res.* 11:201-208.

تأثير استخدام الكميوست والبرولين والسماذ الحيوي (البكتريا المقاومة للملوحة) على نمو محصول الشعير وامتصاصه للعناصر الغذائية تحت ظروف الملوحة العالية

ياسر محمود عباس* - محمد صابر علي عويس**

*معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية - الجيزة - مصر.

**قسم الأراضي والمياه - كلية زراعة - جامعة الفيوم.

اجريت تجربة حقلية على تربة متأثرة بالملوحة (التوصيل الكهربائي لمستخلص عجينة التربة المشبعة = 11,33 ديسمينز/م) في قرية قصر الباسل جنوب منطقة اطسا . محافظة الفيوم. مصر. خلال الموسم الشتوي ٢٠١٣/٢٠١٤ بهدف تحديد التأثيرات الإيجابية لأضافه محسن تربة محلي (كميوست بمعدل ٢٠م^٢/فدان)، حمض اميني (برولين رش بمعدل ٣ مجم/ لتر عند ٦٠، ٤٥، ٢٠ يوم من الزراعة) وسماذ حيوي (بكتريا متحملة الملوحة) اما منفردة او بالجمع بينها على نمو وقياسات الشعير (صنف جيزة ١٢٣). رويت اراض التجربة بمياه مالحة من بحر الغرق (خليط من مياه النيل ومياه الصرف الزراعي) وهي ذات درجة C2S1 طبقا للتقسيم الأمريكي القديم لنوعية مياه الري (التوصيل الكهربائي = ١.٦٦ ديسيمتر/متر، النسبة الامصاصية للصدويوم SAR = ٤.٢٥) وقد اخذ في الاعتبار في هذه الدراسة التحسين المصاحب للمعاملات على خواص التربة (مثل التوصيل الكهربائي لمستخلص عجينة التربة المشبعة، رقم الحموضة، النسبة المئوية للصدويوم المتبادل، ومحتوى التربة من العناصر المغذية الكبرى والصغرى الميسرة للنبات).

تشير النتائج الى ان قيم التوصيل الكهربائي والنسبة المئوية للصدويوم المتبادل ورقم الحموضة قد تناقصت بإضافة الكميوست والبرولين والسماذ الحيوي وتختلف قيمة هذا النقص من معاملة لأخرى وقد تبين أن افضل معاملة كانت (الكميوست + برولين + السماذ الحيوي). حيث كان تأثيرها واضحا على نسبة المادة العضوية والسعة التبادلية والكاتيونية للتربة بالزيادة حيث ادى الجمع بينها الى الحصول على اعلى قيمة للمادة العضوية والسعة التبادلية الكاتيونية كما أدت اضافة الكميوست + برولين + سماذ حيوي الي تقليل قيمة الكثافة الظاهرية للتربة، وزيادة قيم التوصيل الهيدروليكي، والمسامية الكلية ومحتوى التربة من الماء الميسر كذلك أدت هذه المعاملة الي تأثير ايجابي على كل من طول النبات، عدد الحبوب في السنبل، عدد السنابل/ م^٢، وزن المائة حبة، محصول الحبوب والقش وكذلك على محتوى النبات من العناصر المغذية الكبرى والصغرى . وكانت المعاملات افضل بإضافة المحسنات الثلاثة مجتمعة (كميوست+ برولين+ سماذ حيوي) وذلك مقارنة بمعاملة الكنترول او الإضافات الأخرى منفردة.

وعلى أساس نتائج هذه الدراسة فإنه يمكن التوصية باستخدام الكميوست، برولين، السماذ الحيوي (البكتريا المتحملة للملوحة) لتخفيف الأثر الضار لكل من ملوحة التربة وماء الري والتي تؤثر سلباً على نوعية وكمية محصول الشعير وامتصاص العناصر الغذائية.