



**Safety and Quality of Ready to Eat Chicken Meat products**

Taghreed, H. Abbas<sup>(1)</sup>, Tolba, K.S<sup>\*</sup>, Ibrahim, A.M.<sup>\*\*</sup>; and Elmoosalami, M.K. <sup>\*\*</sup>

<sup>\*</sup> Meat Hygiene and Control Department, Faculty of Veterinary Medicine, Cairo University

<sup>\*\*</sup> Food Hygiene Department, Animal Health Research Institute, Dokki-Giza

(1) Corresponding Author: [taghreed7amdy@gmail.com](mailto:taghreed7amdy@gmail.com)

**ABSTRACT**

Contamination of ready-to-eat (RTE) chicken products with food borne pathogens remains an important public health issue, because it can lead to illness if there are mal practices during slaughtering, eviscerating, washing, handling and subsequently preparation by cooking. Therefore, this study focused on the safety of some ready to eat chicken products during the preparation in terms of sensory attributes, chemical parameters for detection of freshness and bacterial load. The samples were collected from different restaurants in Cairo and Giza Governorate. The research study focused on ten specific products including chicken shawarma, grilled chicken, chicken pane, shish tawoak, chicken fajita, chicken burger, chicken salad, chicken mandy, fried chicken and finally pop chicken. The bacteriological criteria used for integrity of the tested products were as follows: Aerobic Plate Count (APC), Enterobacteriaceae, coliforms and Staph. aureus counts. The present study indicated that the examined chicken salad was the most contaminated products in terms of APC, Enterobacteriaceae, coliforms and Staph. aureus counts ( $1.094 \times 10^6 \pm 8.81 \times 10^5$ ,  $1.696 \times 10^2 \pm 4.82 \times 10^1$ ,  $1.183 \times 10^4 \pm 4.65 \times 10^3$  and  $30.0 \pm 21.34$  cfu/g) respectively. On the contrary, the lowest bacterial load of the aforementioned bacterial counts was recorded for fried chicken ( $5.78 \times 10^4 \pm 3.1 \times 10^4$ ,  $< 3$ ,  $3.1 \times 10^2 \pm 43.33$  and  $< 10$  cfu/g), respectively followed by pop chicken products which recorded ( $1.797 \times 10^4 \pm 9.34 \times 10^3$ ,  $< 3$ ,  $2.20 \times 10^2 \pm 72.7$  and  $< 10$  cfu/g), respectively. From these results, it was cleared that five chicken products (Salad, pane, shish tawoak, fajita and chicken burger) had higher mean values of bacteriological analysis, but still within the accepted range concerning APC as defined by Public Health Laboratory Service (PHLS) (2000) regarding judgment of RTE meals. Chemical characters were within the permissible limits. Moreover, sensory attributes and the reflection of microbial counts on the consumer and public health were discussed.

**(Keywords:** RTE chicken meals, PHLS, APC, Coliforms, Enterobacteriaceae)

**Introduction**

Ready-to-eat food products provide a source of readily available and nutrition meals for the consumers. However, questions have been raised about the safety and microbiological quality of these food products. In this respect, 12% of children (aged 2-16 years), 16% of adults (aged 17-69 years) and 13% of people aged 70 and above reported consumption of cooked chicken meat in the 1995 National Nutrition Survey (McLennan and Podger, 1999). Australian National Children's Nutrition and Physical Activity Survey 2007 reported that 33% of children (aged 2-16 years) reported consumption of cooked chicken meat (DOHA, 2008). It has been agreed that the product quality has a multidimensional nature including performance, durability, reliability, conformity, consistency, etc. (Munoz et al., 1992 and Lawless, 1995). After appearance and tenderness characteristics, flavour is considered the most important characteristic feature of meat quality perceived by the consumer (Love, 1994 and Sow and Grongnet, 2010). The importance of sensory evaluation for product development and for quality control has been reported extensively. However, the average acceptability scores of

consumer responses have not been sufficiently meaningful for companies to understand the level of the product's sensory quality despite using different scales such as the 5-point hedonic scale, 7 or 9-point hedonic scale, and so on. (Imm et al., 2009; Etaio et al., 2010 and Imm et al., 2010). Susceptibility of chicken meat and chicken-based meat products to microbial spoilage, presents a potential health hazard, since poultry meat may harbor pathogenic microorganisms (Geornaras et al., 1998). Spoilage is commonly detected by sensory and/or microbiological analysis (Dainty, 1996). Poultry and poultry products have become a common food for humans in developing countries and they are often sold in street restaurants. Ready-to-eat cooked meat products are recognized to be contaminated during slicing which has been associated with several outbreaks (Perez et al., 2010). As any perishable meat, fish or poultry, bacteria can be found on raw or undercooked chicken. They multiply rapidly at temperatures between 4.4 °C and 60 °C out of refrigeration and before thorough cooking, occur. Freezing does not kill bacteria but only thorough cooking destroys them. USDA's Food Safety and Inspection Service (FSIS) (2012) has a zero tolerance for

certain pathogens, including Salmonella and L. monocytogenes, in cooked and ready-to-eat products, such as chicken franks or lunch meat, that can be eaten without further cooking. Most food-borne illness outbreaks are a result of improper handling or contamination when meals are prepared. Sanitary food handling and proper cooking and refrigeration should prevent food-borne illnesses. Aerobic Plate Count and Enterobacteriaceae counts are considered as indication of bacteriological quality, which give an idea about the hygienic measures adopted during further processing and help in assuring the keeping quality of further processed chicken meat products (Aberle et al., 2001 and Mohamed et al., 2015). Detection of freshness is very important through conducting required chemical analysis of cooked and uncooked processed chicken meat products to ensure compliance of such products with national or international standard legal requirements (Beckers, 1998 and Ibrahim, et al., 2014). Staph. aureus are carried on human hands, in nasal passages, or in throats. The bacteria are

found in foods made by hand and then improperly refrigerated, such as chicken salad. Staphylococcus aureus plays a great role in bacterial contamination of fast and RTE foods, because workers during preparation and processing may touch fast foods, which are usually eaten without sufficient cooking or heating (Soliman, 1988). Staphylococcus aureus have been implicated in cases of severe diarrhea as well as the main cause of food poisoning gastroenteritis among consumers (Davies and Board, 1998 and Eid et al., 2014). Mulla (2002) established that the increase in thiobarbituric acid (TBA) value, resulted in the presence of detectable unaccepted flavor and lower degree of acceptability of poultry processed products. Fat oxidation was due to prolonged storage or due to the use of low quality meat in the processing of such products. Therefore, the present work planned out to assure the sensory, chemical and bacteriological quality of some RTE chicken products randomly collected from different supermarkets in Cairo and Giza Governorates.

#### Material and Methods

A grand total of 100 random samples of RTE chicken products including chicken shawarma, grilled chicken, chicken pane, shish tawoak, chicken fajita, chicken burger, chicken salad, chicken mandy, fried chicken and finally pop chicken (ten samples each) were collected from different restaurants in Cairo and Giza Governorates and aseptically transferred in its original containers without delay to the laboratory and subjected to the following examinations:

##### I. Analysis of sensory attributes:

Collected samples were examined in terms of their appearance, color, taste, flavor, consistency and juiciness using 9 points hedonic scale as (1) was dislike extremely and (9) was liked extremely. Panel team formed from 1-12 members (Food Hygiene Department, Animal Health Research Institute) with experience with RTE chicken products were examined the RTE chicken products by using 9 point hedonic scale according to Anna. 1998.

##### II. Bacteriological examination:

Culture media used in this study was prepared, produced and measured its performance to define its efficiency in carrying bacteriological examination according to ISO/TS 11133-2014.

**Sample preparation:** according to ISO 6887-2:2003:

Twenty-five grams of the examined samples were removed by sterile scissors and forceps and stomached using stomacher (Seward stomacher 80 Biomaster, Serial No. 46464. England) with

225ml of sterile buffered peptone water (0.1%) to give a homogenate of 1/10 dilution. One ml from the original dilution was transferred with sterile pipette to another sterile test tube containing 9 ml of sterile peptone water 0.1% and mixed well to make the next dilution from which further decimal dilutions were prepared. The prepared dilutions were subjected to the following examinations:

1-Aerobic Plate Count (APC) according to APHA, (2001), on APC agar at 35°C for 48±2 hrs.

2-Total Coliform count (as a Most Probable Number (MPN) according to FDA, (2002), on Lauryl Sulphate Tetrathionate (LST) broth at 35°C for 24-48hrs.

3-Enterobacteriaceae count according to ISO 21528-2 :( 2004), on Violet Red Bile Glucose Agar (VRBGA) at 37°C for 24±2hrs.

4-Staphylococcus aureus count according to FDA, (2001), on Baird Parker media at 35°C for 24-48 hrs.

##### III. Chemical analysis for detection of freshness

1-Determination of Hydrogen Ion Concentration (pH) according to E.O.S 63/11 (2006) by using pH meter,

2-Determination of Total Volatile Basic Nitrogen (TVB-N) according to E.O.S 63/10 (2006) by distillation method.

3-Determination of TBA value of lipid oxidation according to E.O.S 63/9 (2006) by distillation method.

**IV. Statistical analysis:**

Results were recorded as mean values of 3 replicates for each analysis ±SD/SE. on way


ANOVA were used for the collected data by SPSS statistics 17.0 for Windows. Comparison among different products was performed by LSD and significance was defined at P<0.05.

**Results**

**Table (1)** Overall acceptability of sensory attributes of examined RTE chicken products (n=10).

Products	Sensory attributes						Overall acceptability
	Appearance	Color	Taste	Flavor	Consistency	Juiciness	
Shawerma	4.2±0.33	4.3±0.34	3.8±0.20	3.8±0.25	4.0±0.26	3.6±0.34	3.95±0.29 <sup>A</sup>
Grilled chicken	4.9±0.23	4.7±0.21	4.9±0.18	4.5±0.17	4.7±0.15	4.4±0.16	4.68±0.18 <sup>AB</sup>
Chicken Pane	4.2±0.47	4.4±0.52	3.8±0.44	3.6±0.43	4.1±0.35	3.7±0.65	4.0±0.48 <sup>BC</sup>
Shish tawoak	5.4±0.27	5.1±0.23	5.1±0.23	4.9±0.18	4.7±0.15	5.1±0.23	5.1±0.22 <sup>acD</sup>
Fajita	4.9±0.43	4.8±0.25	5.1±0.32	4.8±0.20	4.8±0.20	5.1±0.32	4.92±0.29 <sup>ac</sup>
Chicken Burger	4.6±0.37	4.8±0.33	4.6±0.22	4.4±0.16	4.7±0.26	4.7±0.26	4.63±0.27 <sup>ac</sup>
Chicken Salad	4.8±0.13	4.8±0.13	5.0±0.15	4.8±0.20	4.8±0.20	5.2±0.13	4.90±0.16 <sup>ac</sup>
Chicken Mandy	4.7±0.15	4.6±0.16	4.3±0.15	4.3±0.15	4.5±0.17	4.5±0.17	4.48±0.16
Fried chicken	4.9±0.18	4.9±0.18	4.5±0.22	4.4±0.22	4.5±0.22	4.3±0.15	4.58±0.20 <sup>ac</sup>
Pop chicken	4.5±0.17	4.6±0.16	4.5±0.17	4.0±0.00	4.4±0.16	4.5±0.17	4.42±0.14 <sup>d</sup>

There are significant differences (P<0.05) between means having the same capital and small letters in the same column. Chicken mandy is the only product that did not show any significant difference with other products under study.



**Fig. (1)** Mean values of overall sensory attribute of examined RTE chicken products.

**Table (2)** Mean statistical value of different bacterial counts in RTE chicken meat products.

product	Bacterial counts (Mean±SE)			
	APC (cfu/g)	Coliforms (MPN/g)	Enterobacteriaceae (cfu/g)	Staph. aureus (cfu/g)
Shawerma	5.21x10 <sup>4</sup> ±2.11x10 <sup>4</sup> <sup>A</sup>	15.24±9.60	8.22x10 <sup>2</sup> ±2.73x10 <sup>2</sup> <sup>A</sup>	<10
Grilled chicken	5.07x10 <sup>4</sup> ±3.9x10 <sup>4</sup> <sup>B</sup>	48.66±45.76	5.56x10 <sup>2</sup> ±4.39x10 <sup>2</sup> <sup>B</sup>	30.00±21.34
Fried chicken	5.78x10 <sup>4</sup> ±3.1x10 <sup>4</sup> <sup>C</sup>	<3	3.1x10 <sup>2</sup> ±43.33 <sup>C</sup>	<10
Chicken Pane	4.244x10 <sup>5</sup> ±3.76x10 <sup>5</sup> <sup>D</sup>	39.26±19.84	1.84x10 <sup>3</sup> ±8.08x10 <sup>2</sup> <sup>BD</sup>	30.0±21.34
Fajita	2.22x10 <sup>5</sup> ±8.4x10 <sup>4</sup> <sup>bcE</sup>	31.90±21.1	3.10x10 <sup>2</sup> ±54.7 <sup>E</sup>	90.0±64
Shish tawoak	2.67x10 <sup>5</sup> ±1.41x10 <sup>5</sup> <sup>bcF</sup>	1.48x10 <sup>2</sup> ±1.08x10 <sup>2</sup>	9.92x10 <sup>2</sup> ±3.25x10 <sup>2</sup> <sup>F</sup>	1.1x10 <sup>2</sup> ±99.39
Chicken Burger	1.49x10 <sup>5</sup> ±8.0x10 <sup>4</sup> <sup>bcG</sup>	1.10±0.79	1.468x10 <sup>3</sup> ±8.64x10 <sup>2</sup> <sup>G</sup>	30±21.34
Chicken Salad	1.094x10 <sup>6</sup> ±8.81x10 <sup>5</sup> <sup>abcdH</sup>	1.696x10 <sup>2</sup> ±4.82x10	1.183x10 <sup>4</sup> ±4.65x10 <sup>3</sup> <sup>abcdefgh</sup>	30.0±21.3
Chicken Mandy	2.11x10 <sup>4</sup> ±8.18x10 <sup>3</sup> <sup>efgh</sup>	1.64±.96	46.0±19.56 <sup>abcdefgh</sup>	60.0±49.89
Pop chicken	1.797x10 <sup>4</sup> ±9.34x10 <sup>3</sup> <sup>efgh</sup>	<3	2.20x10 <sup>2</sup> ±73 <sup>acdefgh</sup>	<10

There are significant differences (P<0.05) between means having the same capital and small letters in the same column.




Fig. (2) Mean of different bacterial counts in examined RTE chicken meat product samples.

Table (3): Mean values of chemical analysis for deterioration of RTE chicken meat products.

Products	Chemical analysis for product freshness		
	pH	TBA	TVB-N
Shawerma	5.33±0.135	0.50±0.077	13.19±0.184
Grilled chicken	5.70±0.133	0.53±0.021	14.70±0.369
Chicken Pane	5.81±0.131	0.57±0.040	13.05±0.717
Shish tawoak	5.79±0.048	0.40±0.026	13.31±0.440
Fajita	5.40±0.087	0.49±0.031	13.61±0.788
Chicken Burger	6.31±0.064	0.57±0.033	13.70±0.535
Salad	5.91±0.118	0.58±0.044	14.57±0.620
Chicken mandy	6.13±0.075	0.48±0.025	12.84±0.156
Fried chicken	5.86±0.184	0.42±0.047	13.14±0.492
Pop chicken	4.95±0.025	0.32±0.025	13.32±0.289




Fig. (3) Chemical analysis for deterioration associated with detection of freshness of examined RTE chicken meat products.

From the results reported in **table (1) and fig. (1)**, the overall acceptability of sensory attributes of RTE chicken meat products including shawerma, grilled chicken, pane, shish tawoak, fajita, burger, chicken salad, mandy, fried chicken and pop chicken were  $3.95 \pm 0.29$ ,  $4.68 \pm 0.18$ ,  $4.0 \pm 0.48$ ,  $5.1 \pm 0.22$ ,  $4.92 \pm 0.29$ ,  $4.63 \pm 0.27$ ,  $4.90 \pm 0.16$ ,  $4.48 \pm 0.16$ ,  $4.58 \pm 0.20$  and  $4.42 \pm 0.14$ , respectively. The statistical analysis of the obtained results regarding the overall acceptability sensory attributes of shawerma meals in comparison with other different RTE chicken meat products revealed that there were a significant differences ( $P < 0.05$ ) between examined shawerma and each of grilled chickens, chicken fajita, chicken burger, chicken salad, shish tawoak and fried chicken. On the other hand, there were no significant difference ( $P > 0.05$ ) between shawerma samples and each of chicken pane, chicken mandy and pop chicken. In this respect, **Ibrahim, et al., (2014)** mentioned higher results than that in the present investigation where mean values of organoleptic scores of examined chicken shawerma samples were ranged from 6.3 and 7.45 with overall acceptability of 7.05 using 9-points hedonic scores. The overall acceptability of examined grilled chicken meat products recorded  $4.68 \pm 0.18$ , meaning that the sensory evaluation was laid between borderline and good area but they tended to shift towards the good. Moreover, the obtained results revealed that there were a significant differences in sensory attributes between examined RTE grilled chicken and both of chicken shawerma and chicken pane ( $P < 0.05$ ), while the data analysis showed no significant difference ( $P > 0.05$ ) between grilled chicken and rest of the products under study. Nearly similar result regarding overall acceptability (6.4) for grilled chicken was reported by **Ibrahim et al. (2014)** which also located in the good area when evaluating the rate of sensory scores. In this respect, **Sow and Grongnet (2010)** stated that RTC broiler was the least preferred one as compared with other treatments. Moreover, table (1) and fig. (1), revealed the overall acceptability of examined chicken pane samples were  $4.0 \pm 0.48$ , it could be concluded that chicken pane samples were more aligned to the border lined area. Otherwise, pane samples were better than shawerma in terms of sensory properties. There were a significant difference ( $P < 0.05$ ) between RTE chicken pane and other products under study except for samples of chicken shawerma, chicken mandy and pop chicken ( $P > 0.05$ ). High significant difference was recorded with RTE

## Discussion

chicken shish tawoak ( $P < 0.001$ ). These results were supported by **Ibrahim, et al. (2014)** as the overall acceptance of organoleptic score was 7.9 that it has been evaluated by the ten panelists. The same table declared the mean  $\pm$  SE of overall acceptance of sensory parameters of examined RTE chicken shish tawoak was  $5.1 \pm 0.22$ , meaning that shish tawoak had good sensory attributes. The results showed that there was a highly significant difference ( $P < 0.001$ ) between examined RTE shish tawoak and each of chicken shawerma and chicken pane samples, while the difference was significant ( $P < 0.05$ ) between shish tawoak and pop chicken. On the contrary, absence of significant differences ( $P > 0.05$ ) between chicken shish tawoak and other products under study. RTE chicken shish tawoak is a traditional marinated chicken shish kebab of Middle Eastern cuisine (**Virginia, 2010 and Ozan, 2013**). It is widely eaten in Turkey, Lebanon and Egyptas well as many cities around the world (**Yvonne, 2002 and Lisa and Lisa 2009**). The results in table and fig. (1) illustrated the mean overall acceptability of examined chicken fajita meals which recorded a score of  $4.92 \pm 0.29$  that more shifted towards the good area. Otherwise, fajita meals were nearly similar to grilled and shish tawoak meals, while they were better in sensory parameters as compared with chicken shawerma and pane meals. The statistical analysis declared that there were significant differences ( $P < 0.05$ ) between examined samples of RTE chicken fajita and each of chicken shawerma and chicken pane. On the other hand, there were no significant differences between chicken fajita and the other products under investigation. In this respect, **Chuaynukool, et al. (2007)** and **Jaturasitha et al., (2002)** concluded that toughness and firmness of chicken products may be attributed to the age and breed of chickens at slaughter, as the lower the age, increased the proportion of muscle collagen which earns a good appearance, texture and consistency of the product. Regarding the overall acceptability of chicken burger sensory attributes, it was found to be  $4.63 \pm 0.27$ . The results declared that chicken burger meals were located between poor and good area but more shifted towards the good area. Otherwise, chicken burger meals looked like grilled, shish tawoak and fajita meals, while better than shawerma and pane meals. The obtained results assured that there were significant differences ( $P < 0.05$ ) between examined samples of RTE chicken burger and both of chicken shawerma and chicken pane. On the other hand, the results proved the absence of

significant differences ( $P>0.05$ ) between burger and the rest of the products under study. Differences in sensory character between the chicken samples may be as a result of many factors including differences in lipid oxidation "Tocopherol and fatty acid profiles". In general the extent of lipid oxidation was higher in burgers than other products (Lawlor et al., 2003). During heat denaturation, insoluble cross-linked collagen shrank and effectively compressed heat-denatured myofibrils, and eventually resulting in moisture loss, decreases in fiber diameter and a tougher texture; Differences in the microstructures and cross-linked collagen content that exist between breeds might account for the varied textural appearance of meat (Wattanachant et al., 2004). The obtained results regarding RTE chicken salad, the overall acceptability of sensory parameters registered  $4.90\pm 0.16$ . Chicken salad meals were located between border line and good area but more shifted towards the good area. Otherwise, chicken salad meals were nearly similar to grilled, shish tawoak, fajita and chicken burger meals, while they were better in sensory attributes than chicken shawerma and pane meals. The obtained results confirmed that there were significant differences ( $P<0.05$ ) between examined samples of RTE chicken salad and both of chicken shawerma and chicken pane. On the other hand, the results proved the absence of significant differences ( $P>0.05$ ) between chicken salad and the rest of the products under study. In this respect, Esperance (2016) concluded that when chicken salad has a strong, unpleasant odor, this means that the salad has gone bad, the objectionable smell is caused by the chicken absorbing off-flavors of the other salad ingredients such as lettuce, onions, tomatoes or apples. Table and Fig. (1) also stated that overall acceptance of sensory parameters of examined RTE chicken mandy assigned to be  $4.48\pm 0.16$ , the results declared that chicken mandy meals were located in the border line area. Otherwise, chicken mandy meals were nearly similar to shawerma, and chicken pane meals, while they had lower sensory attributes than grilled chicken, shish tawoak, fajita, salad and chicken burger meals. The obtained results specify that there were no significant differences ( $P>0.05$ ) between examined RTE chicken mandy samples and the rest of the products in this study. Nearly similar results were reported by Lawlor, et al. (2003) who carried out One-way ANOVA on the sensory scores, the results showed significant ( $P<0.05$ ) differences between roast chicken samples and other chicken products for appearance, flavor,

odor and texture attributes. The authors clarified that the sensory panel was easier to distinguish between chicken samples using appearance and texture rather than odor and flavor attributes. Table (1) and Fig. (1) described also the overall acceptance of sensory parameters of examined RTE fried chicken meals were  $4.58\pm 0.20$ . The results declared that fried meals were located between border line and good area but more shifted towards good area. Otherwise, fried chicken meals were nearly similar to grilled, shish tawoak, fajita, salad and chicken burger meals, while they were better in sensory parameters than chicken shawerma, mandy and chicken pane meals. The results obtained confirmed the existence of significant differences ( $P<0.05$ ) between RTE fried chicken samples and each of chicken shawerma and chicken pane, while there were no significant differences with other examined RTE chicken product in this study. Consumers usually evaluate the fried product as acceptable or not first by its color. Krokida, et al. (2001) stated that oil temperature and sample thickness are the process parameters that affect the color significantly during frying. The difference in texture scores could be due to differences in frying time (Altunakar et al., 2004), where frying in oil with higher degree of hydrogenation resulted in products of lighter color and harder texture (Li, 2005). Pop chicken showed the mean overall acceptability of sensory attributes of  $4.42\pm 0.14$ . Moreover, the obtained results proved the existence of significant difference ( $P<0.05$ ) between examined RTE pop chicken and shish tawoak only, while did not established existence of statistically significant differences with other RTE products in this study. Indumathi and Obula (2015) found that organoleptic properties of chicken popcorn (Mean  $\pm$  S.E) were  $7.16\pm 0.02$  for color,  $6.61\pm 0.05$  for flavor,  $6.44\pm 0.09$  for tenderness and  $6.31\pm 0.04$  for juiciness. The overall acceptability was  $6.43\pm 0.08$ . They added that the spent hen chicken popcorn with different types of enrobing batters has good palatability. Based on the physico-chemical and proximate evaluation, enrobing can be successfully employed to add value to the products and develop a low calorie fried products. The obtained results in Table (2) and Fig. (2) revealed that chicken salad was the most contaminated meals in terms of APC, coliform, Enterobacteriaceae and Staph. aureus counts ( $1.094\times 10^6\pm 8.81\times 10^5$ ,  $1.696\times 10^2\pm 4.82\times 10$ ,  $1.183\times 10^4\pm 4.65\times 10^3$  and  $30.0\pm 21.34$ ) respectively, followed by chicken pane, shish tawoak, fajita, chicken burger, shawerma and grilled chicken

meals. On the contrary, the lowest microbial load of the aforementioned bacterial counts was recorded for fried chicken ( $5.78 \times 10^4 \pm 3.1 \times 10^4$ ,  $<3$ ,  $3.1 \times 10^2 \pm 43.33$  and  $<10$  cfu/g), respectively as well as pop chicken meals which recorded  $1.797 \times 10^4 \pm 9.34 \times 10^3$ ,  $<3$ ,  $2.20 \times 10^2 \pm 72.7$  and  $<3$  cfu/g, respectively. In the present study, shawerma was one of the contaminated RTE meals. This agreed with those reported by **Ibrahim, et al., (2014)**. Nearly similar result for APC was recorded by **Hassanein et al., (2015)** who found that APC of chicken shawerma was  $5.91 \times 10^4 \pm 1.08 \times 10^4$ . Moreover, **Hassanein (2010)** could detect APC count with 5.28 log cfu/g in examined chicken shawerma samples. This agreed with the present study. In this respect, **Gad (2004) and Sharaf and Sabra (2012)** failed to detect Staph. aureus count in examined chicken shawerma samples in Al-Taif Governorate KSA. This agreed with the present study, while **Sharaf and Sabra (2012)** recorded higher APC ( $1.2 \times 10^5$  cfu/g) and Enterobacteriaceae mean count ( $2 \times 10^4$  cfu/g). Meanwhile, mean value of APC, coliforms and Staph. aureus of examined chicken shawerma meals recovered by **Ibrahim, et al., (2014)** represented by  $4.58 \times 10^5 + 0.74 \times 10^5$  (cfu/g),  $9.97 \times 10^3 + 2.53 \times 10^3$  (MPN/g) and  $1.75 \times 10^4 + 0.31 \times 10^4$  (cfu/g) respectively, which were higher than that recorded in the present study. Chicken meals are subjected to be contaminated with several types of microorganisms from different sources during the period elapsed from the time of slaughtering, preparation, processing and cooking to consumption. These microorganisms varied according to the method of manufacture, quality of used non-meat ingredients, and contamination level during the processing chain, packaging and storage. This substantiates the findings of **Narasimha and Ramesh (1988)**. Grilled chicken meals are subjected to be contaminated with several types of microorganisms from different sources during the period elapsed from the time of slaughtering, preparation, processing and cooking to consumption. These microorganisms varied according to the method of manufacture, quality of used non-meat ingredients, and contamination level during the processing chain, packaging and storage. This substantiates the findings of **Narasimha and Ramesh (1988)**. The results regarding mean values of APC, coliforms and Staph. aureus of RTE chicken pane meals in the present study were higher than those reported by **Eid et al., (2014)** ( $21.6 \times 10^3 \pm 3 \times 10^3$ ,  $3.33 \pm 0.6$  and  $2.0 \times 10^3 \pm 1.1 \times 10^2$ ), respectively. Meanwhile, **Abd El-Aziz (2015)** could detect Staph. aureus in

examined half cooked chicken pane ranged from  $<10$  to  $7 \times 10^4$  with a mean value of  $9.29 \times 10^2 \pm 5.54 \times 10^2$  cfu/g. In this respect, **Ibrahim, et al. (2014)** could detect APC, with  $7.35 \times 10^4 + 1.17 \times 10^4$  (cfu/g), which was lower than the present study, while mean values of coliforms and Staph. aureus were higher than those of the present study ( $1.18 \times 10^3 + 0.26 \times 10^3$  (MPN/g) and  $3.01 \times 10^3 + 0.46 \times 10^3$  cfu/g), respectively. Also **Arab (2010)** found higher APC count ( $6.3 \times 10^4 \pm 0.35 \times 10^4$ ) in examined chicken pane. Nearly similar results regarding Min, Max and Mean  $\pm$  SE of Staph. aureus count of examined shish tawoak was recorded by **Eid et al., (2014)** ( $<10$ ,  $3 \times 10^3$  and  $2.6 \times 10^2 \pm 7.4 \times 10^2$ ), respectively. On contrary, higher mean value of Staph. aureus was observed by **Abo El-Enaen, et al., (2012)** who recorded  $2.39 \times 10^3$  cfu/g. and **Sampers, et al., (2010)** who detected mean Staph. aureus count of  $7.9 \times 10^3$  cfu/g. Moreover, **Eid, et al., (2014)** recorded  $2 \times 10^2$ ,  $2.2 \times 10^4$  and  $5.1 \times 10^3 \pm 6.6 \times 10^4$  as min, max and mean  $\pm$  SE of coliform count (MPN/g) of examined RTE chicken shish tawoak, respectively. These results were higher than those reported in the present study. Also higher coliform count was reported by **Sampers et al., (2010) and USDA-FSIS, (2012)** ( $2.51 \times 10^5$  and  $2.5 \times 10^3$ ), respectively. Moreover, **Ibrahim, et al., (2014)** illustrated higher mean values of APC, coliform and Staph. aureus represented by  $1.92 \times 10^5 \pm 0.46 \times 10^5$ ,  $4.32 \times 10^3 \pm 0.85 \times 10^3$  and  $9.84 \times 10^3 \pm 1.68 \times 10^3$  (cfu/g) for examined shish tawoak, respectively. The obtained results of chicken burger meals declared that The mean value of APC and Enterobacteriaceae of examined chicken burger samples in the present study was similar to that obtained by **Ali, 2011** ( $6.33 \times 10^4 \pm 1.84 \times 10^4$  and  $1.1^3 \times 10^3 \pm 0.25 \times 10^3$  cfu/g.) for fried beef burger samples, respectively, Also the obtained result in this study regarding APC was more or less agreed with those reported by **Hassanein, (2010)** (5.53 log cfu/g) in examined beef burger samples. Higher result ( $1.85 \times 10^3 \pm 0.42 \times 10^3$  cfu/g) was recorded for Staph. aureus count. Moreover, the trend in the present study, was lower than that reported by **Hassanein, et al., (2015)** who found that the mean values and incidences of APC, coliforms and Staph. aureus cfu/g were  $3.58 \times 10^3 \pm 0.72 \times 10^5$  (100%),  $2.39 \times 10^3 \pm 0.61 \times 10^3$  (73.33%) and  $2.73 \times 10^3 \pm 0.52 \times 10^3$  (80%) of examined chicken burger respectively. Moreover, Nearly similar results for APC were reported by **Becker, et al. (2002)** who carried out a survey of deli salads at retail and found APC at a range of 5 to almost 8 log cfu/g with a mean of 6.5 log cfu/g

( $3.2 \times 10^6$  cfu/g) and **Christensen and King (1971)** who found chicken salad have bacterial levels ranging from 4 to 7 log cfu/g, with a median of (5.8 log cfu/g). They also reported lower contamination with coleslaw at a range of 2.7 to 4.5, and a median of 3.6, log cfu/g. In another study of deli salads from various manufacturers, egg salads were found to have the poorest initial quality of 4.1 and 6.8 log cfu/g (**Folwer and Clark, 1975**). Regarding fried chicken meals, APC was agreed with that reported by **Mohamed et al., (2015)** ( $7.18 \times 10^4 \pm 1.44 \times 10^4$  cfu/g), while they reported higher Enterobacteriaceae, coliforms and Staphylococcal counts than that in the present study ( $8.73 \times 10^3 \pm 1.96 \times 10^3$ ,  $6.40 \times 10^3 \pm 1.23 \times 10^3$  and  $2.10 \times 10^3 \pm 0.32 \times 10^3$  cfu/g), respectively. The author also detected a significant difference ( $P < 0.01$ ) between boiled and fried chicken regarding APC and Enterobacteriaceae which was compatible with the results of this study, while obtained results of APC was lower than that obtained by **Zaki, et al. (2012)** who mentioned that the mean value of APC in fried chicken sandwiches was  $1.3 \times 10^6 \pm 1.2 \times 10^6$  cfu/g. Fast foods might not only have an impact on health but also could cause serious problem when contaminated with pathogenic microorganisms. Dangerous microorganisms, i.e. Salmonella and Escherichia coli might be transmitted to foods through preparation practices (**Adams and Moss, 2000**). In the recent years; all the fast food service restaurants have added fresh vegetables and seasonings in their meals. The contamination during processing and changes in microbial growth during storage might quantitatively and qualitatively alter microflora of foods. Regardless its inner tissues might be free from microorganisms, its surfaces could be contaminated with a variety of microorganisms depending on condition of the raw product, handling method and time and conditions during storage (**Odumeru et al., 1997 and Pelczar et al., 2006**). The consumption of such served meals has become very serious public health concern (**Meng and Doyle, 1998**). Consumers became more concerned about the food they consume; therefore consumers must attain information about the consumed product. One way ANOVA of APC among meals declared presence of highly significant differences ( $P < 0.01$ ) between pop chicken meals and both of chicken fajita, shish tawoak, burger and chicken salad meals. Enterobacteriaceae using One Way ANOVA, the results revealed a highly significant difference ( $P < 0.01$ ) between pop chicken and all chicken

meals under investigation, except for grilled and chicken mandy meals, which proved absence of significant difference with pop chicken meals. (**Table and Fig. 2**). The obtained results in **Table (3) and Fig. (3)** displayed the chemical analysis of RTE chicken products, pH and TBA value as well as TVB-N. pH of all chicken recorded low value as  $4.95 \pm 0.025$  in pop chicken and as high as  $6.31 \pm 0.064$  and  $6.13 \pm 0.075$  in chicken burger and mandy, respectively. TBA value recorded lower values ( $0.32 \pm 0.025$  mg mal. /kg) in pop chicken and  $0.42 \pm 0.047$  mg Mal. /kg in fried chicken meals, higher value ( $0.58$  mg Mal./kg) was recorded in chicken salad and  $0.57 \pm 0.040$  in chicken pane. Moreover, TVB-N recorded lower value ( $12.84 \pm 0.156$  mg) in examined chicken mandy meals as well as higher values in grilled chicken and chicken salad ( $14.70 \pm 0.369$  mg and  $14.57 \pm 0.620$  mg), respectively. In this object, **Edris, et al., (2013)** recorded pH of chicken pane ( $5.86 \pm 0.01$ ) and chicken fillet of  $5.95 \pm 0.01$  which is similar to the obtained results in the present study. Nearly similar results for pH were obtained by **Shedeed, (1999)** (5.7 to 6.1); **Affi, (2000)** (5.9 to 6.4) and **Fathy, 2012** (5.6 to 6.2). While, higher results were obtained by **Hassanein and Hassan, (2003)** ( $6.57 \pm 0.03$  to  $6.57 \pm 0.02$ ) for chicken pane. Regarding TVB-N, nearly similar values of examined chicken products were obtained by **Affi, (2000)** ( $12.57 \pm 0.222$  mg %). Higher result was recorded by **Hassanein and Hassan, (2003)** ( $30.76 \pm 1.07$  mg %) and **Eid, (2014)** ( $27.4 \pm 6.2$  mg %) for shish tawoak. While lower findings were recorded by **Edris, et al. (2013)** ( $7.06 \pm 0.26$  mg %) and **Fathy, (2012)**, ( $6.57 \pm 0.19$  mg %) as well as for chicken fillet ( $7.89 \pm 0.28$ ). It is important to emphasize that the meat and chicken meat products begin to deteriorate when it contains 30 mg TVN per 100 grams (**Pearson, 1984**). Thus, all examined samples were within the accepted limit. Moreover, TVB-N can be considered as a reliable indicative measure for the quality of various food articles specially poultry and its products. The results obtained by **Eid (2014)** for the mean value of TBA of heat treated poultry products were 0.59 mal./kg, **Hassanein, and Hassan, (2003)** ( $0.352 \pm 0.015$ ) and **Koreeski and Dwiatkiewicz, (2006)** ( $0.454$  mg/kg.) which were similar to that in the present study. Lower TBA values were recorded by **Shedeed, (1999)** (1, 69 to 0.28%), **Affi, (2000)** (0.051% to 0.223 mg %) and **Edris, et al. (2013)** ( $0.05 \pm 0.01$ ) for chicken pane and ( $0.03 \pm 0.01$ ) for chicken fillet. Accurately, TBA-value tends to decrease during frozen storage (**Lai, et al., 1991**). Oxidative rancidity was



occurred at TBA-value more than 0.9 mg % according to (Pearson, 1984). Thus, all samples were within the accepted limit. TBA value is closely related with the sensory characteristics of

### Conclusion

The achieved results in the current study allow concluding that chicken salad was the highly contaminated product, while pop and fried chicken meals were the least contaminated products. The variation in bacterial load in chicken meat products may be due to many causes mainly bad hygiene. Furthermore, the examined samples of chicken meat products as chicken salad was more contaminated with the highest level of microorganisms because such products may receive more handling during preparation as well as absence of heat treatment. Other products may be due to addition of spices which act as a source of bacterial contamination. However, among the control measures applied during chicken processing it was seen that the equipments comes in contact with personnel hands considered the main source of secondary

food article as rancidity (Salem, 1992). TBA values of examined chicken meals in present study were within the acceptable limit (<0.9 mg mal/kg).

microbial contamination. The initial microbial load of chicken carcasses can be controlled as a result of application of hygienic measures including Good Manufacture practices (GMP), Good Hygienic practices (GHP) as well as Risk Management program (RMP) which help identifying the existed hazard to evaluate the risk consequence in order to apply risk mitigation in due time to obtain a safe and healthy food for consumer. Effective control strategies for this diverse array of bacteria that currently limit the shelf life of ready-to-eat chicken meals and may constitute public health hazard and adversely affect on consumer health and this will require comprehensive integrated efforts along the food chain Generally , especially Restaurants manufacture and processing of those meals.

### References

- Aberle, E. D.; Forrest, J.C.; Gerrard, D.E. and Mills, E.W. (2001):** Principles of meat Science. 4<sup>th</sup> Ed. Kendall / Hunt Publishing Co., Dubuque, I.A.
- Abd El-Aziz, Amira, A. F. (2015):** Staphylococcus aureus enterotoxins in some breaded/Half cooked chicken and fish products. M. V. Sc. Thesis, Fac. Vet. Med., Benha Univ.
- Abo El-Enaen, Neveen H.; Eid, Samah and Salah-El-Dein, W.M. (2012):** Prevalence of Salmonella and staph.aureus microorganisms in broiler meat at Zagazig city and the effects f some organic acids on their viabilities. Egypt. J. Agric. Res. 90(1): 235-245.
- Adams MR, Moss MO (2000):** Food microbiology R Soc Chem. Sci Park Cambridge P 447.
- Affi-Jehan, S.A. (2000):** Chemical studies on some poultry meat products. M.V. Sci., Fac. Vet. Med. Zagazig Univ.
- Ali, E.A.M. (2011):** Microbial and chemical evaluation of fast foods. M.V. Sc. Thesis, (Meat Hygiene), Fac. of Vet. Med., Benha Univ. Egypt.
- Altunakar, B.; Sahin, S. and-Sumnu, G. (2004):** Functionality of batters containing different starch types for deep fat frying of chicken nuggets. Eur. Food Res. Technol., 218: 318-322.
- Anna, V. A. Resurreccion (1998):** consumer sensory testing for the product development. A chapman and Hall Food Science book. (Sensory).
- APHA "American Public Health Association" (2001):** Compendium of Methods for Microbiological Examination of food. 4<sup>th</sup> Ed., Washington, DC, USA.
- Arab, W.S.S. (2010):** Quality improvement of meat meals provided by a university student restaurant. Ph.D. Thesis (Meat Hygiene), Fac. Vet. Med., Benha University. Egypt.
- Becker, B.; Trierweiler, B.; Fehler, J.; Bohme, T.; Spiess, W.E.L. and W. H. Holzapfel, W.H. (2002):** Attended chilling cabinets in the food retail sale—3. Hygienic quality of delicatessen salads. Fleischwirtschaft, 82:104–107.
- Beckers, S.A. (1998):** More US consumers prefer chicken "Misset-World Poultry, 9:20-21.
- Christiansen, L.N. and King, N.S. (1971):** The microbial content of some salads and sandwiches at retail outlets. J. Milk Food Technol., 34:289–293.
- Chuaynukool, K.; Wattanachant, S. and Siripongvutikorn, S. (2007):** Chemical and physical properties of raw and cooked spent hen, broiler and Thai indigenous chicken muscles in mixed herbs acidified soup (Tom Yum) J Food Technol., 5: 180–186.
- Dainty, R.H. (1996):** Chemical and biochemical detection of spoilage. Int. J. Food Microbiol., 33: 19–34.
- Davies, A. and Board, R. (1998):** The microbiology of meat and poultry. 1st Ed., Edmundsburg Press, Ltd., Edmunds, London, UK.
- DOHA (2008):** Australian national children's nutrition and physical activity survey – Main

- finding. Department of Health and Ageing, 2007.<http://www.health.gov.au/internet/main/publishing.nsf/Content/health-pubhlth-strateg-foodmonitoring.htm#07survey> Accessed 6 August 2014
- Edris, A.M.; Hassanin, F.S. and Ghanim, S.H. (2013):** Chemical indices of some chicken meat products. *Benha Vet. Med. J.*, 24(1): 248-253.
- Egyptian Organization for Standardization "E.O.S" 63/11 (2006):** Egyptian organization For Standardization and quality control. Egyptian Standards for poultry meat products treated with heat. (Determination of pH value).
- Egyptian Organization for Standardization "E.O.S" 63/9 (2006):** Egyptian Organization For Standardization and quality control. Egyptian Standards for poultry meat products treated with heat. (Determination of Thiobarbituric Acid number (TBA) (mg/kg).
- Egyptian Organization for Standardization "E.O.S"63/10 (2006):** Arab republic of Egypt, Egyptian Organization for Standardization and quality control. Egyptian standards for poultry meat products treated with heat. (Determination of TVB-N, mg %).
- Eid, Amal, M., Eltalawy- Mona F., Zahran-Seham, E. and Khedre, A. Z. (2014):** Bacteriological and chemical evaluation of some heat-treated chicken products. *Benha Vet. Med. J.*, 27(2): 437-443.
- Eid, Samah, Saiid, Nagah, Karam, Hend, Ahmed M. Erfan, Hassan, Wafaa M.M., A.Baset, Afaf and Hassan M.K (2014):** Bacteriological and molecular studies on food poisoning pathogens isolated from chicken meat products. *Animal Health Research J.*, 2(3): 128-143.
- Esperance, B. (2016):** How to Tell If Chicken Salad Is Bad? Texas Agri Life Extension: Identifying and Handling Spoiled/Unsafe Food and Drinks after a Disaster. Kids Health: Food Poisoning, 1999-2016 Demand Media, Inc.
- Etaió, I.; Albisu, M.; Ojeda, M.; Gil, P. F.; Salmeron, J.; and Perez Elortondo, F. J. (2010):** Sensory quality control for food certification: A case study on wine. Panel training and certification, method validation and monitoring. *Food Cont.*, 21:542.
- Fathy, Eman (2012):** Chemical analysis of chicken meat with relation to its quality. Ph.D., Fac. Vet. Med. Banha Univ.
- FDA (Food and Drug Administration), USDA (United States Department of Agriculture), and CDC (Centers for Disease Control and Prevention) (2001):** Draft assessment of the relative risk to public health from foodborne *Listeria monocytogenes* among selected categories of ready-to-eat foods [Online] <http://www.foodsafetv.gov/~dms/lmrisksu.html> (verified July 18, 2002).
- Fowler, J.L. and W. S. Clark, W.S. (1975):** Microbiology of delicatessen salads. *J. Milk Food Technol.*, 38:111-113.
- Gad, M.A. (2004):** Microbiological evaluation of poultry meat and its products. M.V.Sc., Thesis Meat hygiene, Fac. Vet. Med., Sadat branch Minofyia Univ.
- Geornaras, I.; De Jesus, A.; Vanzyl, E. and von Holy, A. (1998):** Bacterial populations associated with the dirty area of a South African poultry abattoir. *J. of Food Prot.*, 61:700-703.
- Hassanein, Fatin, S. and Hassan, M.A. (2003):** Chemical indices of incipient deterioration in chicken cut-up products. *Benha Vet. M. J.*, 14: 54-65.
- Hassanein, Fatin S., El-Shater, Mohamed A. H. and Abd El-Fatah- Rabab R. (2015):** Bacteriological aspect of meat and poultry meat meals. *Benha Veterinary Medical J.*, 28 (2):91-97.
- Hassanein, Saadia M. E. (2010):** The Microbial Quality of Fast Food and Traditional Fast Food. *J. Nature and Sci.*, 8 (10): 117-133.
- Ibrahim, Hemmat M.; Salem -Amany, M and Mahmoud S. S. (2014):** Quality evaluation of some locally manufactured chicken meat products. *Benha Vet. Med. J.*, 26(2): 143-149.
- Imm, B. Y.; Lee, J. H.; and Yeo, I. K. (2009):** Capability analysis of sensory quality of jajang sauce. *Food Science & Biotechnology*, 18(3), 745-748.
- Imm, B. Y.; Shim, J. Y.; and Imm, J. Y. (2010):** Relationships between sensory and instrumental hardness and the estimation of hardness specifications for commercial soybean curd. *Food Science and Biotechnology*, 19(5): 1289-1293.
- Indumathi, J. and Obula Reddy, B. (2015):** Study on process optimization and effect of different types of enrobing on the quality of spent hen chicken popcorn. *International J. of Recent Scientific Res.*, 6 (5): 4070-4073.
- ISO 11133 :( 2014):** Microbiology of food, animal feed and water -- Preparation, production, storage and performance testing of culture media.
- ISO 21528-2(2004):** Microbiology of food and animal feeding stuffs -- Horizontal methods for the detection and enumeration of Enterobacteriaceae -- Part 2: Colony-count method.
- ISO 6887-2 :( 2003):** Microbiology of food and animal feeding stuffs -- Preparation of test samples, initial suspension and decimal dilutions for microbiological examination -- Part

**2:Specific rules for the preparation of meat and meat products**

- Jaturasitha, S.; Leangwunta, V.; Leotaragul, A.;Phongphaew, A.; Apichartsrungskoon, A.; Simasathitkul, N.; Vearasilp, T.; Worachai, L. and Meulen, U.T. (2002):** A comparative study of Thai native chicken and broiler on productive performance, carcass and meat quality. Proceedings of Deutscher Tropentag 2002.
- Kennedy, C.J. (2000):** Managing Frozen Foods. Wood head Publishing Limited and CRC press LLC, Cambridge, UK, and pp.143-144.
- Koreeski, I. and Swiatkiewicz, S. (2006):** Effect of stabilized fish oil supplementation and storage on changes in the fatty acid profile, TBARs content and sensory properties of breast meat from broiler chickens. Polish J. Natural Sci.,3: 421-426.
- Krokida, M.K.; Oreopoulou, V.; Maroulis, Z.B. and MarinouKouris, D. (2001a):** Deep fat frying of potato strips—quality issues. Drying Technol., 19(5): 879-935.
- Lai, S. M.; Gray, J. I.; Smith, D.M.; Booren, A.H.; Crackel, R. L. and Buckley, D. J. (1991):**Effect of oleoresin rosemary, tertiary butyl hydroquinone and sodium tripolyphosphate on the development of oxidative rancidity in restructured chicken nuggets. J. Food Sci., 56: 616-620.
- Lawless, H. T. (1995):** Dimensions of sensory quality: A critique. Food Quality and Preference, 6: 191–199.
- Lawlor, J.B.; Sheehan, E.M.; C.M. Delahunty, C.M.; Kerry, J.P. and Morrissey, P.A. (2003):** Sensory Characteristics and Consumer Preference for Cooked Chicken Breasts from Organic, Corn-fed, Free-range and Conventionally Reared Animals. International J. of Poult. Sci., 2(6): 409-416.
- Li, y. (2005):** Quality changes in chicken nuggets fried in oils with different degrees of hydrogenation. Thesis of Master of Science, Department of Bio resource Engineering Macdonald Campus, McGill University Montreal, Quebec, Canada.
- Lisa, M. R. and Lisa, A.R. (2009):** New American Cuisine for Today's Family: Fresh Ideas to Prepare Healthy Mediterranean Meals in Under 30 Minutes. SDP Publishing LLC. 69– ISBN 978-0-9824461-1-9.
- Love, J. (1994):** Product acceptability evaluation. In A. M. Pearson T. R. Dutson (Eds.) Quality attributes and their measurement in meat, poultry and fish products (pp. 337–358). Glasgow: Blackie Academic and Professional.
- McLennan, W. and Podger, A. (1999):** National nutrition survey. Foods eaten Australia. 1995. ABS Catalogue number 4804.0. Australian Bureau of Statistics and Common wealth Department of Health and Family Services, Canberra.
- Meng, J. and Doyle, M. P. (2002):** Introduction Microbiological food safety. Microbes and Infection, 4(4): 395-397.
- Mohamed, A. H.; Amin- Reham, A.; Emam-Sherien, M. and Abdel Aal -Asmaa, A.A .M. (2015):** Bacterial Status of Food Meals served at Governmental Hospital. Benha Vet. Med. J., 29(1):143-150.
- Mulla, Z. S. (2002):** Studies on the onset of warmed over and the use of M – S based electronic nose. Thesis, the Ohio state University.
- Munoz, A. M.; Civille, G. V. and Carr, B. T. (1992):** Sensory evaluation in quality control New York: Van Nostrand Reinhold.
- Narasimha, R.D. and Ramesh, B.S. (1988):** Microbial profiles of minced meat. Meat Science 23:279-291.
- Odumeru, J.; Mitchell, S.; Alves, D.; Lynch, J.; Yee, A.; Wang, S.; Styliadis, S. and Farber, J. (1997):** Assessment of the microbiological quality of ready to use vegetables for health care food services. J Food Prot., 60: 954-960.
- Ozcan, O. (2013):** The Sultan's Kitchen: A Turkish Cookbook. Tuttle Publishing. 179– ISBN 978-1-4629-0639-0.
- Pearson, D. (1984):** Chemical analysis of foods 8<sup>th</sup> Ed, Publishing Co., Churchill Livingstone, Edinburgh, London, UK.
- Pelczar, M.J.; Chane, C.S. and Kreig, N.R. (2006):** Microbiology 5<sup>th</sup> edition. Tata McGraw-Hill Publishing Company Limited, New Delhi.
- Pérez, F.; Castro, R.; Posada-Izquierdo, G.D.; Valero, Carrasco, E.; García-Gimeno, R.M. and Zurera,G.( 2010):**Evaluation of hygiene practices and microbiological quality of cooked meat products during slicing and handling at retail. Meat Sci., 86(2): 479-485.
- Public Health Laboratory Service (PHLS) (2000):** Guidelines for the microbiological quality of some ready-to-eat foods sampled at the point of sale. Comm. Dis. and Public Health, 3(3): 163-167.
- Sabine, L. and Brian, S. E. (2004):** A handbook of statistical analysis using Statistical Package for the Social Sciences (SPSS). One Way ANOVA, Ver. 20.
- Salem, Amany, M. (1992):** Biological analytical studies on incipient deterioration of beef meat on liver. M. V. Sc., Faculty of Veterinary Medicine, Zagazig Univ. Benha branch.
- Sampers, I.; Jacxsens, L.; Luning, P.A.; Marcelis, W.J.; Dumoulin, A. and Uyttendaele, M.(2010):** Performance of food safety management systems

- in poultry meat preparation processing plants in relation to *Campylobacter* spp. contamination. *J. Food Protec.*, 73(8): 1447-1457.
- Sharaf, Eman, M., and Sabra, Sherifa, M. (2012):** Microbiological loads for some types of cooked chicken meat products at Al-Taif Governorate, KSA. *World Applied Sciences Journal*, 17(5):593-597.
- Shedeed, N.A. (1999):** Evaluation of microwave cooking of chicken meat. M.Sc., Fac. Agric., Cairo University.
- Soliman, M.R. (1988):** Sanitary status of ready to eat meat products and fishes. M.V.Sc., Thesis (Meat hygiene), Fac. Vet. Med. Cairo Univ.
- Sow, T.M.A. and Grongnet, J.F. (2010):** Sensory characteristics and consumer preference for chicken meat in Guinea. *Poult Sci.*, 89(10): 2281-2292.
- United States Department of Agriculture and Food Safety and Inspection Service office of "USDA-FSIS" The Nationwide Microbiological Baseline Data Collection Program (2012):** Raw chicken parts baseline survey (RCPBS). Food Safety Net Service, Ltd., San Antonio.
- Virginia, M. (2010):** Istanbul. Lonely Planet. 145-ISBN 978-1-74220-404-8.
- Warries, P.D. (2000):** Meat Science. 1<sup>st</sup> Ed. CABI Publishing Co. CABI International, Walling Ford, United Kingdom.
- Wattanachant, S.; Benjakul, S. and Ledward, D. A. (2004):** Composition, color, and texture of Thai indigenous and broiler chicken muscles. *Poult Sci.*, 83:123-128.
- Yvonne, Y. H. (2002):** Muslim Minorities in the West: Visible and Invisible. Roman Altamira. 138- ISBN 978-0-7591-0218-7.
- Zaki, R.S.; Hefnawy, Y.A. and El-Khateib, T.S.A. (2012):** Assessment of microbiological quality and safety of home delivery ready to eat foods. Assiut university researches.

### الملخص العربي

امان و جودة منتجات الدجاج الجاهزة للأكل

تغريد حمدي عباس\* (١)، خالد شوقي طلبية\*، عادل محمد إبراهيم\*\* و محمد خالد المسلمي\*\*  
قسم الرقابة على اللحوم و منتجاتها\*\* - كلية الطب البيطري - جامعة القاهرة\*\*  
و معهد بحوث صحة الحيوان\* - الدقي - الجيزة

تلوث منتجات الدجاج الجاهزة للأكل بمسببات الأمراض التي تنقلها الأغذية لا تزال مشكلة صحية عامة مهمة لأنها تؤدي إلى المرض إذا كان هناك سوء ممارسة أثناء الذبح، نزع الأحشاء، المعالجة و بعد ذلك التحضير عن طريق الطهي. لذلك هذه الدراسة ركزت على أمان بعض وجبات الدجاج الجاهزة للأكل أثناء التحضير من خلال النواحي الحسية، و الكميائية لكشف مدى طزاجتها و سلامتها من الميكروبات. وقد تم تجميع هذه المنتجات من مطاعم مختلفة بمحافظة القاهرة و الجيزة، و الدراسة البحثية ركزت على عشرة منتجات تشمل شورما الدجاج، الدجاج المشوي، بانيه الدجاج، الشيش طاووق، فاهيتا الدجاج، برجر الدجاج، سلطة الدجاج، المندي، الدجاج المقلي و أخيراً كور الدجاج. و المعايير البيكتريولوجية المستخدمة هي: عدد الميكروبات الكلية، عدد الميكروبات القولونية، عدد الميكروبات المعوية، عدد ميكروب الاستافيلوكوكاس اورياس. و قد أوضحت الدراسة ان سلطة الدجاج هي أكثر المنتجات تلوثاً من حيث العد البكتيري الكلي، القولوني، المعوي، الاستافيلوكوكاس اورياس ( $1.09410^5 \times 8.81 \pm 10^6$ ،  $1.696 \times 10^2 \pm 10^2$ ،  $1.82 \times 10^4 \pm 10^4$ ،  $1.0 \times 10^6 \pm 4.65 \times 10^6$ ،  $10^3 \pm 3.3 \times 10^3$  وحدة خلية/جم) على التوالي. و بالمقابل أقل عدد ميكروبات وجد في منتج الدجاج المقلي و هي (وحدة خلية/جم) على التوالي و كذلك كور الدجاج ( $10^4 \times 10^5$ ،  $10^3 \pm 10^3$ ،  $10^2 \times 10^3$ ،  $3 >$ ،  $3.33 \times 10^2 \pm 43.33$ ) و على التوالي. و كذلك كور الدجاج ( $10^4 \times 10^7$ ،  $10^3 \times 10^4$ ،  $10^3 \times 10^4$ ،  $20$ ،  $3 >$ ،  $10^3 \times 10^4$ ،  $10^3 \times 10^4$ ،  $20$ ،  $3 >$ ،  $10^3 \times 10^4$ ) من خلال هذه النتائج من الواضح ان خمسة منتجات و هي (السلطة، البانيه، الشيش طاووق، الفاهيتا، البرجر) لديها اعلى متوسط قيمة للفحص البكتيري و لكن مازالت بالحد المسموح به في العد البكتيري الكلي تبعاً لخدمات معمل الصحة العامة. و من حيث الخواص الكميائية فكانت في الحدود المسموح بها. علاوة على ذلك تم فحص السمات الحسية و مناقشة انعكاس عدد الميكروبات على المستهلك و الصحة العامة.