



Isolation and characterization of avian reovirus (Egypt/ARV/Giza 2011) associated with arthritis in broiler breeder flocks

Latifa, Bekheet⁽¹⁾; Hussein, H.A.⁽²⁾; Elsafty, M.M.⁽³⁾; Tantawy, L.A.⁽⁴⁾ and Helal, A.M.⁽³⁾

⁽¹⁾ Specific Pathogen Free Eggs Project, Koum Oshiem, El-Fayoum Governorate; ⁽²⁾ Virology Department, Faculty of Veterinary Medicine, Cairo University; ⁽³⁾ Central Laboratory for Evaluation of Veterinary Biologics, VSVRI, Abbasia, Cairo. ⁽⁴⁾ Animal Health Researches Institute, AHRI, Dokki, Giza.

Abstract

Samples of tibiotarsal tendons, tendon sheaths and fluids around ruptured tendons were collected from broiler breeder flock of 28 weeks old in Giza governorate, showed symptoms of arthritis, lameness and swelling in the hock joints. Trial for isolation of avian reovirus (ARV) was done by inoculation in embryonated chicken eggs (ECE). After propagation and titration of the isolated virus, it had been antigenically characterized by agar gel precipitation test (AGPT) and fluorescent antibody technique (FAT). Pock lesions on the chorioallantoic membrane (CAM) was negatively stained and examined by electron microscope, also histopathological examination was carried out. Molecular characterization was performed by reverse transcriptase polymerase chain reaction (RT-PCR). The results confirm the isolation of avian reovirus from the suspected flock.

Key words: avian reovirus, arthritis, CAM and pock

Introduction

Avian reoviruses belong to genus Orthoreovirus of family Reoviridae (Mertens, 2004). Reoviruses have a double capsid structure comprised of an outer capsid of ~85 nm in diameter and an inner core of ~45 nm in diameter, which accommodates 10-segmented genomic dsRNA (Zhang et al., 2005). The 10 segments encode for at least 12 viral proteins, 8 structural proteins and 4 nonstructural proteins. σ C is one of the most important polypeptides among ARV proteins that induces ARV-specific neutralizing antibodies which makes such protein a target and marker in studies on ARV genetic variations as well vaccine development (Martinez-Costas et al., 1997 and Shapouri et al., 1995). The avian reovirus can induce several manifestations in chickens including arthritis, lameness and swelling in the hock joint (Jones and Guneratne, 1984), inclusion body hepatitis (McFerran et al., 1976), enteritis (Dutta and Pomeroy, 1969), hydro-pericardium (Bains and MacKenzie, 1974), myocarditis (Davis et al., 2012), central nervous system disease (Van de Zande and Kuhn, 2007) and runting-stunting syndrome or malabsorption syndrome (MAS) (Goodwin et al., 1993). ARVs could be inoculated on CAM as it is a successful route for ARV isolation and the main pathological changes are appearance of pock lesions on CAM (Schwartz et al., 1976). The electron microscope has been used as a powerful tool in the characterization of ARV either by negative stain or thin section electron microscopic examination

(Goldsmith, 2014). ARVs were classified into 11 serotypes (Wood et al., 1980), based on antigenic characterization using serological tests like agar gel precipitation test (AGPT), virus neutralization test (VNT), enzyme linked immunosorbent assay (ELISA) and fluorescent antibody technique (FAT). Molecular-based techniques including reverse transcriptase polymerase chain reaction (RT-PCR) (Bruhn et al., 2005), nested PCR (Liu et al., 1997), multiplex PCR (Caterina et al., 2004), real-time PCR (Ke et al., 2006), PCR followed by restriction fragment length polymorphism (RFLP) (Lee et al., 1998), and in situ hybridization (ISH) (Liu and Giambrone, 1997), have been used to detect ARVs. Phylogenetic studies classified isolates of ARVs into various groups and lineages also provided evidences showing frequent reassortments among the circulating lineages which responsible for variation in the ARV genome segments. Previous studies on comparison of sequences of ARVs were focused on σ C-encoding gene (Liu et al., 2003). The aim of the present study is to isolate and characterize ARV associated with arthritis in broiler breeder chickens.

Material and methods

Samples of tibiotarsal tendons, tendon sheaths and fluids around ruptured tendons were collected from chickens of broiler breeder farm of 28 weeks old located in Giza governorate, suffered from rupture of Achilles tendon, swelling in joints and lameness, showed 10% morbidity 0.2% mortality. The chickens were vaccinated with reovirus

S1133 live vaccine at 38 days old and inactivated ARV vaccine at 18 weeks and boosted at 22 weeks old. Homogenate of the samples was prepared in phosphate buffer saline (PBS) pH 7.2 with antibiotic (200 IU/ml of penicillin and 100µg/ml of dehydro-streptomycin) followed by three times freezing and thawing then centrifuged at 5000 r.p.m. for 15 minutes at 4°C. The supernatant was further passed through 0.45 µl filter membrane. The filtrate was used for virus isolation by inoculation on the chorioallantoic membrane of specific pathogen-free embryonated chicken eggs (SPF-ECE) at 11-13 days old. The inoculated eggs were incubated at 37° C and 80% humidity for 5 days with daily candling then the CAMs were harvested (Tantawy, 1999). Propagation of the isolated virus on CAM was carried out for 6 passages in (SPF-ECE) (Wood et al., 1980). Virus titration test was performed in SPF-ECE according to (Neelima et al., 2003) using filtrate of the harvested CAM from the sixth passage that showing pock lesions and the virus titer was calculated by Karber method (Finney, 1978). For the antigenic characterization of the isolated virus, AGPT (Tantawi et al., 1984) and FAT (Adair et al., 1987) were conducted using specific monoclonal antibodies against ARV. Transmission electron microscopic examination was applied on filtrate of pock lesion preparations (Goldsmith, 2014). Images were captured by CCD camera model AMT. Histopathological examination of pock lesions on the harvested CAM was carried out according to Tantawy, (1999). RT-PCR was employed for molecular characterization of the isolated virus using primers designed by Xie et al. (1997). These primers were specific to the conserved fragment of sigma C gene and amplify fragment with molecular size of 532 bp. Sequences of the utilized primers were as follows:

Primer (Sense)	1	5'GGTGCGACTGCTGTATTGGTAAC 3'.
Primer (Antisense)	2	5'AATGGAACGATAGCGTGTGGG 3'.

Results

Isolation, propagation and titration of ARV

Infected CAM of the inoculated eggs showed thickening, opaqueness and swelling forming pock lesion in the third passage. Number of pock lesions increased and markedly appeared between fourth to sixth passages figure (1). The titer of the isolated virus was 10^{6.1} EID₅₀/ml.




Figure (1) showed marked appearance of pock lesion on CAM of ECE in the fifth passage of the virus.

Antigenic characterization of the isolated virus by AGPT and FAT

AGPT showed positive result using specific monoclonal antibodies figure (2). Examination of frozen sections of pock lesions by fluorescent microscope using specific monoclonal antibodies revealed specific fluorescent foci in the sections confirming the antigenic characterization of the isolated virus figure (3).




Figure (2) AGPT showed line of precipitation.




Figure (3) FAT showed positive fluorescent foci on CAM section.

Electron microscopic examination

Appearance of virus particles in aggregates with apparent damage to the outer capsid of the virus with size reached about 45 nm figure (4).




Figure (4) showed aggregates of the virus particles with apparent damage to the outer capsids, by TEM Mode, Direct Mag: 120000 x Print Mag: 252000x@211 mm

histopathological examination of harvested cams

The ectodermal layer of CAMs showed mild to moderate hyperplasia associated with hypertrophy and vacuolation of cells. Very characteristic oesinophilic intracytoplasmic inclusion bodies were also noticed in ectodermal cells figure (5) and (6). Focal necrosis of ectoderm was observed. The mesoderm showed pronounced edema with congested blood vessels, in addition to inflammatory cells and fibroblasts. Few blood vessels occluded with inflammatory cells. The endoderm showed mild hyperplasia with focal necrosis.




Figure (5) & (6) showed the intracytoplasmic inclusion bodies in sections of pock lesions.

RT-PCR

Electrophoresis of the amplified products in agarose gel revealed the expected band with correct molecular size confirming the molecular characterization of the isolated virus figure (7).




Figure (7) showed the expected band with correct molecular size.

1-sample tested. 2-positive control.

Discussion

ARV could be isolated from different organs like kidney, thymus, ceecal tonsils, spleen, pancreas (Van Loon et al., 2001), liver, intestine, heart, joints and tendons (Lu et al., 2015). In the current study, chickens of broiler breeder farm were suffered from rupture of Achilles tendon, swelling in joints and lameness. Samples were collected from the affected joints and tendons as the virus still for long time in the joints and tendons for at least 285 days post infection (Kerr and Olson, 1969). ARV isolation could be performed in tissue culture or in the SPF-ECE. There are different routes were used for inoculation of the virus in SPF-ECE including yolk sac, allantoic sac and chorioallantoic membranes. The most suitable conditions for propagation of the avian reovirus vaccinal strain were on the embryonated chicken eggs and the preferable route for inoculation was on the chorioallantoic membrane (Hassan et al., 1993). In this study, ARV was isolated on the CAM of SPF-ECE and after number of passages; characteristic lesions of reovirus including opaque and thick pocks appeared on the CAM figure (1). Performing the infectivity test revealed that titer of the isolated virus reached $10^{6.1}$ EID₅₀/ml after fourth passages. Antigenic characterization of the isolated virus revealed the appearance of line of

precipitation in AGPT between the isolated virus and the antibodies specific to ARV confirmed the isolation of ARV figure (2). Similar results were obtained by Tantawi et al., (1984) who isolated and characterized two infectious tenosynovitis producing viruses from tendon sheaths and synovial fluids of broiler and broiler breeder flocks in Egypt in 1983. Also, Similar results obtained by Kheir El-Din and El-Sanoussi (1986) who isolated reovirus from broiler breeder chickens suffered from retardation of growth, diarrhea and abnormal gate. The isolated viruses were characterized by AGPT. Fluorescent foci in the frozen sections of the obtained pock lesions on the harvested CAM were observed confirming the antigenic characterization of the isolated virus figure (3). Negative staining of the isolated virus and examination under the electron microscope demonstrated the presence of the characteristic aggregates of the virus particles with apparent damage to the outer capsid figure (4). Size of the virus particles reached about 45 nm and that similar to previous studies achieved by others like Walker et al. (1972); Mustaffa-Babjee et al. (1973) and Tantawy, (1999). The collected CAMs of the inoculated SPF-ECE were macroscopically opaque, thickened and oedematous with engorged and tortuous blood vessels. White pock lesions of different sizes appeared on the CAMS, some of them were small minute (pin-headed) and the others looked large prominent specially at sit of inoculation figure (1). Several researchers have been reported these lesions even by inoculation of local isolate of ARV or vaccinal strain as Tantawi et al., (1984) and Tantawy, (1999). Histopathological changes of the pock lesions were characteristic to ARV with appearance of oesinophilic intracytoplasmic inclusion bodies in the ectodermal layer which considered pathognomonic to ARV and that ensure the replication and maturation of the virus in the cytoplasm of the affected cells. The virus invaded the CAM leading to appearance of another changes like hypertrophy, hyperplasia, vacuolation of the cells and focal necrosis in the ectodermal layer also edema with congested blood vessels and inflammatory cells in the mesoderm. The observed lesions are agree with previous study performed by the reference vaccinal strain of ARV which showed the same picture as

detected by **Tantawy, (1999)**. Different segments were target for amplification in the RT-PCR assay for the detection and molecular characterization of the virus in the clinical samples. RT-PCR could segment (**Lee et al., 1998**) or S4 genome segment (**Roussan et al., 2012**) or L1 genome segment (**Leary et al., 2002**). The sigma C gene of the S1 segment was selected to be applied in the present study utilizing primers specific to the conserved region to obtain fragment of 532 bp (**Xie et al., 1997**). The employed RT-PCR revealed band of corrective size confirming the molecular characterization of the isolated reovirus. Indeed,

References

- Adair, B.M.; Burns, K. and Mckillop, E.R. (1987)**. Serological studies with reoviruses in chickens turkeys and ducks. *J. Comp. Path.* Vol (97): 495-501.
- Bains, B. S. and MacKenzie, M. (1974)**. Reovirus-associated mortality in broiler chickens. *Avian Dis.* 18, 472-476.
- Bruhn, S.; Bruckner, L., and Ottiger, H. P. (2005)**. Application of RT-PCR for the detection of avian reovirus contamination in avian viral vaccines. *J. Virol. Meth.* 123, 179-186.
- Caterina, K. M.; Frasca, S.; Girshick, T. and Khan, M. I. (2004)**. Development of a multiplex PCR for detection of avian adenovirus, avian reovirus, infectious bursal disease virus, and chicken anemia virus. *Mol. Cell. Probe.* 18, 293-298.
- Davis, J.F.; Kulkarni, A. and Fletcher, O. (2012)**. Myocarditis in 9- and 11- day-old broiler breeder chicks associated with a reovirus infection. *Avian Disease.* 56(4): 786-790.
- Dutta, S. K. and Pomeroy, B. S. (1969)**. Isolation and characterization of an enterovirus from baby chicks having an enteric infection. II. physical and chemical characterization and ultrastructure. *Avian Dis.* 11, 9-15.
- Finney, D.J. (1978)**. *Statistical Method in Biological Assay*, 3rd edit., Griffin, London.
- Goodwin, M. A.; Davis, J. F.; McNulty, M. S.; Brown, J. and Player, E. C. (1993)**. Enteritis (so-called runting stunting syndrome) in Georgia broiler chicks. *Avian Dis.* 37, 451-458. associated with arthritis of chickens. *Avian Dis.* 17, 415-424.
- Goldsmith, C.S. (2014)** Morphologic differentiation of viruses beyond the family level. *Viruses* 2014,6,4902-4913; doi:10.3390/v6124902 available at www.mdpi.com/journal/viruses
- Hassan, Nadia; Abou El-Khair, M.; El bordeny, F.; El-Assily, S.; Khashabah, E.; Wahba, S. and Saber, S. (1993)**. Studies on the vaccinal strain of reovirus. *Assuit Vet. Med. Journal.* Vol.29 (No.57): 50-61.
- Jones, R. C. and Guneratne, J. R. M. (1984)**. The pathogenicity of some avian reoviruses with particular reference to tenosynovitis. *Avian Pathol.* 13, 173-189.
- Ke, G. M.; Cheng, H. L.; Ke, L. Y.; Ji, W. T.; Chulu, J. L. C.; Liao, M. H.; Chang, T. J. and Liu, H. J. (2006)**. Development of a quantitative Light Cycler real-time RT-PCR for detection of avian reovirus. *J. Virol. Meth.* 133, 6-13.
- Kerr, K. M. and Olson, N. O. (1969)**. Pathology of chickens experimentally inoculated or contact-infected with an arthritis-producing virus. *Avian Dis.* 13, 729-745.
- Kheir El-Din, A.M. and El-Sanousi, A. (1986)**. Infectious stunting and runting syndrome (helicopter disease) : Clinical investigation ,isolation and identification of the possible etiological agent. *Journal of Egyptian Veterinary Medical Association.* Vol.46 (No.3):283-294.
- Leary, T.P.; Erker, J.C.; Chalmers, M.L.; Wetzel, J.D.; Desai, S.M.; Mushahwar, I.K. and Dermody, T.S. (2002)**. Detection of reovirus by reverse transcription-polymerase chain reaction using primers corresponding to conserved regions of the viral L1 genome segment. *Journal of Virological Methods.* 104: 161-165.
- Lee, L. H.; Shien, J. H. and Shieh, H. K. (1998)**. Detection of avian reovirus RNA and comparison of a portion of genome segment S3 by polymerase chain reaction and restriction

- enzyme fragment length polymorphism. Res. Vet. Sci. 65, 11-15.
- Liu, J. and Giambrone, J.J. (1997). In situ detection of reovirus in formalin-fixed
- Liu, H. J.; Giambrone, J. J. and Nielsen, B. L. (1997). Molecular characterization of avian reoviruses using nested PCR and nucleotide sequence analysis. J. Virol. Meth. 65, 159-167.
- Liu, H. J.; Lee, L. H.; Hsu, H. W.; Kuo, L. C. and Liao, M. H. (2003). Molecular evolution of avian reovirus: evidence for genetic diversity and reassortment of the S-class genome segments and multiple cocirculating lineages. Virology. 314, 336-349.
- Lu, H.; Tang, Y.; Dunn, P.A.; Wallner-Pendleton, E.A. Lin, I. and Knol, E.A. (2015). Isolation and molecular characterization of newly emerging avian reovirus variants and novel strains in Pennsylvania, USA, 2011-2014. Sci Rep. 2015; 5: 14727. doi: 10.1038/srep14727
- Martinez-Costas, J.; Grande, A.; Varela, R.; Garcia-Martinez, C. and Benavente, J. (1997). Protein architecture of avian reovirus S1133 and identification of the cell attachment protein. J. Virol. 71, 59-64.
- McFerran, J. B.; McCracken, R. M.; Connor, T. J. and Evans, R. T. (1976). Isolation of Viruses from Clinical Outbreaks of Inclusion Body Hepatitis. Avian Pathol. 5, 315-324.
- Mertens, P. (2004). The dsRNA viruses. Virus Res. 101, 3-13.
- Mustaffa-Babjee, A.; Spradbrow, P.B. and Omar, A.R. (1973). Characterization of an avian reovirus isolated in Queensland. J. Comp. Path. 1973. Vol.83.387-400.
- Neelima, S.; Ram, G.C.; Kataria, J.M. and Goswami, T.K. (2003). Avian reovirus induces an inhibitory effect on lymphoproliferation in chickens. Veterinary Research Communications. 27:73-85.
- Roussan, D.A.; Shaheen, I.A.; Khawaldeh, G.Y.; Totanji, W.S. and Al-Rifai, R.H. (2012). Simultaneous detection of astrovirus, rotavirus, reovirus and adenovirus type I in boiler chicken flocks. Polish Journal of Veterinary Science, Vol (15), No (2), 337-344.
- Schwartz, L. D.; Gentry, R. F. and Rothenbacher, H. (1976). Infectious tenosynovitis in commercial white Leghorn chickens. Avian Dis. 20, 769-773.
- Shapouri, M. R. S.; Kane, M.; Letarte, M.; Bergeron, J.; Arella, M. and Silim, A. (1995). Cloning, sequencing and expression of paraffine embedded chicken tissues using a digoxigenie-labeled cDNA probe. Avian Diseases 41:374-378.
- the S1 gene of avian reovirus. J. Gen. Virol. 76, 1515-1520.
- Tantawi, H.H.; Amina, N.; yousef, Y.I.; Fawzia, M.; Al-Abdulla, j.M.; El-Batrawi, A.; El-Ghawas, A.; Nasser, A.A. and Reda, I.M. (1984). Infectious tenosynovitis in broilers and broiler breeders in Egypt. Veterinary Research Communications. Vol.8: 229-235.
- Tantawy, L.A. Ph D Thesis (1999). Pathological studies on SPF fertile eggs inoculated with viral strains. Faculty of Veterinary Medicine, Cairo University.
- Van de Zande, S. and Kuhn, E.M. (2007). Central nervous system signs in chickens caused by a new avian reovirus strain: a pathogenesis study. Vet. Microbiol. 120:42-49.
- Van Loon, A.A.; Koopman, H.C. W.; Mumczur, K.J.; Szeleszczuk, O.; Karpinska, E.; Kosowska, G. and Luticken, D. (2001). Isolation of a new serotype of avian reovirus associated with malabsorption syndrome in chickens. Veterinary Quarterly. Volume 23, Issue 3, pages 129-133.
- Walker, E.R.; Friedman, N.O. and Olson, N.O. (1972). Electron microscopic study on an avian reovirus that causes arthritis. J. Ultrastructure Research. 41, 67-79.
- Wood, G. W.; Nicholas, R. A. J.; Hebert, C. N. and Thornton, D. H. (1980). Serological comparisons of avian reoviruses. J. Comp. Pathol. 90, 29-38.
- Xie, Z.; Fadl, A.A.; Grishick, T. and Khan, M.I. (1997). Amplification of avian reovirus RNA using the reverse transcription polymerase chain reaction. Avian Disease 41:654-660.
- Zhang, X.; Tang, J.; Walker, S. B.; O'Hara, D.; Nibert, M. L.; Duncan, R. and Baker, T. S. (2005). Structure of avian orthoreovirus virion by electron cryomicroscopy and image reconstruction. Virology. 343, 25-35.
- Zhang, Y.; Liu, M.; Shuidong, O.; Hu, Q.L.; Guo, D.C.; Chen, H.Y. and Han, Z. (2006). Detection and identification of avian, duck and goose reoviruses by RT-PCR: goose and duck reoviruses are part of the same genogroup in the genus Orthoreovirus. Arch. Virol. (2006) 151: 1525-1538.

المستخلص العربي

عزل و توصيف فيروس ريو الدجاج (Egypt/ARV/GIZA 2011) المصاحب لمرض التهاب المفاصل في قطعان امهات التسمين لطيفة بخيت⁽¹⁾-حسين على حسين⁽²⁾-منير الصفتى⁽³⁾-خليلي طنطاوى⁽⁴⁾-احمد هلال⁽⁵⁾

⁽¹⁾ مشروع انتاج البيض الخالى من المسببات المرضية-كوم اوشيم-الفيوم ⁽²⁾ قسم الفيروسات-كلية الطب البيطرى-جامعة القاهرة ⁽³⁾ المعمل المركزى للرقابة على المستحضرات الطبية البيطرية معهد بحوث الامصال واللقاحات البيطرية-العباسية-القاهرة ⁽⁴⁾ معهد بحوث صحة الحيوان-الدقى-الجيزة

تم جمع عينات من اوتار التيببوتارسال و الاغشيه والسوائل حول الاوتار الممزقة من قطيع امهات التسمين عند عمر ٢٨ اسبوع في محافظة الجيزة حيث كانت تظهر أعراض التهاب المفاصل، العرج وتورم في مفاصل العرقوب و وقد تم اجراء محاولة لعزل فيروس ريو الدجاج بحقنه على الغشاء السقائى المشيمانى فى بيض الدجاج المخصب الخالى من المسببات المرضية و بعد تمرير الفيروس و معايرته تم عمل توصيف انتيجينى للفيروس باستخدام اختبارى الترسيب المناعى و التعادل الفلورسنتى الضوئى. وقد اظهرت النتائج وجود بثرات على الغشاء السقائى المشيمانى بالبيض المحقون و قد تم اخذ عينات منها و صبغها صبغ سالب و فحصها تحت الميكروسكوب الالكترونى و ايضا تم عمل فحص التغيرات المرضية فى انسجة الغشاء السقائى المشيمانى للبيض المحقون. كما تم عمل التوصيف الجزيئى بواسطة اختبار تفاعل البلمرة المتسلسل المسبوق بالنسخ العكسى و قد اسفرت النتائج التاكيد على عزل فيروس ريو الدجاج من القطيع المشتبه به.