



## IMPROVING AGENTS FOR PHYTOREMEDIATION OF SOIL CONTAMINATED WITH PYMETROZINE INSECTICIDE

Aly A.A. Shalaby<sup>1\*</sup>, A.A.A. Aioub<sup>1</sup>, A.A.A. Romeh<sup>2</sup> and M.R.A. Ramadan<sup>2</sup>

1. Plant Prot. Dept., Fac. Agric., Zagazig Univ., Egypt

2. Plant Prod. Dept., Fac. Technol. and Dev., Zagazig Univ., Egypt

Received: 22/02/2017 ; Accepted: 13/03/2017

**ABSTRACT:** The present work was designed to investigate the potential of using plantain (*Plantago major* L.) for the phytoremediation of pymetrozine contaminated soil. The use of soluble silicon dioxide (SiO<sub>2</sub>), Tween 80, hydroxypropyl-β-cyclodextrin (HPβCD) and liquid humic acid (HA) for enhancing the availability and uptake of pymetrozine contaminated soil by *P. major* were evaluated. Results revealed that pymetrozine concentrations in soil with *P. major* reduced by 30.00 – 83.25% throughout 1 to 12 days exposure, compared with 12.50–61.90% in the soil of control. Pymetrozine uptake in the roots and translocated in the leaves of *P. major* to reach the maximum levels, 53.41 mg/kg and 58.08 mg/kg, respectively, after 4 days. The phytoremediation efficiency of *P. major* amended with SiO<sub>2</sub> was greater than that of other solubility-enhancing agents with respect to the removal of pymetrozine from contaminated soil within 1-12 days of treatment. The addition of SiO<sub>2</sub> increased pymetrozine uptake in roots and translocation in leaves by about 170.84% and 322.83% compared with *P. major* roots and leaves alone, respectively within 4 days. The most-effective to least-effective supplements to use in combination with *P. major* for the purpose of pymetrozine phytoremediation in roots and leaves were found to be as follows: SiO<sub>2</sub> > HA > HPβCD > Tween 80. This study indicates that SiO<sub>2</sub> can improve the efficiency of phytoremediation of pymetrozine.

**Key words:** Phytoremediation, *Plantago major*, pymetrozine, improving agents, soil.

## INTRODUCTION

Pymetrozine was the compound of the pyridine azomethine family, representing a newly developed chemical class of insecticides (Shen *et al.*, 2009). Pymetrozine is efficient toward aphids, whiteflies and plant hoppers in pest control programmes (Lashkari *et al.*, 2007). Pymetrozine acts by interfering in the regulation of the nervous system for feeding behavior, which results in death of the insect due to starvation a few days after application (Guoqing *et al.*, 2009). EFSA (2014) showed that pymetrozine exhibits low to moderate persistence in soil under laboratory aerobic conditions. It degraded by hydroxylation of the methylene group of the triazine ring and by oxidation. Degradation under field conditions

from 8 sites in Europe (Switzerland, France and Germany) resulted in pymetrozine dissipating with single first order DT<sub>50</sub> of 19.6 – 183 d (5sites) and biphasic DT<sub>50</sub> 3.81-10.3 d with associated DT<sub>90</sub> 167 ≥1000 d. Pymetrozine degraded from water mainly by distribution to the sediment. Single first order DT<sub>50</sub> of pymetrozine in the whole systems ranged between 289 and 495 d.

Phytoremediation is an environmentally sound technology for pollution prevention, control and remediation. One medicinal herb is plantain (*Plantago major* L., Plantaginaceae family). *Plantago major* has a wide geographic distribution through the temperate grasslands of the world and grows in a wide area of Europe, temperate regions of Asia and South Australia, North Africa and North America (Velasco-

\*Corresponding author: Tel. : +2001278194994

E-mail address: aaashalaby@zu.edu.eg :

Lezama *et al.*, 2006). It is a familiar perennial weed and may be found at roadsides, meadows, cultivated fields, waste areas, and even cracks in sidewalks and canal banks. The seeds and husks contain high levels of fiber; they expand and become highly gelatinous when soaked in water (Samuelsen, 2000; Sharifa *et al.*, 2008). *P. major* can accumulate a variety of inorganic metals or metabolize a variety of organic compounds including imidacloprid, chlorpyrifos, diethyl and dioctyl phthalates, azoxystrobin and cyanophos (Romeh, 2010; Romeh, 2013; Romeh and Hendawi, 2013; Romeh, 2014; Romeh, 2015a&b). Therefore, the goal of this work was to assess the capability of using *P. major* for the phytoremediation of pymetrozine -contaminated soil. The utilization of soluble silicon dioxide (SiO<sub>2</sub>) and improving agents such as the surfactants, Tween 80, hydroxypropyl-β-cyclodextrin (HPβCD) and liquid humic acid (HA) for enhancing the bioavailability and uptake of pymetrozine contaminated soil by *P. major* were evaluated.

## MATERIALS AND METHODS

### Pesticide and Plant

Pymetrozine (Technical grade 95.00%), 6-methyl-4-[(E)-pyridin-3-ylmethylideneamino]-2, 5-dihydro-1, 2, 4-triazin-3-one was acquired from Central Agricultural Pesticides Laboratory, Agricultural Research Center, Dokki – Giza, Egypt. The normal broadleaf plantain (*P. major*) was acquired as seedlings in phytoremediation experiment. Seeds of *P. major* grow naturally on meadow land in Zagazig University, Zagazig, Sharkia Governorate. After the seeds germinated, the seedlings (the age of the seedlings is 30 days and 10–12 cm height with 4 – 6 leaves) were collected for the experiment.

### Experimental Design

To assess the elimination of pymetrozine from the soil, two treatments were performed in this experiment, and each treatment consisted of five replicates: pymetrozine contaminated soil without plants, pymetrozine contaminated soil with *P. major* only (each pot contained one seedling of *P. major*). Experimental uptake was performed on soil in a pot experiment for 21-day exposure. Air-dried sieved clay loam soil (organic matter, 1.79%, pH 7.8, electric

conductivity 2.36) was obtained from Kamrona Village, Menia EL-Kamh district, Sharkia Governorate, Egypt, and then placed in plastic pots. The pots were provided with 0.5 kg of air dried soil. After planting, pymetrozine dissolved in water was spiked into the 150 ml of distilled water used for irrigation to obtain the original concentration of 20 mg/kg. The irrigation water containing pymetrozine was dropped into the pots with a caution to avoid the direct contact of plant leaves. Samples from exposed and control plants were collected through 1, 3, 7, 10, 14 and 21 days. Roots of plant from soil were rinsed in running tap water for 2 min and were blotted dry. The plants were dissected into individual roots and leaves then, 10 g of each leaves and roots and 20 g of soil were analyzed for the pesticide. All pots were watered with 50 ml tap water every 4 days or additionally watered when essential.

### Enhancing Agents for Phytoremediation of Soil Contaminated by Pymetrozine

To assess the removal of pymetrozine from the soil, eight treatments each consisting of five replicates were performed, as follows: Autoclaved soil contaminated with pymetrozine without plants. Soil contaminated by pymetrozine without plants. Contaminated soil with pymetrozine plus *P. major* only. Contaminated autoclaved soil with Pymetrozine containing *P. major* only. Contaminated soil with pymetrozine plus *P. major* and amended with soluble silicon dioxide (SiO<sub>2</sub>), these called silica, at 750 mg/l for a total concentration of 187.5 mg/kg. Contaminated soil with pymetrozine plus *P. major* and amended with 2-Hydroxypropyl-beta-cyclodextrin (HPβCD) at 1.0% (Chen *et al.*, 2010). Contaminated soil with pymetrozine plus *P. major* and amended with humic acid solution (HA) at 10 mg/l (humus WSG 90, produced by organist-Hungary). The latter concentration is reported to be the critical micelle concentration of HA (Guetzloff and Rice, 1994). Contaminated soil with pymetrozine plus *P. major* and amended with polyoxyethylene sorbitan monooleate (Tween 80) at 9.2 mg/l, corresponding to 0.5 critical micelle concentration (CMC), where the CMC of Tween 80 was determined as 13–45 mg/l (Edwards *et al.*, 1991; Mitton *et al.*, 2012).

In treatments (3)–(8), each pot contained one seedling of *P. major*. Each whole plant uptake experiment was performed in potted soil for 12 days as described above.

### Residues Analysis

The system followed in this work depended on QuEChERS strategy depicted by Anastassiades *et al.* (2003) and Lehotay *et al.* (2005). Soil tests were homogenized sieved (2 mm mesh) and air-dried at room temperature. A 10 g of homogenised soil was weighed into a 50 ml polypropylene tube, then 5 ml of HPLC water was included and the mixture was shaken for 1 min with a vortex apparatus. After that 10 ml of acetonitrile was included (acidified with acetic acid 1%) and the mixture was shaken for 1 min by hand and for 1 min with a vortex apparatus. Four grams of anhydrous magnesium sulfate, and 1 g sodium chloride were added, and the mixture was instantly hand-shaken for 30 sec., and centrifuged for 5 min at 4000 rpm in a Sigma 2–5 rotator (Sigma, Steinheim, Germany). At that point, a clean-up dispersive solid phase extraction step was implemented by including the supernatant (7.5 ml, *i.e.* 1.33 g of soil for every ml), to a 15 ml polypropylene tube that contained 1.125 g of MgSO<sub>4</sub> (150 mg MgSO<sub>4</sub> per ml of concentrate) and 0.225 g of C<sub>18</sub> (30 mg C<sub>18</sub> per ml of concentrate), hand-shaken for 30 sec. and centrifuged for 5 min at 4000 rpm (Asensio-Ramos *et al.*, 2010; Padilla-Sanchez *et al.*, 2010). For the determination of pymetrozine, 1.0 ml of the final concentrate was determined by HPLC.

A 10 g of fine macerated plant tissue (roots or leaves) was weighed into a 50 ml polypropylene tube. Then, 10 ml of acetonitrile was added (acidified with acetic acid 1%) and the mixture was hand-shaken for 2 min. Four grams anhydrous magnesium sulfate, and 1 g sodium chloride, was added, and the mixture was immediately hand-shaken for 30 sec. and centrifuged for 5 min at 4000 rpm. The clean-up step was implemented by adding the supernatant (7.5 ml, *i.e.* 1.33 g of plant tissue per ml), to a 15 ml polypropylene tube that included 1.125 g of MgSO<sub>4</sub> (150 mg MgSO<sub>4</sub> per ml of extract), and 0.188 g of PSA (25 mg PSA per ml of extract), hand-shaken for 30 sec. and centrifuged for 5 min at 4000 rpm (Anastassiades *et al.*,

2003; Lehotay *et al.*, 2005). For the determination of pymetrozine residues, 1.0 ml of the final extract was analyzed by HPLC.

### HPLC analysis

Soil, root, and leaf samples were analyzed for pymetrozine using high-performance liquid chromatography (HPLC) according to (Shen *et al.*, 2009). A 10 µl aliquot of final extract was injected into HPLC system and determined using a C<sub>18</sub> reversed-phase column [250 mm x 4.6 mm (i.d.)] and eluted isocratically with a mobile phase of water and acetonitrile (15:85, *V/V*) at the rate of 1 ml/min. The UV detection was adjusted at 299 nm. The retention time (RT's) was 2.63 minutes under these conditions. The performance of HPLC method was tested by evaluating quality parameters, such as recovery values.

### Recovery Samples

The extraction effectiveness of the analytical procedure was assessed by recovery experiments prepared in triplicate using the fortified blank leaves and roots of *P. major*, and soil samples at 0.5 mg/kg. The percent recoveries were respectively, 92.33, 90.12, and 88.5% in roots, leaves, and soils.

### Data Analysis

The rate of degradation ( $K_r$ ) and half-life ( $t_{1/2}$ ) was obtained according to Gomaa and Belal (1975) and Ashour (1976). The rate of degradation ( $K_r$ ) = 2.303 x slope. Half-life ( $t_{1/2}$ ) = 0.693/ $K_r$ .

## RESULTS AND DISCUSSION

### Phytoremediation of Soil Contaminated with Pymetrozine

Results obtained during the present investigation revealed that all experimental sets containing plantain removed a high amount of pymetrozine. As shown in Table 1, pymetrozine concentrations in soil containing *P. major* declined by 30.00–83.25% during 1 to 12 days of the experimental period, compared with 12.50–61.90% in un-planted soil. The half-life value ( $t_{1/2}$ ) of pymetrozine, estimated by first-order reaction, for soil plus *P. major* was found to be 5.65 days, compared with 8.58 days for

Table 1. Dissipation of pymetrozine in soil planted with *Plantago major* L

| Treatment                           | Days after application |              |               |              |              |              |      | T <sub>1/2</sub><br>(days) | K <sub>r</sub><br>(days) | AUC <sub>s</sub> mg/kg<br>(days) |
|-------------------------------------|------------------------|--------------|---------------|--------------|--------------|--------------|------|----------------------------|--------------------------|----------------------------------|
|                                     | 1                      | 2            | 4             | 6            | 8            | 12           |      |                            |                          |                                  |
| <b>In soil</b>                      |                        |              |               |              |              |              |      |                            |                          |                                  |
| mg/kg                               | 17.50                  | 15.30        | 13.52         | 11.40        | 9.05         | 7.62         | 8.58 | 0.08                       | 142.19                   |                                  |
| Loss (%)                            | 12.50                  | 23.50        | 32.40         | 43.00        | 54.75        | 61.90        |      |                            |                          |                                  |
| <b>In soil with <i>P. major</i></b> |                        |              |               |              |              |              |      |                            |                          |                                  |
| mg/kg                               | 14.00                  | 11.52        | 9.71          | 7.87         | 5.66         | 3.35         | 5.65 | 0.12                       | 97.45                    |                                  |
| Loss (%)                            | 30.00                  | 42.40        | 51.45         | 60.65        | 71.70        | 83.25        |      |                            |                          |                                  |
| <b>In <i>P. major</i> roots</b>     |                        |              |               |              |              |              |      |                            |                          |                                  |
| mg/kg                               | 38.74                  | 45.56        | 53.41         | 42.91        | 37.69        | 22.58        |      |                            |                          |                                  |
| <b>In <i>P. major</i> leaves</b>    |                        |              |               |              |              |              |      |                            |                          |                                  |
| mg/kg                               | 13.58                  | 44.95        | 58.08         | 37.72        | 28.13        | 16.06        |      |                            |                          |                                  |
| <b>Total uptake</b>                 | <b>52.32</b>           | <b>90.51</b> | <b>111.49</b> | <b>80.63</b> | <b>65.82</b> | <b>38.64</b> |      |                            |                          |                                  |

T<sub>1/2</sub>, half-life; k<sub>r</sub>, disappearance rate constant; AUC<sub>s</sub>, areas under the curve represent compound concentration during the period of study

soil alone (Table 1). Results with the disappearance rate constant (k<sub>r</sub>) values showed that pymetrozine had the highest k<sub>r</sub> value and lowest t<sub>1/2</sub> in soil with *P. major*, while pymetrozine had the shortest k<sub>r</sub> and longest t<sub>1/2</sub> in unplanted soil.

Pymetrozine concentrations (mg/kg) in various parts of *P. major* are found in Fig. 1. Concentrations of pymetrozine in the root were always higher than those in the leaves of the plant, with the exception of 4 days period; the translocation ratio was about 1.08 times higher than for pymetrozine contaminated soil with *P. major* roots (Fig. 1). In the roots of *P. major*, pymetrozine accumulated to reach the maximum levels through 4 days (53.41 mg/kg). Afterwards, concentration decreased gradually during the experiment (Table 1). Pymetrozine translocated into the leaves of *P. major* and reached the maximum through 4 days of treatment (58.08 mg/kg) then decreased until the end of exposure. Pymetrozine is highly moved in plants. It can be taken up from the soil as well as through the leaves (Fliückige *et al.*, 1992). Pymetrozine is both systemic and translaminar, making it highly mobile within plants (Wyss and Bolsinger, 1997). From bioassays and autoradiographic techniques of pymetrozine, it

has been shown that this systemic behavior originates not only from xylem but also from phloem mobility. After foliar application, the growing points of plants are protected by pymetrozine imports mainly from leaves. This indicates a high importance of phloem mobility for the systemic activity of pymetrozine for plant-sucking insects (Wyss and Bolsinger, 1997).

### Combination of *P. major* and Agents that Increase the Pymetrozine Availability in Soil

Changes in the levels of pymetrozine in soils subject to various treatments were measured following 1 to 12 days (Table 2) to estimate (1) plant's capacity to remove pymetrozine, (2) the role of various agents in increasing pymetrozine accessibility, microorganisms, plants and combination of plants with microorganisms, which lead to the dissipation of pymetrozine in the soil. Different agents contributed in increasing pymetrozine availability and reached the maximum enhancing through 4 - 8 days; therefore, the removal percentage of pymetrozine in a control group (C) was compared with removal percentages in experimental treatments after 4 days.




Fig. 1. Uptake and translocation of pymetrozine by *Plantago major* L. in soil

In C, 24.70% of the pymetrozine was degraded through natural biotic losses, whereas degradation percentages in treatments T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> were 32.40%, 49.50%, and 51.45%, respectively. We examined the contribution of three factors in the degradation of pymetrozine: biotic dissipation or natural hydrolyzation, as in the control group (C), microbial dissipation (T<sub>1</sub>), remediation by *P. major* (T<sub>2</sub>), and a combination of microbial degradation and *P. major* remediation (T<sub>3</sub>) (Table 2). Abiotic degradation was not considered in the treatments. The degradation of pymetrozine in the soil due to contribution by microorganisms (T<sub>1</sub>) was 7.70%, while contribution by plant alone (T<sub>2</sub>) and a combination of microorganisms and plants (T<sub>3</sub>) resulted in dissipation amounts of 24.80 % and 26.75%, respectively. The results showed that a combination of microorganisms plus plants (T<sub>3</sub>) is the most effective treatment for the dissipation of pymetrozine in soil, followed by plant alone (T<sub>2</sub>) and then microorganisms (T<sub>1</sub>), compared with natural dissipation processes.

It could be concluded that enhancement in pymetrozine dissipation in phytoremediation system could be achieved possibly due to the degradation induced by effects of plant with microorganisms in rhizosphere (Cheng *et al.*, 2007). The microbial-enhanced phytoremediation offers much potential for the remediation of organic pollutants in the soil (Chen *et al.*, 2010).

Pollutant-degrading bacteria may accelerate plants adaptation to contaminants by detoxifying contaminated soils during direct mineralization of these organic contaminants (Escalante-Espinosa *et al.*, 2005). In addition to, plant exudates increase the density and activity of potential pollutant-degrading bacteria in the area surrounding the roots (Siciliano and Germida, 1998). Romeh (2010) found that short-rod gram-negative bacteria that isolated from the water solution containing *P. major* was able to induce 93.34% loss of imidacloprid as a source of both nitrogen and carbon through 48 hr.

**Table 2. Contribution of agents that increase the pymetrozine availability in soil**

| Treatment                                | Days after treatments               |       |   |                  |
|--|-------------------------------------|-------|---|------------------|
|  | Amount added to the soil (20 mg/kg) | mg/kg | Removal (%)   | Contribution (%) |
| <b>1 day</b>                             |                                     |       |   |                  |
| C:In autoclaved soil                     | 18.55                               | 7.25  | 0.0   |                  |
| T1: In soil                              | 17.50                               | 12.50 | 5.25 microorganisms (T <sub>1</sub> -C)                   |                  |
| T2: Autoclaved soil with plantain        | 14.30                               | 28.50 | 21.25 Plant (T <sub>2</sub> -C)                           |                  |
| T3: Soil with plantain                   | 14.00                               | 30.00 | 22.75 Combination (T <sub>3</sub> -C)                     |                  |
| T4: Soil with plantain+ SiO <sub>2</sub> | 11.10                               | 44.50 | 14.50 SiO <sub>2</sub> (T <sub>4</sub> - T <sub>3</sub> ) |                  |
| T5: Soil with plantain+ HPβCD            | 11.20                               | 44.00 | 14.00 HPβCD (T <sub>5</sub> - T <sub>3</sub> )            |                  |
| T6: Soil with plantain+ HA               | 11.16                               | 44.20 | 14.20 Humic acid (T <sub>6</sub> - T <sub>3</sub> )       |                  |
| T7: Soil with plantain+ Tw 80            | 11.60                               | 42.00 | 12.00 Tw 80 (T <sub>7</sub> - T <sub>3</sub> )            |                  |
| <b>2 days</b>                            |                                     |       |   |                  |
| C:In autoclaved soil                     | 17.39                               | 13.05 | 0.0   |                  |
| T1: In soil                              | 15.30                               | 23.50 | 10.45 microorganisms (T <sub>1</sub> -C)                  |                  |
| T2: Autoclaved soil with plantain        | 12.00                               | 40.00 | 26.95 Plant (T <sub>2</sub> -C)                           |                  |
| T3: Soil with plantain                   | 11.52                               | 42.40 | 29.35 Combination (T <sub>3</sub> -C)                     |                  |
| T4: Soil with plantain+ SiO <sub>2</sub> | 7.42                                | 62.90 | 20.50 SiO <sub>2</sub> (T <sub>4</sub> - T <sub>3</sub> ) |                  |
| T5: Soil with plantain+ HPβCD            | 8.24                                | 58.80 | 16.40 HPβCD (T <sub>5</sub> - T <sub>3</sub> )            |                  |
| T6: Soil with plantain+ HA               | 7.71                                | 61.45 | 19.05 Humic acid (T <sub>6</sub> - T <sub>3</sub> )       |                  |
| T7: Soil with plantain+ Tw 80            | 8.80                                | 56.00 | 13.60 Tw 80 (T <sub>7</sub> - T <sub>3</sub> )            |                  |
| <b>4 days</b>                            |                                     |       |   |                  |
| C:In autoclaved soil                     | 15.06                               | 24.70 | 0.0   |                  |
| T1: In soil                              | 13.52                               | 32.40 | 7.70 microorganisms (T <sub>1</sub> -C)                   |                  |
| T2: Autoclaved soil with plantain        | 10.10                               | 49.50 | 24.80 Plant (T <sub>2</sub> -C)                           |                  |
| T3: Soil with plantain                   | 9.71                                | 51.45 | 26.75 Combination (T <sub>3</sub> -C)                     |                  |
| T4: Soil with plantain+ SiO <sub>2</sub> | 5.30                                | 73.50 | 22.05 SiO <sub>2</sub> (T <sub>4</sub> - T <sub>3</sub> ) |                  |
| T5: Soil with plantain+ HPβCD            | 6.45                                | 67.75 | 16.30 HPβCD (T <sub>5</sub> - T <sub>3</sub> )            |                  |
| T6: Soil with plantain+ HA               | 6.01                                | 69.95 | 18.50 Humic acid (T <sub>6</sub> - T <sub>3</sub> )       |                  |
| T7: Soil with plantain+ Tw 80            | 6.90                                | 65.50 | 14.05 Tw 80 (T <sub>7</sub> - T <sub>3</sub> )            |                  |
| <b>6 days</b>                            |                                     |       |   |                  |
| C:In autoclaved soil                     | 13.07                               | 34.65 | 0.0   |                  |
| T1: In soil                              | 11.40                               | 43.00 | 8.35 microorganisms (T <sub>1</sub> -C)                   |                  |
| T2: Autoclaved soil with plantain        | 8.29                                | 58.55 | 23.90 Plant (T <sub>2</sub> -C)                           |                  |
| T3: Soil with plantain                   | 7.87                                | 60.65 | 26.00 Combination (T <sub>3</sub> -C)                     |                  |
| T4: Soil with plantain+SiO <sub>2</sub>  | 3.47                                | 82.65 | 22.00 SiO <sub>2</sub> (T <sub>4</sub> - T <sub>3</sub> ) |                  |
| T5: Soil with plantain+ HPβCD            | 4.20                                | 79.00 | 18.35 HPβCD (T <sub>5</sub> - T <sub>3</sub> )            |                  |
| T6: Soil with plantain+ HA               | 3.90                                | 80.50 | 19.85 Humic acid (T <sub>6</sub> - T <sub>3</sub> )       |                  |
| T7: Soil with plantain+ Tw 80            | 4.70                                | 76.50 | 15.85 Tw 80 (T <sub>7</sub> - T <sub>3</sub> )            |                  |
| <b>8 days</b>                            |                                     |       |   |                  |
| C:In autoclaved soil                     | 11.50                               | 42.50 | 0.0   |                  |
| T1: In soil                              | 9.05                                | 54.75 | 12.25 microorganisms (T <sub>1</sub> -C)                  |                  |
| T2: Autoclaved soil with plantain        | 6.00                                | 70.00 | 27.50 Plant (T <sub>2</sub> -C)                           |                  |
| T3: Soil with plantain                   | 5.66                                | 71.70 | 29.20 Combination (T <sub>3</sub> -C)                     |                  |
| T4: Soil with plantain+SiO <sub>2</sub>  | 1.50                                | 92.50 | 20.80 SiO <sub>2</sub> (T <sub>4</sub> - T <sub>3</sub> ) |                  |
| T5: Soil with plantain+ HPβCD            | 1.72                                | 91.40 | 19.70 HPβCD (T <sub>5</sub> - T <sub>3</sub> )            |                  |
| T6: Soil with plantain+ HA               | 1.60                                | 92.50 | 20.80 Humic acid (T <sub>6</sub> - T <sub>3</sub> )       |                  |
| T7: Soil with plantain+ Tw 80            | 2.30                                | 88.50 | 16.80 Tw 80 (T <sub>7</sub> - T <sub>3</sub> )            |                  |

The potential of *P. major* for phytoremediation of pymetrozine in a the solution amended with SiO<sub>2</sub> (T<sub>4</sub>) was greater than those amended by other solubility enhancing agents, as measured by the removal of pymetrozine from contaminated soil at all experimental periods (Table 2). The percentage removal of pymetrozine through 4 days was ~73.50% in T<sub>4</sub>. In addition to, the removal percentages of pymetrozine in T<sub>6</sub>, T<sub>5</sub>, and T<sub>7</sub> were 69.95%, 67.75%, and 65.50%. The contribution of T<sub>4</sub> to the release of pymetrozine from soil was 22.05% while the contributions of T<sub>6</sub>, T<sub>5</sub>, and T<sub>7</sub> were 18.50%, 16.30%, and 14.05% (Table 2). Results in Table 2 indicate that, the maximum contribution of T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>5</sub>, T<sub>6</sub>, and, T<sub>7</sub> (12.25%, 27.50%, 29.20%, 19.70%, 20.80%, and 16.80%) to the release of pymetrozine from soil during 8 days, while, T<sub>4</sub> during 4 days cleared (22.05%) (Table 2).

The above results showed that, most of the pymetrozine disappearance in T<sub>2</sub> - T<sub>8</sub>, may be attributed to bioavailability of pymetrozine by enhancing agents. Romeh (2015b) found that the phytoremediation potential of *P. major* plus liquid silicon dioxide, SiO<sub>2</sub> was more potent than other solubility enhancing agents in removing cyanophos from the contaminated soil, improving removal percentage to 74.05% from 45.90% in soil with *P. major* only.

### Improvement the Phytoremediation of Soil Contaminated with Pymetrozine Using Soluble-Enhancing Agents

Pymetrozine amounts (mg/kg) in the roots and leaves of *P. major* are shown in Table 3. Pymetrozine accumulated in the roots of *P. major* to reach the maximum levels after 4 days (53.41 mg/kg). Afterwards, the concentration decreased gradually throughout the test (Table 3). In the leaves, pymetrozine translocated into the *P. major* leaves and reached the maximum after 4 days of exposure (58.08 mg/kg), then decreased until the end of testing. In autoclaved soil with *P. major*, pymetrozine accumulated in the roots of *P. major* to reach the maximum level after 4 days (52.67 mg/kg). Afterwards, concentration decreased gradually throughout the test (Table 3). Pymetrozine translocated into the leaves of *P. major* and reached the maximum through 4 days of exposure (57.72


mg/kg) then declined until the end of testing. Liquid silicon dioxide (SiO<sub>2</sub>) produced a synergistic effect on pymetrozine uptake and translocation. The phytoremediation potential of *P. major* plus SiO<sub>2</sub> was greater than that of other solubility-enhancing agents with respect to the removal of pymetrozine from contaminated soil within 1-12 days exposure (Table 3). Amending the soil with *P. major* containing SiO<sub>2</sub> resulted in a decrease in pymetrozine, half-life T<sub>1/2</sub>; k<sub>d</sub>, disappearance rate constant; AUCs, areas under the curve represent compound concentration during the period of study in the soil, and increased pymetrozine concentrations in plant leaves and roots (Table 3); it was detected in roots after one day then increased gradually and reached the maximum after 4 days (91.25 mg/kg) then decreased to the end of experiment. The pymetrozine concentration in *P. major* roots amended with SiO<sub>2</sub> increased by about 170.84% compared to those treated with pymetrozine in *P. major* roots alone through 4 days. The pymetrozine concentrations in *P. major* roots amended with SiO<sub>2</sub>, HA, HPβCD and Tween 80 achieved 91.25 mg/kg, 87.50 mg/kg, 86.12 mg/kg, and 58.04 mg/kg within 4 days, respectively. Pymetrozine concentrations in *P. major* roots alone reached 53.41 mg/kg (Fig. 2). Combination of *P. major* with enhancing agents for the purpose of pymetrozine phytoremediation in roots and leaves followed the order: SiO<sub>2</sub> > HA > HPβCD > Tween. SiO<sub>2</sub> addition caused increased pymetrozine translocation to leaves by about 322.83% compared with *P. major* leaves alone through 4 days. The concentrations of pymetrozine in *P. major* enhanced with SiO<sub>2</sub>, HA, HPβCD and Tween 80 reached 187.50 mg/kg, 176.87 mg/kg, 105.12 mg/kg, and 94.25 mg/kg through 4 days, respectively. Pymetrozine concentrations in *P. major* leaves alone reached 58.08 mg/kg.

The total pymetrozine accumulation in entire plant amended with SiO<sub>2</sub>, HA, HPβCD and Tween 80 arrived 278.75 mg/kg, 264.37 mg/kg, 191.24 mg/kg, and 152.29 mg/kg during 4 days compared with 111.49 mg/kg in *P. major* alone. Soil containing *P. major* and amended with SiO<sub>2</sub> caused a decrease in pymetrozine in soil and increases in plant leaves and roots, which is explained by the silicic acid Si (OH)<sub>4</sub> that enhance availability of compound from the soil to the roots and leaves of plant (Ma and Yamaji,

Table 3. Efficiency of solubility-enhancing agents in phytoremediation of pymetrozine contaminated soil

| Treatment  | Days after application |        |        |        |        |       | $t_{1/2}$<br>(days) | $K_r$<br>(days) | AUC, mg/kg<br>(days) |
|--|------------------------|--------|--------|--------|--------|-------|---------------------|-----------------|----------------------|
|  | 1                      | 2      | 4      | 6      | 8      | 12    |                     |                 |                      |
| In soil<br>mg/kg   | 17.50                  | 15.30  | 13.52  | 11.40  | 9.05   | 7.62  | 8.58                | 0.08            | 142.19               |
| Loss (%)   | 12.50                  | 23.50  | 32.40  | 43.00  | 54.75  | 61.90 |                     |                 |                      |
| In soil with <i>P. major</i><br>mg/kg                    | 14.00                  | 11.52  | 9.71   | 7.87   | 5.66   | 3.35  | 5.65                | 0.12            | 97.45                |
| Loss (%)   | 30.00                  | 42.40  | 51.45  | 60.65  | 71.70  | 83.25 |                     |                 |                      |
| In <i>P. major</i> roots<br>mg/kg                        | 38.74                  | 45.56  | 53.41  | 42.91  | 37.69  | 22.58 |                     |                 |                      |
| In <i>P. major</i> leaves<br>mg/kg                       | 13.58                  | 44.95  | 58.08  | 37.72  | 28.13  | 16.06 |                     |                 |                      |
| Total uptake   | 52.32                  | 90.51  | 111.49 | 80.63  | 65.82  | 38.64 |                     |                 |                      |
| In soil with <i>P. major</i> + HP $\beta$ CD<br>mg/kg    | 11.20                  | 8.24   | 6.45   | 4.20   | 1.72   | 0.16  | 3.10                | 0.22            | 57.84                |
| Loss (%)   | 44.00                  | 58.80  | 67.75  | 79.00  | 91.40  | 99.20 |                     |                 |                      |
| In <i>P. major</i> roots<br>mg/kg                        | 56.22                  | 72.63  | 86.12  | 70.19  | 60.05  | 35.09 |                     |                 |                      |
| In <i>P. major</i> leaves<br>mg/kg                       | 43.15                  | 80.12  | 105.12 | 65.43  | 47.66  | 26.85 |                     |                 |                      |
| Total uptake   | 99.37                  | 152.75 | 191.24 | 135.62 | 107.71 | 61.94 |                     |                 |                      |
| In soil with <i>P. major</i> + Tween 80<br>mg/kg         | 11.60                  | 8.80   | 6.90   | 4.70   | 2.30   | 0.60  | 3.41                | 0.20            | 63.28                |
| Loss (%)   | 42.00                  | 56.00  | 65.50  | 76.50  | 88.50  | 97.00 |                     |                 |                      |
| In <i>P. major</i> roots<br>mg/kg                        | 46.26                  | 53.80  | 58.04  | 47.61  | 40.42  | 30.92 |                     |                 |                      |
| In <i>P. major</i> leaves<br>mg/kg                       | 37.09                  | 64.74  | 94.25  | 57.62  | 36.82  | 20.13 |                     |                 |                      |
| Total uptake   | 83.35                  | 118.54 | 152.29 | 105.23 | 77.24  | 51.05 |                     |                 |                      |
| In soil with <i>p. major</i> + SiO <sub>2</sub><br>mg/kg | 11.10                  | 7.42   | 5.30   | 3.47   | 1.50   | 0.01  | 2.66                | 0.26            | 51.05                |
| Loss (%)   | 44.50                  | 62.90  | 73.50  | 82.65  | 92.50  | 99.95 |                     |                 |                      |
| In <i>P. major</i> roots<br>mg/kg                        | 45.41                  | 65.62  | 91.25  | 56.25  | 35.00  | 0.00  |                     |                 |                      |
| In <i>P. major</i> leaves<br>mg/kg                       | 125.25                 | 156.21 | 187.50 | 133.75 | 90.37  | 6.25  |                     |                 |                      |
| Total uptake   | 170.66                 | 221.83 | 278.75 | 190.00 | 125.37 | 6.25  |                     |                 |                      |
| In soil with <i>P. major</i> + HA<br>mg/kg               | 11.16                  | 7.71   | 6.01   | 3.90   | 1.60   | 0.05  | 2.92                | 0.23            | 54.65                |
| Loss (%)   | 44.20                  | 61.45  | 69.95  | 80.50  | 92.00  | 99.75 |                     |                 |                      |
| In <i>P. major</i> roots<br>mg/kg                        | 55.61                  | 68.75  | 87.50  | 22.50  | 15.00  | 0.00  |                     |                 |                      |
| In <i>P. major</i> leaves<br>mg/kg                       | 36.25                  | 126.25 | 176.87 | 131.25 | 56.25  | 12.50 |                     |                 |                      |
| Total uptake   | 91.86                  | 195.00 | 264.37 | 153.75 | 71.25  | 12.50 |                     |                 |                      |
| In autoclaved soil with <i>P. major</i><br>mg/kg         | 14.30                  | 12.00  | 10.10  | 8.29   | 6.00   | 3.90  | 6.01                | 0.11            | 102.44               |
| Loss (%)   | 28.50                  | 40.00  | 49.50  | 58.55  | 70.00  | 80.50 |                     |                 |                      |
| In <i>P. major</i> roots<br>mg/kg                        | 37.31                  | 44.86  | 52.67  | 41.70  | 36.46  | 21.99 |                     |                 |                      |
| In <i>P. major</i> leaves<br>mg/kg                       | 17.34                  | 43.46  | 57.72  | 36.88  | 27.26  | 15.68 |                     |                 |                      |
| Total uptake   | 54.65                  | 88.32  | 110.39 | 78.58  | 63.72  | 37.67 |                     |                 |                      |
| In autoclaved soil<br>mg/kg                              | 18.55                  | 17.39  | 15.06  | 13.07  | 11.50  | 9.20  | 10.48               | 0.06            | 163.52               |
| Loss (%)   | 7.25                   | 13.05  | 24.70  | 34.65  | 42.50  | 54.00 |                     |                 |                      |





**Fig. 2.** Efficiency of solubility enhancing agents in phytoremediation of pymetrozine contaminated soil through 1-12 days of exposure; HPβCD: 3- hydroxyl, B-cyclodextrin, Tw 80: tween 80, SiO<sub>2</sub>: silicon dioxide, HA: humic acid; (A): 1 day, (B): 2 days, (C): 4 days, (D): 6 days, (E): 8 days, (F): 12 days

2006). In addition to, humic acid (HA) caused a synergistic effect on pymetrozine uptake and translocation. Humic acid (HA) could act as a natural surfactant for enhancing the bioavailability of pymetrozine -contaminated soil. The combined effects of plants plus HA resulted in increasing the removal amount of pymetrozine from the soil, enhancing percentage degradation of 69.95% from 51.45% in soil with *P. major* only through 4 days of treatment (Table 3). The enhanced degradation performance for pymetrozine observed might be due to an increase in microbial activities and bioavailable in soils caused by the combined effects of plants and HA. Also, results in Table 3 show that HA was a little bit better than Tween 80 in decreasing pymetrozine - polluted soil. The surfactant activity of HA was found to increase solubility of organic contaminants on soils, hence enabling desorption-remediation of polycyclic aromatic hydrocarbon (PAH) (Holman *et al.*, 2002), In addition to, the fraction of humic substances remaining in the soil cause a favorable role in the growth of plant and microbes and thus useful in the full recuperation of treated soils (Nardi *et al.*, 2002). Several reports discussed the use of surfactant-enhanced phytoremediation (Mitton *et al.*, 2012). This work also showed that enhancing agents such as the surfactants, HP $\beta$ CD, natural HA and Tween 80 removes amounts convergent of pymetrozine from a contaminated soil (Fig. 2). HP $\beta$ CD helped in recovering 86.12 mg/kg and 105.12 mg/kg pymetrozine from contaminated soil by *P. major* roots and leaves within 4 days of treatment (Fig. 2). This increase in the removal of pymetrozine in soils amended with HP $\beta$ CD may be due to the formation of an inclusion complex with pymetrozine (Villaverde *et al.*, 2006). The use of plants plus surfactants has been proposed for improving phytoremediation strategies. These methods are based on the ability of agents to enhance the water solubility of hydrophobic organic compounds (HOCs) and to promote desorption, bio-degradation and phytoremediation processes (Wang and Keller, 2009).

## Conclusion

The results showed that *P. major* removes efficiently of pymetrozine residues in soil and

has a potential activity for pesticides phytoremediation. Also, this study indicates that SiO<sub>2</sub> can enhance the phytoremediation effectiveness for pymetrozine.

## REFERENCES

- Anastassiades, M., S.J. Lehotay, D. Stajnbaher and F. Schenck (2003). Fast and easy multiresidue method employing acetonitrile extraction/partitioning and "dispersive solid-phase extraction" for the determination of pesticide residues in produce. *J. AOAC. Int.*, 86: 412–431.
- Asensio-Ramos, M., J. Hernández-Borges, L.M. Ravelo-Pérez and M.A. Rodríguez-Delgado (2010). Evaluation of a modified QuEChERS method for the extraction of pesticides from agricultural, ornamental and forestal soils. *Anal. Bioanal. Chem.*, 396 (6): 2307-2319.
- Ashour, M.B.A. (1976). Residual effect of certain soil pesticides on some field crops. M.Sc. Thesis, Fac. Agric., Zagazig Univ., Egypt.
- Chen, Y., X. Tang, S.A. Cheema, W. Liu and C. Shen (2010). Beta cyclodextrin enhanced phytoremediation of aged PCBs-contaminated soil from waste recycling area. *J. Environ. Monit.*, 12: 1482–1489.
- Cheng, S., X. Jin, X. Huiping, Z. Liping and W. Zhenbin (2007). Technical note phytoremediation of triazophos by *canna indica* L. in a hydroponic system. *Int. J. Phytoremediation*, 9: 453–463.
- Edwards, D.A., R.G. Luthy and Z. Liu (1991). Solubilization of polycyclic aromatic hydrocarbons in micellar nonionic surfactant solutions. *Environ. Sci. Technol.*, 25: 127–133.
- EFSA (European Food Safety Authority) (2014). Conclusion on the peer review of the pesticide risk assessment of the active substance pymetrozine. *EFSA J.*, 12 (9): 3817.
- Escalante-Espinosa, E., M.E. Gallegos-Martinez, E. Favela-Torres and M. Gutiérrez-Rojas (2005). Improvement of the hydrocarbon phytoremediation rate by

- Cyperus laxus* Lam. inoculated with a microbial consortium in a model system, *Chemosphere*, 59: 405–413.
- Fliickige, C.R., H. Kristinson, R. Senn, A. Rindlisbacher, H. Buholzer and G. Voss (1992). A novel agent to control aphids and whiteflies. *Brighton Crop Protection Conf. - Pests and Diseases*, 1: 43-50.
- Gomaa, E.A. and M.H. Belal (1975). Determination of dimethoate residues in some vegetables and cotton plant. *Zagazig J. Agric. Res.*, 2: 215–221.
- Guetzloff, T.F. and J.A. Rice (1994). Does humic acid form amicelle?. *Sci. Total Environ.*, 152: 31–35.
- Guoqing, S., H. Xuan and H. Yinan (2009). Kinetic study of the degradation of the insecticide pymetrozine in a vegetable-field ecosystem. *J. Hazard Mater.*, 164: 497–501.
- Holman, H., K. Nieman, D. Sorensen, C. Miller, M.C. Martin, T. Borch, W. McKinney and R. Sims (2002). Catalysis of PAH biodegradation by humic acid shown in synchrotron infrared studies. *Environ. Sci. Technol.*, 36: 1276–1280.
- Lashkari, M.R., A. Sshragard and M. Ghadamyari (2007). Sublethal effects of imidacloprid and pymetrozine on population growth parameters of cabbage aphid, *Brevicoryne brassicae* on rapeseed, *Brassica napus* L. *Insect Sci.*, 14: 207.
- Lehotay, S.J., K. Mas̃tovska'and S.J. Yun (2005). Evaluation of two fast and easy methods for pesticide residue analysis in fatty food matrixes. *J. AOAC. Int.*, 88: 630–638.
- Ma, J.F. and N. Yamaji (2006). Silicon uptake and accumulation in higher plants. *Trends Plant Sci.*, 11: 392–397.
- Mitton, F.M., G. Mariana, P. Aranzazu and S. Karina (2012). Effects of amendments on soil availability and phytoremediation potential of aged p, p-DDT, p, p<sub>-</sub>-DDE and p, p-DDD residues by willow plants (*Salix* sp.). *J. Hazard. Mater.*, 203–204: 62–68.
- Nardi, S., E. Sessi, D. Pizzeghello, A. Sturaro, R. Rella and G. Pargoli (2002). Biological activity of soil organic matter mobilized by root exudates. *Chemosphere*, 46: 1075–1081.
- Padilla-Sanchez, J.A., P. Plaza-Bolanos, R. Romero-Gonzalez, A. Garrido-Frenich and J.L.M. Vidal (2010). Application of a quick, easy, cheap, effective, rugged and safe-based method for the simultaneous extraction of chlorophenols, alkylphenols, nitrophenols and cresols in agricultural soils, analyzed by using gas chromatography–triple quadrupole-mass spectrometry/mass spectrometry. *J. Chromatography A*. 1217 (36): 5724 – 5731.
- Romeh, A.A. (2010). Phytoremediation of water and soil contaminated with imidacloprid pesticide by *plantago major* L. *Int. J. Phytoremediation*, 12: 188–199.
- Romeh, A.A. (2013). Diethyl phthalate and dioctyl phthalate in *Plantago major* L. *Afri. J. Agric. Res.*, 8: 4360-4364.
- Romeh, A.A. (2014). Phytoremediation of cyanophos insecticide by *Plantago major* L. in water. *J. Environ. Health Sci. and Eng.*, 12: 38.
- Romeh, A.A. (2015a). Evaluation of the phytoremediation potential of three plant species for azoxystrobin-contaminated soil. *Int. J. Environ. Sci. Technol.*, 12 : 3509-3518.
- Romeh, A.A. (2015b). Enhancing agents for phytoremediation of soil contaminated by cyanophos. *Ecotoxicol. and Environ. Safety* 117: 124–131.
- Romeh, A.A. and M.Y. Hendawi (2013). Chlorpyrifos insecticide uptake by plantain from polluted water and soil. *Environ. Chem. Lett.*, 11:163–170.
- Samuelsen, A.B. (2000). The traditional uses, chemical constituents and biological activities of *Plantago major* L.: a review. *J Ethnopharmacol*, 71: 1–21.
- Sharifa, A.A., Y.L. Neoh, M.I. Iswadi, O. Khairul, M.M. Abdul Halim, A. Jamaludin Mohamed and H.L. Hing (2008). Effects of methanol, ethanol and aqueous extract of *Plantago major* on gram positive bacteria, gram negative bacteria and yeast. *Ann. Microsc.*, 8: 42–44.

- Shen, G., X. Hu and Y. Hu (2009). Kinetic study of the degradation of the insecticide pymetrozine in a vegetable-field ecosystem. *J. Hazard. Mater.*, 164 (2-3): 497–501.
- Siciliano, S.D. and J.J. Germida (1998). Mechanisms of phytoremediation: biochemical and ecological interactions between plants and bacteria. *Environ. Rev.*, 6 (1): 65–79.
- Velasco-Lezama, R., R. Tapia-Aguilar, R. Román-Ramos, E. Vega-Avila and M.S. Perez-Gutierrez (2006). Effect of *Plantago major* on cell proliferation *in vitro*. *J. Ethnopharmacol.*, 103 (1): 36–42.
- Villaverde, J., C. Maqueda and E. Morillo (2006). Effect of the simultaneous addition of cyclodextrin and the herbicide norflurazon on its adsorption and movement on soils. *J. Agric. and Food Chem.*, 54: 4766–4772.
- Wang, P. and A.A. Keller (2009). Partitioning of hydrophobic pesticides within a soil–water–anionic surfactant system, *Water Res.*, 43: 706–714.
- Wyss, P. and M. Bolsinger (1997). Translocation of pymetrozine in plants. *Pestic. Sci.*, 50: 195–202.

### المواد المحسنة للعلاج النباتي للتربة الملوثة بالمبيد الحشري بيميتروزين

على عطا على شلبي<sup>١</sup> - على أحمد علي أيوب<sup>١</sup>

أحمد علي رميح<sup>٢</sup> - محمود راضي عبد الرحمن رمضان<sup>٢</sup>

١- قسم وقاية النبات - كلية الزراعة - جامعة الزقازيق - مصر

٢- قسم الإنتاج النباتي - كلية التكنولوجيا والتنمية - جامعة الزقازيق - مصر

تم تصميم هذا البحث لدراسة أهمية استخدام نبات لسان الحمل (*Plantago major* L.) في المعالجة النباتية للتربة الملوثة بالمبيد الحشري بيميتروزين، وتقييم تأثير إضافة hydroxypropyl -  $\beta$  - cyclodextrin (HP $\beta$ CD) ، silicon dioxide (SiO<sub>2</sub>) ، Tween 80 و liquid humic acid (HA) لتحسين كفاءة التقاط المبيد من التربة الملوثة باستخدام نبات لسان الحمل، وأوضحت النتائج أن تركيز المبيد قد انخفض في التربة بواسطة النبات إلى ٣٠,٠٠ - ٨٣,٢٥% خلال فترة التجربة ١ - ١٢ يوماً مقارنةً بانخفاض قدره ١٢,٥٠ - ٦١,٩٠% في تربة الكنترول (بدون النبات) ، كما أمتص المبيد بجذور وأوراق نبات لسان الحمل ليصل إلى ٥٣,٤١ مجم/كجم ، ٥٨,٠٨ مجم/كجم بعد ٤ أيام، أدت إضافة أكسيد السيليكون إلى زيادة التقاط الجذور للمبيد وانتقاله إلى الأوراق بحوالي ١٧٠,٨٤% و ٣٢٢,٨٣% على التوالي مقارنة بتجربة استخدام النبات فقط، وأمكن ترتيب تأثير إضافة المواد المحسنة مع النبات تنازلياً كما يلي: SiO<sub>2</sub> > HA > HP $\beta$ CD > Tween80 ، ولقد أوضحت هذه الدراسة أنه يمكن إضافة SiO<sub>2</sub> لزيادة المعالجة النباتية للتربة الملوثة بهذا المبيد.

#### المحكمون :

- ١- أ.د. محمد عبد الله صالح رئيس بحوث متفرغ بالمعمل المركزي للمبيدات - مركز البحوث الزراعية - وزارة الزراعة.
- ٢- أ.د. رفعت مصطفى محمد شريف أستاذ المبيدات المتفرغ - كلية الزراعة - جامعة الزقازيق.