

# Impact of Different Application Methods and Concentrations of Ascorbic Acid on Sugar Beet under Salinity Stress Conditions

Eman M. Abdel Fatah<sup>1</sup> and K.A. Sadek<sup>2</sup>

<sup>1</sup>Physiol. and Chem. Res. Dept., Sugar Crops Res. Inst., Agric. Res. Center, Giza, Egypt

<sup>2</sup>Agron. Res. Dept., Sugar Crops Res. Inst., Agric. Res. Center, Giza, Egypt

## ABSTRACT

Salinity is one of the major abiotic stresses which affected on plant growth and reduces plant productivity. Antioxidants compounds considered as one of the methods and strategies to ameliorate the deleterious effects of salinity stress on plants to reduce the effect of oxidative stress. Field experiment was carried out at Sinnuris District which located mainly adjacent to Lake Qarun (saline soil) at El Fayoum Governorate (latitude 29°20' and 29°30' N and longitude 30°43' and 30°56' E) during 2015/2016 and 2016/2017 growing seasons to study the effect of relation between three application methods (seed soaking, foliar spraying and seed soaking plus foliar spraying together) and four concentrations (without treatment, tap water, 100 ppm and 200 ppm of ascorbic acid) of ascorbic acid on some biochemical and physiological characteristics, antioxidant enzyme activity as well as growth, yield and quality of sugar beet (*Beta vulgaris* var. *saccharifera*, L.) under salt stress conditions. The treatments were arranged in a strip-split plot design with three replications. The vertical plots were occupied with three application methods while, the horizontal plots were devoted to four ascorbic acid treatments.

Results showed that application of soaking sugar beet seed in ascorbic acid (AsA) plus foliar spraying at 50 and 70 days from sowing significantly increased chlorophyll a, b and carotenoides, LAI, catalase (CAT) and superoxide dismutase (SOD) activities, yield components and sucrose % in both seasons while juice impurities (K, Na and  $\alpha$ -Amino-N) were decreased and significantly increased root yield by (9.19, 9.63%) and sugar yield by (13.43, 15.59%) compared to AsA applied as a seed soaking or foliar spray in the 1<sup>st</sup> season and the same trend was attained in the 2<sup>nd</sup> season.

Applying 200 ppm of ascorbic acid was more effective and significantly increased root and sugar yields and sucrose % and caused more reduction in juice impurities in comparison to the control, in both seasons.

Higher stomata area, lower stomata density and stomata closure% were recorded by seed soaking plus foliar spraying with 200 ppm AsA treatment for upper and lower surface of sugar beet leaf.

Applying seed soaking plus foliar spraying by 200 ppm of ascorbic acid significantly increased root yield by (6.99 and 4.54 ton/fed) and sugar yield by (2.19 and 2.06 ton/fed) over that gained by untreated plants in the 1<sup>st</sup> and 2<sup>nd</sup> seasons.

**Keywords:** Sugar beet, salinity, ascorbic acid, application methods, enzyme activities, stomata.

## INTRODUCTION

Salt stress is one of the world wide abiotic factors which caused inhibition of soil micro-organisms' and cause damage in plant growth and its yield. In most of the cases, the negative effects of salinity have been attributed to increase in Na<sup>+</sup> and Cl<sup>-</sup> ions. In different plants, salinity at higher levels may cause membrane damage, nutrient imbalance, altered levels of growth regulators, enzymatic inhibition and metabolic dysfunction, including photosynthesis which ultimately leads to plant death (Hasanuzzaman *et al.* 2012). Biochemical and molecular studies of salt stress responses in plants have revealed significant increases of reactive oxygen species (ROS), including singlet oxygen ( $^1O_2$ ), superoxide ( $O_2^-$ ), hydroxyl radical (OH $\cdot$ ) and hydrogen peroxide ( $H_2O_2$ ) (Ahmad *et al.* 2012). These radicals cause peroxidation of membrane lipids, destruction of skeleton structure and dysfunction of cell (Yin, *et.al* 2011). El-Fayoum Governorate is a natural depression in the Eocene limestone plateau located at about 90-km southwest of Cairo. Sinnuris District in the eastern-north part of Fayoum depression. These areas are located mainly adjacent to Lake Qarun which covers an area of about 210 km<sup>2</sup> of salty water of average 33.9–37.6 g.L<sup>-1</sup> (FWMP, 1999) however the increase in salinity indicates annual salt accumulation in Qarun is still as much as 70–85 mln.kg/year (Shadrin *et al.* 2016) and

thus the capacity of the salt extraction process must be increased to prevent further salinity increases in the future ,where, poor drainage conditions, shallow saline groundwater, and clayey soils with low permeability occurred, soils are also very saline. The storing capacity of the lake and its salinity are considered main limiting factors for agriculture development in Fayoum Governorate (Euroconsult, 1992). In order to tackle salinity problem, choice of crop that can utilize or tolerate high salt levels in soil and/or irrigation water should be included in agricultural programs. Sugar beet is one of the most salt tolerant crops, but with less tolerance during germination, emergence and seedling stage (Maas, 1986). Rhoades and Loveday (1990) indicated that, sugar yield of sugar beet was not affected by salinity up to an electrical conductivity value of soil paste extract (ECe) of 7dSm.

Over-production of reactive oxygen species (ROS) in plants under stress conditions is a common phenomenon. Plants tend to counter this problem through their ability to synthesize ROS neutralizing substances including non-enzymatic and enzymatic antioxidants. In this context, ascorbic acid (Vitamins C) is one of the universal non-enzymatic antioxidants having substantial potential of not only scavenging ROS, but also modulating a number of fundamental functions in plants both under stress and non-stress conditions. Most efficient role of AsA is to protect lipids and proteins against salinity -induced oxidative

adversaries (Naz *et al.*, 2016), enhancing plant growth, rate of photosynthesis, transpiration, and oxidative defense potential and photosynthetic pigments, it also has a role in cell division and cell enlargement, stomatal regulations (Barth *et al.*, 2006), as well as acting as a co-factor for many enzymes. Additionally, ascorbic acid protects metabolic processes against H<sub>2</sub>O<sub>2</sub> and toxic derivatives of oxygen which affect various enzyme activities and decreases the damage caused through oxidative processes by functioning in synergy with other stabilizing membranes and antioxidants (Shao *et al.*, 2008). Foliar application of AA at 200 mg l<sup>-1</sup> decreased the adverse effect of salinity that was accompanied by a significant increase in plant growth of flax cultivars (El-Hariri *et al.* 2010). Applied ASA (250 mg/L) as (seed soaking, foliar spraying or seed soaking and foliar spraying together) were the most effective in increasing all yield and its parameters compared with water as control and decreased the harmful effects of salinity stress of faba bean plants (Sakr *et al.* 2014). Abido *et al.* (2015) showed that foliar spraying with 150 ppm of ascorbic acid improved growth attributes, yield and its components of sugar beet. Samin *et al.* (2016) observed that treated *Solanum melongena* with ascorbic acid (100ppm and 200ppm) showed increase in plant height, root length, in saline environments as compared to control plants. Elsayed (2017) reported that, the interaction between salinity stress and seed soaking (0.5 mM ASA) was significant for whole plant and root fresh and dry weights, and chlorophyll content index of sugar beet. Jagdish *et al.* (2017) recorded that, foliar spraying 1.0 mM ascorbic acid, improved the SOD activity, total chlorophyll, yield attributes, of mungbean (*Vigna radiata* L.) under salt stress condition.

The current study took-place to study the effects of different Ascorbic acid (ASA) concentrations and methods of their application on yield and quality of sugar beet as under Sinnuris District conditions, Fayoum governorate, Egypt.

### Materials and Methods

Field experiments was conducted at Sinnuris District, Fayoum Governorate (latitude 29°20 and

29°30 N and longitude 30°43 and 30°56 E) during 2015/2016 and 2016/2017 seasons to evaluated different ascorbic acid (AsA) application methods and concentrations and their interactions on some Biochemical and physiological Characteristics, antioxidant enzyme activity as well as growth, yield and quality of sugar beet (*Beta vulgaris* var. saccharifera, L.) grown under salinity stress condition. This area is mainly located adjacent to Lake Qarun where poor drainage conditions, shallow saline groundwater, and clayey soils with low permeability occurred. In the study area, soil is also very saline (Fig. 1). The present work included 12 treatments, represented the combinations of three application methods (seed soaking, foliar spraying and seed soaking plus foliar spraying) and four ascorbic acid treatments (without treatments (control), tap water, 100 and 200 ppm). The treatments were arranged in a strip plot design in three replicates. The vertical plots were occupied with three application methods while, the horizontal plots were devoted to the four ascorbic acid treatments. Each experiment basic unit area was 15.00 m<sup>2</sup> including five ridges of 0.60 m apart and 5.0 m long, with 20-cm hill spacing.

Seeds soaking treatments were done for 12 hours in water or ASA, and then air dried before sowing, whereas foliar spraying treatment or seeds soaking and foliar spraying together treatment were sprayed with foliar solution volume (350 L/faddan) by hand sprayer (for experimental plots) after 50 and 70 days from sowing (DFS) until saturation point. Ascorbic acid was obtained from El- Gomhouria CO, for Trading Chemicals and Medications. Seeds of sugar beet multi-germ variety "Magribel" were sown at 20<sup>th</sup> and 26<sup>th</sup> October, while harvesting was done 7 months later in the first and second seasons. Soil samples were taken at random from the experimental site at a depth of 0-30 cm from soil surface. Mechanical and chemical analyses were carried out according to the standard methods (Ryan *et al.*, 1996) (Table 1). The recommended agricultural practices for growing sugar beet were followed according recommendation.

**Table.1 Physical properties and some chemical analysis of experimental soils for 2015/2016 and 2016/2017 seasons**

| Physical properties              |                  |                  |                 |                                  | Chemical properties           |                 |                              |                              |
|----------------------------------|------------------|------------------|-----------------|----------------------------------|-------------------------------|-----------------|------------------------------|------------------------------|
| Seasons                          | Fine sand %      | Silt %           | Clay %          | Texture                          | pH (1:2:5)                    | EC (ds/m)       |                              |                              |
| 2015/2016                        | 24.70            | 37.10            | 38.20           | Clay loam                        | 8.16                          | 8.60            |                              |                              |
| 2016/2017                        | 25.30            | 36.90            | 37.80           |                                  | 8.14                          | 8.50            |                              |                              |
| Soluble Cations (meq/l)          |                  |                  |                 |                                  | Soluble Anions (meq/l)        |                 |                              |                              |
|                                  | Ca <sup>++</sup> | Mg <sup>++</sup> | Na <sup>+</sup> | K <sup>+</sup>                   | HCO <sub>3</sub> <sup>-</sup> | Cl <sup>-</sup> | SO <sub>4</sub> <sup>-</sup> | CO <sub>3</sub> <sup>-</sup> |
| 2015/2016                        | 25.21            | 19.19            | 40.20           | 1.25                             | 6.69                          | 70.90           | 8.26                         | -                            |
| 2016/2017                        | 25.70            | 19.55            | 39.60           | 1.22                             | 6.72                          | 70.71           | 8.64                         | -                            |
| Available Macronutrients (mg/kg) |                  |                  |                 | Available Micronutrients (mg/kg) |                               |                 |                              |                              |
|                                  | N                | P                | K               | Fe                               | Mn                            | Zn              | Cu                           |                              |
| 2015/2016                        | 39.60            | 5.25             | 147.50          | 12.25                            | 6.50                          | 1.20            | 4.50                         |                              |
| 2016/2017                        | 42.50            | 4.75             | 145.00          | 11.50                            | 7.20                          | 0.85            | 4.25                         |                              |




Fig. 1: Location map of the study area

### Studied traits

After 120 days from sowing, random samples of sugar beet plants were taken from each sub plot to determine the following traits:

1. **Leaf area index (LAI)** = leaf area per plant ( $\text{cm}^2$ ) / Plant ground area ( $\text{cm}^2$ ).
2. **Photosynthetic pigments**, *i.e.* chlorophyll a, b and carotenoides (mg/g leaf fresh weight), which were determined according to the method of Wettstein (1957).
3. **Antioxidant Enzyme Activities:** Super oxide dismutase (SOD) and Catalase (CAT) activities (Unit  $\text{g}^{-1}$  protein) were estimated by the method of Beyer and Fridovich (1987).
4. **Leaf stomata measurements:** The morphological changes of stomata (stomata density (No.  $\text{mm}^2$ ), area ( $\mu\text{m}$ ) and stomatal closure (%) for adaxial (upper) and abaxial (lower) surface of fully expanded leaves from mature leaf from different treatments were measured through Transmission Electronic Microscope (TEM) Model JEOL JEM-1400 (Jeol Ltd., Tokyo, Japan) linked with the software program at TEM lab (FA-CURP) Faculty of Agriculture, Cairo University Research Park.

At harvest time (210 day from sowing), ten plants were randomly chosen from each plot to determine the following traits:

- 1- Root length (cm).
- 2- Root diameter (cm).
- 3- Root fresh weight g/plant
- 4- Weight per plot was obtained and used to calculate root yield per-faddan.

5- Sugar yield ( $\text{ton fad}^{-1}$ ) = Root yield (ton) x Extracted sugar %.

### Quality measurements

1. Sucrose percentage was determined by using Saccharometer on an extract of fresh macerated root according to the procedure of the El-Fayoum Sugar Company (Le-Docte,1972), juice impurities contents (K, Na and  $\alpha$ -amino N) were estimated as meq/100 g beet were determined by using Analyzer –HG in reception laboratory in El-Fayoum Company.
2. Sugar lost to molasses percentage (SLM %) was calculated by the following formula, according to Devillers (1988):  

$$\text{SLM}\% = 0.29 + (\text{Na} + \text{K}) 0.343 + 0.094 (\alpha\text{-amino N}).$$

### Statistical analysis:

The collected data of the studied traits were statistically analyzed as shown by Snedecor and Cochran (1980). Treatments means were compared using LSD test at 5% of probability.

## RESULTS AND DISCUSSIONS

### Photosynthetic pigments, leaf area index and antioxidant enzymes activities:

Results illustrated in Table (2) showed that, different application methods (seed soaking, foliar spraying and seed soaking plus foliar spraying) significantly affected leaf area index (LAI), photosynthetic pigments, antioxidant enzyme activities, in both seasons. Under saline soil conditions, it was noticed that, treatment of soaking sugar beet seeds before sowing in ascorbic acid

**Table 2: Photosynthetic pigments and antioxidant enzymes activities of sugar beet as affected by different application methods and concentrations of ascorbic acid during 2015/2016 and 2016/2017 seasons**

| Treatments                       | Photosynthetic pigments<br>(mg/g f.w.) |                 |                 |                 |                 |                 |                 |                 | Antioxidant enzymes activities<br>(U g <sup>-1</sup> protein) |                 |                            |                 |  |
|----------------------------------|--|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|---|-----------------|----------------------------|-----------------|--|
|                                  | Chl. a                                 |                 | Chl. b          |                 | Carot.          |                 | LAI             |                 | Catalase (CAT)  |                 | Superoxide dismutase (SOD) |                 |  |
|                                  | 1 <sup>st</sup>                        | 2 <sup>nd</sup> | 1 <sup>st</sup> | 2 <sup>nd</sup> | 1 <sup>st</sup> | 2 <sup>nd</sup> | 1 <sup>st</sup> | 2 <sup>nd</sup> | 1 <sup>st</sup>   | 2 <sup>nd</sup> | 1 <sup>st</sup>            | 2 <sup>nd</sup> |  |
| Application Methods              |  |                 |                 |                 |                 |                 |                 |                 |   |                 |                            |                 |  |
| Seed soaking                     | 2.53                                   | 2.77            | 1.57            | 1.52            | 0.551           | 0.559           | 3.32            | 3.90            | 0.184   | 0.188           | 49.82                      | 53.87           |  |
| Foliar spraying                  | 2.82                                   | 3.23            | 1.66            | 1.71            | 0.600           | 0.608           | 3.57            | 4.13            | 0.181   | 0.176           | 51.30                      | 55.47           |  |
| Seed soaking and foliar spraying | 3.18                                   | 3.26            | 1.78            | 1.84            | 0.632           | 0.641           | 3.68            | 4.64            | 0.188   | 0.193           | 54.42                      | 56.23           |  |
| LSD at 5%                        | 0.01                                   | 0.04            | 0.03            | 0.01            | 0.034           | 0.035           | 0.30            | 0.33            | 0.004   | 0.006           | 0.51                       | 0.22            |  |
| Ascorbic acid concentrations     |  |                 |                 |                 |                 |                 |                 |                 |   |                 |                            |                 |  |
| Untreated (control)              | 2.44                                   | 2.56            | 1.34            | 1.38            | 0.512           | 0.520           | 3.24            | 3.84            | 0.181   | 0.183           | 49.91                      | 52.29           |  |
| Water                            | 2.48                                   | 2.60            | 1.37            | 1.41            | 0.525           | 0.533           | 3.49            | 4.00            | 0.179   | 0.179           | 49.51                      | 52.03           |  |
| 100 (ppm)                        | 3.21                                   | 3.60            | 2.01            | 2.02            | 0.633           | 0.642           | 3.51            | 4.42            | 0.187   | 0.188           | 53.35                      | 57.91           |  |
| 200(ppm)                         | 3.25                                   | 3.59            | 1.96            | 1.94            | 0.706           | 0.717           | 3.85            | 4.63            | 0.190   | 0.193           | 54.62                      | 58.52           |  |
| LSD at 5%                        | 0.01                                   | 0.04            | 0.03            | NS              | 0.018           | 0.018           | 0.35            | 0.34            | 0.002   | 0.003           | 0.40                       | 0.31            |  |
| Interactions                     | *                                      | *               | *               | *               | *               | *               | NS              | NS              | NS  | NS              | *                          | *               |  |

Chl.a= Chlorophyll a; Chl.b= Chlorophyll b; LAI=Leaf area index; Carot.= Carotenoid 1<sup>st</sup> = First season; 2<sup>nd</sup> = Second season

(AsA) solution plus foliar spraying at 50 and 70 days from sowing (DFS) enhanced and produced the maximum LAI (3.68 and 4.64), chlorophyll a (3.18 and 3.26), chlorophyll b (1.78 and 1.84) and carotenoids (0.632 and 0.641) as well as catalase activity (0.188 and 0.193) and superoxide dismutase activity (54.42 and 56.23) compared to the other two application methods in the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively. This observation coincides with those found by Samin *et al.* (2016), Elsayed (2017) and Jagdish *et al.* (2017).

Concerning ascorbic acid concentrations, it was noticed that plants treated with 100 and/or 200 ppm concentrations of ascorbic acid (AsA) significantly affected chlorophyll a, b and carotenoids, LAI, catalase activity (CAT), superoxide dismutase activity (SOD) in both seasons with exception, chlorophyll b which was insignificant in the 2<sup>nd</sup> season as compared with untreated plant (control) and water tap treatment (Table 2). The maximum increases were observed with 200 ppm ascorbic acid, which enhanced chl.a by (33.20 and 40.23%), Carotenoids by (37.89 and 37.88%), LAI by (18.83 and 20.57%), catalase activity (CAT) by (4.97 and 5.46%), Superoxide dismutase activity (SOD) by (9.44 and 11.91 %), more than the control (untreated) in the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively. No significant differences were observed between 100 and 200 ppm of AsA concentrations for chl.a in the 2<sup>nd</sup> season. Several studies were reported about the AsA-regulation of antioxidant defense metabolism in different plants grown under salinity stress conditions, e.g., canola by (Bybordi, 2012), *Abelmoschus esculentus* (Raza *et al.* 2013), barley (*Hordeum vulgare*) (Agami, 2014). Furthermore, Athar *et al.* (2009) found that application of AsA at 100 mg L<sup>-1</sup> improved growth, and activities of superoxide dismutase (SOD), and catalase (CAT) antioxidative enzymes in wheat. The stimulation effect of ascorbic acid on plant under stress condition might be due to the substantial role of ascorbic acid in many metabolic and physiological processes and counteracting the adverse effects of salinity (Shaddad *et al.*, 1990), regulation of photosynthesis, hormone biosynthesis and regeneration of other antioxidants, plant defense against oxidization by scavenger of the toxic species of activated oxygen, cofactor of key enzyme as well as the role of ascorbate in photosynthesis and involvement both in the electron transport of photosystems and oxidized system of chloroplasts. Panda and Khan (2003) added that, AsA is absorbed readily after exogenous application and moves within the plant Hameed *et al.* (2012).

### **Root length, diameter and fresh weight and root and sugar yields:**

As shown in Table (3) root length, diameter and fresh weight, root and sugar yields/fad significantly affected by the three application methods of AsA acid in both seasons with exception root diameter in the 1<sup>st</sup> season. Soaking sugar beet seeds before sowing in ascorbic acid (AsA) solution plus foliar spraying at 50 and 70 days from sowing, significantly increased root yield by (9.19, 9.63% and 9.88, 11.02%) and sugar yield by (13.43, 15.59% and 5.41, 9.24%) as compared with soaking seed and/or foliar spray in the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively with insignificant differences between them in both seasons. While foliar spraying came in the second rank. On the other hand, seed soaking application recorded the minimum values of all last mention traits, except for, root and sugar yields which recorded the lowest values with exogenous application in both seasons. Similar trends could be seen in the results of Abido *et al.* (2015) and Elsayed (2017). By contrast, Azooz *et al.* (2013) found that pretreatment with ASA resulted in a significant increment of yield of broad bean (*vicia faba*, L.).

Data in Table (3) indicated that, root length, diameter and fresh weight, root and sugar yields /fad significantly increased due to plants treatment with 100 and/or 200 ppm of AsA compared to untreated plant (control) and water tap treatment either soaking or foliar spray in both seasons. The most beneficial AsA concentration was 200 ppm, at which the root were longest (23.47 and 24.33 cm) and the widest (11.00 and 11.08 cm), produced the greatest fresh weigh per plant (0.875 and 0.941g/plant) and enhanced root yield by (32.89 and 23.27 %) and sugar yield by (20.87 and 53.85%) more than the control in the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively. Similar results were obtained by Jagdish *et al.* (2017), Elsayed (2017) and Samin *et al.* (2016). This is might be attributed to the role of AsA in plant cell division, cell expansion, growth and development and senescence (Zhang 2013). Ascorbic acid beneficial effect on plant growth and root length could be attributed to the involvement of ascorbic acid in root elongation regulation, cell expansion and cell vacuolation (Smirnoff, 1996).

### **Sucrose %, potassium, sodium, alpha amino contents and sugar lost to molasses %:**

Soaking sugar beet seed plus foliar spraying, significantly increased sucrose % by 1.37, 2.08 % and 1.49, 2.35% over the other applications methods, i.e.; soaking seed and/or foliar spray in the 1<sup>st</sup> and 2<sup>nd</sup> season, respectively, and gave the best result for the reduction of juice impurities (K, Na and  $\alpha$ -Amino -N) and sugar lost to molasses % in both seasons (Table 4).

**Table 3: Root length, diameter and fresh weight, root and sugar yields of sugar beet as affected by different application methods and concentrations of ascorbic acid and their interaction during 2015/2016 and 2016/2017 seasons.**

| Treatments                       | Root length(cm) |           | Root diameter(cm) |           | Root fresh weight (g/plant) |           | Root yield(ton/fed) |           | Sugar yield(ton/fed) |           |
|----------------------------------|-----------------|-----------|-------------------|-----------|-----------------------------|-----------|---------------------|-----------|----------------------|-----------|
|                                  | 2015/2016       | 2016/2017 | 2015/2016         | 2016/2017 | 2015/2016                   | 2016/2017 | 2015/2016           | 2016/2017 | 2015/2016            | 2016/2017 |
| Application Methods              |                 |           |                   |           |                             |           |                     |           |                      |           |
| Seed soaking                     | 20.93           | 21.64     | 10.28             | 10.41     | 0.736                       | 0.848     | 17.20               | 17.61     | 2.68                 | 3.14      |
| Foliar spraying                  | 20.95           | 21.71     | 10.33             | 10.49     | 0.762                       | 0.890     | 17.13               | 17.43     | 2.63                 | 3.03      |
| Seed soaking and foliar spraying | 23.23           | 24.33     | 10.47             | 10.65     | 0.824                       | 0.936     | 18.78               | 19.35     | 3.04                 | 3.31      |
| LSD at 5%                        | 0.14            | 0.28      | NS                | 0.04      | 0.002                       | 0.009     | 0.20                | 0.29      | 0.06                 | 0.14      |
| Ascorbic acid concentrations     |                 |           |                   |           |                             |           |                     |           |                      |           |
| Untreated (control)              | 19.96           | 20.85     | 9.82              | 10.00     | 0.698                       | 0.848     | 15.23               | 16.33     | 2.54                 | 2.60      |
| Water                            | 20.49           | 21.21     | 9.86              | 10.04     | 0.701                       | 0.852     | 15.31               | 16.40     | 2.48                 | 2.55      |
| 100 (ppm)                        | 22.88           | 23.84     | 10.76             | 10.94     | 0.823                       | 0.925     | 20.03               | 19.66     | 3.04                 | 3.49      |
| 200 (ppm)                        | 23.47           | 24.33     | 11.00             | 11.08     | 0.875                       | 0.941     | 20.24               | 20.13     | 3.07                 | 4.00      |
| LSD at 5%                        | 0.07            | 0.13      | 0.05              | 0.02      | 0.001                       | 0.004     | 0.09                | 0.20      | 0.17                 | 0.14      |
| Interactions                     | *               | *         | NS                | *         | *                           | *         | *                   | *         | *                    | *         |

Data in Table (4) also showed that, treated sugar beet plant by 100 and /or 200 ppm ascorbic acid significantly affected sucrose %, impurities (meq/100 g beet) i.e. K and Na in both seasons and sugar lost to molasses% with exception  $\alpha$ -Amino-N in both seasons and sugar lost to molasses% in the 1<sup>st</sup> season. The maximum increases were observed by treated plant with 200 ppm ascorbic acid, which enhanced

sucrose % by 1.34 and 1.83% more than the control and caused significant reduction in juice impurities potassium and sodium in both seasons and sugar lost to molasses % as compared to control. The same trend was recorded by Abido *et al.* (2015) whose showed that, spraying sugar beet with 150 ppm ascorbic acid recorded significant increase in sucrose % than control (without treatment).

**Table 4: Sucrose%, Impurities (Na, K and  $\alpha$ -amino-N) and sugar lost to molasses (SLM %) of sugar beet juices as affected by different application methods and concentrations of ascorbic acid and their interaction during 2015/2016 and 2016/2017 seasons.**


| Treatments                       | Sucrose%        |                 | Impurities (meq/100 g beet) |                 |                 |                 |                   |                 |                 |                 |
|----------------------------------|-----------------|-----------------|-----------------------------|-----------------|-----------------|-----------------|-------------------|-----------------|-----------------|-----------------|
|                                  |                 |                 | K                           |                 | Na              |                 | $\alpha$ -Amino-N |                 | SLM%            |                 |
| Application Methods              | 1 <sup>st</sup> | 2 <sup>nd</sup> | 1 <sup>st</sup>             | 2 <sup>nd</sup> | 1 <sup>st</sup> | 2 <sup>nd</sup> | 1 <sup>st</sup>   | 2 <sup>nd</sup> | 1 <sup>st</sup> | 2 <sup>nd</sup> |
| Seed soaking                     | 17.53           | 17.87           | 4.01                        | 3.96            | 2.23            | 2.33-15         | 1.34              | 1.35            | 2.56            | 2.57            |
| Foliar spraying                  | 16.82           | 17.01           | 4.32                        | 4.33            | 2.34            | 2.18            | 1.31              | 1.33            | 2.70            | 2.65            |
| Seed soaking and foliar spraying | 18.90           | 19.36           | 3.55                        | 3.62            | 2.09            | 2.07            | 1.19              | 1.21            | 2.34            | 2.36            |
| LSD at 5%                        | 0.21            | 0.14            | 0.26                        | 0.42            | 0.19            | 0.21            | 0.09              | 0.08            | 0.20            | 0.19            |
| Ascorbic acid treatments         |                 |                 |                             |                 |                 |                 |                   |                 |                 |                 |
| Untreated (control)              | 17.15           | 17.20           | 4.10                        | 4.08            | 2.32            | 2.29            | 1.38              | 1.40            | 2.62            | 2.61            |
| Water                            | 17.19           | 17.41           | 3.98                        | 4.04            | 2.26            | 2.25            | 1.35              | 1.38            | 2.56            | 2.58            |
| 100 (ppm)                        | 18.17           | 18.68           | 3.95                        | 3.91            | 2.12            | 2.13            | 1.22              | 1.20            | 2.49            | 2.47            |
| 200 (ppm)                        | 18.49           | 19.03           | 3.83                        | 3.83            | 2.19            | 2.10            | 1.18              | 1.21            | 2.46            | 2.44            |
| LSD at 5%                        | 0.17            | 0.26            | 0.16                        | 0.12            | 0.09            | 0.07            | NS                | NS              | NS              | 0.05            |
| Interactions                     | *               | *               | NS                          | NS              | NS              | NS              | NS                | NS              | NS              | NS              |

K= Potassium; Na= Sodium ; SLM= Sugar lost to molasses; 1<sup>st</sup> = First season; 2<sup>nd</sup> = Second season


#### Analysis of stoma morphological parameters:

The effect of different application methods and concentrations of ascorbic acid (AsA) on morphological changes of stomatal characteristics (Stomata area, density and closure %) for upper (adaxial) and lower (abaxial) surfaces of sugar beet leaf are shown in Fig. (1 and 2). Individual response to different application method of AsA concentrations was observed for each parameter of stomata. Largest area or size and lower density of stomata for both adaxial and abaxial leaf surfaces, as well as, lower closure% were recorded by application of seed soaking with AsA and foliar spraying together followed by foliar spraying treatment while the higher density and smaller size of stomata was observed by seed soaking treatment at all AsA concentrations. Treated sugar beet plants with (100 and 200 ppm) of AsA or distilled water as seed soaking ,foliar spraying or seed soaking plus foliar spraying gave the highest stomata untreated plants-size and lower density and closure% than those untreated plant (control) for upper and lower surfaces of sugar beet. Increasing concentrations of ascorbic acid from 100 to 200 ppm lead to an increase in stomata area for both leaf surfaces. The microscopic analysis (Image a, b, c, and d) showed that: higher stomata area, lower stomata density and stomata closure % were recorded by seed soaking plus foliar spraying application with 200 ppm AsA for upper and lower surfaces of sugar beet leaf (Images a

and b). On the other hand, untreated sugar beet plants by AsA (control) affected all stomata parameter and recorded the higher stomata closure %, and reduced dimensions of stomata in both beet leaf surfaces (Images c and d) this might be due to Salinity that caused both water stress and osmotic stress in plants and that accumulated salt ions had a toxic effect on plants and decreased leaf turgor, further causing stomata closure and decreases of stomata size (Chaves *et al.* 2009). Our results also confirm that, stomatal density is inversely related to leaf area and that salinity might increase stomatal density by causing reduction of leaf area. Higher density and smaller size of stomata is a form of adaptation to salinity, because it allowed plants to be more efficient in water transport regulation and transpiration (Dickison, 2000) added that, stomatal density might increase with leaf size reduction to overcome a simple geometric practicality of fitting enough functional stomatal units per unit of leaf surface area to meet the desired CO<sub>2</sub> flux and to service photosynthetic capacity (Franks and Farquhar, 2007). Ascorbic acid (AsA) is the major antioxidant that scavenges H<sub>2</sub>O<sub>2</sub> which an important stress signaling function and promotes stomata closure (Chen and Gallie 2004). These results are in harmony with those achieved by Arafa *et al.* (2014) whose concluded that, soaking seeds of sweet pepper plant in ascorbic acid at 50 mg/L might be useful for overcoming the harmful effects of salinity on the stomata density on both leaf surfaces.




**Fig. (1): Stomatal area ( $\mu\text{m}$ ) and density ( $\text{No. mm}^{-2}$ ) for upper (U) and lower (L) surfaces of sugar beet leaf as affected by different application methods of AsA concentrations in 2015/2016 season.**




**Fig. (2): Stomatal closure% for upper and lower surfaces of sugar beet leaf as affected by different application methods of AsA concentrations in 2015/2016**





**Image a and b: Electron microscopic images (magnification 300 x) of adaxial (image a) and abaxial (image b) surface of sugar beet leaf showing the effect of seed soaking plus foliar spraying treatment with 200 ppm AsA**



**Image c and d: Electron microscopic images (magnification 300 x) of adaxial (image c) and abaxial (image d) surface of sugar beet leaf showing the effect of untreated sugar beet plants with AsA (control)**

#### Interaction effects:

The effect of the interaction between application methods and different treatments of AsA were significant in their effect on chlorophyll a, b and carotenoids, superoxide dismutase (SOD) activity, root and sugar yields and its components and sucrose % in both seasons as shown in Tables (5 and 6) which cleared that, ascorbic acid at 200 ppm when applied as a seed soaking plus foliar spraying proved most effective to reduce salt stress inhibition followed by ascorbic acid at 100 ppm. The aforementioned treatment significantly increased root yield by (6.99, 6.80 and 4.54, 3.95 ton/fad) and sugar yield by (2.19, 1.02 and 2.06, 1.11 ton/fed) over that gained by untreated plants in the 1<sup>st</sup> and 2<sup>nd</sup> seasons, respectively.

#### Conclusion

It could be concluded that application of ascorbic acid at 200 and or 100 ppm as a seed soaking plus foliar spraying had proved as the most effective treatment to reduce salt stress inhibition and resulted in sugar beet maximum root and sugar yields/fad of under this experimental conditions.

#### References

- Abido, W. A. E., M. A. E. Ibrahim and Maha M. El-Zeny (2015). Growth, productivity and quality of sugar beet as affected by antioxidants foliar application and potassium fertilizer top dressing. *Asian J. Crops Sci.* 7(2):113-127.
- Agami, R.A. (2014) Applications of ascorbic acid or proline increase resistance to salt stress in barley seedlings. *Bio. Plan.* 58 (2): 341-347.
- Ahmad, P., M. Ozturk and S. Gucel (2012). Oxidative damage and antioxidants induced by heavy metal stress in two cultivars of mustard (*Brassica juncea*, L.) plants. *Fresenius Environ. Bull.* 21:2953-2961
- Arafa, A. A., M. A. Khafagy, A. M. Abo-El Kheer, R. A. Fouda and M.F. El-Banna, (2014). Stomatal density in the leaves of sweet pepper plant as affected by certain bio-stimulants under salt stress conditions. *J. Plant Production, Mansoura Univ., Vol.5 (4): 649 - 662.*
- Athar, H., A. Khan and M. Ashraf (2009). Including salt tolerance in wheat by exogenously applied ascorbic acid through different modes. *J. Plant Nutr.* 32(11): 1799–1817.

**Table (5): Root length and fresh weight, root and sugar yields, Chlorophyll a and b, carotenoides, Superoxide dismutase (SOD) activity and sucrose % as affected by the interaction between application methods and concentrations of AsA during 2015/2016 seasons.**

| Application methods              | AsA Concentrations (ppm) | RL (cm)* | RFW (g/plant) | RY (t/fed) | SY (t/fed) | Photosynthetic pigments (mg/g F W) |        |        | SOD (U g <sup>-1</sup> protein) | Suc. % |
|----------------------------------|--------------------------|----------|---------------|------------|------------|------------------------------------|--------|--------|---------------------------------|--------|
|                                  |                          |          |               |            |            | Chl. a                             | Chl. b | Carot. |                                 |        |
| Seed soaking                     | Untreated (control)      | 20.31    | 0.672         | 15.23      | 2.45       | 2.40                               | 1.35   | 0.51   | 45.99                           | 16.87  |
|                                  | Water                    | 20.55    | 0.683         | 15.27      | 2.37       | 2.48                               | 1.40   | 0.52   | 46.37                           | 16.81  |
|                                  | 100                      | 21.22    | 0.776         | 18.89      | 3.00       | 2.61                               | 1.70   | 0.53   | 54.78                           | 17.94  |
|                                  | 200                      | 21.65    | 0.813         | 19.12      | 3.04       | 2.63                               | 1.81   | 0.64   | 52.13                           | 18.51  |
| Foliar spraying                  | Untreated (control)      | 19.54    | 0.706         | 15.15      | 2.34       | 2.46                               | 1.33   | 0.51   | 49.01                           | 16.35  |
|                                  | Water                    | 20.36    | 0.706         | 15.28      | 2.25       | 2.48                               | 1.35   | 0.52   | 49.65                           | 16.44  |
|                                  | 100                      | 21.38    | 0.816         | 19.10      | 2.88       | 3.22                               | 1.95   | 0.73   | 53.55                           | 17.13  |
|                                  | 200                      | 22.43    | 0.821         | 19.27      | 2.91       | 3.12                               | 2.00   | 0.77   | 52.97                           | 17.35  |
| Seed soaking and Foliar spraying | Untreated (control)      | 20.04    | 0.714         | 15.31      | 2.22       | 2.46                               | 1.33   | 0.52   | 53.52                           | 18.22  |
|                                  | Water                    | 20.57    | 0.714         | 15.38      | 2.23       | 2.48                               | 1.35   | 0.54   | 53.70                           | 18.33  |
|                                  | 100                      | 24.04    | 0.878         | 22.11      | 3.24       | 3.81                               | 2.24   | 0.64   | 55.53                           | 19.44  |
|                                  | 200                      | 26.33    | 0.990         | 22.30      | 4.41       | 3.98                               | 2.20   | 0.71   | 54.95                           | 19.62  |
| LSD at 5%                        |                          | 0.12     | 0.002         | 0.16       | 0.30       | 0.02                               | 0.01   | 0.03   | 0.69                            | 0.29   |

RL=Root length; RFW= Root fresh weight; RY=Root yield; SY=Sugar yield; Chl.a= Chlorophyll a; Chl.b = Chlorophyll b; Carot.= Carotenoid; SOD= Superoxide dismutase; Suc.% =Sucrose %.

**Table (6):** Root length, diameter and fresh weight, root and sugar yields, Chlorophyll a and b, carotenoids, Superoxide dismutase (SOD) activity and sucrose % as affected by the interaction between application methods and concentrations of AsA during 2016/2017 seasons

| Application methods              | AsA Concentrations (ppm) | RL (cm) | RD (cm) | RFW (g/plant) | RY (t/fed) | SY (t/fed) | Photosynthetic pigments(mg/g FW) |        |        | (SOD) (U g <sup>-1</sup> protein) | c. %  |
|----------------------------------|--------------------------|---------|---------|---------------|------------|------------|----------------------------------|--------|--------|-----------------------------------|-------|
|                                  |                          |         |         |               |            |            | Chl. a                           | Chl. b | Carot. |                                   |       |
| Seed soaking                     | Untreated (control)      | 21.42   | 9.88    | 0.871         | 15.68      | 2.59       | 2.61                             | 1.21   | 0.52   | 49.86                             | 16.83 |
|                                  | Water                    | 21.11   | 9.92    | 0.877         | 15.73      | 2.50       | 2.63                             | 1.25   | 0.53   | 50.38                             | 17.00 |
|                                  | 100                      | 21.96   | 10.82   | 0.906         | 19.20      | 3.46       | 2.91                             | 1.77   | 0.54   | 58.66                             | 18.67 |
|                                  | 200                      | 22.33   | 11.02   | 0.907         | 19.83      | 3.99       | 2.95                             | 1.83   | 0.65   | 56.56                             | 18.97 |
| Foliar spraying                  | Untreated (control)      | 20.21   | 9.96    | 0.798         | 16.13      | 2.42       | 2.53                             | 1.39   | 0.52   | 52.05                             | 16.36 |
|                                  | Water                    | 21.41   | 10.02   | 0.801         | 16.15      | 2.32       | 2.58                             | 1.40   | 0.53   | 52.27                             | 16.68 |
|                                  | 100                      | 22.03   | 10.92   | 0.883         | 18.66      | 3.40       | 3.88                             | 1.98   | 0.74   | 58.41                             | 17.23 |
|                                  | 200                      | 22.91   | 11.06   | 0.912         | 18.80      | 3.97       | 3.91                             | 2.07   | 0.78   | 59.15                             | 17.75 |
| Seed soaking and Foliar spraying | Untreated (control)      | 20.93   | 10.15   | 0.875         | 17.18      | 2.49       | 2.53                             | 1.55   | 0.53   | 54.17                             | 18.41 |
|                                  | Water                    | 21.11   | 10.19   | 0.880         | 17.34      | 2.52       | 2.58                             | 1.58   | 0.55   | 54.22                             | 18.54 |
|                                  | 100                      | 27.53   | 11.08   | 0.987         | 21.13      | 3.60       | 3.99                             | 2.08   | 0.64   | 58.48                             | 20.13 |
|                                  | 200                      | 27.76   | 11.16   | 1.004         | 21.72      | 4.55       | 3.92                             | 2.15   | 0.72   | 58.03                             | 20.37 |
| LSD at 5%                        |                          | 0.23    | 0.03    | 0.007         | 0.34       | 0.25       | 0.06                             | 0.01   | 0.03   | 0.54                              | 0.45  |

RL=Root length; RD=Root diameter; RFW= Root fresh weight; RY=Root yield; SY= Sugar yield; Chl.a= Chlorophyll a; Chl.b= Chlorophyll b; Carot.= Carotenoid; SOD= Superoxide dismutase; Suc.% =Sucrose %.

- Azooz, M.M., A.M. Alzahrani and M.M. Youssef (2013). The potential role of seed priming with ascorbic acid and nicotinamide and their interactions to enhance salt tolerance in broad bean (*vicia faba L.*). *Aust. J. Crop Sci.* 7(13):2091-2100.
- Barth, C., M. De Tullio, and P. L. Conklin (2006). The role of ascorbic acid in the control of flowering time and the onset of senescence. *J. Exp. Bot.* 57(8):1657-1665.
- Beyer, w.f. and i. Fridovich (1987). Assaying for superoxide dismutase activity: some large consequences of minor changes in condition. *Anal. Biochem.*, 16 (2):559-566.
- Bybordi, A.(2012).Effect of ascorbic acid and silicium on photosynthesis, antioxidant enzyme activity, and fatty acid contents in canola exposure to salt stress. *J. Integr. Agric.* 11, 1610–1620.
- Chaves, M. M., J. Flexas, and C. Pinheiro (2009). Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Ann. Bot.* 103(4):551–560.
- Chen Z. and D. R. Gallie (2004). The Ascorbic Acid Redox State Controls Guard Cell Signaling and Stomatal Movement. *The Plant Cell*, Vol. 16, 1143–1162
- Devillers, P. (1988). Previsioin du sucre melasse Scurries francases 129: 190-200. In: Cooke, D.A. and R. K. Scott (1993). The sugar beet crop. Chapman and Hall London, pp.262-265.
- Dickison W.C. (2000) Integrative Plant Anatomy. San Diego: Academic Press.
- El-Hariri, D. M., M. S. Sadak and H. M. S. El-Bassiouny (2010). Response of flax cultivars to ascorbic acid and  $\alpha$ -tocopherol under salinity stress condition. *Inter. J. of Academic Res.* 2(6): 101-109.
- Elsayed, A. A. Abdelraouf (2017). Effect of Seed soaking Sugar Beet (*Beta vulgaris*, L.) Seeds with Gibberellic, Abscisic or Ascorbic Acids on Alleviation of Salinity Stress. *Alex. Sci., Exchange J.* 38(1): 74-81.
- Euroconsult (1992). Environmental profile, Fayoum Governorate, Egypt. Al-Shorouk Press, Cairo.
- Franks P.J. and G.D. Farquhar (2007). The mechanical diversity of stomata and its significance in gas-exchange control. *Plant Physiol* 143:78–87.
- FWMP. (1999). Fayoum Water Management Project II "Salinisation Monitoring of Lake Qarun Between 1901 and 1998". Technical Report No. 55: 16pp.
- Hameed, A., T. Hussain, S. Gulzar, I. Aziz, B. Gul and M.A. Khan (2012).Salt tolerance of a cash crop halophyte *Suaeda fruticosa*: biochemical responses to salt and exogenous chemical treatments. *Acta Physiologiae Plantarum* 34:2331 –2340.
- Hasanuzzaman, M., M. A. Hossain, J. A. Teixeira da Silva and M. Fujita (2012). "Plant responses and tolerance to abiotic oxidative stress: antioxidant defense is a key factor," in Crop Stress and Its Management: Perspectives and Strategies, V. Bandi, A. K. Shanker, C. Shanker, and M. Mandapaka, Eds., pp. 261–316, Springer, Berlin, Germany,.
- Jagdish, K. N., A. B. Nishant, K. Deepmala, H. Akhouri and K. Y. Dinesh (2017). Mitigating effect of foliar applied ascorbic acid on morpho-physiological, biochemical changes and yield attributes induced by salt stress in *Vigna radiate*. *Agric. Sci. Digest.*, 37(2): 112-116.
- Le-Docte A. (1927). Commercial determination of sugar beet in the beet roots using Sachs Le-Docte process. *International Sugar J.* 29: 488-492.
- Maas, E.V (1986). Salt tolerance of plants. *Appl. Agric. Res.* 1(1): 12-26.
- Naz, H., N. A. Akram and M. Ashraf (2016). Impact of ascorbic acid on growth and some physiological attributes of cucumber (*Cucumis sativus*) plants under water-deficit conditions. *Pak. J. Bot.*, 48(3): 877-883.
- Panda, S.K. and M.H. Khan (2003). Salt stress influences lipid peroxidation and antioxidants in the leaf of an indica rice (*Oryza sativa*, L.). *Physiol. Mol. Biol. Plants*, 9: 273-278.
- Raza, S. H., F. Shafiq, M. Chaudhary and I. Khan (2013). Seed invigoration with water, ascorbic and salicylic acid stimulates development and biochemical characters of okra (*Ablemoschus esculentus*) under normal and saline conditions. *Int. J. Agric. Biol.* 15, 486–492.
- Rhoades, J. and J. Loveday (1990). Salinity in Irrigated Agriculture. In Stewart, B.A. and D.R Nielsen (eds.); Irrigation of Agric. Crops. Agron.No. 30, American Soc. pp. 1089–1142.
- Ryan, J., S. Garabet, K. Harmsen and A. Rashid (1996). A soil and plant analysis. Manual Adapted for the West Asia and North Africa Region. ICARDA, Aleppo, Syria. 140 pp
- Sakr, M.T., Z.A. Mohamed, I. A. Marouah and M.T. Zalama (2014). Response of faba bean plants to application of some growth promoters under salinity stress conditions. *J. Plant Production, Mansoura Univ.*, 5 (1): 79-94.

- Samin j., M. Hamayun, S. Wali, A. Bibi, H. Gul and F. Rahim (2016). Foliar application of ascorbic acid mitigates sodium Chloride induced stress in eggplant (*solanum melongena* l.). *Pak. J. Bot.*, 48(3): 869-876.
- Shaddad, M.A., A.F. Radi, A.M. Abdel-Rahman and M.M. Azooz. (1990). Response of seeds of *lupinus termis* and *Vicia faba* to the interactive effect of salinity and ascorbic acid or pyridoxine. *Plant & Soil*, 122(2): 177-183.
- Shadrin N.V., G.M. EL-Shabrawy, E.V. Anufriieva, M.E. Goher and E. Ragab (2016). 'Long-term changes of physicochemical parameters and benthos in Lake Qarun (Egypt): Can we make a correct forecast of ecosystem future, *Knowl. Manag. Aquat. Ecosyst.* 417, 18:1-11.
- Shao, H. B., L. Y.Chu, Z. H. Lu, and C. M. Kang (2008). Primary antioxidant free radical scavenging and redox signaling pathways in higher plant cells. *Int. J. Biol. Sci.* 4(1): 8–14.
- Smirnoff, N. (1996). The function and metabolism of ascorbic acid in plants. *Ann. Bot.*, 78: 661-669.
- Snedecor G.W. and Cochran W.G. (1980). *Statistical Methods*. 7 ed., Iowa State Univ. Press, Ames, Iowa, USA, pp 255-269.
- Wettstein, D. (1957). Chlorophyll, Letal under submikro svopische formmech, sall-plastiden- Exptl Cell Ser., 12: 427-433.
- Yin H., L. Xu and N. A. Porter (2011). "Free radical lipid peroxidation: Mechanisms and analysis," *Chem. Rev.*, 111 (10): pp. 5944–5972.
- Zhang, Y. (2013). Biological role of ascorbate in plants (Chapter 2). Springer Briefs in Plant Science.

## تأثير طرق إضافة وتركيزات لحمض الأسكوربيك على بنجر السكر تحت ظروف الاجهاد الملحي ايمان محمد عبد الفتاح<sup>١</sup> - كرم عبد الصادق<sup>٢</sup>

١ قسم بحوث الفسيولوجي والكيمياء بمعهد بحوث المحاصيل السكرية- مركز البحوث الزراعية

٢ قسم بحوث المعاملات الزراعية بمعهد بحوث المحاصيل السكرية- مركز البحوث الزراعية

الاجهاد الملحي هي احد عوامل الإجهاد البيئي التي تؤثر على نمو النبات ويقلل من إنتاجيتها. تعتبر مضادات الأكسدة واحدة من الأساليب والاستراتيجيات المتبعة لتخفيف الآثار الضارة لإجهاد الملوحة على النباتات و لتقليل تأثير الإجهاد التأكسدي. اقيمت تجربتان حقليتان بمنطقة سنورس التي تقع بشكل رئيسي بجوار بحيرة قارون (تربة مالحة) في محافظة الفيوم (خطي العرض ٢٩ ° ٢٠ و ٢٩ ° ٣٠ N وخط الطول ٣٠ ° ٤٣ و ٣٠ ° ٥٦ E) خلال موسمي الزراعة ٢٠١٦/٢٠١٥ و ٢٠١٦/٢٠١٧. لدراسة تأثير ثلاث طرق لإضافة حمض الاسكوربيك (نقع البذور ، رش الأوراق ، نقع البذور بالإضافة إلى رش الأوراق معاً) و أربعة تركيزات (بدون معاملة (كنترول) ، ماء الصنبور ، ١٠٠ و ٢٠٠ جزء في المليون) على بعض الخصائص الكيميائية الحيوية والفسيولوجية ونشاط بعض إنزيمات مضادات الأكسدة وكذلك على بعض صفات النمو وحاصل وجودة بنجر السكر المنزرع تحت ظروف الاجهاد الملحي. تم استخدام تصميم الشرائح المتعامدة في ثلاث مكررات ، اوضحت النتائج التالي:

- ادى نقع بذور بنجر السكر في حمض الأسكوربيك (AsA) بالإضافة إلى الرش الورقي عند عمر ٥٠ و ٧٠ يوماً من الزراعة إلى تخفيف التأثير الضار لملوحة التربة و اعطى زيادة معنوية في متوسطات الكلوروفيل (أ) ، (ب) والكاروتينيدات و دليل مساحة الاوراق و زيادة معنوية في فعالية مضادات الأكسدة الإنزيمية السوبرأوكسيد دسميوتيز (SOD) و الكاتاليز (CAT) و مكونات المحصول والنسب المئوية للسكروز % وانخفاض الشونب في كلا الموسمي ،فضلاً عن زيادة حاصل الجذور والسكر/فدان بنسبة (٩,١٩ ، ٩,٦٣ ، %) و (١٣,٤٣ ، ١٥,٥٩ ، % ) مقارنة بمعاملتى نقع البذور أو الرش الورقي في الموسم الأول. نفس الاتجاه كان في الموسم الثاني.
- دلت النتائج على ان تطبيق ٢٠٠ جزء في المليون من حمض الأسكوربيك كان أكثر فاعلية وزاد حاصل الجذور و حاصل السكر والنسب المئوية للسكروز وانخفاض الشونب مقارنة بالكنترول في كلا الموسمين.
- تم تسجيل أعلى مساحة و أقل كثافة للثغور و نسبة الثغور المغلقة للأسطح العلوية والسفلية لأوراق بنجر السكر من خلال نقع البذور مع الرش الورقي بحمض الاسكوربيك بتركيز ٢٠٠ جزء في المليون.
- تحقق اعلي حاصل جذور وحاصل سكر عند نقع البذور مع الرش الورقي لحمض الاسكوربيك بتركيز ٢٠٠ جزء في المليون حيث زاد حاصل الجذور بمقدار (٦,٩٩ و ٤,٥٤ طن / فدان) حاصل السكر بمقدار (٢,١٩ و ٢,٠٦ طن / فدان) مقارنة بالكنترول في في الموسمين الأول والثاني على التوالي.
- يتضح من هذا البحث انه يمكن التغلب على الآثار الضارة للإجهاد الناشئ عن ملوحة التربة على صفات النمو والمحصول ومكوناته والمحتويات البيوكيميائية والجودة لنباتات بنجر السكر باستخدام المواد المضادة للأكسدة (حمض الاسكوربيك) حيث أدى استخدامه نقعاً ورشاً معاً بتركيز ٢٠٠ جزء في المليون ، إلى مقاومة إجهاد ملوحة التربة وزيادة إنتاجيتها من محصول بنجر السكر. وبذلك يمكن زيادة مساحة رقعة الأرض الزراعية وذلك بإضافة مزيد من الأراضي التي بها نسبة ملوحة إلى المساحة المنزرعة مما ينعكس بالإيجاب على زيادة محصول بنجر السكر.