

Journal of Plant Protection and Pathology

Journal homepage: www.jppp.mans.edu.eg
Available online at: www.jppp.journals.ekb.eg

Expected Reasons of Population Decline in Honey Bee (*Apis mellifera*) Colonies

Ayoub, Z. N. ^{*1}; R. J. Rashed²; T. H. Shekh Faraj² and O. M. Ahmad³

^{*1} Department of Plant Protection, College of Agricultural Engineering sciences, University of Duhok, Kurdistan Region, Iraq.

² Department of Horticulture, College of Agricultural Engineering sciences, University of Sulaimani, Kurdistan Region, Iraq.

³ General Directorate of Agriculture, Ministry of Agriculture and Water Resources, Kurdistan Region, Iraq.



Cross Mark

ABSTRACT

The study aimed to investigate the relationship between colony survival and some preventive procedures. The experiments were conducted in Sulemani governorate during August, September and October 2020. Seventy-two colonies were tested in four localities (Qaradax , Sulaimani center, Mergapan, and Sartake bamo); (18 colonies for each location). The tested colonies were treated with; probiotics, formic acid, eucalyptus, Tetracycline and Terramycin. Larger brood area was found in all treated colonies compared to untreated ones. Colonies provided with probiotics showed significantly more brood than untreated colonies in the four locations. Maximum brood area was 203.667 (inch)² in Sartake bamo followed by 199.667 (inch)² in Mergapan; then 179.000 (inch)² in Sulaimani center. While the brood area was not more than 15.667 (inch)² in all untreated colonies in the four tested apiaries. Colonies provided with probiotics showed significantly more density of adult workers than untreated colonies in the four locations. Maximum area covered with adult workers (density) was 6.667 Lf. (Langstroth frame) in Sartake bamo followed by 6.333 Lf. in Mergapan; then 6.000 Lf. in those treated with Formic acid in Sartake bamo. While the density of adult workers was not more than 1.000 Lf. in all untreated colonies in the four tested apiaries. Using probiotics and organic acid treatments were the best preventive measures.

Keywords: Honey Bee, Probiotics, Formic Acid, Brood Area

INTRODUCTION

Huge number of colony losses of managed honey bees in the recent years have disrupted beekeeping industry and bee researchers. The honeybee is not only pollinators of agricultural crops and wild plants, but also provide abundant bee products. In recent years, the dramatic reductions in bee colonies, causing significant economic losses, were reported from all over the world. The main reasons were diverse; agrochemicals, parasites, viruses, methods of planting structure and distribution, or their interaction factors (Vanengelsdorp *et al.*, 2009).

Domestic apiculture industry in Iraq was destroyed during gulf war only feral colonies existed in the mountains. After 1991 beekeeping process began again and a large number of infested honey bee colonies were illegally imported from neighboring countries. *Varroa* mite infestation was first detected in Iraq in the mid-1980s (FAO). Many beekeepers, particularly those with traditional hives lost almost all their colonies. In 1990, *Varroa* mite was reported in all Arab countries (Haddad 2011). Although different kinds of acaricides from various sources were applied by beekeepers but still remains threat to the bee hives of the area. (Ayoub *et al.*, 2014)

The single greatest threat to honey bee populations worldwide is the invasive mite *Varroa destructor* Andersen & Trueman. The life cycle of the varroa mites is tightly adapted to the development of the honey bees. *Varroa* mites are serious and devastating ectoparasites of the honey bee. During the phoretic phase, the varroa mites live on the bodies of honey bees and feed on their haemolymph. The reproductive phase of

varroa mites happens exclusively in the capped cells of developing bee pupae (Ifantidis, 1983). Several studies have documented the negative effects of varroa infestation on honey bees including reduced lifespan (Kralj *et al.*, 2007), decreased survivorship (Yang *et al.*, 2007) and weight loss in drones (Duay *et al.*, 2002).

The second threat to honey bees is American Foulbrood disease which classified on list B of the Office International de Epizootic (OIE), the world organization for animal health. List B diseases are those diseases which have significant impact on the socio-economic and/or public health of the countries as well as international trade of animals and animal products (de Graaf *et al.*, 2006). Colony Collapse Disorder (CCD) is a syndrome describing the large-scale loss of managed honey bees worldwide first reported in 2006–2007 (Vanengelsdorp *et al.*, 2009). Several studies have investigated and reported various causes for this sudden decline in bees such as viruses (Dainat *et al.*, 2012), varroa mites (Le Conte *et al.*, 2010), microsporidean *Nosema* spp. and bacterial brood diseases (Paxton, 2010).


Due to the lack of definite causal agent of colony losses, it is being investigated extensively and it is becoming clear that a single causal agent is difficult to identify and these causes are possibly multiple and very complex (Dainat *et al.*, 2012; Nazzi *et al.*, 2012; Evans *et al.*, 2012).

Beekeepers have shown their interest in chemical treatments that instantly show their effect on the pests rather than using natural products. Natural chemicals, such as formic acid, oxalic acid, lactic acid, thymol eucalyptus, camphor, menthol, thyme oil, laurel, and lavender oils have been recommended and used in recent years. Since organic acids

* Corresponding author.

E-mail address: zahra.naef@uod.ac

DOI: 10.21608/jppp.2021.198010



appear to be less harmful to human health and are naturally found in the chemical structure of honey, they are suggested in the process of producing organic and healthy hive products (Esmen, Dodoloğlu, & Genc., 2010).

According to European Union regulation 1804/1899 on organic production, the use of formic acid is authorized as a natural compound in organic apiculture standard management (Mato et al., 2006).

Aims of the study;

- This study aimed to investigate the relationship between colony survival and some preventive procedures.
- To find out and confirm the most obvious reasons of population decline in order to perform suitable solutions to preserve honey bee (*Apis mellifera*) colonies in our area.

MATERIALS AND METHODS

All experiments were conducted in Sulemani governorate during August, September and October 2020. Seventy-two colonies were tested in four localities (Qaradax , Sulaimani center, Mergapan, and Sartake bamo). Fifteen colonies were differently treated and compared to three untreated colonies in each locality (apiary). The treatments can be illustrated as the following;

The 1st treatment: three colonies were provided with probiotic, doses for probiotics were calculated from the manufacturer’s instructions, considering an average body weight of 100 mg for individual bee. All doses were prepared in 50% sucrose syrup. This treatment was applied weekly for six weeks.

The 2nd treatment: three colonies were treated with formic acid; the evaporator pad was placed on a top of the frames in the brood chamber.

The 3rd treatment: three colonies were treated continuously with eucalyptus smoking during colony inspection.

The 4th treatment: three colonies were treated with Tetracycline (Bee tetracycline Alfarabi), 100 g of the product was mixed with 200 g of powdered sugar; each hive received 20 g of the mixture.

The 5th treatment: three colonies were treated with Terramycin, the product was combined with powdered sugar to make a dust and applied to honey bee colonies for six weeks.

The 6th treatment (control): three untreated colonies compared to the previously mentioned colonies.

The total area of capped brood and the density of adult workers (area covered with bees) on both sides of the combs were considered as parameters of colony strength. These parameters were weekly measured during experimental period using standard Langstroth frame.

The results were analyzed statistically using factorial RCBD design with triple replicates and performed using XLSTA program (2017) m, Duncan’s multiple range Test was used to determine the differences between means at P = 0.05.

RESULTS

Larger brood area was found in all treated colonies compared to untreated ones. Colonies treated with probiotic showed larger brood area in the four locations. Colonies provided with probiotics showed significantly more brood than untreated colonies. Maximum brood area was 203.667 (inch)² in Sartake bamo followed by 199.667 (inch)² in Mergapan; then 179.000 (inch)² in Sulaimani center. While the brood area was

not more than 15.667 (inch)² in all untreated colonies in the four tested apiaries, table (1); fig (1,2,3, and 4).

Table 1. Effect of five different treatments on honey bee brood area (inch)² compared to untreated colonies in four locations.

Treatments	Locations (apiaries)			
	Qaradax	Sulaimani center	Mergapan	Sartake bamo
Probiotic	167.333 a	179.000 a	199.667 a	203.667 a
Formic acid	144.667 ab	159.333 ab	167.667 ab	159.667 ab
Eucalyptus	144.000 ab	129.667 b	144.000 b	139.667 ab
Tetracycline	133.000 bc	153.000 ab	166.667 ab	166.000 ab
Terramycin	114.667 c	164.333 a	167.333 ab	165.000 b
control	15.667 d	7.667 c	13.330 c	9.333 c

Means with the same letter are not significantly different.




Fig 1. Effect of five different treatments on honey bee brood area (inch)² compared to untreated colonies in Qaradax.




Fig 2. Effect of five different treatments on honey bee brood area (inch)² compared to untreated colonies in Sulaimani center.




Fig 3. Effect of five different treatments on honey bee brood area (inch)² compared to untreated colonies in Mergapan.




Fig 4. Effect of five different treatments on honey bee brood area (inch)² compared to untreated colonies in sartaki bamo.

More density of adult workers was found in all treated colonies compared to untreated ones. Colonies treated with probiotic showed higher density of adult workers in the four locations. Colonies provided with probiotics showed significantly more density of adult workers than untreated colonies. Maximum brood area was 6.667 Lf. (Langstroth frame) in Sartake bamo followed by 6.333 Lf. in Mergapan; then 6.000 Lf. in those treated with Formic acid in Sartake bamo. While the density of adult workers was not more than 1.000 Lf. in all untreated colonies in the four tested apiaries, table (2); fig (5,6,7and 8).

Using probiotics and formic acid treatments were the best preventive measures in all tested apiaries.

Table 2. Effect of five different treatments on honey bee density (Langstroth frame) compared to untreated colonies in four locations.

Treatments	Locations (apiaries)			
	Qaradax	Sulaimani center	Mergapan	Sartake bamo
Probiotic	5.667 a	5.667 a	6.333 a	6.667 a
Formic acid	4.667 ab	5.000 ab	4.667 bc	6.000 a
Eucalyptus	4.333 b	4.000 b	3.667 c	5.000 b
Tetracycline	3.667 b	4.333 b	5.000 b	4.000 c
Terramycine	4.333 b	5.000 ab	5.667 ab	6.000 a
control	0.333 c	1.000 c	1.000 c	0.667 d

Means with the same letter are not significantly different.




Fig 5. Effect of five different treatments on honey bee density (Langstroth frame) compared to untreated colonies in Qaradax.




Fig 6. Effect of five different treatments on honey bee density (Langstroth frame) compared to untreated colonies in Sulaimani center.




Fig 7. Effect of five different treatments on honey bee density (Langstroth frame) compared to untreated colonies in Mergapan.




Fig 8. Effect of five different treatments on honey bee density (Langstroth frame) compared to untreated colonies in Sartake bamo.

DISCUSSION

Many pest and disease problems in managed honeybee colonies can be avoided by performing good sanitation and cultural controls. Prevention is the best way of defense against organisms that can harm our colonies.

There is no agreement on the definite reason(s) of the problem of honey bee colony losses and the inhibition of the population growth of the colonies (Zoran *et al.*, 2019). Scientific community has been confirmed that colony losses and the decrease in population growth of the colony are a multifactorial issue (vanEngelsdorp *et al.*, 2010; Neumann *et al.*, 2010; Goulson *et al.*, 2015).

Various non-specific factors (e.g., climate changes, agrochemisation and inadequate food) decrease the strength of the colonies; such colonies easily become susceptible for bacterial, and many microbial infections. Then these unhealthy situations diminish the immune system of the bee (Evans *et al.*, 2004; Gätschenberger *et al.*, 2013). Inadequate anti-varroa Management lead to significant health problems in honey bee colonies and cause the spread of viruses.

Our results are in agreement with recent studies that have found other natural compounds such as natural plant extracts, probiotics bacteria, and some organic acids that are very useful for maintaining the survival of honey bee colonies and supporting the population growth of the colonies. For example, the probiotics bacteria have antimicrobial activity against a large number of bacteria and fungi and produces large amounts of lactic acid (Kulhanek *et al.*, 2017).

Because of this is a multifactorial issue, therefore the recommended solutions to the problem includes a sequence of activities and preventive procedures during beekeeping management leads to the protection of the large number of the susceptible colonies from expected pests and pathogens in our area.

REFERENCES

- Aydın, L., Cakmak, I., & Cakmak, S. 2007. Efficacy of Ecostop (Thymol+ Menthol) and perizin (coumaphos) against Varroa destructor found on naturally infested honey bee colonies. *Uludag Bee Journal*, 59–62.
- Aydın, L., Senlik, B., & Girisgin, A.O. 2009. Efficacy of Obeson (Thymol) against Varroa destructor found on naturally infested honey bee colonies. *Uludag Bee Journal*, 9, 72–75.
- Ayoub, Z.N., Ahmed, D.S., Ismael, H.R. 2014. *Varroa* mite infestation in apiaries of Duhok province, Kurdistan, Iraq. *Acarina*, 22 (1):47–52.
- Bahreini, R. 2003. A comparison of two methods of applying oxalic acid for control of varroa. *Journal of Apicultural Research*, 42, 82–83. doi:10.1080/00218839.2003.11101098

- Cox-Foster DL, Conlan S, Holmes EC, Palacios G, Evans JD, et al. 2007. A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* 318: 283–287.
- Dainat B, Evans JD, Chen YP, Gauthier L, Neumann P 2012. Predictive markers of honey bee colony collapse. *Plos One* 7: e32151.
- de Graaf, A.M. Alippi, M. Brown, J.D. Evans, M. Feldlaufer, A. Gregorc, M. Hornitzky, S.F. Peral, D.M.T. Schauch, D. Titera, V. Tomkies, and W. Ritter, 2006. Diagnosis of American foulbrood in honey bees: a synthetic and proposed analytical protocols, *Letters in Applied Microbiology*, 43, 583-590.
- Duay P, De Jong D, Engels W 2002. Decreased flight performance and sperm production in drones of the honey bee (*Apis mellifera*) slightly infested by Varroa destructor mites during pupal development. *Genetics and Molecular Research* 1: 227–232.
- Esmen, B., Dodoglu, A., & Genc, F. 2010. Natural compounds for the control of Varroa destructor in honey bees (*Apis mellifera* L.) and an economic assessment of these products. *Mellifera*, 10–20, 32–36.
- Evans JD, Lopez DL, 2004. Bacterial probiotics induce an immune response in the honey bee (Hymenoptera: Apidae). *J Econ Entomol*, 97:752-756.
- Gätschenberger H, Azzami K, Tautz J, Beier H, 2013. Antibacterial immune competence of honey bees (*Apis mellifera*) is adapted to different life stages and environmental risks. *PLoS ONE*, 8: e66415.
- Girisgin, A.O. 2008. Using and efficacy of organic acids to Varroa destructor on naturally infested honey bee (*Apis mellifera*) colonies (PhD thesis) (65 pp.). Uludag University, Health Sciences Institute, Turkey.
- Goulson D, Nicholls E, Botías C, Rotheray EL, 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, 347:1255957.
- Haddad, N. 2011. Honey bee viruses, diseases and hive management in the Middle East and their relation to the colony collapse disorder and bee losses. *Uludag Bee Journal*, 11 (1): 17–24.
- Highfield AC, El Nagar A, Mackinder LC, Noel LM, Hall MJ, et al. 2009. Deformed wing virus implicated in overwintering honeybee colony losses. *Appl Environ Microbiol* 75: 7212–7220.
- Higes M, Martin-Hernandez R, Botías C, Bailon EG, Gonzalez-Porto AV, 2008. How natural infection by Nosema ceranae causes honeybee colony collapse. *Environmental Microbiology* 10: 2659–2669.
- Higes M, Martin-Hernandez R, Garrido-Bailon E, Gonzalez-Porto AV, Garcia- Palencia P, 2009. Honeybee colony collapse due to Nosema ceranae in professional apiaries. *Environmental Microbiology Reports* 1: 110–113.
- Paxton RJ, 2010. Does infection by Nosema ceranae cause “Colony Collapse Disorder” in honey bees (*Apis mellifera*)? *Journal of Apicultural Research* 49: 80–84.
- Ifantidis MD, 1983. Ontogenesis of the Mite Varroa-Jacobsoni in Worker and Drone honeybee Brood Cells. *Journal of Apicultural Research* 22: 200–206.
- Kralj J, Brockmann A, Fuchs S, Tautz J, 2007. The parasitic mite Varroa destructor affects non-associative learning in honey bee foragers, *Apis mellifera* L. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology* 193: 363–370.
- Kulhanek K, Steinhauer N, Rennich K, Caron DM, Sagili RR, Pettis JS, Ellis JD, Wilson ME, Wilkes JT, Tarpay DR, Rose R, Lee K, Rangel J, vanEngelsdorp D, 2017. A national survey of managed honey bee 2015–2016 annual colony losses in the USA. *J Apicult Res* 2017, 56:328-340.
- Le Conte Y, Ellis M, Ritter W, 2010. Varroa mites and honey bee health: can varroa explain part of the colony losses? *Apidologie* 41: 353–363.
- Mato, I., Huidobro, J.F., Simal-Lozano, J., & Sancho, M.T, 2006. Analytical methods for the determination of organic acids in honey. *Critical Reviews in Analytical Chemistry*, 36, 3–11.
- Nazzi F, Brown SP, Annoscia D, Del Piccolo F, Di Prisco G, 2012. Synergistic Parasite-Pathogen Interactions Mediated by Host Immunity Can Drive the Collapse of Honeybee Colonies. *PLoS Pathogens* 8: e1002735.
- Neumann P, Carreck NL, 2010. Honey bee colony losses. *J Apiculture Res*, 49:1-6.
- Runckel C, Flenniken ML, Engel JC, Ruby JG, Ganem D, 2011. Temporal analysis of the honey bee microbiome reveals four novel viruses and seasonal prevalence of known viruses, Nosema, and Crithidia. *Plos One* 6: e20656.
- VanEngelsdorp D, Evans JD, Saegerman C, Mullin C, Haubruge E, 2009. Colony collapse disorder: a descriptive study. *Plos One* 4: e6481.
- VanEngelsdorp D, Speybroeck N, Evans JD, Nguyen BK, Mullin C, 2010. Weighing risk factors associated with bee colony collapse disorder by classification and regression tree analysis. *Journal of Economic Entomology* 103: 1517–1523.
- Yang X, Cox-Foster D, 2007. Effects of parasitization by Varroa destructor on survivorship and physiological traits of *Apis mellifera* in correlation with viral incidence and microbial challenge. *Parasitology* 134: 405–412.

الأسباب المتوقعة لانخفاض تعداد شغالات نحل العسل في طوائف *Apis mellifera*

زهرة نايف أيوب¹ و روخوش جوهر رشيد² و تيشك حسن شيخ فرج² و عثمان مجيد أحمد³

¹ قسم وقاية النبات، كلية علوم الهندسة الزراعية، جامعة دهوك، إقليم كردستان، العراق

² قسم البستنة، كلية علوم الهندسة الزراعية، جامعة السليمانية، إقليم كردستان، العراق

³ المديرية العامة للزراعة، وزارة الزراعة والموارد المائية، إقليم كردستان، العراق

استهدفت الدراسة التحري عن العلاقة بين استمرار بقاء طوائف النحل وبعض الإجراءات الوقائية خلال التربية. تم تنفيذ التجارب في محافظة السليمانية خلال الأشهر؛ اب؛ أيلول وتشرين الأول 2020. تم اختبار 72 طائفة نحل في اربعة مواقع (قرداغ؛ مركز السليمانية؛ ميركه بان؛ ومنطقة سرتاكي بامو)؛ (18 طائفة في كل موقع). تم استخدام البكتريا المفيدة؛ حامض الفورميك؛ البيوكالبتوس؛ تتراسايكلين؛ و تيراميسين؛ مع طوائف التجربة. كانت مساحة الحضنة اكبر في جميع الطوائف الخاضعة للاختبار مقارنة بالطوائف غير المعاملة (طوائف السيطرة). الطوائف المزودة بالبكتريا المفيدة أظهرت زيادة معنوية في مساحة الحضنة مقارنة بطوائف السيطرة. اكبر مساحة للحضنة 203.66 انج² كانت في منطقة سرتاكي بامو؛ و بعدها 199.667 انج² في ميركه بان؛ ثم 179.00 انج² في مركز السليمانية؛ بينما لم تتجاوز مساحة الحضنة 15.66 انج² في جميع الطوائف غير المعاملة (السيطرة) في المواقع الأربعة الطوائف المعاملة بالبكتريا المفيدة أظهرت زيادة معنوية في المساحة المغطاة بشغالات النحل مقارنة بطوائف السيطرة في المواقع الأربعة. أعلى قيمة للمساحة المغطاة بالنحل كانت 6.667 اطار لانكستروث في منطقة سرتاكي بامو؛ و بعدها 6.333 اطار لانكستروث في منطقة ميركه بان؛ ثم 6.000 اطار لانكستروث في الطوائف المعاملة بحامض الفورميك في منطقة سرتاكي بامو. بينما لم تتجاوز كثافة النحل عن 1.000 اطار لانكستروث في الطوائف غير المعاملة (السيطرة) في المواقع الأربعة لكل من البكتريا المفيدة و الحامض العضوي اعطت افضل النتائج عند استخدامها كأجارات وقائية للمحافظة على طوائف النحل.