

Role of Environmental Conditions on the Epidemic of Wheat Yellow Rust in Gharbia Governorate of Egypt

Moustafa M. El-Shamy*¹  Mona E. Mohamed¹, Shaimaa E. Ibrahim², and Samir M. Saleh³

Address:

¹ Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt

² Field Crops Research Institute, Agricultural Research Center, Giza, Egypt

³ Central Laboratory for Agricultural Climate, Agricultural Research Center, Giza, Egypt

*Corresponding author: **Moustafa M. El-Shamy**; email dr.elshamy@yahoo.com

Received: 27-01-2022; Accepted: 01-05-2022; Published: 17-05-2022

doi: [10.21608/ejar.2022.118682.1201](https://doi.org/10.21608/ejar.2022.118682.1201)



ABSTRACT

This work aimed to evaluate one hundred and fifty-five commercial wheat fields including six varieties of yellow rust (stripe rust) epidemics caused by *Puccinia striiformis* f. sp. *tritici*, in Gharbia governorate during the 2019 to 2021 seasons. Also, study the weather variables associated with yellow rust outbreaks in wheat. Temperature, relative humidity, and rain variables were obtained from the regional climate model imeto2 (<http://meto.at/farmview/>) as well as the World Climate Research Programme (WCRP) Global Energy and Water Cycle Experiment's (GEWEX) Surface Radiation Budget (NASA/GEWEX SRB) during the infection period from January to March. Results revealed that the high disease epidemic linked to rain precipitation and number of precipitation days combined with the minimum temperatures (5–10°C) and relative humidity (RH) of at least 60%. Levels of yellow rust severity varied according to the studied year, and wheat variety. Gemmeiza-11, Sids-12, Misr-1, and Misr-2 were the most susceptible varieties, while both Gemmeiza12 and Sakha95 were resistant to yellow rust disease during the studied period. In the 2021 season, all the wheat varieties showed a resistance response. The loss % ranged between 36.00, 40.00 % (Gemmeiza-11); 36.00, 37.25% (Sids-12); 25.00, 33.33% (Misr-1) and 21.21, 24.84% (Misr-2) in 2019 and 2020 growing seasons, respectively compared to the grain yield in 2021 season. Low disease severities were detected ranging from zero to trace infection (2.00%) in the 2021 season. A high and positive correlation coefficient (R^2) was found between disease severity % and loss in grain yield of the tested wheat varieties.

Keywords: Yellow rust, *Puccinia striiformis*, disease severity, rain precipitation, relative humidity, temperature.

INTRODUCTION

Wheat (*Triticum aestivum*) is produced in almost all the governorates of Egypt, which is considered the first field crop followed by maize. Stripe (yellow) rust caused by *Puccinia striiformis* f. sp. *tritici* (Pst), is one of the most important diseases of wheat that incurred great yield losses around the world (Hovmøller *et al.*, 2011; de Vallavieille-Pope *et al.*, 2012; Chen *et al.*, 2014). In the last decade, epidemics of yellow rust destroyed most of the wheat area grown with Giza 144 variety in the Northern Egyptian provinces, particularly in Manzala district in 1967 (Abdel-Hak *et al.*, 1972). Also, in 1995, the disease attacked most of the commercial wheat varieties including Gemmeiza-1, Giza-163, Sakha-69, Giza-157, Sakha-92, Giza-166, and Giza-164, causing severe infection, particularly in the north and south Delta region. The national average loss in grain yield ranged from 14.00 –20.50% in the Nile delta region (El-Daoudi *et al.*, 1996; El-Shamy and Hamada, 1997; Abu El-Naga *et al.*, 1998). Then, yellow rust epidemics occurred regularly in all the governorates of Egypt including Gharbia governorate until the 2020 season. During this period, the recent wheat varieties i.e. Gemmeiza-11, Sids-12, Misr-1, Misr-2, Giza-171 showed susceptible responses to stripe rust under field conditions and new virulent stripe rust races were appeared (Abou-Zeid and Omara, 2018; Omar, 2020; Shahin, 2020; Shahin *et al.*, 2020). Favorable climatic conditions include different variables i.e. temperature, relative humidity (RH), and rain precipitation, which leads to an increasing threat of the disease (El Jarroudi *et al.*, 2017; Bitar and Farhad, 2019; Draz and Abd El-Kreem, 2021). The duration of the favorable temperature and moisture during wheat growth plays an important role in increasing yellow rust epidemics (van den Berg and van den Bosch, 2007). On the role of temperature, severe epidemics of stripe rust were linked to optimal temperatures from 7 to 12°C (Grabow *et al.*, 2016). This work aims to study the relationship between weather factors and occurrence yellow rust disease epidemics. In addition to investigate the reaction of six wheat varieties to the disease and its effect on grain yield in Gharbia governorate during the 2019 –2021 growing seasons.

MATERIAL AND METHODS

Selected locations:

This study was carried out in five districts in Gharbia governorate during 2019 – 2021 growing seasons. The surveys for yellow rust were carried out across the wheat-growing areas of El-Santa, El-Mahalla El-Kubra, Samnoud, Zifta, and Tanta districts during the National Campaign of wheat. The district's extension engineers shared in selecting the wheat fields. Six commercial bread wheat varieties i.e. Gemmeiza-11, Gemmeiza-12, Sids-12, Sakha-94, Misr-1, and Misr-2 were observed from January–March during the 2019 – 2021 growing seasons. Three to six fields per site were surveyed depending on the year. The number of observed fields ranged from 47 to 54 each year and the total number was 155 fields during the period of study [Fig.1 and Table\(1\)](#). Bread wheat is usually sown from mid-November to first December.




Fig. 1. Selected five districts for the observed wheat fields during the 2019 – 2021 cropping seasons in Gharbia governorate

Table 1. Distribution of 155 bread wheat fields selected in Gharbia governorate during 2019 - 2021 growing seasons.

| District | Location | No. of fields in | | |
|--------------------------------|----------------|------------------|-----------|-----------|
| | | 2019 | 2020 | 2021 |
| El-Santa | El-Gemmeiza | 4 | 4 | 6 |
| | Shinraq | 5 | 4 | 5 |
| | Shubra Qas | 3 | 4 | 4 |
| El-Mahalla El-Kubra | Saft Turab | 4 | 5 | 4 |
| | Shubra Malakan | 3 | 5 | 4 |
| | EL Hyatem | 4 | 4 | 5 |
| Sammanoud | Sammanoud | 4 | 4 | 4 |
| | Aziziyah | 4 | 5 | 5 |
| Zifta | Sunbat | 3 | 6 | 5 |
| | Hanout | 4 | 4 | 4 |
| Tanta | Ragdia | 4 | 5 | 4 |
| | Shabsheir | 5 | 4 | 4 |
| Total no. of fields 155 | | 47 | 54 | 54 |

Meteorological data.

Temperature (°C), relative humidity (RH %), and precipitation (mm), and precipitation days were computed using the regional climate model imeto2 (<http://meto.at/farmview/>). This weather station is located in Gemmeiza Agricultural Research Station which is based on its location relative to wheat-producing areas in Gharbia governorate, the station covers about 25 km in all directions and is away from 7 and 25 km from the selected sites. Obtained data are present in (Table 2).

Table 2. Mean of Metrological data obtained by the regional climate model imeto2 (<http://meto.at/farmview/>) in Gharbia governorate during 2019 up to 2021 seasons.

| Variable | 2019 | | | 2020 | | | 2021 | | |
|--------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | Jan. | Feb. | Mar. | Jan. | Feb. | Mar. | Jan. | Feb. | Mar. |
| Precipitation(mm) | 1.13 | 0.37 | 0.65 | 2.03 | 0.77 | 3.05 | 0.77 | 0.79 | 0.12 |
| Precipitation days | 19 | 10 | 7 | 11 | 22 | 15 | 9 | 4 | 6 |
| RH% | 55.34 | 58.79 | 58.3 | 67.24 | 66.94 | 57.9 | 58.56 | 57.77 | 52.28 |
| TMAX | 18 | 20.41 | 21.36 | 17.23 | 20.29 | 25.68 | 20.56 | 25.06 | 21.62 |
| TMIN | 6.5 | 7.37 | 7.87 | 7.82 | 8.5 | 10.92 | 9.4 | 11.68 | 9.78 |

Rain: Precipitation (mm day⁻¹) RH: Relative Humidity at (%) TMAX: Maximum Temperature(C) TMIN: Minimum Temperature

To confirm the meteorological data, we obtained the weather data per each district from the World Climate Research Programme (WCRP) Global Energy and Water Cycle Experiment's (GEWEX) Surface Radiation Budget (NASA/GEWEX SRB), Table (3).

Table 3. Mean of Metrological data obtained by the World Climate Research Programme (WCRP) Global Energy and Water Cycle Experiment's (GEWEX) Surface Radiation Budget (NASA/GEWEX SRB) in Gharbia governorate during 2019 up to 2021 seasons

| Variable | 2019 | | | 2020 | | | 2021 | | |
|-------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | Jan. | Feb. | Mar. | Jan. | Feb. | Mar. | Jan. | Feb. | Mar. |
| Precipitation(mm) | 1.23 | 0.43 | 0.82 | 2.10 | 0.60 | 2.74 | 0.27 | 1.16 | 0.23 |
| RH% | 55.29 | 58.54 | 56.2 | 69.07 | 65.91 | 59.81 | 62.02 | 62.75 | 60.40 |
| TMAX | 19.72 | 21.39 | 23.84 | 18.36 | 20.80 | 24.65 | 22.08 | 19.81 | 30.78 |
| TMIN | 7.25 | 8.17 | 9.70 | 8.47 | 8.72 | 10.35 | 9.67 | 9.52 | 10.09 |

Assessment of wheat yellow rust severity:

Visual disease severity estimates were done using the 0 -100 scale proposed by (Peterson *et al.*, 1948) and infection type adopted by (Johnston, 1961). Yellow rust severity was detected as percent leaf area affected by the disease, while Infection types were classified into five categories, 0 (immune), R (resistance), MR (moderate resistance), MS (moderately susceptible), and S (susceptible). The first record started with appearance of yellow rust symptoms in any variety and location. Assessment of the disease was carried out at least on ten sites of each field. Four assessments were carried out along the season to detect the final disease severity.

Grain yield affected by yellow rust epidemics:

Estimation of the grain yield for each variety/ site and season was carried out in cooperation with farmers as a ton per acre. The loss % of grain yield/acre was estimated by comparing the yield in 2019 and 2020 and 2021 seasons, which relatively no yellow rust infection was detected, using the equation adopted by (Calpouzos *et al.*, 1976)

$$\text{Loss\%} = (1 - Y_d / Y_h) \times 100$$

Y_d = Yield of diseased field Y_h = yield of healthy field

Least significant differences (L.S.D. at 0.05%) were used to comparisons yield components parameters under study according to Snedcor (1957). The correlation coefficient was used to expos the relation disease severity % and grain yield loss

RESULTS**Yellow rust mean disease severity:**

Mean disease severity of yellow rust of 6 bread wheat varieties evaluated in 155 fields in Gharbia governorate from the 2019 - 2021 seasons Table (4). The first observation of yellow rust varied depending on the year, and wheat variety. First symptoms were generally observed between the 1st January and 5th February 2019 and 2020 in most of the sites. The varieties Gemmeiza-11, Sids-12, Misr-1, and Misr-2 showed high mean disease severity percentages ranging from 46 to 94 S in 2019 and 2020 seasons at all the tested sites, while the Gemmeiza-12 variety showed low mean disease severity % (4S) in 2019 and high (55S) in 2020. Slight disease severities were detected on the evaluated varieties ranging from zero to trace infection in the 2021 season.

Table 4. Mean disease severity of yellow rust of 6 bread wheat varieties evaluated in 156 fields in Gharbia governorate during 2019- 2021 growing seasons.

| Variety | 2019 | 2020 | 2021 |
|-------------|-------|-------|------|
| Gemmeiza-11 | 76S | 94S | 1.8S |
| Gemmeiza-12 | 4S | 5.5S | 0 |
| Sakha-94 | 2.33S | 0 | 0 |
| Sids-12 | 76S | 77.5S | 2.0S |
| Misr-1 | 48S | 80S | 2.2S |
| Misr-2 | 46S | 55S | 2.0S |

Mean disease severity % was assessed according to the modified 0 -100 scale proposed by Peterson *et al.* (1948) and infection type (Johanston, 1961).

Metrological analysis:

Precipitation (mm/ day) and Precipitation days, humidity, and maximum and minimum temperature during the critical months of infection (January – March) in 2019 up to 2021 seasons were obtained from the regional climate model imeto2 (Table 2). Data clear that mean precipitation (mm) and precipitation days in 2019 and 2020 seasons (1.3, 0.37, 0.65mm and 2.03, 0.77, 3.05 mm/ day) were more than those in 2021(0.77, 0.79 and 0.12 mm/ day (Fig. 2). The associated degrees of minimum temperature in 2019 and 2020 were relatively less than those in 2021. The minimum temperatures (5–10°C) and humidity percentages(RH%) ranged from 52 up to 67% were relatively more in 2019, 2020 seasons than those in 2021 seasons Fig. (3). In general, slight differences in metrological data were observed between the regional climate model imeto2 and those obtained from NASA/GEWEX SRB Table (3).




Fig. 2. Rain precipitation (mm/day) and precipitation days during the critical months in 2019 up to 2021 growing seasons.




Fig. 3. Association between precipitation days (mm/day), minimum temperature (TMIN), and relative humidity (RH%) in the critical months during 2019 up to 2021 seasons.

Grain yield of the wheat varieties:

Data in the Table (5) revealed that the loss in grain yield was associated with yellow rust disease severity of each wheat variety / growing season. Significant differences were found between the tested wheat varieties. The wheat variety Gemmeiza-11 showed the lowest grain yield (1.725 and 1.600 ton /feddan) followed by Sids-12 variety (1.763 and 1.600 ton / feddan), Misr-1 (1.800 and 1.600 ton /acre) then Misr-2 variety (1.950 and 1.860 ton / feddan) in 2019 and 2020 seasons, respectively. While in 2021 season, all the wheat varieties gave high grain yields ranging from 2.400 ton / feddan (Misr-1) to 2,850ton/ feddan (Sakha-94). The loss % of grain yield/variety was assessed as compared between those in the 2019, 2020 seasons and the yield in the 2021 season. Gemmeiza-11, Sids-12, Misr-1 and Misr-2 gave the highest losses of grain yield, respectively. The loss % ranged between 36.00, 40.00 %(Gemmeiza-11), 36.00, 37.25% (Sids-12), 25.00, 33.33% (Misr-1) and 21.21, 24.84% (Misr-2) in 2019 and 2020 growing seasons, respectively compared to the obtained grain yield in 2021 season Table 5. Mean grain yield (ton/feddan) and grain loss% of 6 bread wheat varieties evaluated in 155 fields in El-Gharbia governorate during 2019- 2021 growing seasons.

| Wheat variety | Grain yield/ ton/ feddan | | | Loss % | |
|----------------|--------------------------|--------|---------|--------|-------|
| | 2019 | 2020 | 2021 | 2019 | 2020 |
| Gemmeiza-11 | 1.725 | 1.600b | 2.700b | 36.00 | 40.00 |
| Gemmeiza-12 | 2.750a | 2.700a | 2.775ab | 0.90 | 2.00 |
| Sakha-94 | 2.840a | 2.800a | 2.850a | 0.35 | 1.75 |
| Sids-12 | 1.763b | 1.600b | 2.550c | 30.86 | 37.25 |
| Misr-1 | 1.800b | 1.600b | 2,400d | 25.00 | 33.33 |
| Misr-2 | 1.950b | 1.860b | 2.475cd | 21.21 | 24.84 |
| L.S.D. at 0.05 | 0.62 | 0.43 | 0.12 | | |

Correlation between disease severity and grain yield loss.

A positive and high significant correlation coefficient (R2) was found between disease rust severity (%) and loss (%) in grain yield (0.988**) in 2019 season and (0.999) in 2020 season Table (6)

Table 6. Correlation coefficient (R^2) between disease severity% and grain yield loss in 2019 and 2021 seasons.

| Wheat variety | 2019 season | | | 2020 season | | |
|---------------|------------------|--------|---------|------------------|--------|---------|
| | Disease severity | Loss % | R^2 | Disease severity | Loss % | R^2 |
| Gemmeiza-11 | 76S | 36.00 | 0.988** | 94S | 40.00 | 0.999** |
| Gemmeiza-12 | 4S | 0.90 | | 5.5S | 2.00 | |
| Sakha-94 | 2.33S | 0.35 | | 0 | 1.75 | |
| Sids-12 | 76S | 30.86 | | 77.5S | 37.25 | |
| Misr-1 | 48S | 25.00 | | 80S | 33.33 | |
| Misr-2 | 46S | 21.21 | | 55S | 24.84 | |

DISCUSSION

Yellow rust (*Puccinia striiformis* f. sp. *tritici*) is considered the most effective disease on wheat and causes a great reduction in grain yield worldwide. There is no prediction model for yellow rust epidemics now in Egypt. Only, the prediction depends on observations of the wheat trap nursery for yellow rust, which is cultivated in almost all the governorates of Egypt. These trap nurseries include all the resistance genes of yellow rust as well as the local wheat varieties. Evaluation of wheat varieties revealed that Gemmeiza-11, Sids-12, Misr-1, and Misr-2 varieties had high disease severities of yellow rust, respectively at all sites in 2019 and 2020 years Table (4). (Shahin, 2020) stated that recent commercial wheat varieties, Misr-1, Misr-2, Giza-171 became susceptible under field conditions. Also, infections of stripe rust were observed on some wheat lines carrying the *Yr* genes *Yr1*, *Yr17*, *Yr27*, *Yr32* and *YrSp* which previously known to be resistant in trap nurseries. In addition to prevalence of new virulent races of the fungus lead to breakdown rust resistance in wheat varieties. (Esmail et al., 2021) reported that Six new aggressive races i.e. 151E80, 135E16, 160E173, 224E191, 238E143 and 72E8 during 2019 and 2020 seasons. There is no doubt that the cultivation of the highly susceptible variety, Gemmeiza-11 in about 60% of the wheat area of the governorate in the 2019 and 2020 seasons leads to an increase in yellow rust disease infection and considered as spreader of rust spores to the other varieties. (Solh et al., 2012) stated that individual crop losses of up to 80% were reported in the widespread epidemic in the Middle East and North Africa in 2010 when initial infection occurred on susceptible wheat varieties at early growth stages. Also, the late discovery of the disease and the using the wrong machine for managing the disease leads to an increase in the disease severity. In contrast, in the 2021 year the same varieties were relatively resistant or showed very low disease severities to yellow rust due to the unsuitable weather conditions for the fungus.

Yellow rust epidemics are the outcome of interactions between susceptible host varieties, favorable environmental conditions, and sufficient quantities of virulent pathogen inoculum (Zadoks, 1985). Early field observations of yellow rust have been shown to have predictive value, so a forecast system combining weather and disease observations may be sufficiently accurate to be useful. (Young et al., 2003). In our results, a low temperature (5–10 °C) associated with great precipitation (mm) and more precipitation days and relative humidity ranging from 55 to more than 66% from January - March period were associated with subsequent higher disease severity. (Bita et al., 2019) stated that occurrence of stripe rust epidemic was linked to the number of rainy days, the number of days with minimum temperatures within the range of 7–8°C and relative humidity (RH) above 60%. (El Jarroudi et al., 2017) found that high humidity, precipitation, or dew is the requirement for the development of the yellow rust epidemic. (Papastamati and Van den Bosch, 2007) found that the temperature in January and February was the most influential variable for the progress of yellow rust disease. The temperature range from 10 - 15°C was suitable for the development of yellow rust while temperatures higher than 23°C slow down the epidemic (Milus et al., 2009). (In contrast, Te Beest et al., 2008) found that the best model describing the disease epidemic of yellow rust consisted of maximum temperature and rain, with an RH of 86.4%. It could be stated that the regional climate model imeto2 presented relatively the same patterns of meteorological conditions obtained by NASA/GEWEX SRB.

The data revealed that the wheat varieties Gemmeiza-11, Sids-12, Misr-1, and Misr-2 highest disease severity consequently gave the highest percentage of grain loss in the 2019 and 2020 seasons. Sahin et al (2020) reported that the cultivars Gemmeiza 11 recorded 64.20% grain yield losses followed by Misr- 1 (62.38%), Misr-2 (57.66%) and Sids-12 (50.89%). The obtained results showed a positive and high significant correlation coefficient (R^2) between stripe rust disease severity and grain loss/ feddan releasing (988** and 999**) in the two seasons, respectively. Many papers reflected the great relationship between final stripe rust severity and grain yield loss % (Ahmad et al., 2010; Ashmawy et al., 2013; Taye et al., 2015). Yield loss was correlated with the proportion of leaf area affected by yellow rust development. In most wheat-producing areas, yield losses caused by yellow rust ranged from 10-70% depending upon the susceptibility of cultivar, earliness of the initial infection, rate of disease development, and duration of the disease (Chen, 2005).

CONCLUSION

From our study, we found that low temperature (5-10°C) associated with more precipitation quantities (mm/day), number of rainy days and humidity (RH %) more than 50% are the most suitable prediction for the occurrence of the yellow rust

epidemics in Egypt. The increase in grain losses % may be related to cultivating the highly sensitive wheat variety Gemmeiza-11 on a large scale as well as the late discovery of the disease and emergence of new virulent stripe rust races. To reduce yellow rust epidemics, the following steps must be followed, 4-5 resistant wheat varieties should be cultivated, early discovery of the disease, use the appropriate fungicide and spray machine. High significance and positive correlation coefficient (R^2) were found between stripe rust severity % and loss % of grain yield.


ACKNOWLEDGEMENT

The Prediction of Worldwide Energy Resource (POWER) Project is funded through the National Aeronautics and Space Administration (NASA) Applied Sciences Program within the Earth Science Division of the Science Mission Directorate. The POWER team could not have completed this task without both technical and scientific inputs from the following Earth Science Division teams: The World Climate Research Programme (WCRP) Global Energy and Water Cycle Experiment's (GEWEX) Surface Radiation Budget (SRB) and the Clouds and the Earth's Radiant Energy System (CERES) projects at NASA LaRC and the Global Modeling and Assimilation Office at the NASA Goddard Space Flight Center. The data obtained through the POWER web services was made possible with collaboration from the NASA Langley Research Center (LaRC) Atmospheric Science Data Center (ASDC).

REFERENCES

- Abdel-Hak, T., Stewart, D.M., and Kamel, A.H. (1972). The Current Stripe Rust Situation Near East Countries. *Regional Wheat Workshop, Beirut, Lebanon*, pp: 50 -55.
- Abou-Zeid, M., and Omara, R. (2018). Characterization of partial resistance stripe rust (*Puccinia striiformis* f. sp. *tritici*) cultivars under in some Egyptian wheat. *Journal of Plant Protection and Pathology, Mansoura University*,9: 111 - 119.
- Abu El-Naga, S.A., Khalifa M. M., Youssef, W. A., and Abd El-Latif, H. A. (1998). Stripe rust situation during the period (1994-1996) with special reference to designating genes conferring resistance in certain Egyptian wheat germplasm. *Journal of Agricultural Science, Mansoura University*. 23 :1127-1136.
- Ahmad, S., Afzal, M., Noorka, I. R., Iqbal, Z., Akhtar, N., Iftkhar, Y., and Kamran, M. (2010). Prediction of yield losses in wheat (*Triticum aestivum* L.) caused by yellow rust in relation to epidemiological factors in Faisalabad. *Pakistan Journal of Botany*, 42(1), 401-407.
- Ashmawy, M. A., El-Orabey, W. M., Nazim, M., and Shahin, A. A. (2013). Effect of stem rust infection on grain yield and yield components of some wheat cultivars in Egypt. *International Journal of Phytopathology*, 2(3), 171-178.
- Calpouzus, J., Roelfs, A.P., Madson, M.E., Martin, F.B., Wilcoxon, R.D., (1976). A new model to measure yield losses caused by stem rust in spring wheat. *Technical Bulletin - Agricultural Experiment Station, University of Minnesota*. 307, 1-23.
- Chen, W., Wellings, C., Chen, X., Kang, Z., and Liu, T. (2014). Wheat stripe (yellow) rust caused by *Puccinia striiformis* f. sp. *tritici*. *Molecular Plant Pathology*, 15(5), 433-446.
- Chen, X. M. (2005). Epidemiology and control of stripe rust (*Puccinia striiformis* f. sp. *tritici*) on wheat. *Canadian Journal of Plant Pathology*, 27(3), 314-337.
- de Vallavieille-Pope, C., Ali, S., Leconte, M., Enjalbert, J., Delos, M., and Rouzet, J. (2012). Virulence dynamics and regional structuring of *Puccinia striiformis* f. sp. *tritici* in France between 1984 and 2009. *Plant Disease*, 96(1), 131-140.
- Draz, I., Abd El-Kreem, T. (2021). Partial resistance to powdery mildew and leaf rust of wheat in Egyptian and CIMMYT genotypes. *Egyptian Journal of Agricultural Research*, 99(1), 61-76. doi: 10.21608/ejar.2021.56575.1070
- El-Daoudi, Y. H., Shafik, I., Ghanem, H. E., Abu El-Naga, S., Mitkees, R., Sherif, S., ... and Bassiouni, A. A. (1996, November). Stripe rust occurrence in Egypt and assessment of grain yield loss in 1995. In *Proceedings, Symposium Regional sur les Maladies des Cjr Jales at des II gumineuses Alimentaires* (pp. 11-14).
- El Jarroudi, M., Kouadio, L., Bock, C. H., El Jarroudi, M., Junk, J., Pasquali, M., ... and Delfosse, P. (2017). A threshold-based weather model for predicting stripe rust infection in winter wheat. *Plant Disease*, 101(5), 693-703.
- El-Shamy, M. M., and Hamada, A. A. (1997). Yellow rust occurrence in Egypt on some Egyptian and exotic wheat germplasm in 1997 growing season. *Annals of Agricultural Science, Moshtohor (Egypt)*.
- Esmail, S. M., Draz, I. S., Ashmawy, M. A., and El-Orabey, W. M. (2021). Emergence of new aggressive races of *Puccinia striiformis* f. sp. *tritici* causing yellow rust epiphytotic in Egypt. *Physiological and Molecular Plant Pathology*, 114, 101612.
- Grabow, B. S., Shah, D. A., and DeWolf, E. D. (2016). Environmental conditions associated with stripe rust in Kansas winter wheat. *Plant Disease*, 100(11), 2306-2312.
- Hovmøller, M. S., Sørensen, C. K., Walter, S., and Justesen, A. F. (2011). Diversity of *Puccinia striiformis* on cereals and grasses. *Annual review of phytopathology*, 49, 197-217.
- Johnston, C. O. (1961). Sixth revision of the international register of physiologic races of *Puccinia recondita* rob. ex desm. (formerly *P. rubigo-vera tritici*). *ARS* 34; 27.
- Milus, E. A., Kristensen, K., and Hovmøller, M. S. (2009). Evidence for increased aggressiveness in a recent widespread strain of *Puccinia striiformis* f. sp. *tritici* causing stripe rust of wheat. *Phytopathology*, 99(1), 89-94.
- Omar, E.G. (2020). Genetic analysis of *Puccinia striiformis tritici*, the causal pathogen of stripe rust disease and its effect on wheat cultivars resistance in Egypt. Ph.D., Kafrelsheikh Uni., Faculty of Agric., pp159.
- Papastamati, K., and van den Bosch, F. (2007). The sensitivity of the epidemic growth rate to weather variables, with an application to yellow rust on wheat. *Phytopathology*, 97(2), 202-210.

- Peterson, R. F., Campbell, A. B., and Hannah, A. E. (1948). A diagrammatic scale for estimating rust intensity on leaves and stems of cereals. *Canadian Journal of research*, 26(5), 496-500.
- Shahin, A. A. (2020). Occurrence of new races and virulence changes of the wheat stripe rust pathogen (*Puccinia striiformis* f. sp. *tritici*) in Egypt. *Archives of Phytopathology and Plant Protection*, 53(11-12), 552-569.
- Shahin, A., Ashmawy, M., El-Orabey, W., and Esmail, S. (2020). Yield losses in wheat caused by stripe rust (*Puccinia striiformis*) in Egypt. *American Journal of Life Sciences*, 8(5), 127-134.
- Snedicor, G.w. (1957). *Statistical Methods*, 5th eds. Iowa State College Press, Amelia, Wang.
- Solh, M., Nazari, K., Tadesse, W., and Wellings, C. R. (2012, September). The growing threat of stripe rust worldwide. In *Borlaug Global Rust Initiative (BGRI) conference, Beijing, China* (pp. 1-4).
- Taye, T., Fininsa, C., and Woldeab, G. (2015). Yield variability of bread wheat under wheat stem rust pressure at bore field condition of Southern Oromia. *Journal of Agricultural Science and Food Technology*, 1, 11-15.
- Te Beest, D. E., Paveley, N. D., Shaw, M. W., and Van Den Bosch, F. (2008). Disease–weather relationships for powdery mildew and yellow rust on winter wheat. *Phytopathology*, 98(5), 609-617.
- Van den Berg, F., and Van den Bosch, F. (2007). The elasticity of the epidemic growth rate to observed weather patterns with an application to yellow rust. *Phytopathology*, 97(11), 1512-1518.
- Young, C. S., Paveley, N. D., Vaughan, T. B., Thomas, J. M., and Lockley, K. D. (2003). Predicting epidemics of yellow rust (*Puccinia striiformis*) on the upper canopy of wheat from disease observations on lower leaves. *Plant Pathology*, 52(3), 338-349.
- Zadoks, J. C., and Bouwman, J. J. (1985). Epidemiology in Europe. In 'The cereal rusts Vol. 2: Diseases, distribution, epidemiology and control'. (Eds AP Roelfs, WR Bushnell) pp. 329–369.



Copyright: © 2022 by the authors. Licensee EJAR, EKB, Egypt. EJAR offers immediate open access to its material on the grounds that making research accessible freely to the public facilitates a more global knowledge exchange. Users can read, download, copy, distribute, print or share a link to the complete text of the application under [Creative Commons BY-NC-SA 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/).



دور الظروف الجوية في تطور وبائية مرض الصدأ الأصفر على القمح في محافظة الغربية بمصر

مصطفى محمود الشامى¹، منى السيد محمد¹، شيماء الدسوقي ابراهيم²، سمير محمود صالح³

¹معهد بحوث أمراض النباتات، قسم بحوث أمراض القمح، مركز البحوث الزراعية

²معهد بحوث المحاصيل الحقلية، قسم بحوث القمح، مركز البحوث الزراعية

³المعمل المركزي للمناخ الزراعي، مركز البحوث الزراعية

*بريد المؤلف المراسل: dr.elshamy@yahoo.com

الملخص

تهدف هذه الدراسة الى تقييم مائة وخمسة وخمسين حقلاً تجارياً ممثله في ستة أصناف من القمح لوبائيات مرض الصدأ الأصفر الناجم عن *Puccinia striiformis* f. sp. *tritici* في محافظة الغربية خلال المواسم من 2019 إلى 2021. وأيضاً دراسة الظروف البيئية المناسبة لحدوث وبائيه بمرض الصدأ الأصفر. تم الحصول على درجات الحرارة والرطوبة النسبية والأمطار من محطة التنبؤ imeto2 بمحطة بحوث الجميزة وكذلك من برنامج أبحاث المناخ العالمي (NASA / GEWEX SRB) خلال الفترة من يناير إلى مارس خلال مدة الدراسة. أظهرت النتائج أن حدوث وبائيه شديده بالمرض كان مرتبط بكمية وعدد أيام الأمطار مع درجات حراره منخفضه (5-10 درجات مئوية) ورطوبة نسبية (RH) على الأقل 60%. وقد أظهرت نتائج الدراسة الى تفاوت شدة الاصابه بالصدأ الأصفر تبعاً للسنة وصنف القمح المنزرع. وكانت الأصناف جميزة 11 - سدس 12 - مصر 1 - مصر 2 على التوالي الأكثر شدة اصابه خلال موسمي الزراعه 2019 و 2020. بينما أظهرت أصناف القمح جميزة 12 و سخا 94 أثار من الاصابه بالمرض. تراوحت نسبة الخسارة ما بين 36.00 ، 40.00% (جميزة 11) - 36,00 ، 37,25% (سدس 12) - 25,00 ، 33,33% (مصر 1) و 21,21 ، 24.84% (مصر 2) في موسمي 2019 و 2020 على التوالي بالمقارنة بمحصول الحبوب لنفس الأصناف في موسم 2021. والتي كانت تقريبا مقاومة للمرض. أظهرت الدراسة وجود معنويه عاليه موجب له معامل الارتباط بين شدة الاصابه والخساره في محصول الحبوب للأصناف تحت الدراسة.

الكلمات المفتاحية: صدأ أصفر، بكسينيا ستريفورمز، شدة الاصابه، سقوط الامطار، الرطوبه النسبيه، درجة الحرارة