

CONTENTS

	Page
I. INTRODUCTION.....	1
II. REVIEW OF LITERATURE.....	3
1. Seasonal abundance of predatory spiders.....	3
2. Seasonal abundance of <i>S. littoralis</i> (Boisd.), <i>A. craccivora</i> (Koch.), <i>A. gossypii</i> (Glover) and <i>T. urticae</i> (Koch.).....	10
3. Biological studies.....	14
4. Toxicity of some alternative pesticides on predatory spiders.....	15
5. Effect of some alternative pesticides on total protein and protein bands.....	17
III. MATERIAL AND METHODS.....	20
1. Seasonal abundance of the spiders and their preys during successive seasons of cotton and broad bean plants.....	20
A. Estimation of spider populations.....	20
B. Estimation of <i>S. littoralis</i> populations.....	20
C. Estimation of Aphids populations: <i>Aphis craccivora</i> in broad bean plants and <i>Aphis gossypii</i> in cotton plants.	21
D. Estimation of phytophagous mite populations.....	21
2. Biological studies.....	21
A. Spider's rearing.....	21
B. Spider's food.....	21
C. Prey's rearing.....	22
3. Toxicological of alternative pesticides.....	23
4. Biochemical studies.....	24
1. Tissue preparation.....	24

2. Determination of total protein.....	24
3. Separation of protein bands by electrophoresis (using SDS-PAGE).....	25
5. Statistical analysis.....	25
IV. RESULTS AND DISCUSSION.....	26
1. Ecological studies.....	26
1.1. Occurrence of spider families collected from broad bean and cotton crops.....	26
1.1.1. Occurrence of spiders families in Qaha Station.....	26
1.1.2. Occurrence of spiders families in Seds Station.....	31
1.2. The percentage of spider families and occurrence of their species their families during two successive years...	31
1.2.1. In Qaha Station (2003/2004).....	31
1.2.2. In Qaha Station (2004/2005).....	39
1.2.3. In Seds Station (2003/2004).....	42
1.2.4. In Seds Station (2004/2005).....	45
Seasonal abundance.....	49
1.3.1. Population density of predatory spider and their preys (<i>Spodoptera littoralis</i> , <i>Aphis craccivora</i> and <i>Tetranychus urticae</i>) which collected from broad bean plants at Qaha Station during two successive years (2003-2004) and (2004-2005).....	49
1.3.2. Population density of predatory spider and their preys (<i>Spodoptera littoralis</i> , <i>Aphis gossypii</i> and <i>Tetranychus urticae</i>) which collected from cotton plants at Qaha Station during two successive years (2004) and (2005).....	53
1.3.3. Population density of predatory spider and their	

preys (<i>Spodoptera littoralis</i> , <i>Aphis craccivora</i> and <i>Tetranychus urticae</i>) which collected from broad bean plants at Seds Station during two successive years (2003-2004) and (2004-2005).....	53
1.3.4. Population density of predatory spider and their preys (<i>Spodoptera littoralis</i> , <i>Aphis gossypii</i> and <i>Tetranychus urticae</i>) which collected from cotton plants at Seds Station during two successive years (2004) and (2005)).....	57
1.4. Association of spider families and preys on broad bean and cotton plants during two successive years (2003-2004) and (2004-2005) at two stations (Qaha and Seds).....	65
1.4.1. Association of spider families and preys on broad bean and cotton plants during (2003-2004) at Qaha Station.....	65
a) Broad bean.....	65
b) Cotton.....	65
1.4.2. Association of spider families and preys on broad bean and cotton plants during (2004-2005) at Qaha Station.....	69
a) Broad bean.....	69
b) Cotton.....	69
1.4.3. Association of spider families and preys on broad bean and cotton plants during (2004-2005) at Kaha Station.....	73
a) Broad bean.....	73
b) Cotton.....	73
1.4.4. Association of spider families and preys on broad	

bean and cotton plants during (2004-2005) at Seds Station.....	77
a) Broad bean.....	77
b) Cotton.....	77
1.5.Average number of spiders' predators and their families from broad bean and cotton plants during two successive years (2003-2004) and (2004-2205) at Kaha and Seds Stations	82
2. Biological studies.....	90
3. Toxicity of some alternative pesticides on adults (females and male).....	103
4.Biochemical study to investigate the effect of nutrition and alternative pesticide on total protein and protein bands in adult spiders.....	109
4.1. Total protein content.....	109
4.2. Refraction of protein patterns.....	109
4.2.1. The effect of treated the adult of (<i>T. abini</i>) with alternative pesticides on its protein bands molecular weight.....	112
4.2.2. Molecular weight of protein bands for adult (<i>T. abini</i>) which feeding on different prey.....	112
V. SUMMARY.....	123
VI. REFERENCES.....	127
ARABIC SUMMARY	

V. SUMMARY

Spiders are considered important and world wide predators of many agricultural pests. There are about 30.000 species of spiders identified around the world till now. Spiders spread in many habitats such as caves, cervices, above mountains and defied desert i.e. spider can be found in most region of earth (Rod & Ken, 1984).

This study included the following points:

1. **Seasonal abundance of the three pests (cotton leafworm, *Spodoptera littoralis*, aphids, *Aphis craccivora* and *Aphis gossypii* and two spotted spider mite, *Tetranychus urticae*) and their associated predatory spiders:**

This study occurred for determination of population density of three pest and spiders at Kaha Station, Qalubiya Governorate and Seds Station at Beni-Sueif Governorate during two successive years (3003-2004 and 2004-2005), on two host plants (cotton and broad bean). The results revealed that the occurrences of 13 families including 22 species these families are Lycosidae, Philodromidae, Linyphiidae, Theridiidae, Miturgidae, Ulobroridae, Oonopidae, Dictynidae, Gnaphosidae, Pholocidae, Araneidae, Thomisidae and Salticidae some of these families present in 2 seasons (summer and winter) as Lycosidae, Philodromidae, Linyphidae, Theridiidae, Miturgidae, Uloboridae, Dictynidae, Thomisidae and Salticidae. Studying the population density and seasonal abundance of the spiders associated with different three pests, temperature and relative humidity. Data in the tables showed that the increasing of temperature and relative humidity followed by the increase of spider population and that is due to the abundance of their prey population during the hot months and there was positive relationship

On the other hand, it was observed that the third group rate of consumption was more than second and first group relative to *S. littoralis*, they were increasing from G₄, G₁ and G₃, respectively. In case *T. urticae*, but the food consumption of *A. crassivora* increasing from G₄, G₂ and G₁, respectively.

Toxicity of some alternative pesticides (KZ oil, Vertimic and *P. nigrum*) on spider adults:

This study aims to demonstrate the effect of some alternative pesticides application on three pests on cotton or broad bean plants and on spider adults (*T. albini*) which play an important role in biological control for different pests.

Duration of female and male of spider (*T. albini*) increasing in case treated with *P. nigrum* than Vertimic and KZ oil.

The mortality percentages of spider were decreasing after 24, 48 and 72 h, respectively when applicated with three alternative pesticides. Toxicity of Vertimic was more affected on spider after 24h than KZ oil and *P. nigrum*.

Biochemical study:

Investigate the effect of these elements (alternative pesticides) and different feeding group on the biochemical change.

Results revealed some difference between the control and the treated samples and they indicated that:

- 1) Total protein was increased with third group followed by first and fourth group
- 2) Total protein was decreased with *P. nigrum* followed by KZ oil and Vertimic.
- 3) Electrophoretic bioassay analysis of protein showed that, protein bands were varied and there were difference between proteins in tissue in case of G₁, G₃ and G₄ and proteins in tissue which treated

with alternative pesticides both in their concentration and their molecular weights.

- 4) In other hand, the protein bands which appeared in the tissue adult of *T. albibi* when treated with alternative pesticides from *P. nigrum*, KZ oil and Vertemic. Protein band 9, 19 and 34 presented only in case treated with KZ oil, (14, 33) were presented in tissue which treated with *P. nigrum* only.
- 5) There were some of protein bands present 26 bands (8, 32) they absent in G₃ and G₄ and 2 bands present in G₃, (27, 33) they absent in other groups.