

Abstract

Khaled Abd El-Fattah Mahmoud El-Mangoury, Biotechnological Studies on Some Wheat Cultivars. Unpublished Ph. D. thesis, Ain Shams University, Faculty of Agriculture, Department of Agronomy, 2005.

This study was carried out in the laboratory of Plant Cellular and Molecular Genetics (PCMG), Agricultural Genetic Engineering Research Institute (AGERI), Agricultural Research Center (ARC), Ministry of Agriculture and Land Reclamation, Giza, during the period from 2001 till 2004.

The aim of the present investigation was two main goals, the first, was to optimize high efficient regeneration system and the second was to optimize reporter gene expression in callus using the biolistic method, causing highest gene expression and minimal tissues damage.

To efficiently complement traditional wheat breeding with genetic transformation technology, it will be desirable to introduce transgenes into the ideal genetic background. Poor tissue culture performance is limiting the number of wheat cultivars that can be stably transformed. Therefore, tissue culture response of three current Egyptian wheat cultivars cv. Sids 1, Sakha 69 and Giza 168 were statistically analyzed in this research. Regenerable callus cultures were initiated from immature embryos of the three wheat cultivars and regeneration characteristics differed highly significantly among the mentioned cultivars.

On the other hand, this research announced a rapid and facile method to investigate both osmotic conditioning treatments of the target tissues (mixture as 0.2 M mannitol and 0.2 M sorbitol or 0.4 M mannitol), and bombardment

pressure (1100 psi or 1350 psi) affecting beta-glucuronidase

(GUS) expression in the three Egyptian wheat cultivars. Immature embryo derived calli were bombarded with gold particles coated with *pBA6* plasmid, harboring *gus* and *bar* gene. GUS activity was observed after two and seven days from bombardment. At 1100 psi, number of blue spots and clusters per bombarded callus (GUS activity) was higher compared with 1350 psi across the three tested cultivars and reached 5% level of significance. Mixture osmotic treatment had resulted high levels of *gus* gene expression in both Sakha 69 and Sids 1 cultivars, while, mannitol was favorable for Giza 168 cultivar. Applying biological and physical parameters (osmotic treatment and bombardment pressure; respectively), high levels of *gus* gene expression were obtained.

In this study, we report one procedure that drastically reduce bombardment damage by optimizing the osmotic treatments and bombardment pressure protocol and the combination of these procedures with improving regeneration protocol for each cultivar, causes high frequency in regeneration capabilities of the bombarded calli.

Transgenic plants were recovered from the cultured immature embryo-derived callus of the three wheat cultivars after biolistic transfer and bialaphos selection. Transgene integration and expression were confirmed by herbicide application and polymerase chain reaction. The introduction of transgenes into the ideal genetic background will allow an evaluation of their crop improvement potential

Key Words: biolistic, osmotic, bombardment pressure, regeneration, transformation, *gus* gene, *bar* gene.

List of Contents

1. INTRODUCTION.....	1
2. REVIEW OF LITERATURE	
2.1. Plant Regeneration.....	4
2.1.1. Effect of cultivars.....	4
2.1.2. Effect of growth regulators.....	14
2.2. Plant transformation.....	26
2.2.1. Effect of cultivars.....	27
2.2.2. Effect of osmotic treatments.....	33
2.2.3. Effect of bombardment pressures	40
3. MATERIALS AND METHODS	
Part One	
3.1 Optimization of regeneration system of Egyptian wheat cultivars; Sakha 68, Sids 1 and Giza 168.	48
Experiment One	
Effect of cultivars, BA and NAA concentrations on regeneration characteristics.....	48
Experiment Two	
Effect of cultivars, growth regulator types, and concentrations on regeneration characteristics.....	49
Part Two	
3.2. Optimization of transformation system of Egyptian wheat cultivars; Sakha 68, Sids 1 and Giza 168.....	50
3.2.1. Plasmid Preparation	
3.2.1.1. Preparation of bacterial cells.....	51
3.2.1.2. Plasmid transformation into <i>E. coli</i> competent cells.....	51
3.2.1.3. Plasmid purification.....	51
3.2.2. Wheat transformation <i>via</i> particle bombardment technique.....	51
3.2.2.1. Preparation and coating of gold particles with plasmid DNA.....	52

3.2.2.2. Optimizing parameters of gene gun delivery.....	52
Experiment One	
Effect of cultivars, osmotic treatments, and bombardment pressures on <i>gus</i> gene expression.....	52
Experiment Two	
1. Biolistic bombardment.....	54
2. Selection and regeneration of transformed calli.....	55
3. Verification of gene integration.....	57
3.1. Isolation of plant genomic DNA.....	57
3.2. Detection of transformed DNA in genetically modified plants using Polymerase Chain Reaction (PCR).....	57
4. RESULTS AND DISCUSSION	
Part One	
Optimization of regeneration system of Egyptian wheat cultivars; Sakha 68, Sids 1 and Giza 168.....	60
Experiment One	
4.1. Effect of cultivars, BA and NAA concentrations on regeneration characteristics.....	60
4.1.1. Effect of cultivars.....	60
4.1.2. Effect of BA concentrations.....	60
4.1.3. Effect of NAA concentrations.....	63
4.1.4. Effect of interaction between cultivars and BA ...	63
4.1.5. Effect of interaction between cultivars and NAA.....	66
4.1.6. Effect of interaction between BA and NAA.....	66
4.1.7. Effect of interaction between cultivars, BA, and NAA.....	69
Experiment Two	
4.2. Effect of cultivars, growth regulator types, and concentrations on regeneration characteristics.....	76
4.2.1. Effect of cultivars.....	76
4.2.2. Effect of growth regulator types.....	76
4.2.2. Effect of growth regulator concentrations.....	79

4.2.4. Effect of interaction between growth regulator types and cultivars.....	79
4.2.5. Effect of interaction between growth regulator concentrations and cultivars.....	82
4.2.6. Effect of interaction between growth regulator types and growth regulator concentrations.....	82
4.2.7. Effect of interaction between cultivars, growth regulator types, and growth regulator concentrations.....	84

Part Two

Experiment I

Optimization of Transformation System in Egyptian Wheat Cultivars; Sakha 68, Sids 1 and Giza 168.....	93
---	----

4.3. Effect of cultivars, osmotic treatments, and bombardment pressures on <i>gus</i> gene expression.....	93
4.3.1. Effect of cultivars.....	93
4.3.2. Effect of osmotic treatments	94
4.3.3. Effect of bombardment pressures.....	95
4.3.4. Effect of interaction between cultivars and osmotic treatments	96
4.3.5. Effect of interaction between cultivars and bombardment pressures	97
4.3.6. Effect of interaction between osmotic treatments and bombardment pressures.....	98
4.3.7. Effect of interaction between cultivars, osmotic treatments, and bombardment pressures.....	100

Experiment II

Production of transgenic plants.....	105
--------------------------------------	-----

5. SUMMARY.....	112
6. LIST OF REFERENCES.....	117